Published in final edited form as:

J Dev Orig Health Dis. 2019 December ; 10(6): 636–644. doi:10.1017/S2040174419000102.

Seasonality of births in horizontal strabismus: comparison with birth seasonality in schizophrenia and other disease conditions

Andrea B. Agarwal¹, Kellie Cassinelli², L. Alan Johnson^{1,2}, Koji Matsuda³, Brian Kirkpatrick⁴, Wei Yang⁵, Christopher S. von Bartheld^{1,*}

¹Department of Physiology & Cell Biology, University of Nevada, Reno School of Medicine, Reno, NV, USA

²Sierra Eye Associates, Reno, NV, USA

³Matsuda Eye Clinic, Sennan, Osaka, Japan

⁴Department of Psychiatry & Behavioral Sciences, University of Nevada, Reno School of Medicine, Reno, NV, USA

⁵School of Community Health Sciences, Nevada Center for Health Statistics and Informatics, University of Nevada, Reno, NV, USA

Abstract

Recent work has implicated one type of horizontal strabismus (exotropia) as a risk factor for schizophrenia. This new insight raises questions about a potential common developmental origin of the two diseases. Seasonality of births is well established for schizophrenia. Seasonal factors such as light exposure affect eye growth and can cause vision abnormalities, but little is known about seasonality of births in strabismus. We examined birth seasonality in people with horizontal strabismus in a retrospective study in Washoe County, Nevada, and re-examined similar, previously obtained data from Osaka, Japan. We then compared seasonal patterns of births between strabismus, refractive error, schizophrenia, and congenital toxoplasmosis. Patients with esotropia had a significant seasonality of births, with a deficit in March, then increasing to an excess in September, while patients with exotropia had a distinctly different pattern, with an excess of births in July, gradually decreasing to a deficit in November. These seasonalities were statistically significant with either chi-square or Kolmogorov-Smirnov type statistics. The birth seasonality of esotropia resembled that for hyperopia, with an increase in amplitude, while the

*Corresponding author: cvonbartheld@med.unr.edu (CSvB).

Conflicts of Interest

Except for Brian Kirkpatrick, the authors have no conflict of interests or potentially conflicting interests. Brian Kirkpatrick receives licensing royalties and travel support from ProPhase for use of the brief negative symptoms scale (BNSS) by for-profit groups; these fees will be donated to the Brain and Behavior Research Foundation. He has also received fees and travel support from ProPhase for teaching the BNSS, consulting fees and travel support from Genentech/Roche, and consulting fees to anonymized pharmaceutical companies through Decision Resources, Inc. He also receives fees from Walsh Medical Media for editorial services, and received fees for editorial services from Physicians Postgraduate Press, Inc.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guidelines on human experimentation (Code of Federal Regulations on the Protection of Human Subjects, 45 CFR) and with the Helsinki Declaration of 1975, as revised in 2008, and has been reviewed by the institutional committee (University of Nevada, Reno Biomedical IRB). The IRB determined that this project is exempt from IRB review according to federal regulations (Exemption category #4; Decision Date: December 29, 2014; Reference Number 699359–1).

seasonality for myopia involved a phase-shift. There was no correlation between seasonality of births between strabismus and congenital toxoplasmosis. The pattern of an excess of summer births for people with exotropia was remarkably similar to the well-established birth seasonality of one schizophrenia subtype, the deficit syndrome, but not schizophrenia as a whole. This suggests a testable hypothesis: that exotropia may be a risk factor primarily for the deficit type of schizophrenia.

Keywords

strabismus; seasonality of births; schizophrenia; refractive error; toxoplasmosis

Introduction

In many diseases, developmental factors contribute to pathophysiology.^{1–3} Therefore, information about seasonality of births can provide insights into the etiology and possible prevention of diseases.⁴ Birth seasonality has been examined and reported for several diseases and conditions, including allergies, Alzheimer's disease, anorexia, arthritis, autism, autoimmune thyroid disease, birth defects, breast cancer, cerebral palsy, childhood cancers, Crohn's disease, depression, diabetes, dyslexia, epilepsy, hyperopia, multiple sclerosis, myopia, narcolepsy, schizophrenia and sudden infant death syndrome. It is well established that people with schizophrenia have an altered seasonality of births, with nearly 300 studies conducted on this topic in about 30 countries.^{4–6}

Recent work has suggested that one type of strabismus, exotropia, is a risk factor for mental illness,^{7, 8} and, in particular, schizophrenia.^{9–13} Strabismus is a deviation of eye alignment that can be vertical, but most often is horizontal. An inward deviation is called an esotropia, and an outward deviation is an exotropia.¹⁴ If seasonality of births in exotropia resembles that in schizophrenia, this would provide further support for an association between the two conditions and may lead to new pathophysiological insights. Seasonal effects on eye growth and refractive error have been documented.^{15–17} However, only a single study from Japan (published in Japanese) in the 1980s has reported on the seasonality of births for people with horizontal strabismus, but without adjustments for the length of months, or consideration of normal seasonal variation in the local population.¹⁸ The paucity of information regarding birth seasonality in strabismus is astounding when one considers the fact that there are about 5–9 times more people affected by strabismus¹⁹ than schizophrenia.²⁰

We determined whether there is a seasonality of births in patients with horizontal strabismus and explored how seasonalities of esotropia and exotropia compare with those previously reported for vision-related conditions. These include refractive errors (hyperopia, myopia),^{15–17} schizophrenia (which has frequent vision comorbidities),^{4–6, 21} and congenital toxoplasmosis. Toxoplasmosis has been implicated in ocular disease/vision problems²² as well as in schizophrenia.^{23, 24} We examined whether esotropia and exotropia have distinct patterns of birth seasonalities, and re-examined previous reports that hyperopia¹⁵ and myopia¹⁶ have birth seasonalities. Furthermore, we assessed how seasonality of births in horizontal strabismus compares with schizophrenia, and with one specific type of

schizophrenia: deficit schizophrenia.^{25–27} Deficit schizophrenia is a well validated subtype and may be a separate disease within the syndrome of schizophrenia.^{25–27} It is thought to involve about 20% of all schizophrenia cases and has previously been proposed to have particular vision and eye abnormalities when compared to non-deficit schizophrenia.^{28–30} Parts of our study have been presented in abstract form.³¹

Methods

Medical records were retrospectively reviewed for all patients who were diagnosed with either esotropia or exotropia at a private ophthalmology practice in Reno, Nevada, between 2001 and 2014 (n= 2,513 subjects with esotropia; n= 2,332 subjects with exotropia). The total number of patients with diagnoses of horizontal strabismus was 4,845. Although clinic-based studies do not provide information about prevalence, the ratio of esotropia to exotropia (51.9% esotropia vs. 48.1% exotropia) is as would be expected for a population-based study with a significant hispanic component.³² The clinic's computer programs were used to compile diagnoses (esotropia, exotropia) and birth months and years. The youngest patient was 13 months and the oldest was over 90 years old. The practice in Reno provides eye care for a large fraction of the population in Washoe County and surrounding areas in Northern Nevada, with little alternative care available within a wide radius (200 km to the west, across the Sierra mountain passes, and 500–600 km distance in all other directions). We used the Washoe County birth records from 1975 through 2014 as our control population to account for the normal seasonality of births in this region.³³ Nevada does not currently have, and has not previously shown, any confounding biannual fluctuations in births, as demonstrated for most southeastern states in the United States.³³ The University of Nevada Biomedical IRB determined that this study was exempt from IRB review. Data from Osaka, Japan (n=3,507 subjects with esotropia; n=3,188 with exotropia) that were analyzed for this study had been de-identified,¹⁸ and therefore did not require IRB oversight. The total number of patients with horizontal strabismus in the Osaka clinic was 6,695, with a ratio of 52.4% esotropia vs. 47.6% exotropia. Such a ratio is similar to those from population-based studies in Japan.³⁴ The normal seasonal variation of births in Osaka was examined for comparison, based on the report by Matsuda and Kahyo, 1994.³⁵

Births were compiled for each month of each year in spreadsheets and analyzed using two types of statistical tests for seasonality. The first test was the chi-square test which is the most popular test for comparing birth rates in specific diseases.^{36–38} Since the chi-square test is not ideal for the study of seasonal patterns such as unimodal, bimodal and multiple peak patterns,³⁷ we applied a second test, the Kolmogorov-Smirnov (KS) non-parametric method which was designed to test hypotheses about seasonal variation.^{37, 39} Since much of our data were spread over 80 years of birth months, we could not apply a time-series as an alternative approach.^{36, 38} For both the chi-square and the KS test, all data were corrected for the difference in numbers of days between months (x 1.0775 for the month of February = 28.25 days/year; x 1.0147 for months with 30 days per year, and x 0.9819 for months with 31 days per year), so that February was not by default the month with the lowest numbers due to being the shortest month. We graphed the distribution of births of people with strabismus relative to the slight seasonal variations among births of the normal local population. Monthly birth excess or reductions (seasonal birth amplitudes) were calculated

by taking one-half the difference between the maxima and minima, measured as percent of the annual mean,³³ after correction for lengths of months and taking into account the normal local seasonality of births. We not only compared the Nevada cohorts with the Osaka cohorts,^{15, 18} but, because the data turned out to be very similar (r-values of 0.714 and 0.686, Pearson's R correlation test), we combined the two data sets to increase the statistical power. To compare the resulting combined data with a normal seasonal variation of births, we constructed a normal curve by calculating the mean of each month for the Washoe County and the Osaka births.³⁵ Since the two sites are located at a similar latitude (39.5° for Reno, 34.7° for Osaka), the normal seasonal variation in births is expected to be,³³ and indeed was, similar between the two sites.

We used both the chi-square test and the KS test to determine whether differences from the expected values were statistically significant, which is the same methodology applied in previous seasonality studies.^{22, 36, 38, 39} A power analysis⁴⁰ predicted that with 2,500–5,000 subjects and an amplitude of 10–20%, there was a 90–100% confidence of detecting any existing trends. We compared our data with previous reports on seasonality in strabismus,¹⁸ hyperopia,^{15, 17} myopia,^{16, 17} as well as published reports about seasonality of births for people with schizophrenia^{4, 5} (well established for the Northern hemisphere),^{4–6, 21} and with three studies reporting on seasonality of congenital toxoplasmosis.^{22, 41, 42} For the precise definitions and inclusion criteria for schizophrenia diagnoses, we refer the reader to the published meta-analyses and original studies.^{4, 5, 21} The meta-analysis by Torrey et al. (1997) provides a detailed discussion of methodological problems.⁴ We applied error bars only to graphs when data from more than two cohorts were combined. Error bars were not reported in the schizophrenia meta-analyses of schizophrenia births.^{4, 5} Because the universality of an excess of summer births among people with deficit schizophrenia has been questioned,⁴³ we compiled and examined all relevant studies, including the most recent ones, to gain an up-to-date perspective on this issue. This provided a stronger confirmation of the association of deficit schizophrenia with summer births than was possible in previous studies.²¹

Results

In our study, we report on data obtained from cohorts at two locations. Data from Reno, Nevada were newly obtained, while some of the data from Osaka, Japan had been reported previously.¹⁸ The births in the Washoe County population (n=173,635) showed seasonality in births similar to data published for the entire United States (1942–1989),⁴⁴ (1931–2008),³³ with a deficit in October through February, and an increase in May through August (peak in June), amounting to a seasonal birth amplitude (= percent deviation from the annual mean)³³ of 6.0% (Fig. 1a,b). This pattern is consistent with the known increase in summer births in the central Western United States (latitude of Washoe County is 39.5°),^{33, 44} with an expected modest increase between July and September in the state of Nevada.³³ Therefore, these data provided a reliable local baseline for comparison with seasonal birth amplitudes from the patient diagnoses. Normal seasonal variation of births in Osaka was obtained from the work of a 1994 study.³⁵ The normal population of Osaka had a seasonal birth amplitude of 4.85% (Fig. 1c,d). Because of the latitudinal gradient in peak birth timing

and amplitude,³³ local rather than national birth data have to be used to establish the normal seasonal variation (baseline).

Strabismus.

In Nevada, patients with esotropia (n=2,513) showed a reduced number of births in March through July and an increased number of births in August through October. This dataset did not reach statistical significance ($p=0.2926$, chi-square test; $p>0.05$, KS test) (Fig. 1a). Patients with exotropia (n=2,333) showed a deficit in winter and spring births, and an excess number of births in July/August. The seasonality was statistically significant with the chi-square test ($p=0.0070$), but not with the KS test ($p>0.05$) (Fig. 1b).

The only previous study on the seasonality of births in strabismus - from Osaka, Japan^{15, 18} - did not correct data for length of month and normal seasonal variation. We therefore recalculated the original data and plotted the births together with the normal local seasonal variation (using the information from the 1994 study)³⁵ (Fig. 1c, d). Esotropia births from Osaka showed a deficit of births from March to June, and a peak from August to November (Fig. 1c) - which is similar to the deficit seen in Nevada from March to July, and the peak in August through October (Fig. 1a). Exotropia births from Osaka showed a peak in July/August (Fig. 1d) that was virtually identical to Nevada's peak in the same two months (Fig. 1b), and the winter months had the lowest numbers of births. At both sites (Nevada and Osaka), there was an abrupt increase in exotropia births in July and August, followed by a sudden decrease in October/November (Fig. 1b,d). The esotropia seasonality for the Osaka cohort was statistically significant with the chi-square test ($p<0.0001$) and with the KS method ($p<0.01$), while the exotropia seasonality was borderline significant with the chi-square test ($p=0.0544$), and did not reach statistical significance with the KS test ($p>0.05$). To assess the similarity of the data from the two cohorts, we calculated r-values based on Pearson's R correlation test. The r-values for the two graphs (Washoe County and Osaka) were 0.714 and 0.686, indicating strong linear correlations. Therefore, we considered it justified to combine the data from the two cohorts to achieve a stronger statistical power.

The combined data from the cohorts in Nevada and Osaka resulted in a data set of 6,020 subjects with esotropia, and 5,520 with exotropia. The combined numbers resulted in birth seasonality curves with less noise than those of the individual cohorts, thus making it easier to recognize patterns and trends. When compared with the expected seasonal variation, esotropia showed the greatest deficit of births in March, with a gradually increasing number of births until a peak in September (Fig. 1e), while exotropia showed an abrupt peak in July, followed by a gradual decrease in births until the greatest deficit was reached in November (Fig. 1f). Thus, the patterns and trends for esotropia and exotropia were distinctly different in their timing and directionality (see black arrows in Fig. 1e,f). The seasonality of esotropia was statistically significant with the chi-square test ($p<0.0001$) and the KS test ($p<0.01$), and that of exotropia was significant with the chi-square test ($p=0.0002$), but it did not reach significance with the KS test ($p>0.05$). The seasonal birth amplitude for esotropia was 19.55%, and the amplitude for exotropia was 14.65%. Both of these values greatly exceeded the seasonal birth amplitude of the combined normal population (which was 4.95%).

Refractive Error.

Patients with esotropia often have hyperopia, and patients with exotropia often have myopia, as has been reviewed.¹⁴ Therefore, we compared our strabismus births with previous reports of birth seasonality for severe or moderate hyperopia^{15, 17} and severe or moderate myopia.¹⁶ We plotted the birth frequency from the Osaka study¹⁷ against the normal local birth seasonality³⁵ (Fig. 2c). In hyperopia, the seasonal birth amplitude was enhanced more than fourfold, from 4.85% to 23.0% (Fig. 2c). When compared with esotropia births (Fig. 2a), it becomes apparent that the esotropia seasonality is essentially identical to hyperopia ($r=0.8936$, Pearson's R correlation test, Fig. 2a,c), confirming previous conclusions.¹⁵ To compare esotropia and hyperopia with myopia births, we plotted the birth frequency from the 2008 Israel myopia study¹⁶ against the normal seasonal variation of the local population, as reported in 2009.⁴⁵ The resulting graph (Fig. 2d) shows a phase-shift of approximately one month (Fig. 2d), but no change in seasonal birth amplitude (as was observed in hyperopia). The myopia seasonality does not resemble the exotropia seasonality (Fig. 2b,d). The data for myopia from a 2016 study¹⁷ also showed a birth deficit in October through December, but there was no sustained peak in spring and early summer births (data not shown).¹⁷ However, unlike the Israeli study,¹⁶ the 2016 study¹⁷ did not distinguish severe, moderate and mild myopia, and mild myopia showed very little seasonality.¹⁶ We fitted sine waves to the birth seasonalities of refractive error, in order to emphasize the differences with regard to amplitude and phase shift (Fig. 2e,f).

Schizophrenia.

When compared with previously published data for births of patients with schizophrenia, there was no correlation between schizophrenia patients and exotropia patients ($r = 0.01$), and a negative month-by-month correlation between exotropia patients and the majority of schizophrenia patients who have an excess of births in winter and spring months ($r = -0.43$)^{4, 5} (Fig. 3 a). However, the pattern of winter births holds only for patients with schizophrenia who have positive symptoms without enduring negative symptoms (about 80% of schizophrenia cases), but not those with enduring, primary negative symptoms, which has been termed "deficit schizophrenia."^{25–27} This subtype within schizophrenia has a well-replicated excess of births in summer months (pooled odds ratio of 1.95)²¹ (see Table 1; Fig. 3b).^{46–53} When we compared seasonality of births between patients with exotropia and people with schizophrenia of the deficit type, we found that the peak in these summer months was remarkably similar to that of patients with exotropia (Figs. 3a,b). There was a large difference in the similarity of exotropia with deficit and non-deficit schizophrenia births: non-deficit schizophrenia births were negatively correlated with exotropia births (r -values of -0.14 to -0.47);^{21, 46} whereas deficit schizophrenia correlated positively and strongly (r -values ranging from $+0.39$ to $+0.61$),^{21, 46} with an overall r -value of $+0.62$, with a z -score of 2.52, and $p=0.01$, which is highly significant. This is a remarkably robust result, because of the limitation of $n=12$, due to the monthly data collection scheme. To directly show the similarity of birth seasonality in people with exotropia and those with deficit schizophrenia, we present the two curves within the same panel (Fig. 3c).

Regarding the association between exotropia and schizophrenia, the significance of our finding of birth seasonalities in exotropia depends on deficit schizophrenia indeed having an

excess of summer births. Since most work on summer births in deficit schizophrenia originated from one group or network (Dr. Kirkpatrick and his colleagues, see Table 1) and since one of the early studies failed to replicate this pattern,⁴³ we examined the entire literature on this topic, including the most recent studies from Tunisia and Turkey.^{46, 53} As shown in Table 1, nine out of the ten studies support the proposed association of deficit schizophrenia with summer, rather than winter births, while non-deficit schizophrenia is associated with a peak of winter and early spring births in the Northern hemisphere, as has been established for schizophrenia as a whole in multiple meta-analyses.^{4, 5}

Toxoplasmosis.

Since toxoplasmosis has been implicated as a risk factor for both schizophrenia^{23, 24} and eye disease,²² and since toxoplasmosis birth seasonality is known to be correlated with seasonality of births among people with schizophrenia as a whole,²³ we compared the seasonality of births for congenital toxoplasmosis from three published studies^{22, 41, 42} with our strabismus (exotropia) data. These studies either collected data on births²² or maternal seroconversion (typically in the last trimester).^{41, 42} The three studies agree on a peak of maternal infection in winter months.^{22, 23} The peak of toxoplasmosis births is in May (Fig. 3d),²² which precedes the excess exotropia births in July/August by 2–3 months (there is no correlation: r -value=0.0556, Pearson's R correlation test).

Discussion

Developmental factors can affect disease expression and prevalence.^{1–3, 33} Such influences may manifest through seasonal risk factors and may be visible by fluctuations in birth frequencies. Seasonality of births has been extensively studied for schizophrenia,^{4–6} but one, relatively frequent condition that is a risk factor for schizophrenia, exotropic (outwards) horizontal strabismus,¹³ has remained virtually unstudied for seasonal factors. Only one previous study examined seasonality of births in patient populations with horizontal strabismus: a Japanese study in the 1980s.¹⁸ This older and the current study both found that patients with exotropia had a distinct excess of births in the summer months of July and August, while esotropia showed birth months with a later and broader peak, from August to October (our Nevada data) or August to December.¹⁸ The similarity of patterns between the two studies suggests that this seasonality is a global, rather than a local phenomenon. Importantly, the timing of the summer births is very similar to that seen in one type of schizophrenia: deficit schizophrenia, which has previously been associated with distinct oculomotor and vision abnormalities.^{28–30}

Multiple factors may contribute to seasonality of births, including seasonal viral and parasitic maternal infections and immune reactions,^{4, 22, 33, 54} temperature, rainfall,^{55, 56} dietary fluctuations, differences in sperm quality, external toxins,⁴ cultural variation of procreation,^{57, 58} effects of religious rules and customs (including Christmas, the Hajj and Ramadan),^{45, 59} and vitamin D and light (sunshine) exposure.⁵⁵ Interestingly, variation in the length of daylight appears to be a relevant factor that affects eye growth and therefore pertains to myopia/ hyperopia,¹⁶ with a peak sensitive period that was calculated to be at about five months of age.¹⁷

Refractive Errors.

We are intrigued by our finding of a seemingly reverse pattern of esotropia and exotropia seasonality: esotropia shows a gradually *increasing* number of births in the first half of the year, from March until September, while exotropia shows a gradually *decreasing* number of births in the second half of the year, from July through November (compare Fig. 2e with Fig. 2f). This pattern - resembling in some ways a mirror image - suggests a gradually advancing factor in spring/summer for esotropia, possibly due to an increasing period of light or increasing deviation from the average length of the light period,¹⁷ until a distinct peak and reversal point is reached in September (Fig. 1e). On the other hand, exotropia births display an abrupt peak and reversal point in July, followed by a gradual decrease to a minimum in November (Fig. 1f), possibly due to a decreasing period of light or decreasing deviation from the average length of the light period.¹⁷ We found that seasonality of hyperopia and myopia involves a distinct phase-shifting of the sine-wave between the two conditions, as illustrated in Fig. 2e-f. Recognition of these distinct seasonal patterns constitutes an extension of previous work,¹⁷ and strongly suggests that changes in light periods and light exposure are involved, since the peak and turning point between the two conditions' seasonality marks the time of maximal light in the Northern hemisphere. Elucidation of the relevant parameters requires further studies, since the current conclusions rely on only one or two reports for each condition. A third myopia study⁶⁰ generally confirmed the conclusions of the previous two studies,^{16, 18} but did not report monthly births (and did not compare with normal seasonal variation), and therefore could not be evaluated for the apparent phase shift of 1–2 months.⁶⁰ The seasonal and daily changes in the eye and eye growth that occur in response to light and darkness are complex and, despite significant progress, have to date eluded any simple or even comprehensive explanations.⁶¹

Schizophrenia.

Regarding the hypothesis of a similar pattern of births for patients with exotropia and patients with schizophrenia, we did not find a pattern in exotropia patients similar to that of subjects with schizophrenia as a whole (that is, with and without primary negative symptoms), who have an excess of winter/spring births.^{4, 5} This shows that patients with the main type of schizophrenia do not have a birth seasonality similar to patients with exotropia. However, a sizable fraction of patients with schizophrenia (about 20%) have a well-validated, distinct syndrome with primary negative symptoms. Patients with these clinical features differ from those without such features in multiple ways, including other signs and symptoms, course of illness, biological correlates, etiological risk factors, and treatment response.^{25–27} People with this type of schizophrenia (deficit schizophrenia with primary negative symptoms) have a replicated excess of births in summer months (June-July^{21, 47–49, 51} or May-August,^{48, 50, 52} Table 1). It has been concluded that the size of the season-of-birth effect in schizophrenia as a whole is larger in sites further away from the equator, and therefore geographic locations of similar latitudes need to be compared.^{5, 49, 62} Multiple studies examining the seasonality of births in deficit schizophrenia have been conducted at 34–55° latitude in the Northern hemisphere (Table 1), which includes the latitudes of our cohorts in Washoe County (39.5°) and in Osaka (34.7°).¹⁸ The season of birth effect reveals that etiologically distinct subgroups within schizophrenia may be differentially influenced by genes and the environment.⁶³ We wish to emphasize that

similarity in seasonality of births does not prove disease association or any causality between diseases - its presence (similarity) only makes a developmental association more likely, while a dissimilarity makes an association less likely.

Regarding the magnitude of the winter/spring birth excess for schizophrenia as a whole, this is often stated to be about 5–8%.^{4, 5} It should be noted that this excess would be considerably larger for people with schizophrenia of the non-deficit type when one subtracts people with schizophrenia of the deficit type.²¹ Since this subgroup has a substantial excess in summer births (Table 1), this would increase the effect size of winter/spring births for people with non-deficit schizophrenia.

The nature of the association between exotropia and schizophrenia remains unclear, but vision appears to be fundamental for the development of schizophrenia. Congenital blindness (or very early loss of vision) has been reported to protect against developing schizophrenia,^{64, 65} while abnormal vision and/or abnormal early visual experiences (as would be frequent in children with exotropia) appears to be a significant risk factor for schizophrenia.¹³

Our findings raise the possibility that exotropia is a risk factor specifically for deficit schizophrenia. The question whether people with deficit schizophrenia have exotropia more often than would otherwise be expected has not yet been tested, but should be explored. If exotropia is indeed a risk factor primarily for development of schizophrenia with negative symptoms, this could provide new insights into the pathophysiological etiology of this type of schizophrenia.

It has been recommended that children diagnosed with exotropia should be monitored for mental illness and in particular signs of psychosis and development of schizophrenia.^{7, 8, 12} Recent work did not find that surgical correction of childhood exotropia significantly affected the rate of mental illness in younger people (mental illness diagnosed at 8–18 years of age),⁶⁶ but the question for schizophrenia, which has a peak for first diagnosis at 18–25 years of age for males and 25–35 for females,⁶⁷ remains unexplored.

Limitations of our study include that our subject numbers, even when cohorts were combined (n=6,020 for esotropia, and n=5,520 for exotropia), were smaller and therefore statistically less robust than some previous studies, e.g., on myopia (~30,000 subjects with severe or moderate myopia).¹⁶ We investigated only strabismus, but not schizophrenia and its different subtypes, in the same population, so we had to compare our data with those from previous studies at other sites (Table 1). Future research with additional cohorts is needed to confirm that exotropia is associated with a peak of births in summer months, to explore possible reasons for the seasonality of exotropia births, and to determine whether exotropia is indeed linked primarily to the deficit type of schizophrenia rather than schizophrenia with positive symptoms. This line of research may identify strategies for disease prevention, since some developmental risk factors such as light exposure and misalignment of eyes during infancy can be manipulated or controlled.

Conclusions

People with esotropia and exotropia have seasonality of births, as do people with hyperopia or myopia, but with distinctly different patterns when compared with the seasonal variation in the normal population. Exotropia showed significant excess in summer births that correlated with the seasonality of births for deficit schizophrenia, but not schizophrenia as a whole. Further studies are needed to confirm the link between exotropia and deficit schizophrenia, and explore its etiology.

Acknowledgments

We thank Kyra Morgan, Chief Biostatistician, Nevada Division of Public and Behavioral Health, Carson City, Nevada, and the office staff of Sierra Eye Associates, Reno, Nevada for help with data collection and processing. We also thank Dr. Fuller Torrey (Stanley Medical Research Institute, Kensington, MD) for helpful comments on this project.

Financial Support

This work was supported by grants from the National Institutes of Health (C.S.v.B, GM103554 and GM104944).

References

1. Wadhwa PD, Buss C, Entringer S, Swanson JM. Developmental origins of health and disease: brief history of the approach and current focus on epigenetic mechanisms. *Semin Reprod Med.* 2009; 27, 358–368. [PubMed: 19711246]
2. Gluckman PD, Hanson MA, Buklijas T. A conceptual framework for the developmental origins of health and disease. *J Dev Orig Health Dis.* 2010; 1, 6–18. [PubMed: 25142928]
3. Almond D, Currie J. Killing me softly: The fetal origins hypothesis. *J Econ Perspect.* 2011; 25, 153–172. [PubMed: 25152565]
4. Torrey EF, Miller J, Rawlings R, Yolken RH. Seasonality of births in schizophrenia and bipolar disorder: a review of the literature. *Schizophr Res.* 1997; 28, 1–38. [PubMed: 9428062]
5. Davies G, Welham J, Chant D, Torrey EF, McGrath J. A systematic review and meta-analysis of Northern Hemisphere season of birth studies in schizophrenia. *Schizophr Bull.* 2003; 29, 587–593. [PubMed: 14609251]
6. Bembeneck A Seasonality of birth in schizophrenia patients. Literature review. *Psychiatr Pol.* 2005; 39, 259–270. [PubMed: 15881621]
7. Schiffman J, Maeda JA, Hayashi K, et al. Premorbid childhood ocular alignment abnormalities and adult schizophrenia-spectrum disorder. *Schizophr Res.* 2006; 81, 253–260. [PubMed: 16242918]
8. Mohny BG, McKenzie JA, Capo JA, et al. Mental illness in young adults who had strabismus as children. *Pediatrics.* 2008; 122, 1033–1038. [PubMed: 18977984]
9. Korn H Schizophrenia and eye movement - a new diagnostic and therapeutic concept. *Med Hypotheses.* 2004; 62, 29–34. [PubMed: 14729001]
10. Toyota T, Yoshitsugu K, Ebihara M, et al. Association between schizophrenia with ocular misalignment and polyalanine length variation in PMX2B. *Hum Mol Genet.* 2004; 13, 551–561. [PubMed: 14709596]
11. Yoshitsugu K, Yamada K, Toyota T, et al. A novel scale including strabismus and ‘cuspidal ear’ for distinguishing schizophrenia patients from controls using minor physical anomalies. *Psychiatry Res.* 2006; 145, 249–258. [PubMed: 17070930]
12. Ndlovu D, Nhleko S, Pillay Y, et al. The prevalence of strabismus in schizophrenic patients in Durban, KwaZulu Natal. *S Afr Optom.* 2011; 70, 101–108.
13. Agarwal AB, Christensen AJ, Feng CY, et al. Expression of schizophrenia biomarkers in extraocular muscles from patients with strabismus: an explanation for the link between exotropia and schizophrenia? *PeerJ.* 2017; 5, e4214. [PubMed: 29302405]

14. von Noorden GK, Campos EC. Binocular Vision and Ocular Motility, 6th edn, 2002 Mosby: St. Louis.
15. Matsuda K, Yokoyama T, Nakagawa M, et al. Statistical analysis of the incidence of hyperopia by month of birth. *Nippon Ganka Kiyo (Folia Ophthalmol Jpn)* 1986; 37, 440–443.
16. Mandel Y, Grotto I, El-Yaniv R, et al. Season of birth, natural light, and myopia. *Ophthalmology*. 2008;115, 686–692. [PubMed: 17698195]
17. Matsuda K, Yokoyama T, Fukui M, Morita T, Kozaki M. Inverse Analysis of the Relationship between Ocular Refraction and Birth Month Using Matrix Equations. *Nippon Ganka Gakkai Zasshi (J Jpn Ophthalmol Soc)* 2016; 120, 533–539.
18. Matsuda K, Nakagawa M, Yokoyama T, et al. Incidence of strabismus by season of birth. *Ganka Rinsho Iho (Jpn Rev Clin Ophthalmol)* 1985; 79, 2115–2118.
19. Chew E, Remaley NA, Tamboli A, et al. Risk factors for esotropia and exotropia. *Arch Ophthalmol*. 1994; 112, 1349–1355. [PubMed: 7945039]
20. Saha S, Chant D, Welham J, McGrath J. A systematic review of the prevalence of schizophrenia. *PLoS Med*. 2005; 2:e141.
21. Messias E, Kirkpatrick B, Bromet E, et al. Summer birth and deficit schizophrenia: a pooled analysis from 6 countries. *Arch Gen Psychiatry*. 2004; 61, 985–989. [PubMed: 15466671]
22. Meenken C, Rothova A, Kijlstra A, Oosting J. Seasonal variation in congenital toxoplasmosis. *Br J Ophthalmol*. 1991; 75, 639.
23. The Stanley Medical Research Institute. Epidemiological similarities and differences between toxoplasmosis and schizophrenia. Updated July, 2017 Retrieved from <http://www.stanleyresearch.org/patient-and-provider-resources/toxoplasmosis-schizophrenia-research/epidemiological-similarities-and-differences-between-toxoplasmosis-and-schizophrenia/>.
24. Torrey EF, Bartko JJ, Yolken RH. *Toxoplasma gondii* and other risk factors for schizophrenia: an update. *Schizophr Bull*. 2012; 38, 642–647. [PubMed: 22446566]
25. Carpenter WT Jr, Heinrichs DW, Wagman AM. Deficit and nondeficit forms of schizophrenia: the concept. *Am J Psychiatr*. 1988; 145, 578–583. [PubMed: 3358462]
26. Kirkpatrick B, Buchanan RW, Ross DE, Carpenter WT Jr. A separate disease within the syndrome of schizophrenia. *Arch Gen Psychiatry*. 2001; 58, 165–171. [PubMed: 11177118]
27. Kirkpatrick B, Galderisi S. Deficit schizophrenia: an update. *World Psychiatry*. 2008; 7, 143–147. [PubMed: 18836581]
28. Ross DE, Thaker GK, Buchanan RW, et al. Eye tracking disorder in schizophrenia is characterized by specific ocular motor defects and is associated with the deficit syndrome. *Biol Psychiatry*. 1997; 42, 781–796. [PubMed: 9347127]
29. Ross DE. The deficit syndrome and eye tracking disorder may reflect a distinct subtype within the syndrome of schizophrenia. *Schizophrenia Bull*. 2000; 26, 855–866.
30. Nkam I, Thibaut F, Denise P, et al. Saccadic and smooth-pursuit eye movements in deficit and non-deficit schizophrenia. *Schizophr Res*. 2001; 48, 145–153. [PubMed: 11278161]
31. von Bartheld CS, Agarwal AB, Feng CY, et al. Why is exotropia (one specific type of strabismus) a significant risk factor for schizophrenia? 5th Biennial Schizophrenia International Research Society Conference, Florence, Italy, April 2–6, 2016 *npj Schizophrenia*. 2016; 78,
32. Choi TB, Lee DA, Oelrich FO, et al. A retrospective study of eye disease among first grade children in Los Angeles. *J Am Optom Assoc*. 1995; 66, 484–488. [PubMed: 7494083]
33. Martinez-Bakker M, Bakker KM, King AA, Rohani P. Human birth seasonality: latitudinal gradient and interplay with childhood disease dynamics. *Proc Biol Sci*. 2014; 281, 20132438. [PubMed: 24695423]
34. Maruo T, Kubota N, Arimoto H. The results of examinations for strabismus and amblyopia in elementary and secondary school pupils (in Japanese). *Ganka Rinsho Iho (Jpn Rev Clin Ophthalmol)*. 1977; 71, 712–714.
35. Matsuda S, Kahyo H. Geographical differences and time trends in the seasonality of birth in Japan. *Int J Epidemiol*. 1994; 23, 107–118. [PubMed: 8194905]
36. Bradbury TN, Miller GA. Season of birth in schizophrenia: a review of evidence, methodology, and etiology. *Psychol Bull*. 1985; 98, 569–594. [PubMed: 4080898]

37. Verdoux H, Takei N, Cassou de Saint-Mathurin R, Bourgeois M. Analysis of the seasonal variation of schizophrenic births using a Kolmogorov-Smirnov type statistic. *Eur Psychiatry*. 1997; 12, 111–116. [PubMed: 19698516]
38. Nwogu EC, Iwueze IS, Nlebedim VU. Some tests for seasonality in time series data. *J Mod Appl Stat Meth*. 2016; 15, 382–399.
39. Freedman LS. The use of a Kolmogorov--Smirnov type statistic in testing hypotheses about seasonal variation. *J Epidemiol Community Health*. 1979; 33, 223–228. [PubMed: 509003]
40. Walter SD. The power of a test for seasonality. *Br J Prev Soc Med*. 1977; 31, 137–140. [PubMed: 884398]
41. Logar J, Soba B, Premru-Srsen T, Novak-Antolic Z. Seasonal variations in acute toxoplasmosis in pregnant women in Slovenia. *Clin Microbiol Infect*. 2005; 11, 852–855. [PubMed: 16153265]
42. Sagel U, Mikolajczyk RT, Krämer A. Seasonal trends in acute toxoplasmosis in pregnancy in the federal state of Upper Austria. *Clin Microbiol Infect*. 2010; 16, 515–517. [PubMed: 19622079]
43. Dollfus S, Brazo P, Langlois S, et al. Month of birth in deficit and non-deficit schizophrenic patients. *Eur Psychiatry*. 1999; 14, 349–351. [PubMed: 10572368]
44. Lam DA, Miron JA. Global patterns of seasonal variation in human fertility. *Ann N Y Acad Sci*. 1994; 709, 9–28. [PubMed: 8154738]
45. Friger M, Shoham-Vardi I, Abu-Saad K. Trends and seasonality in birth frequency: a comparison of Muslim and Jewish populations in southern Israel: daily time series analysis of 200 009 births, 1988–2005. *Hum Reprod*. 2009; 24, 1492–1500. [PubMed: 19246466]
46. Kallel L, Brunelin J, Zghal A, et al. Summer birth and deficit schizophrenia in Tunisia. *Psychiatry Res*. 2007; 152, 273–275. [PubMed: 17482273]
47. Kirkpatrick B, Ram R, Amador XF, et al. Summer birth and the deficit syndrome of schizophrenia. *Am J Psychiatry*. 1998; 155, 1221–1226. [PubMed: 9734546]
48. Kirkpatrick B, Castle D, Murray RM, Carpenter WT Jr. Risk factors for the deficit syndrome of schizophrenia. *Schizophr Bull*. 2000; 26, 233–242. [PubMed: 10755684]
49. Messias E, Kirkpatrick B. Summer birth and deficit schizophrenia in the epidemiological catchment area study. *J Nerv Ment Dis*. 2001; 189, 608–612. [PubMed: 11580004]
50. Tek C, Kirkpatrick B, Kelly C, McCreadie RG. Summer birth and deficit schizophrenia in Nithsdale, Scotland. *J Nerv Ment Dis*. 2001; 189, 613–617. [PubMed: 11580005]
51. Kirkpatrick B, Tek C, Allardyce J, Morrison G, McCreadie RG. Summer birth and deficit schizophrenia in Dumfries and Galloway, southwestern Scotland. *Am J Psychiatry*. 2002; 159, 1382–1387. [PubMed: 12153832]
52. Kirkpatrick B, Herrera Castanedo S, Vazquez-Barquero JL. Summer birth and deficit schizophrenia: Cantabria, Spain. *J Nerv Ment Dis*. 2002; 190, 526–532. [PubMed: 12193837]
53. Mete L, Sarikaya Ö, Erol A. The relationship of deficit syndrome with clinical symptoms, summer births and heritability in patients with schizophrenia. *Turk Psikiyatri Derg*. 2015; 26, 229–235. [PubMed: 26731019]
54. Brown AS, Patterson PH. Maternal infection and schizophrenia: implications for prevention. *Schizophr Bull*. 2011; 37, 284–290. [PubMed: 21134972]
55. McGrath J, Selten JP, Chant D. Long-term trends in sunshine duration and its association with schizophrenia birth rates and age at first registration—data from Australia and the Netherlands. *Schizophr Res*. 2002; 54, 199–212. [PubMed: 11950544]
56. Messias E, Mourao C, Maia J, et al. Season of birth and schizophrenia in Northeast Brazil: relationship to rainfall. *J Nerv Ment Dis*. 2006; 194, 870–873. [PubMed: 17102713]
57. Häfner H, Haas S, Pfeifer-Kurda M, Eichhorn S, Michitsuji S. Abnormal seasonality of schizophrenic births. A specific finding? *Eur Arch Psychiatry Neurol Sci*. 1987; 236, 333–342. [PubMed: 3678292]
58. Tochigi M, Okazaki Y, Kato N, Sasaki T. What causes seasonality of birth in schizophrenia? *Neurosci Res*. 2004; 48, 1–11. [PubMed: 14687876]
59. Odegård O. Season of birth in the population of Norway, with particular reference to the September birth maximum. *Br J Psychiatry*. 1977; 131, 339–344. [PubMed: 922262]

60. McMahon G, Zayats T, Chen YP, et al. Season of birth, daylight hours at birth, and high myopia. *Ophthalmology*. 2009; 116, 468–473. [PubMed: 19157564]
61. Nickla DL. Ocular diurnal rhythms and eye growth regulation: where we are 50 years after Lauber. *Exp Eye Res*. 2013; 114, 25–34. [PubMed: 23298452]
62. Welham J, Chant D, Saha S, et al. No association between the deficit syndrome in psychosis and summer birth in a Southern hemisphere country. *Aust N Z J Psychiatry*. 2006; 40, 935–936. [PubMed: 16959020]
63. Franzek E, Beckmann H. Gene-environment interaction in schizophrenia: season-of-birth effect reveals etiologically different subgroups. *Psychopathology*. 1996; 29, 14–26. [PubMed: 8711071]
64. Silverstein SM, Wang Y, Keane BP. 2013 Cognitive and neuroplasticity mechanisms by which congenital or early blindness may confer a protective effect against schizophrenia. *Frontiers Psychol*. 3, Article 624.
65. Landgraf S, Osterheider M. 2013 “To see or not to see: that is the question.” The “Protection-Against-Schizophrenia” (PaSZ) model: evidence from congenital blindness and visuo-cognitive aberrations. *Frontiers Psychol*. 4, Article 352.
66. Kilgore KP, Barraza RA, Hodge DO, McKenzie JA, Mohny BG. Surgical correction of childhood intermittent exotropia and the risk of developing mental illness. *Am J Ophthalmol*. 2014; 158, 788–792. [PubMed: 24954680]
67. Ochoa S, Usall J, Cobo J, Labad X, Kulkarni J. Gender differences in schizophrenia and first-episode psychosis: a comprehensive literature review. *Schizophr Res Treatment*. 2012; 916198. [PubMed: 22966451]

Fig. 1.

Comparison of seasonality of births for horizontal strabismus from two different sites: data from Washoe County, Nevada (a, b), and Osaka, Japan (c, d), as previously reported,¹⁸ but without normalization to the local population. Esotropia (a, c, e); Exotropia (b, d, f). Note the similarities in the timings of peaks and troughs between the two sites. (Panel a) Esotropia births ($n=2,513$, red squares) are shown relative to normal Washoe County births (blue dots). (Panel b) Exotropia births ($n=2,332$, red squares) are shown relative to births in the normal Washoe County population (blue dots). (Panel c) Esotropia births from the Osaka cohort ($n=3,507$, orange dots)¹⁸ are shown relative to the normal local birth seasonality (blue dots).³⁵ (Panel d) Exotropia births from the Osaka cohort ($n=3,188$, orange dots)¹⁸ are shown relative to the normal local birth seasonality (blue dots).³⁵ Note the similarity of patterns in seasonality of births between cohorts. (Panel e) The distribution of births from the *combined* esotropia cohorts (Nevada and Osaka, $n=6,020$, orange dots), are shown relative to the combined normal local variation (blue dots). The black arrow denotes the gradually increasing number of esotropia births from a trough in March to the peak in September. (Panel f) The distribution of births from the *combined* exotropia cohorts (Nevada and Osaka, $n=5,520$, orange dots) are shown relative to the combined normal local variation (blue dots). The black arrow denotes the gradually decreasing number of exotropia births from a peak in July to the trough in November. Note the “reverse” or “mirror image” seasonality pattern between esotropia (e) and exotropia (f).

Fig. 2.

Comparison of seasonality of births in horizontal strabismus and refractive errors: esotropia (a), exotropia (b), hyperopia (c, e), myopia (d, f). (Panels a-d) Data from subjects with strabismus or with refractive errors are shown in orange, relative to the normal local population (blue). Data for esotropia and exotropia are the combined cohorts from Nevada and Osaka.¹⁸ Data for hyperopia are from Osaka,^{17, 35} and data for myopia are from Israel.^{16, 45} Note that the birth distribution for esotropia (a) and hyperopia (c) is very similar, while the birth distribution for exotropia (b) and myopia (d) is not. Hyperopia and myopia birth distributions form amplitude- or phase-altered sine waves, with hyperopia increasing the amplitude of the normal seasonal variation (c, e), while myopia has the same amplitude, but is phase-shifted (precocious) by about 1 month compared with the normal seasonal variation (d, f). Hyperopia births have the minimum in spring (c, e), while myopia has the minimum in winter (d, f). The seasonality curves were modeled as sine waves in panels e (hyperopia) and f (myopia). A , amplitude; A_h , amplitude for hyperopia; A_n , Amplitude for normal population; α_m , phase shift for myopia; α_n , phase shift for normal population, P , period (=wavelength), $1/2P$, one half period. Note that a sine wave's amplitude is defined differently than a birth seasonality's amplitude.³³

Fig. 3.

Comparison of seasonality of births for exotropia, two types of schizophrenia, and congenital toxoplasmosis. (Panel a) Exotropia births (orange squares) compared with schizophrenia births (blue diamonds) for similar latitudes (30–39°).⁵ (Panel b) Comparison of births for deficit schizophrenia and non-deficit schizophrenia, compiled from four data sets.^{21, 46} Data from Messias et al., 2004²¹ were redrawn with permission. Note that the summer peak extends into August in at least 4 of the 9 studies showing an excess of summer births (Table 1). Note also that the period of excess births in schizophrenia as a whole

extends between December and May, with some variation, but always occurs in winter/spring. Error bars = SEM. *, data points are significantly different, $p < 0.01$, t-test. (Panel c) Comparison of the birth seasonalities of people with exotropia (combined cohorts from Nevada and Osaka) and people with deficit schizophrenia (combined cohorts from UK, Ireland, Spain, New York, and Tunisia). The graph for deficit schizophrenia was plotted as percent births per month (purple scale on left side, as in the original studies), while the exotropia graph was plotted as deviation from the mean (orange scale on right side). Note the similarities between the two seasonality graphs, and thus between the two disease conditions: exotropia and deficit schizophrenia. (Panel d) Comparison of seasonality of toxoplasmosis births²² (purple dots) and the current study's data on exotropia (orange dots). A similar seasonality was reported in two other studies for toxoplasmosis, supporting maternal infection in winter, and an increased number of affected offspring born in spring.^{41, 42} Note that there is no similarity between the timing of toxoplasmosis infection and exotropia births (r-value=0.0556, Pearson's R correlation test).

Table 1.

Studies on seasonality of births comparing deficit and non-deficit schizophrenia (SZ) in the Northern hemisphere.

Authors and Year	Reference Number	Location	Latitude [°]	Number of Subjects with Deficit SZ	Number of Subjects with nonDeficit SZ	Months with Excess Births (Deficit SZ)
Kirkpatrick et al., 1998	47	Maryland, USA	39.3	26	65	June
Kirkpatrick et al., 1998	47	“Northern Hemisphere”		55	101	July
Kirkpatrick et al., 1998	47	Long Island, NY, USA	40.8	32	52	June/July
Dollfus et al., 1999	43	France	43.3–49.4	53	158	Jan/Feb/Mar
Kirkpatrick et al., 2000	48	UK, Caribbean, Africa, India		26	118	May-August
Messias & Kirkpatrick, 2001	49	Five sites (ECA, USA) *	34.1–41.3	77	146	June/July
Tek et al., 2001	50	Nithsdale, UK	55.4	51	57	June-August
Kirkpatrick et al., 2002	51	Dumfries/Galloway, UK	55.1	65	277	June-August
Kirkpatrick et al., 2002	52	Cantabria, Spain	43	22	55	May-August
Messias et al., 2004	21	UK, Spain, USA	41–55	401	1,193	June/July
Kallel et al., 2007	46	North Tunisia, Africa	35–37	34	46	June/July
Mete et al., 2015	53	Izmir, Turkey	38	35	65	“Summer”
Sums				877	2,333	

* ECA, Epidemiological Catchment Area of the National Institute of Mental Health