Published in final edited form as:

Curr Opin Anaesthesiol. 2019 June ; 32(3): 405–411. doi:10.1097/ACO.0000000000000722.

The Malnourished Surgery Patient: A Silent Epidemic in Perioperative Outcomes?

David G.A. Williams, MD, MPH¹, Jeroen Molinger, MSc^{1,2,3}, Paul E. Wischmeyer, MD, EDIC¹

¹Department of Anesthesiology, Duke University School of Medicine, Durham, North Carolina ²-
Erasmus MC, Erasmus University Rotterdam Dept of Intensive Care, Rotterdam, Netherlands ³-
Clinical Human Performance Center BeLife, Rotterdam, Netherlands

Abstract

Purpose of Review: As many as 2 of every 3 major surgery patients are malnourished preoperatively – a diagnosis rarely made and treated even less frequently. Unfortunately, perioperative malnutrition is perhaps the least often identified surgical risk factor and is among the most treatable to improve outcomes.

Recent findings: Two important perioperative nutrition guidelines were published recently. Both emphasize nutrition assessment as an essential component of preoperative screening. The recently published Perioperative Nutrition Screen (PONS) readily identifies patients at malnutrition risk, allowing for preoperative nutritional optimization. The use of CT scan and ultrasound lean body mass (LBM) evaluation to identify sarcopenia associated with surgical risk and guide nutrition intervention is garnering further support. Preoperative nutrition optimization in malnourished patients, use of immunonutrition in all major surgery, avoidance of preoperative fasting, inclusion of post-operative high-protein nutritional supplements, and early postoperative oral intake have all recently been shown to improve outcomes and should be utilized.

Summary: The recent publication of new surgical nutrition guidelines, the PONS score and use of LBM assessments will allow better identification and earlier intervention on perioperative malnutrition. It is essential that in the future no patient undergoes elective surgery without nutrition screening and nutrition intervention when malnutrition risk is identified.

Keywords

nutrition; nutrition screening; malnutrition; protein; oral nutrition supplement; immunonutrition; lean body mass; ultrasound; sarcopenia; muscle

Background

Perioperative malnutrition is a known independent predictor of poor postoperative outcomes. In fact, it has been determined that malnourished surgical patients experience higher

postoperative mortality, morbidity, length of stay (LOS), hospital readmission rates, and hospital costs [1,2]. Data from the *National Surgical Quality Improvement Program* (NSQIP) demonstrates malnutrition is among the only major readily-modifiable preoperative risk factors associated with poor surgical outcomes, including mortality [2]. While much attention has been paid to the physiological and pharmacological management of the surgical patient, the importance of perioperative nutrition optimization continues to be poorly appreciated as shown by our recent survey of perioperative nutrition practices [3]. It is estimated that greater than 1 in every 3 hospitalized patients is malnourished at hospital admission and the Healthcare Cost and Utilization Project (HCUP) indicates that only 3% of these patients are being properly identified and diagnosed during their hospitalization [4]. Consequently, only 1 in 10 malnourished patients are ever diagnosed and even fewer are effectively treated. Also concerning is that a diagnosis of malnutrition rarely leads to adequate intervention. Large dataset analysis of ~2 million hospitalized U.S. patients who were diagnosed with malnutrition reveals that fewer than 7% of these malnutrition-related hospital stays included nutrition intervention aimed at improving nutrition in a meaningful way [4]. Accordingly, only 1 in 100 malnourished patients in U.S. hospitals are treated for malnutrition. Truly, malnutrition is a silent epidemic occurring daily in our care of patients.

Perioperative medicine is an emerging specialty that emphasizes medical care of patients from the time of contemplation of surgery through the operative period to full recovery [5**]. As the role of the anesthesiologist is being redefined, fundamental to perioperative medicine is a coordinated, multidisciplinary care team model aimed at utilizing evidenced-based best practice techniques to optimize patient comorbidities in order to maximize successful patient outcomes [5**]. Recently published key evidenced-based consensus surgical guidelines indicate malnutrition is a key modifiable preoperative risk factor associated with poor surgical outcomes that we must address [6**,7**,8]. We must, therefore, improve our ability to preoperatively identify the patient at risk for perioperative malnutrition and utilize evidence-based nutrition optimization techniques if we are to improve surgical outcomes and deliver cost effective medical care.

How Can We Improve Identification of the Malnourished Surgical Patient: The Perioperative Nutrition Screen (PONS) Score

Preoperative identification of the malnourished surgical patient is paramount to improving nutrition status during the perioperative period [6**,7**,8]. Screening for malnutrition prior to elective surgery can identify patients at risk for malnutrition who may benefit from preoperative nutritional intervention. The risk of perioperative malnutrition is often most significant for oncologic and gastrointestinal surgeries, where 2 out every 3 patients presenting for surgery are malnourished pre-operatively [9,10]. These high-risk surgeries are also often targeted by enhanced recovery programs allowing for more meaningful optimization to occur [6, 11**]. While several screening tools have been validated for use in hospitalized patients, no universally accepted screening tool for preoperative malnutrition risk has been available. A key recent publication from the Perioperative Quality Initiative (POQI) recommends use of the Perioperative Nutrition Screen (PONS) for preoperative assessment of malnutrition [6**]. (See Figure 1) PONS was developed as a modified version

of the well-validated malnutrition universal screening tool (MUST) and identifies nutrition risk based on commonly utilized malnutrition questions. Each question, including an albumin level < 3.0 is assigned 1 point for a “positive” response (maximum PONS score of 3). Perioperative malnutrition risk is further assessed based on the patient’s preoperative vitamin D and albumin levels (Figure 1). Any patient with PONS ≥ 1 (any positive response to first 3 questions) and/or an albumin < 3.0 (and/or a vitamin D < 20) is considered high risk for perioperative malnutrition [6**] and should receive pre-operative nutrition intervention prior to surgery as described below. If available, referral to a registered dietician for further preoperative nutritional evaluation is also suggested. It has been shown that underweight patients (BMI levels < 18.5 kg/m² for adults < 65 years old) undergoing major surgery have an increase in postoperative complications [12,13]. A longitudinal study by Sergi et al described the nonsurgical risk of malnutrition based on age and found that the risk for all-cause mortality increases starting at a BMI of 24 kg/m² for patients > 65 and doubles when BMI is < 22 kg/m² for men and < 20 kg/m² for women [12]. The PONS, therefore, is based on this association between BMI and all-cause risk, and uses a higher BMI (< 20 kg/m²) for adults > 65 years as it is suggested that higher BMI thresholds should be used for older adults. Irrespective of BMI, unintentional weight loss has been associated with negative postoperative outcomes including morbidity and functional decline [14]. Although limited as a nutritional marker in post-operative setting and states of systemic inflammation, albumin has been long utilized as malnutrition indicator and preoperative albumin levels are a strong predictor of postoperative complications, including mortality [15,16]. Collectively, these components contribute to a thorough preoperative malnutrition screening tool.

The PONS allows quick and efficient preoperative malnutritional risk identification and should be routinely performed on all patients coming for major surgery during preoperative assessment. Delaying major elective surgery should be considered in patients with any positive finding on PONS score, which identifies patient as being at risk for malnutrition, to allow for adequate nutrition optimization [6**]. Although the optimal time period for preoperative nutrition optimization is unclear, it appears a minimum of 7–10 days should occur, and the risk of disease progress from delaying surgery should be weighed against the significant risk of operating on a malnourished patient.

Role of Lean Body Mass and Sarcopenia in Surgical Outcomes

Sustained malnutrition can lead to the reduction in quantity and quality of lean body mass (LBM) – sarcopenia. Within the last 10 years, this syndrome has received increasing attention as a possible predictor of adverse outcomes after surgery [17,18]. In contrast to traditional perioperative nutritional assessments that typically focus on weight loss and serum markers of nutrition (i.e. – serum albumin levels), sarcopenia is a potentially much more objective indicator of nutritional and metabolic reserve prior to surgery [19**]. Currently, sarcopenia largely remains undiagnosed [20,21] as it is often difficult to differentiate from overall weight loss, especially in obese patients [22]. Computerized tomography (CT) scan LBM analysis has begun to be utilized and studied around the world to assess preoperative “metabolic reserve” and “fitness for surgery” [19**]. This innovative technique may effectively identify patients at higher risk for malnutrition and poor surgical outcomes [19**].

In surgical patients, a rapidly growing body of literature demonstrates low baseline LBM may be an independent risk factor for complications in patients undergoing hepatic [23] colorectal [24,25], diverticular [26*], and pancreatic [27] oncological surgery. These findings are supported by a recent meta-analysis evaluating sarcopenia as a predictor of postoperative complication risk post-gastrointestinal cancer surgery [28**]. Twenty-nine studies (n = 7176) were evaluated utilizing a range of CT-scan sarcopenia measures and found perioperative sarcopenia prevalence ranges from 12%–78%. Preoperative sarcopenia was associated with increased risk of major complications (risk ratio 1.40; 95% C.I. 1.20–1.64; $P<0.001$) and total complications (risk ratio 1.35; 95% C.I. 1.12–1.61; $P=0.001$). While sarcopenia was associated with an increased risk of complications after gastrointestinal tumor resection, these retrospective studies lack methodological consensus, limiting interpretation and clinical utilization of these findings [28**].

Sarcopenia carries a significant health cost burden. For patients undergoing major general or vascular surgery, decreasing LBM is associated with increased insurer costs. Sarcopenic patients had a mean payer cost of \$34,796, versus \$21,380.6 in non-sarcopenic patients [29]. Further, CT analysis of LBM in 452 patients (median age 65, 61.5% males) undergoing surgery for colorectal cancer (38.9%), colorectal liver metastases (27.4%), primary liver tumors (23.2%), and pancreatic/periampullary cancer (10.4%) showed 45.6% had sarcopenia and post-adjustment for confounders, low LBM was associated with a cost increase of €4,061 ($P=0.015$) [30**]. Thus, initial data indicate in some surgical populations sarcopenia is independently associated with increased costs. Sarcopenia treatment with nutrition and exercise interventions may reduce hospital costs in an era of escalating health-care costs and an increasingly ageing population [31**].

Given observational evidence for role of LBM in perioperative outcome, we currently require prospective trials examining LBM and surgical outcomes. We urgently need objective methods to preoperatively measure sarcopenia and define objective risk cut-offs for sarcopenia-based measures based on magnitude and type of surgical intervention. Further, as CT Scan is not practical for all patients to undergo for LBM assessment alone and not realistic for longitudinal measures, the use of bedside ultrasound for LBM measurement has become an ideal option. To address the potential role of low LBM in perioperative optimization, a highly innovative and novel muscle-specific ultrasound (U/S) device has been developed (Musclesound, Denver, CO). This new U/S technology provides bedside measures of LBM, intramuscular glycogen and intramuscular adipose tissue (IMAT) (See Figure 2 for example). IMAT has recently been correlated to muscle strength [32*]. Intramuscular glycogen U/S measures have been validated via muscle biopsy [33] and we showed acutely ill patients have significant intramuscular glycogen deficits [34]. Intramuscular glycogen is known to change daily based on adequacy of nutrition intake and “physical stress”, and could prove useful in monitoring nutrition delivery and utilization in perioperative patients [35**]. This promising technology requires investigation in the perioperative setting. In closing, LBM could serve as a key predictor of preoperative risk and aid in identifying patients in need for preoperative optimization.

Preoperative Nutrition Optimization Improves Surgical Outcomes

A summary of our structured perioperative nutrition protocol is shown in Figure 3. Undergoing surgery elicits a state of metabolic and physiologic stress on the human body and there is an increase in production of hepatic acute phase protein synthesis, proteins involved in immune function, and proteins required for wound healing [6**,7**,8]. Perioperative fasting can exacerbate the surgical stress response and intensify protein loss so preoperative fasting should be avoided.

Protein intake requirements are therefore increased and when protein intake is insufficient to meet increased demands of protein synthesis, LBM breakdown becomes the source of amino acid. While optimal protein intake for surgical patients has yet to be definitively determined, current nutrition guidelines suggest intake of a minimum of 1.2–2.0 g of protein/kg/d for stressed patients [6**]. Several studies show consuming 25–35 g of protein in a single meal maximally stimulates muscle protein synthesis, therefore daily protein requirements for patients at malnutrition risk should be addressed by consumption of 25–35 g of protein – particularly whey protein and casein – at every meal to provide high quality amino acids required to stimulate LBM synthesis [36, 37].

The time required for adequate preoperative nutrition optimization and a gold standard marker to quantify optimization progress has not yet been identified. Recent consensus recommendations from the North American Surgical Nutrition Summit emphasizes that nutritional care should be established preoperatively for both malnourished and normal nourished patients in order to foster optimal nutritional status throughout perioperative period [38].

In patients with low risk of perioperative malnutrition (i.e. – PONS <1 and Albumin [ALB] >3.0) high-protein, complex carbohydrate-rich diets should be consumed preoperatively [6**]. Many patients will not be able to meet the suggested perioperative energy goals of 25 kcal/kg/d and 1.5–2 g/kg/d of protein solely from food intake. Thus, it is recommended to encourage patients to take high-protein oral nutritional supplements (HP-ONSs). Immunonutrition (IMN, containing arginine/fish oil) has numerous trials over many years supporting its use in all patients having major GI, cardiac, and ENT surgery. This benefit is independent of malnutrition risk and should be utilized in all major surgery patients where it has been shown to reduce infections and complications by ~40% and significantly shorten length of stay [39]. This association has shown to be true not only when IMN is used in ERAS pathways [40] but also in a large scale prospective cohort study with a propensity score-matched comparative effectiveness evaluation [41*].

Patients determined to be at risk of malnutrition (i.e. – PONS >1 or ALB <3.0) should be prescribed HP-ONS prior to any elective surgery for 2–6 weeks [6**,42**]. These patients should consume a minimum of 1.2 g/kg/d total of protein and the HP-ONS should contain >18 g/protein/serving given at least twice a day [6**]. Follow-up of patient compliance with ONS intake is essential for benefit [6**].

As the role of the anesthesiologist expands to include perioperative nutrition optimization, dietitians are integral to the perioperative care team. For malnourished patients who cannot

meet protein/calorie requirements via oral nutrition, a dietician should be consulted and home enteral nutrition (EN) initiated for a period of at least 7 days pre-operatively. Preoperative parenteral nutrition (PN) should be utilized in patients with malnutrition when >50% of recommended kcal/protein requirement cannot be adequately met by ON or EN [6**].

Management of Postoperative Nutrition

Early oral feeding immediately post-major surgery, including GI surgery, is associated with a decrease in postoperative complications, LOS, and hospital costs [6**,7**,42**]. Specifically, convincing evidence shows feeding within 24 hours of GI surgery decreases mortality and major morbidities [43]. High protein diets are essential to assist in maintaining lean muscle mass in the postoperative period [6**]. Therefore, except for patients with bowel discontinuity, ischemia, or obstruction, a high-protein diet should be initiated within 24 hours post-surgery and overall protein intake goals are more important than total calorie intake in the postoperative period [6**]. Patients tolerating 50%–100% of nutrition goals should receive HP-ONS (2xday) to meet protein needs. A high-impact recent trial conducted in colorectal surgery patients within an ERAS/ERP pathway demonstrated in patients receiving HP-ONS post-operatively to achieve consumption of >60% of protein needs over first 3 post-operative days was associated with a 4.4-day reduction in length of stay ($p<0.001$) [44*]. For patients consuming <50% via the oral route, a dietician should be consulted and EN should be administered. Any patient who cannot achieve >50% of protein/calories requirement by ONS or EN should receive PN for >7 days, in combination with EN where feasible [6**].

Post-Operative Nutrition is Essential to Continued Recovery after Hospital Discharge

Perioperative nutrition optimization does not end at hospital discharge. Recovering postoperative patients, particular the elderly, often experience decreased appetites, persistent nausea, constipation from opiate use, and lack of education about diet optimization [6**,45*]. An observational study of patients after ICU discharge showed an average spontaneous calorie intake of 700 kcal/d, which is far insufficient to support anabolism since a caloric intake of 1.2–1.5×resting energy expenditure is recommended and thought to be required [6**,45*,46]. To facilitate continued recovery from the physiological stress of surgery, it is important to continue nutrition optimization therapy post-discharge [45*]. This is supported by meta-analysis data demonstrating ONS-use reduces mortality, hospital complications, hospital readmissions, LOS, and hospital costs [47,48]. In a retrospective analysis of a large hospital dataset, Philipson et al matched 724,000 patients who received ONS with controls not receiving ONS and demonstrated a 21% reduction in hospital LOS. In fact, for every \$1 USD spent on ONS, \$52.63 was saved in hospital costs [49]. Similar findings are echoed in major randomized trial of 652 patients in 78 centers studying effect of HP-ONS with β -hydroxy β -methylbutyrate versus placebo ONS [50]. Results demonstrate elderly hospitalized patients at risk for malnutrition benefitted from HP-ONS, with reduced 90-day mortality by ~50% relative to placebo (4.8% vs 9.7%; RR, 0.49; 95% CI, 0.27–0.90; $P=$.

018) [49]. Therefore, strong societal guideline recommendations indicate, HP-ONS should be consumed in all patients having major surgery for at least 4–8 weeks postoperatively⁶, and should be continued for up to 3–6 months postoperatively in more severely malnourished patients or patients with prolonged postoperative or ICU courses.

Conclusion

The malnourished surgical patient is underdiagnosed and undertreated - truly a silent epidemic in perioperative outcomes. HP-ONS and immunonutrition can mitigate the surgically induced metabolic response, immune-suppression, and support optimal postoperative recovery. We need further key prospective RCTs to determine the optimal method to identify the surgical patient at nutrition risk using tools like the PONS score (Figure 1) and LBM sarcopenia analysis. We also need study of ideal length of time required for preoperative nutrition optimization and large trials of the structured POET nutrition care pathways (Figure 3). In total, it is now abundantly clear we have a call-to-action to ensure that in the future no patient undergoes elective surgery without nutrition screening and structured nutrition intervention.

Acknowledgements

Financial support and sponsorship

Grant support from NIH T32 Anesthesiology Department Research Training grant to Dr. David Williams

Conflicts of interest

PEW- Is an associate editor of Clinical Nutrition (Elsevier). Has received grant funding related to this work from NIH, Canadian Institutes of Health Research, Abbott, Baxter, Fresenius, and Takeda. Dr. Wischmeyer has served as a consultant to Abbott, Fresenius, Baxter, Cardinal Health Nutricia, and Takeda for research related to this work. Dr. Wischmeyer has received unrestricted gift donation for surgical research from Musclesound. Dr. Wischmeyer has received honoraria or travel expenses for lectures on improving nutrition care in surgery from Abbott, Baxter and Nutricia.

J.M.- Consultant for Musclesound Inc and Nutricia.

DGAW- Grant support from NIH T32 Anesthesiology Department Research Training grant

References

1. Correia MI & Waitzberg DL The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through a multivariate model analysis. *Clinical nutrition* (Edinburgh, Scotland) 22, 235–239 (2003).
2. Vaid S, Bell T, Grim R & Ahuja V Predicting Risk of Death in General Surgery Patients on the Basis of Preoperative Variables Using American College of Surgeons National Surgical Quality Improvement Program Data. *The Permanente Journal* 16, 10–17 (2012).
3. Williams JD & Wischmeyer PE Assessment of perioperative nutrition practices and Attitudes-A national survey of colorectal and GI surgical oncology programs. *Am J Surg*, doi:10.1016/j.amjsurg.2016.10.008 (2016).
4. Weiss AJ (Truven Health Analytics), F. K. T. H. A., Barrett ML (M.L. Barrett, Inc.), Elixhauser A (AHRQ), Steiner CA (AHRQ), Guenter P (American Society for Parenteral and Enteral Nutrition), Brown MH (Baxter International, Inc.). Characteristics of Hospital Stays Involving Malnutrition, 2013. HCUP Statistical Brief #210 Agency for Healthcare Research and Quality, Rockville, MD (9 2016.).

- **5.** Aronson S et al. A Perioperative Medicine Model for Population Health: An Integrated Approach for an Evolving Clinical Science. *Anesth Analg* 126, 682–690, doi:10.1213/ANE.0000000000002606 (2018). [PubMed: 29200059] Key paper that describes integrated multidisciplinary perioperative optimization services and population health strategies in the context of a transitioning healthcare system.
- **6.** Wischmeyer PE et al. American Society for Enhanced Recovery and Perioperative Quality Initiative Joint Consensus Statement on Nutrition Screening and Therapy Within a Surgical Enhanced Recovery Pathway. *Anesth Analg* 126, 1883–1895, doi:10.1213/ANE.0000000000002743 (2018). [PubMed: 29369092] Key consensus statement paper that introduces the Perioperative Nutrition Screen (PONS) and describes perioperative optimization strategies for malnourished surgical patients.
- **7.** Weimann A et al. ESPEN guideline: Clinical nutrition in surgery. *Clinical nutrition* (Edinburgh, Scotland) 36, 623–650, doi:10.1016/j.clnu.2017.02.013 (2017). Key paper that describes clinical best practice guidelines for management of nutrition in surgical patients based on the current evidence of available literature.
8. Gillis C & Carli F Promoting Perioperative Metabolic and Nutritional Care. *Anesthesiology* 123, 1455–1472, doi:10.1097/ALN.0000000000000795 (2015). [PubMed: 26248016]
9. Awad S & Lobo DN What's new in perioperative nutritional support? *Curr Opin Anaesthesiol* 24, 339–348, doi:10.1097/ACO.0b013e328345865e (2011). [PubMed: 21451404]
10. Drover JW et al. Nutrition therapy for the critically ill surgical patient: we need to do better! *JPEN. Journal of parenteral and enteral nutrition* 34, 644–652, doi:10.1177/0148607110372391 (2010). [PubMed: 21097764]
- **11.** Ljungqvist O, Scott M & Fearon KC Enhanced Recovery After Surgery: A Review. *JAMA Surg* 152, 292–298, doi:10.1001/jamasurg.2016.4952 (2017). [PubMed: 28097305] Key review paper that describes major improvements in clinical outcomes and cost secondary to the paradigm shift following implementation of Enhanced Recovery After Surgery (ERAS).
12. Sergi G et al. An adequate threshold for body mass index to detect underweight condition in elderly persons: the Italian Longitudinal Study on Aging (ILSA). *The journals of gerontology. Series A, Biological sciences and medical sciences* 60, 866–871 (2005).
13. Kim JM et al. Relationship between low body mass index and morbidity after gastrectomy for gastric cancer. *Annals of surgical treatment and research* 90, 207–212, doi:10.4174/ast.2016.90.4.207 (2016). [PubMed: 27073791]
14. Truong A, Hanna MH, Moghadamyeghaneh Z & Stamos MJ Implications of preoperative hypoalbuminemia in colorectal surgery. *World journal of gastrointestinal surgery* 8, 353–362, doi:10.4240/wjgs.v8.i5.353 (2016). [PubMed: 27231513]
15. Nelson CL, Elkassabany NM, Kamath AF & Liu J Low Albumin Levels, More Than Morbid Obesity, Are Associated With Complications After TKA. *Clinical orthopaedics and related research* 473, 3163–3172, doi:10.1007/s11999-015-4333-7 (2015). [PubMed: 25995174]
16. Bohl DD, Shen MR, Kayupov E & Della Valle CJ Hypoalbuminemia Independently Predicts Surgical Site Infection, Pneumonia, Length of Stay, and Readmission After Total Joint Arthroplasty. *The Journal of arthroplasty* 31, 15–21, doi:10.1016/j.arth.2015.08.028 (2016). [PubMed: 26427941]
17. Lentine KL, Axelrod D & Abbott KC Interpreting body composition in kidney transplantation: weighing candidate selection, prognostication, and interventional strategies to optimize health. *Clin J Am Soc Nephrol* 6, 1238–1240, doi:10.2215/CJN.02510311 (2011). [PubMed: 21566111]
18. Mueller N et al. Can Sarcopenia Quantified by Ultrasound of the Rectus Femoris Muscle Predict Adverse Outcome of Surgical Intensive Care Unit Patients as well as Frailty? A Prospective, Observational Cohort Study. *Ann Surg* 264, 1116–1124, doi:10.1097/SLA.0000000000001546 (2016). [PubMed: 26655919]
- **19.** Wischmeyer PE, Puthucherry Z, San Milan I, Butz D & Grocott Muscle Mass and Physical Recovery in ICU: Innovations for Targeting of Nutrition and Exercise. *Curr Opin Crit Care*, Accpeted for Publication., doi:10.1097/MCC.0000000000000431 (2017). Key paper describing new innovative techniques targeted at recovery from post-ICU syndrome. These techniques include the use computerized tomography scan and ultrasound analysis of lean body mass to predict metabolic reserve and effectiveness of nutrition/exercise interventions, and 13C-Breath

testing to predict earlier detection of infection and predict over-feeding and under-feeding during nutritional optimization interventions.

20. Fielding RA et al. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc* 12, 249–256, doi:10.1016/j.jamda.2011.01.003 (2011). [PubMed: 21527165]
21. Prado CM et al. Prevalence and clinical implications of sarcopenic obesity in patients with solid tumours of the respiratory and gastrointestinal tracts: a population-based study. *Lancet Oncol* 9, 629–635, doi:10.1016/S1470-2045(08)70153-0 (2008). [PubMed: 18539529]
22. Batsis JA, Sahakyan KR, Rodriguez-Escudero JP, Bartels SJ & Lopez-Jimenez F Normal weight obesity and functional outcomes in older adults. *Eur J Intern Med* 25, 517–522, doi:10.1016/j.ejim.2014.05.008 (2014). [PubMed: 24909976]
23. Okumura S et al. Impact of Skeletal Muscle Mass, Muscle Quality, and Visceral Adiposity on Outcomes Following Resection of Intrahepatic Cholangiocarcinoma. *Ann Surg Oncol* 24, 1037–1045, doi:10.1245/s10434-016-5668-3 (2017). [PubMed: 27837297]
24. Boer BC et al. Skeletal muscle mass and quality as risk factors for postoperative outcome after open colon resection for cancer. *Int J Colorectal Dis* 31, 1117–1124, doi:10.1007/s00384-016-2538-1 (2016). [PubMed: 26876070]
25. Margadant CC et al. Lower muscle density is associated with major postoperative complications in older patients after surgery for colorectal cancer. *Eur J Surg Oncol* 42, 1654–1659, doi:10.1016/j.ejso.2016.05.040 (2016). [PubMed: 27554247]
- *26. Matsushima K et al. Loss of muscle mass: a significant predictor of postoperative complications in acute diverticulitis. *Journal of Surgical Research* 211, 39–44 (2017). [PubMed: 28501129] Key paper that studied the association between low muscle mass (LMM) and outcomes after emergency operations for acute diverticulitis. In univariable analysis, significantly higher rates of postoperative major complications (63% versus 37%, $P = 0.027$) and surgical site infection (47% versus 19%, $P = 0.008$) were identified in the LMM group.
27. Sandini M et al. A high visceral adipose tissue-to-skeletal muscle ratio as a determinant of major complications after pancreatoduodenectomy for cancer. *Nutrition* 32, 1231–1237, doi:10.1016/j.nut.2016.04.002 (2016). [PubMed: 27261062]
- **28. Simonsen C et al. Sarcopenia and Postoperative Complication Risk in Gastrointestinal Surgical Oncology: A Meta-analysis. *Ann Surg* 268, 58–69, doi:10.1097/SLA.0000000000002679 (2018). [PubMed: 29373365] Key meta-analysis that investigated sarcopenia as a predictor of postoperative risk of major and total complications after surgery for gastrointestinal cancer. Preoperative incidence of sarcopenia was associated with increased risk of major complications (risk ratio 1.40; 95% confidence interval, 1.20–1.64; $P < 0.001$; $I = 52\%$) and total complications (risk ratio 1.35; 95% confidence interval, 1.12–1.61; $P = 0.001$; $I = 60\%$).
29. Friedman J, Lussiez A, Sullivan J, Wang S & Englesbe M Implications of sarcopenia in major surgery. *Nutr Clin Pract* 30, 175–179, doi:10.1177/0884533615569888 (2015). [PubMed: 25681482]
- **30. van Vugt JLA et al. Low skeletal muscle mass is associated with increased hospital expenditure in patients undergoing cancer surgery of the alimentary tract. *PLoS One* 12, e0186547, doi: 10.1371/journal.pone.0186547 (2017). [PubMed: 29088245] Key paper that describes the association of health care expenditure with low skeletal muscle mass in patients undergoing gastrointestinal surgery. Hospital costs incrementally increased with lower sex-specific skeletal muscle mass quartiles ($P = 0.029$). After adjustment for confounders, low skeletal muscle mass was associated with a cost increase of €4,061 ($P = 0.015$).
- **31. Whittle J, Wischmeyer PE, Grocott MPW & Miller TE Surgical Prehabilitation: Nutrition and Exercise. *Anesthesiol Clin* 36, 567–580, doi:10.1016/j.anclin.2018.07.013 (2018). [PubMed: 30390779] Key recent review of exercise and nutrition interventions for prehabilitation in surgical patients.
- *32. Akazawa N, Harada K, Okawa N, Tamura K & Moriyama H Muscle mass and intramuscular fat of the quadriceps are related to muscle strength in non-ambulatory chronic stroke survivors: A cross-sectional study. *PLoS One* 13, e0201789, doi:10.1371/journal.pone.0201789 (2018). [PubMed: 30071100] Recent publications describing the relationship of muscle quality with muscle strength in stroke survivors.

33. Hill JC & Millan IS Validation of musculoskeletal ultrasound to assess and quantify muscle glycogen content. A novel approach. *Phys Sportsmed* 42, 45–52, doi:10.3810/psm.2014.09.2075 (2014).
- **34. Wischmeyer PE, Puthuchear Z, San Millan I, Butz D & Grocott MPW Muscle mass and physical recovery in ICU: innovations for targeting of nutrition and exercise. *Curr Opin Crit Care* 23, 269–278, doi:10.1097/MCC.0000000000000431 (2017). [PubMed: 28661414] Key recent paper examining metabolic, nutritional and anabolic analysis techniques and therapeutic strategies to improve recovery of muscle mass and function after acute illness. Describes use of CPET testing, ultrasound LBM measures and other anabolic and nutritional strategies.
- **35. Looijaard W, Molinger J & Weijs PJM Measuring and monitoring lean body mass in critical illness. *Curr Opin Crit Care* 24, 241–247, doi:10.1097/MCC.0000000000000511 (2018). [PubMed: 29847342] Key paper that describes the new innovative use of computed tomography scan analysis, musculoskeletal ultrasound, and bioelectrical impedance analysis to monitor lean body mass during ICU stay and can be used to guide nutrition interventions.
36. McClave SA et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN. Journal of parenteral and enteral nutrition* 40, 159–211, doi:10.1177/0148607115621863 (2016). [PubMed: 26773077]
37. Paddon-Jones D et al. Protein and healthy aging. *The American journal of clinical nutrition*, doi: 10.3945/ajcn.114.084061 (2015).
38. McClave SA et al. Summary Points and Consensus Recommendations From the North American Surgical Nutrition Summit. *Jpen-Parenter Enter* 37, 99s–105s, doi:10.1177/0148607113495892 (2013).
39. Drover JW et al. Perioperative use of arginine-supplemented diets: a systematic review of the evidence. *Journal of the American College of Surgeons* 212, 385–399 (2011). [PubMed: 21247782]
40. Moya P et al. Perioperative Standard Oral Nutrition Supplements Versus Immunonutrition in Patients Undergoing Colorectal Resection in an Enhanced Recovery (ERAS) Protocol: A Multicenter Randomized Clinical Trial (SONVI Study). *Medicine (Baltimore)* 95, e3704, doi: 10.1097/MD.00000000000003704 (2016). [PubMed: 27227930]
- *41. Thornblade LW et al. Preoperative Immunonutrition and Elective Colorectal Resection Outcomes. *Dis Colon Rectum* 60, 68–75, doi:10.1097/DCR.0000000000000740 (2017). [PubMed: 27926559] Key prospective cohort study that aimed to determine whether immunonutrition before elective colorectal surgery improves outcomes in the overall community. Results demonstrate reductions in prolonged length of stay and fewer complications and support the use of immunonutrition in quality improvement initiatives related to elective colorectal surgery.
- **42. Gillis C & Wischmeyer PE Pre-operative nutrition and the elective surgical patient: why, how and what? *Anaesthesia* 74 Suppl 1, 27–35, doi:10.1111/anae.14506 (2019). [PubMed: 30604414] Key review paper that describes the pathophysiology of malnutrition in the context of the surgically stressed patient and provides a knowledge base for the understanding why avoiding preoperative malnutrition and supporting protein anabolism are fundamental goals for perioperative nutritional optimization.
43. Osland E, Yunus RM, Khan S & Memon MA Early versus traditional postoperative feeding in patients undergoing resectional gastrointestinal surgery: a meta-analysis. *JPEN. Journal of parenteral and enteral nutrition* 35, 473–487, doi:10.1177/0148607110385698 (2011). [PubMed: 21628607]
- *44. Yeung SE, Hilkewich L, Gillis C, Heine JA & Fenton TR Protein intakes are associated with reduced length of stay: a comparison between Enhanced Recovery After Surgery (ERAS) and conventional care after elective colorectal surgery. *The American journal of clinical nutrition*, doi:10.3945/ajcn.116.148619 (2017). This key prospective cohort study compared protein adequacy as well as energy intakes, gut function, clinical outcomes, and how well nutritional variables predict length of hospital stay (LOS) in adult patients receiving enhanced recovery after surgery (ERAS) protocols and conventional care for elective colorectal resection. Nutrition variables were independent predictors of earlier discharge after potential confounders were controlled for.

- *45. Wischmeyer PE Tailoring nutrition therapy to illness and recovery. *Crit Care* 21, 316, doi: 10.1186/s13054-017-1906-8 (2017). [PubMed: 29297385] Key paper that describes nutrition optimization in ICU patients by emphasizing early enteral nutrition during the acute phase to correct micronutrient/vitamin deficiencies, deliver adequate protein, and moderate nonprotein calories in well-nourished patients. Significant protein/calorie delivery with high-protein oral supplements being essential to achieve adequate nutrition for months or years is required to facilitate functional and LBM recovery following discharge from ICU.
46. Peterson SJ et al. Adequacy of oral intake in critically ill patients 1 week after extubation. *Journal of the American Dietetic Association* 110, 427–433, doi:10.1016/j.jada.2009.11.020 (2010). [PubMed: 20184993]
47. Elia M, Normand C, Norman K & Laviano A A systematic review of the cost and cost effectiveness of using standard oral nutritional supplements in the hospital setting. *Clinical nutrition (Edinburgh, Scotland)* 35, 370–380, doi:10.1016/j.clnu.2015.05.010 (2016).
48. Stratton RJ, Hebuterne X & Elia M A systematic review and meta-analysis of the impact of oral nutritional supplements on hospital readmissions. *Ageing Res Rev* 12, 884–897, doi:10.1016/j.arr.2013.07.002 (2013). [PubMed: 23891685]
49. Philipson TJ, Snider JT, Lakdawalla DN, Stryckman B & Goldman DP Impact of oral nutritional supplementation on hospital outcomes. *Am J Manag Care* 19, 121–128 (2013). [PubMed: 23448109]
50. Deutz NE et al. Readmission and mortality in malnourished, older, hospitalized adults treated with a specialized oral nutritional supplement: A randomized clinical trial. *Clinical nutrition (Edinburgh, Scotland)* 35, 18–26, doi:10.1016/j.clnu.2015.12.010 (2016).

Key Points:

- Perioperative malnutrition is a critically underdiagnosed modifiable surgical risk factor that is associated with preventable adverse surgical outcomes.
- It is essential that all patients undergo preoperative nutrition screening with the Perioperative Nutrition Screen (PONS) in order to identify patients in need of perioperative nutrition therapy.
- Measurement of lean body mass (LBM) via CT scan and ultrasound demonstrates sarcopenia, shows associations with poor surgical outcome/increased hospital costs and may be used to guide nutrition intervention
- Preoperative nutrition optimization in malnourished patients for > 10 days, use of immunonutrition in all major surgery, and early postoperative oral intake have all recently been shown to improve surgical outcomes.
- All major surgical patients should consume high protein oral nutrition supplements post-operatively to reduce length of stay and continue for at least 1-month post-hospital discharge to optimize recovery

Figure 1.

Pre-Operative Nutrition Score (PONS) Assessment Tool (PONS; adapted from reference [6]). PONS utilizes questions from the validated Malnutrition Universal Screening Tool to assess for malnutrition risk in perioperative patients. A PONS score ≥ 1 and/or an albumin < 3.0 or vitamin D < 20 signifies malnutrition risk, and the patient should receive preoperative nutrition therapy before undergoing surgery.

Musclesound Skeletal Lean Body Muscle Assessment

ICU Admit

ICU Day 10

Figure 2.

Example of MuscleSound® Lean Body Mass Loss Over Hospital Stay in ICU Patient- A) ICU Admit, B)-ICU Day10. Note loss of muscle mass/size and lower muscle glycogen content at Day 10. Rectus femoris muscle shown: Pink areas (red arrows) = low muscle glycogen content; Purple areas (blue arrows) = high muscle glycogen content; White areas = intermuscular connective tissue; Grey areas = areas of muscle injury; Yellow areas = fascia

Figure 3. Perioperative Nutritional Care Pathways used in Duke Perioperative Enhancement Team (POET) Nutrition Program. Includes Nutrition Pathway for Low Nutrition Risk Patients with PONS score < 1 and Nutrition Pathway for High Nutrition Risk Patients— as defined by any positive response on the PONS score.