Strategies for Detecting Colorectal Cancer in Patients with Inflammatory Bowel Disease: A Cochrane Systematic Review and Meta-Analysis

William A. Bye, B. Com, MBBS, FRACP¹, Christopher Ma, MD, FRCPC^{2,3}, Tran M. Nguyen, MSc^{3,4}, Claire E. Parker, MA, MLIS^{3,4}, Vipul Jairath, MD, PhD^{3,5,6} and James E. East, MD (Res), FRCP⁷

OBJECTIVES: Patients with longstanding ulcerative colitis (UC) and colonic Crohn's disease (CD) have an increased risk of colorectal cancer (CRC). We assess the effectiveness of endoscopic surveillance in patients with inflammatory bowel disease (IBD) for diagnosing CRC and reducing CRC-related mortality.

METHODS: MEDLINE, EMBASE, and CENTRAL were searched from inception to 19 September 2016. Randomized controlled trials (RCTs), observational cohorts, or case-control studies assessing any form of endoscopic surveillance for early CRC detection were eligible for inclusion; studies without a comparison non-surveillance group were excluded. The primary outcome was rate of CRC detection. Secondary outcomes were rate of early (Duke stage A and B) versus late (Duke stage C & D) CRC detection and rate of CRC-related death. Data were pooled using fixed or random effects models based on the degree of heterogeneity; pooled odds ratios (ORs) with 95% confidence intervals (CIs) were calculated using the Mantel-Haenszel method.

RESULTS: Five observational studies evaluating 7199 IBD patients were included; no RCTs met criteria for inclusion. There are limited new studies evaluating this clinical question (last included study published 2014). There was a significantly higher rate of cancer detection in the non-surveillance group (3.2%, 135/4256) compared to the surveillance group (1.8%, 53/2895) (OR 0.58 (95% CI: 0.42–0.80), $p < 0.001$). In four pooled studies, there was a significantly lower rate of CRC-associated death in the surveillance group (8.5%, 15/176) compared to the non-surveillance group (22.3%, 79/354) (OR 0.36 (95% CI: 0.19–0.69), $p = 0.002$). In two pooled studies, there was a significantly higher rate of early-stage CRC detection in the surveillance group (15.5%, 17/110) compared to the non-surveillance group (7.7%, 9/117) (OR 5.40 (95% CI: 1.51–19.30), $p = 0.009$).

CONCLUSIONS: Colonoscopic surveillance in IBD is associated with a reduction in CRC development and CRC-associated death, as well as increased detection of early-stage CRC.

SUPPLEMENTARY MATERIAL accompanies this paper at <https://doi.org/10.1038/s41395-018-0354-7>

Am J Gastroenterol (2018) 113:1801–1809. <https://doi.org/10.1038/s41395-018-0354-7>

INTRODUCTION

Patients with longstanding ulcerative colitis (UC) and colonic Crohn's disease (CD) are at increased risk of developing colorectal cancer (CRC), likely as sequelae of chronic inflammation [1]. Inflammatory bowel disease (IBD)-associated CRC has a unique phenotype compared to sporadic CRC: it typically occurs at a younger age [2, 3], in more proximal locations [4], with aggressive mucinous or signet ring cell histology [5], and frequently

with multiple synchronous malignancies [5]. In a meta-analysis of population-based studies, the risk of CRC in IBD patients was increased over twofold (standardized incidence ratio (SIR) 2.4 (95% confidence interval (CI): 2.1–2.7)) compared to the general population [6]. Furthermore, CRC is estimated to account for approximately 15% of all deaths in IBD patients [2].

The goal of endoscopic surveillance is to detect and intervene on precursor dysplasia either through endoscopic or surgical therapy,

¹Department of Gastroenterology and Hepatology, Prince of Wales Hospital, Sydney, NSW, Australia. ²Division of Gastroenterology and Hepatology, University of Calgary, Calgary, AB, Canada. ³Robarts Clinical Trials, Western University, London, ON, Canada. ⁴Cochrane Inflammatory Bowel Disease Group, Robarts Clinical Trials, London, ON, Canada. ⁵Department of Medicine, Western University, London, ON, Canada. ⁶Department of Epidemiology and Biostatistics, Western University, London, ON, Canada. ⁷Translational Gastroenterology Unit, Nuffield Department of Medicine, John Radcliffe Hospital, University of Oxford, Oxford, UK. These authors contributed equally: William A. Bye and Christopher Ma. **Correspondence:** J.E.E. (email: james.east@ndm.ox.ac.uk)

Received 10 January 2018; accepted 21 August 2018; Published online 23 October 2018

or to detect CRC at an early, curative stage, which may lead to an improved prognosis and survival [7, 8]. Multiple international societies have endorsed endoscopic surveillance for patients with chronic UC or colonic CD [9–15] and it has become a cornerstone of IBD-related care, despite paucity of direct evidence for mortality reduction. Understanding the effectiveness of surveillance is crucial: with the increasing incidence of IBD, the costs associated with ongoing surveillance will also continue to rise [16]. Not only are surveillance programs expensive, they result in invasive and recurrent patient interventions that are not without risk.

In a previous meta-analysis, Collins et al. [7] did not demonstrate a statistically significant reduction in CRC-related death in IBD patients undergoing colonoscopy surveillance (pooled relative risk (RR) 0.81 (95% CI: 0.17–3.83)). However, this study was published over a decade ago. Strategies for both IBD management and surveillance have evolved since that time and the introduction of highly effective biologic therapies capable of inducing clinical, endoscopic, and histologic remission in IBD may have altered the natural history of the disease [17]. Currently, the optimal strategy for endoscopic surveillance with respect to timing of colonoscopy, interval between colonoscopies, management of complex lesions using advanced endoscopic resection techniques, and use of chromoendoscopy versus high-definition white light endoscopy (HD-WLE) remains controversial.

In this systematic review and meta-analysis, we update the previous study by Collins et al. [7]. We evaluate the literature with respect to the efficacy of colonoscopy surveillance for CRC detection, stage of CRC diagnosis, and CRC-related mortality in IBD patients with chronic colonic inflammation.

METHODS

Search strategy

We searched MEDLINE (1948–2016), EMBASE (1947–2016), the Cochrane Central Register of Controlled Trials (1994–2016), and the Cochrane IBD Group Specialized Register from inception to 19 September 2016. This was supplemented with hand-searches of reference lists of potentially relevant papers and conference proceedings from Digestive Disease Week, United European Gastroenterology Week, and the European Crohn's and Colitis Organization Congress. The search strategy is detailed in Supplemental Materials and Methods.

Study eligibility criteria

Randomized controlled trials (RCTs) and observational (cohort or case-control) studies evaluating any form of endoscopic surveillance aimed at early detection of CRC in patients of any age with a diagnosis of UC or colonic CD defined by conventional clinical, endoscopic, and histologic criteria selected for surveillance based solely on the duration and extent of disease were eligible for inclusion. Studies without a non-surveillance comparison group were excluded.

Outcomes

The primary outcome was the comparative rate of CRC diagnosis between the surveillance and non-surveillance groups.

Comparative rates of the following secondary outcomes between the surveillance and non-surveillance groups were also evaluated: (1) early-stage (Duke stage [18] A and B) CRC diagnosis; (2) late-stage (Duke stage C and D) CRC diagnosis; (3) colectomy for CRC; and (4) death from CRC. In the original study protocol, time to cancer detection, time to death, and proportion of patients with adverse events (AEs), serious AEs, and study withdrawal due to AEs were a priori defined secondary outcomes; however, the included studies did not sufficiently report these outcomes for inclusion in meta-analysis [19].

Data extraction

Potentially relevant articles were reviewed independently by three authors (WAB, TMN, CEP) to determine eligibility; disagreements were resolved by consensus. Results from the included studies were then extracted independently by two authors (WAB and TMN) into a standardized extraction form. The proportion of patients dying from CRC in the surveillance and control groups of each study was derived from life tables, survival curves, or where possible, by calculating life tables from the data provided.

Assessment risk of bias and quality of evidence

Methodological quality of each included study was independently evaluated by two authors (WAB and TMN) using the Newcastle–Ottawa Quality Assessment Scale (NOS) [20]. The NOS uses a star system to assess quality of cohort and case-control studies, with a maximum score of nine stars. For cohort studies, quality is evaluated on three domains: selection of study groups, comparability between the study and control group, and ascertainment of exposure or outcome of interest. For case-control studies, study quality is assessed on selection of study groups, comparability between cases and controls, and ascertainment of exposure.

The Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach was used to evaluate the overall quality of evidence supporting the primary and secondary outcomes [21]. Evidence from RCTs is considered high quality and evidence from observational studies is considered low quality; quality of evidence can be downgraded due to risk of bias, indirect evidence, inconsistency/unexplained heterogeneity, imprecision, and publication bias. Overall evidence quality is classified as high, moderate, low, and very low quality. Potential reporting bias was assessed by comparison of outcomes listed in protocols to published manuscripts; if protocols were not available, outcomes listed in the Methods section were compared to those reported in the Results section of published manuscripts. Funnel plots for evaluation of potential publication bias were not constructed due to the small number of included studies ($n < 10$).

Data synthesis and analysis

For dichotomous outcomes, we calculated the odds ratio (OR) and corresponding 95% CI. Data from individual studies were combined for meta-analysis and the pooled OR and 95% CI was calculated using the Mantel–Haenszel method. Heterogeneity was assessed among studies using the χ^2 test (p value > 0.10 was considered statistically significant) and the I^2 statistic. An I^2 value of 25% indicates low heterogeneity, 50% indicates moderate

heterogeneity, and 75% indicates high heterogeneity [22]. A fixed effects model was used to pool data unless significant heterogeneity existed between studies; a random-effects model was used if heterogeneity existed (I^2 50–75%). Data were not pooled in meta-analysis if a high degree of heterogeneity ($I^2 > 75\%$) existed. Data analysis was conducted using Review Manager (RevMan 5.3.5).

RESULTS

Search results and study characteristics

The literature search identified 12,896 records; 9499 records were screened after removal of duplicates (Fig. 1). 41 full-text articles were evaluated; most were excluded due to the absence of a non-surveillance control cohort. No RCTs met criteria for inclusion. Seven reports of five observational studies from 1993 to 2014 were included [23–27]. Of these five studies, four [23, 25–27] were cohort studies and one [24] was a case-control design. Four studies [24–27] were retrospective and one study [23] was prospectively performed. In total, 7199 patients with UC or colonic CD were evaluated. The characteristics of the included studies is sum-

marized in Table 1 and patient characteristics from the included studies are reported in Table 2.

Cancer detection

Three studies had available data for rates of CRC detection [24, 25, 27], enrolling 7151 patients (2895 in the surveillance group vs. 4256 in the non-surveillance group). CRC was detected in 53/2895 (1.8%) patients in the surveillance group compared to 135/4256 (3.2%) patients in the non-surveillance group. Using a fixed effects model, surveillance was associated with a 42% reduction in the odds of CRC detection, which was statistically significant (OR 0.58 (95% CI: 0.42–0.80), $p < 0.001$) (Fig. 2a). In a sensitivity analysis excluding the most heavily weighted study (Ananthakrishnan et al. [27]), the magnitude and direction of effect of surveillance colonoscopy on CRC detection was similar, but the results were no longer statistically significant (OR 0.67 (95% CI: 0.30–1.50)).

CRC-related death

Death due to CRC was reported in 4 studies [23, 25–27], enrolling 530 patients (176 in the surveillance group vs. 354 in the non-surveillance group). CRC-related death occurred in 8.5% (15/176) of patients in the surveillance group compared to 22.3% (79/354) of patients in the non-surveillance group. Using a fixed effects model, surveillance was associated with a 64% reduction in the odds of CRC-related death, which was statistically significant (OR 0.36 (95% CI: 0.19–0.69), $p = 0.002$) (Fig. 2b). This effect was maintained even in sensitivity analysis excluding the study by Ananthakrishnan et al. [27] (OR 0.40 (95% CI: 0.17–0.95)).

Tumor stage

Two studies had available data to examine detection of early- and late-stage CRC [23, 25] (Table 3). A higher rate of early-stage CRC was detected in the surveillance group (15.5%, 17/110) compared to the non-surveillance group (7.7%, 9/117) using a fixed effects model (OR 5.40 (95% CI: 1.51–19.30), $p = 0.009$) (Fig. 2c), which was statistically significant. In comparison, there was a nominally but not statistically significant lower rate of detection of late-stage CRC in patients undergoing surveillance (9.1%, 10/110) compared to patients not undergoing surveillance (16.2%, 19/117) (OR 0.46 (95% CI: 0.08–2.51), $p = 0.37$); a random-effects model was used for this analysis due to significant heterogeneity ($I^2 = 71\%$) (Fig. 2d). One study [26] described CRC stage at diagnosis using the American Joint Committee on Cancer 6th Edition Tumor Node Metastasis (TNM) staging system: surveillance was associated with a higher rate of diagnosing stage 1 CRC compared to non-surveillance (43.5%, 10/23 vs. 15.7%, 19/121, $p = 0.008$).

Colectomy

In one study, surveillance was associated with a lower risk of colectomy (36.3%, 33/91) compared to non-surveillance (53.7%, 51/95) (OR 0.49 (95% CI: 0.27–0.88)) [25].

Risk of bias and quality of evidence

The risk of bias in the five included non-randomized trials was assessed using the NOS; all studies scored well based on selection

Fig. 1 Study flow diagram

Table 1 Characteristics of included studies

Study	Sample size (n)		Study population	Study design	Outcomes	Statistical methods
	Surveillance (n)	Non-surveillance (n)				
Lashner et al. [25]	186		UC (disease duration ≥ 9 years)	Retrospective cohort study (1984–1986), annual colonoscopy with q10cm biopsies	Survival, deathCRC detectionColec-tomy rate	Kaplan–Meier survival analysis (log-rank test)Multivariate Cox regression analysis
	91	95				
Choi et al. [23]	41		UC (disease duration ≥ 8 years)	Prospective cohort study (1974–1991), UC patients who developed CRC, colonoscopy q2 years with biopsies	5-Year survivalTumor stageDeath	Kaplan–Meier survival analysis (Tarone–Ware test)
	19	22				
Karlen et al. [24]	142		UC (disease duration ≥ 5 years)	Nested population-based case–control study (1958–1988), 40 patients with UC and 102 controls, colonoscopy with multiple biopsies	Exposure to surveillance colonoscopy	Conditional logistic regression analysis (matched analysis)
	20	122				
Lutgens et al. [26]	149		IBD (UC and CD)	Retrospective, pathology database-derived cohort study of IBD patients with CRC (1990–2006), colonoscopy q<3 years with q10cm biopsies	CRC-related deathTumor stage	Kaplan–Meier survival analysis (Tarone–Ware test)Multivariate Cox regression analysis
	23	126				
Ananthakrishnan et al. [27]	6823		IBD (UC and CD)	Retrospective, population-based cohort study (1994–2014)	Diagnosis of CRC (by diagnostic coding for colon or rectal cancer)	Univariate and multivariate logistic regression modeling
	2754	4509				

CD Crohn's disease, CRC colorectal cancer, UC ulcerative colitis, IBD inflammatory bowel disease

of study groups, comparability between intervention and control group, and outcome and exposure ascertainment (Supplemental Table 1). A GRADE analysis indicated that the quality of evidence supporting the outcomes of cancer detection, CRC-related death, and tumor stage was very low due to the observational nature of the included studies and imprecision of the effect estimates. No potential reporting bias was detected.

DISCUSSION

As the incidence of IBD increases worldwide, a greater resource burden will be placed on endoscopic surveillance programs for CRC detection [16]. A prospective, long-term, RCT directly assessing survival benefit with endoscopic surveillance in IBD patients would be challenging to perform due to low event rates; thus, the best available evidence arises from observational data. Recognizing the inherent limitations of observational studies for drawing conclusions regarding causality, we updated a previous Cochrane systematic review and meta-analysis to evaluate the impact of surveillance colonoscopy on detection of CRC and death attributable to CRC in IBD patients. This updated review suggests that the ongoing use of colonoscopy surveillance may reduce both CRC development and CRC-associated death through detection of early-stage cancers. Although the quality of evidence is very low, an RCT that answers these questions is unlikely to be performed, particularly given the ethical considerations precluding randomization of IBD patients to a non-surveillance arm. Thus, we consider this pooled analysis to be the best available evidence at present to inform the benefit of colonoscopic surveillance.

Drawing conclusions regarding causality are limited by potential confounding bias in the included observational studies. Most authors attempted to control for confounders such as patient age and disease duration in multivariable logistic or Cox regression models (Table 1). However, important covariates such as concomitant therapy, disease phenotype, and frequency of assessments were not always adjusted for (Table 2). The largest study in this meta-analysis by Ananthakrishnan et al. [27] included 6823 IBD patients: the authors retrospectively examined the effect of colonoscopy within 6–36 months of CRC diagnosis (for cases) or end of follow-up (for controls) in a well-validated electronic IBD database capturing two tertiary care referral centers. In sensitivity analyses excluding this study, our conclusions of a significant benefit to surveillance colonoscopy on mortality were unchanged and surveillance was associated with a 33% non-statistically significant reduction in CRC detection. Ananthakrishnan et al. [27] further demonstrated the beneficial effect of surveillance colonoscopy which was robust even after adjusting for disease phenotype, extent, primary sclerosing cholangitis, erythrocyte sedimentation rate, follow-up duration, and intensity of health care utilization.

Since IBD-associated CRC is postulated to arise from precursor dysplasia [28], the theoretical benefit of surveillance colonoscopy is to detect and remove dysplasia, either endoscopically or surgically, prior to the development of CRC. Screening programs for sporadic CRC leverage the therapeutic window between the onset of dysplasia and development of CRC as an opportunity for endoscopic removal of pre-cancerous lesions [29]. Carcinogenesis in sporadic CRC typically results from the “adenoma-carcinoma sequence”, with the progressive accumulation of somatic mutations

in a dysplastic focus; this process can take years and can be arrested by endoscopic polypectomy [30].

This surveillance model may not translate to the IBD population. First, strategies and tools for dysplasia detection in IBD patients have evolved substantially. Previous recommendations for random four-quadrant biopsies every 10 cm have come under scrutiny with multiple studies demonstrating that targeted biopsies may be more effective for dysplasia detection than random biopsies [31–33]. Second, chromoendoscopy may be superior to white light endoscopy for dysplasia detection [15]. However, this was informed by studies conducted before widespread use of HD-WLE and a recent RCT of HD-WLE versus chromoendoscopy demonstrated no significant difference in dysplasia or neoplasia detection [34]. Three studies included in our meta-analysis specified the use of random four-quadrant biopsy surveillance and none identified the use of either HD-WLE or chromoendoscopy. Updated studies evaluating the efficacy of these different strategies on CRC detection and mortality are required.

Although dysplasia detection is an important surrogate for future CRC risk, it does not necessarily confer improved quality of life or increased survival in patients with IBD. While resecting polypoid dysplasia is associated with low future CRC risk [35], it does not obviate the high risk of CRC associated with invisible dysplasia or non-polypoid dysplasia [36]. This highlights the difference in tumorigenesis in patients with IBD compared to sporadic CRC. In IBD, chronic inflammation with repeated epithelial wounding and repair results in selection for mutant clones resistant to apoptosis and with accelerated growth better suited to a hostile microenvironment [37]. Not only does this predispose to synchronous lesions but it also renders large areas of the colonic mucosa genetically unstable (“field cancerization”) [38]. Resection of circumscribed focal dysplastic lesions does not address the underlying predisposition to cancer development in histologically inflamed tissue and may not mitigate future CRC risk [39, 40].

Despite these challenges, we demonstrate in this meta-analysis that surveillance colonoscopy is associated with a decrease in both CRC development and death secondary to CRC. This mirrors epidemiologic data that suggests the incidence of IBD-associated CRC is decreasing, although this cannot be causally attributed to surveillance programs since other factors may influence this, such as better treatments to control inflammation. Initial estimates of cumulative CRC risk in UC patients were as high as 2% at 10 years, 8% at 20 years, and 18% at 30 years of disease [1]. This risk has been recently downgraded: in a meta-analysis of nine population-based studies, Lutgens et al. [41] report a SIR for IBD-associated CRC of 1.7 (95% CI: 1.2–2.2) [41]. Advances in endoscopic surveillance and dysplasia detection are postulated mechanisms contributing to this decrease in CRC incidence. However, enhanced medical therapies for control of active inflammation and possible chemoprophylaxis protection from 5-aminosalicylates are potential confounders [42].

Although the incidence of IBD-associated CRC may be decreasing, the number of patients undergoing surveillance is increasing due to the rising incidence of IBD [16]. Given the substantial financial and resource costs associated with colonoscopy, identifying IBD patients at highest risk for CRC development for targeting

Table 2 Baseline reported patient characteristics

Study	Baseline covariates	Surveillance	Non-surveillance
Lashner et al. [25]	Sex (<i>n</i> , male:female)	53:38	43:52
	Mean age at symptom onset, years (SD)	25.5 (±11.3)	26.5 (±13.1)
	Family history of CRC	1	4
Choi et al. [23]	Mean age at UC onset, years (±SD)	24.9 (±11.9)	30.2 (±12.1)
	Mean disease duration, years (±SD)	18.5 (±5.8)	17.6 (±8.6)
	Pancolitis disease extent (%)	16 (84.2)	11 (50.0)
	Mean follow-up, years (±SD)	4.5 (±2.9)	2.6 (±3.2)
Karlen et al. ^a [24]	Sex (<i>n</i> , male:female)	NR	NR
	Age at UC diagnosis (years)	NR	NR
	UC disease extent at diagnosis	NR	NR
Lutgens et al. [26]	IBD (<i>n</i> , UC:CD)	18:5	71:54
	Sex (<i>n</i> , male:female)	17:6	72:54
	Median age at IBD diagnosis (years, range)	26 (9–50)	30 (6–83)
	Median age at CRC diagnosis (years, range)	48 (38–71)	49 (21–85)
	Primary sclerosing cholangitis (%)	2 (9%)	17 (14%)
	Mean follow-up (months, range)	57 (0–188)	51 (0–235)
Ananthakrishnan et al. [27]	Median age (years, IQR)	47 (32–61)	49 (35–63)
	Median age at IBD diagnosis (years, IQR)	37 (24–51)	39 (27–54)
	Mean follow-up (years, IQR)	8 (5–12)	8 (5–11)
	Sex (% , male:female)	48:52	43:57
	Race (% , white:non-white)	87:13	87:13
	IBD (% , UC:CD)	49:51	54:46
	PSC (%)	3	2
	Any immunomodulator use (%)	40	21
	Any biologic use (%)	21	9
	Median C-reactive protein (mg/dl, IQR)	3.7 (1.4–10.4)	4.8 (1.6–16.7)

^aFor Karlen et al. [24], baseline characteristics reported between cases and controls, not surveillance vs. non-surveillance
 CD Crohn's disease, CRC colorectal cancer, IBD inflammatory bowel disease, IQR interquartile range, NR not reported, UC ulcerative colitis

screening is critical. Current guidelines for the timing of surveillance colonoscopy are primarily based on disease extent and duration which are demonstrated risk factors for IBD-associated CRC

Fig. 2 Forest plots for effect of surveillance compared to non-surveillance for detection of colorectal cancer (a), death from colorectal cancer (b), early-stage colorectal cancer diagnosis (c), and late-stage colorectal cancer diagnosis (d) in patients with inflammatory bowel disease

[37]. Other risk factors for IBD-associated CRC include increasing age, male sex, and strong family history [43]. However, these are relatively crude predictors for identifying specific patients who would benefit from surveillance. Future research should focus on improving the precision of CRC risk estimation, potentially through incorporation of less invasive tests such as fecal DNA testing or rectal mucosal fluorescence in situ hybridization analysis for chromosomal instability [44]. Fecal immunochemical testing has been adopted for sporadic CRC screening in average risk populations; however, its application to IBD patients is limited by false positives due to detection of fecal hemoglobin from mucosal ulcerations in active inflammation [45].

The effect of surveillance colonoscopy on colectomy was investigated by Lashner et al. [25], who reported significantly lower rates

of colectomy for CRC, dysplasia, or active disease (OR 0.49 (95% CI: 0.27–0.88), $p = 0.02$) in patients receiving surveillance colonoscopy compared to those in the non-surveillance group [25]. However, these results should be re-examined given that endoscopic techniques for both lesion identification and excision have advanced significantly since this study. Furthermore, a finding of dysplasia no longer absolutely necessitates colectomy. Indeed, in patients with invisible low-grade dysplasia, the decision for ongoing surveillance colonoscopy versus colectomy as the most cost-effective management strategy is dependent on multiple factors including age, comorbidity, and expected postoperative morbidity [46].

Our study has several limitations. First and most importantly, no RCTs were eligible for inclusion which limited the quality of evidence synthesis in GRADE analysis. However, RCTs that directly

Table 3 Surveillance compared to non-surveillance for detection of colorectal cancer and death from colorectal cancer in patients with inflammatory bowel disease

Outcomes	Anticipated absolute effects ^a (95% CI)		Relative effect OR (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Risk with non-surveillance ^a	Risk with surveillance			
Cancer detection (per patient)	32 Per 1000 ^b	19 Per 1000 (14 to 26)	0.58 (0.42 to 0.80)	7151 (3 Studies)	⊕⊕⊕⊕ VERY LOW ^b
Death from colorectal cancer (per cancer)	223 Per 1000 ^b	94 Per 1000 (52 to 165)	0.36 (0.19 to 0.69)	530 (4 Studies)	⊕⊕⊕⊕ VERY LOW ^d
Cancer stage: Duke A or B (per cancer)	77 Per 1000 ^b	310 Per 1000 (112 to 617)	5.40 (1.51 to 19.30)	227 (2 Studies)	⊕⊕⊕⊕ VERY LOW ^a
Cancer stage: Duke C or D (per cancer)	162 Per 1000 ^b	82 Per 1000 (15 to 327)	0.46 (0.08 to 2.51)	227 (2 Studies)	⊕⊕⊕⊕ VERY LOW ^f

CI confidence interval, GRADE Grading of Recommendations Assessment, Development, and Evaluation, OR odds ratio

^aThe risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI)

^bControl group risk comes from control arm of meta-analysis, based on included trials

^cDowngraded one level due to sparse data (188 events)

^dDowngraded one level due to sparse data (94 events)

^eDowngraded due to very sparse data (26 events)

^fDowngraded due to very sparse data (29 events) and heterogeneity ($I^2 = 71\%$)

GRADE Working Group Grades of Evidence

High quality: we are very confident that the true effect lies close to that of the estimate of the effect

Moderate quality: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low quality: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect

Very low quality: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect

evaluate the effect of colonoscopy surveillance on CRC in IBD patients are unlikely to ever be performed due to ethical and logistical considerations. Even retrospective studies that evaluate this question are difficult to perform because the pool of IBD patients not undergoing surveillance is limited. Second, limitations of the included studies need to be considered when interpreting pooled results from meta-analysis. Such limitations include: (1) the potential for volunteer bias among patients seeking out surveillance colonoscopy; (2) possible referral bias when data are derived from tertiary care centers that treat a higher prevalence of patients with refractory or severe disease; and (3) heterogeneous surveillance practices with respect to interval of repeat surveillance colonoscopy and biopsy strategy. Although we included all studies evaluating endoscopic surveillance, there were insufficient data to compare different surveillance strategies, and comparative studies of HD-WLE, chromoendoscopy, and digital image enhancement colonoscopy are needed. Third, we evaluated the effect of surveillance on stage of CRC diagnosis by the Duke staging system; this has largely been replaced by the American Joint Committee on Cancer TNM staging system [47]. However, many of the included studies were published before the TNM system was widely adopted and both classifications have distinct similarities with respect to depth of tumor penetration. Fourth, we did not evaluate the effect of polypectomy on mortality as it was outside the scope of this review. However, additional data linking dysplasia detection and endoscopic resection to reduction in IBD-associated CRC outcomes are needed.

In conclusion, in this Cochrane systematic review and meta-analysis, we demonstrate in pooled analysis from observational studies that colonoscopy surveillance results in a reduction in IBD-associated CRC development by 42% and CRC-associated death by 64% compared to patients who do not undergo surveillance. We speculate that this reduction in CRC-associated death is due to increased detection of early-stage cancers associated with better prognosis. Updated studies are required to evaluate the benefits of modern surveillance techniques, including advancements in endoscopic technology and dysplasia resection. We further hypothesize that additional benefits may be derived from surveillance: as therapeutic targets for both UC and CD shift from symptom resolution towards normalization of histologic and endoscopic endpoints, surveillance colonoscopy offers a further opportunity to tailor treatment towards targeting more robust objective outcomes and improve long-term disease prognosis.

ACKNOWLEDGEMENTS

This review was supported by the Cochrane Inflammatory Bowel Disease Group.

CONFLICT OF INTEREST

Guarantor of the article: James E. East is the guarantor of the article and accepts full responsibility for the conduct of the study.

Specific author contributions: WAB, VJ, and JEE conceived and designed the study. WAB, TMN, and CEP were involved in data collection and analysis. WAB and CM drafted the initial manuscript.

VJ and JEE contributed to manuscript editing. All authors approve the final draft.

Financial support: CM is supported by the Canadian Institutes of Health Research and the Canadian Association of Gastroenterology. Partial funding for the Cochrane IBD Group (1 April 2016–31 March 2018) has been provided by Crohn's and Colitis Canada (CCC). JEE was funded by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health.

Potential competing interests: WAB, CM, TMN, and CEP: none; VJ: advisory board fees (AbbVie, Sandoz, Takeda, Janssen) and speaker's fees (Takeda, Janssen, Shire, Ferring); JEE: advisory board fees (Lumendi, Boston Scientific)

Study Highlights

WHAT IS CURRENT KNOWLEDGE

- ✓ Patients with ulcerative colitis and colonic Crohn's disease have an increased risk of colorectal cancer.
- ✓ Surveillance colonoscopy for dysplasia and early CRC detection is recommended by multiple international societies for patients with chronic colonic inflammation.
- ✓ In a previous Cochrane review (Collins 2006), there was no definitive evidence that surveillance colonoscopy prolonged survival in IBD patients.

WHAT IS NEW HERE

- ✓ In this contemporary systematic review and meta-analysis, we identified five relevant observational studies enrolling 7199 IBD patients, but no randomized controlled trials.
- ✓ On pooled analysis, a 42% reduction in the odds of CRC with surveillance colonoscopy was observed.
- ✓ On pooled analysis, a 64% reduction in the odds of CRC-related death with surveillance colonoscopy was observed.
- ✓ A greater than 5-fold increase in the odds of early stage CRC detection with surveillance colonoscopy.

REFERENCES

- Eaden JA, Abrams KR, Mayberry JF. The risk of colorectal cancer in ulcerative colitis: a meta-analysis. *Gut*. 2001;48:526–35.
- Munkholm P. Review article: the incidence and prevalence of colorectal cancer in inflammatory bowel disease. *Aliment Pharmacol Ther*. 2003;18(Suppl 2):1–5.
- Rutter MD, Saunders BP, Wilkinson KH, et al. Thirty-year analysis of a colonoscopic surveillance program for neoplasia in ulcerative colitis. *Gastroenterology*. 2006;130:1030–8.
- Bansal P, Sonnenberg A, Risk factors of colorectal cancer in inflammatory bowel disease. *Am J Gastroenterol*. 1996;91:44–8.
- Itzkowitz SH, Yio X. Inflammation and cancer IV. Colorectal cancer in inflammatory bowel disease: the role of inflammation. *Am J Physiol Gastrointest Liver Physiol*. 2004;287:G7–17.
- Jess T, Rungoe C, Peyrin-Biroulet L. Risk of colorectal cancer in patients with ulcerative colitis: a meta-analysis of population-based cohort studies. *Clin Gastroenterol Hepatol*. 2012;10:639–45.
- Collins PD, Mpofu C, Watson AJ, et al. Strategies for detecting colon cancer and/or dysplasia in patients with inflammatory bowel disease. *Cochrane Database Syst Rev*. 2006;2:CD000279
- Farraye FA, Odze RD, Eaden J, et al. AGA technical review on the diagnosis and management of colorectal neoplasia in inflammatory bowel disease. *Gastroenterology*. 2010;138:746–74.
- Farraye FA, Odze RD, Eaden J, et al. AGA medical position statement on the diagnosis and management of colorectal neoplasia in inflammatory bowel disease. *Gastroenterology*. 2010;138:738–45.
- American Society for Gastrointestinal Endoscopy Standards of Practice Committee, Shergill AK, Lightdale JR, et al. The role of endoscopy in inflammatory bowel disease. *Gastrointest Endosc*. 2015;81:1101–21 e1–13.
- Annese V, Daperno M, Rutter MD, et al. European evidence based consensus for endoscopy in inflammatory bowel disease. *J Crohns Colitis*. 2013;7:982–1018.
- Cairns SR, Scholefield JH, Steele RJ, et al. Guidelines for colorectal cancer screening and surveillance in moderate and high risk groups (update from 2002). *Gut*. 2010;59:666–89.
- Eaden JA, Mayberry JF; British Society for Gastroenterology; Association of Coloproctology for Great Britain and Ireland. Guidelines for screening and surveillance of asymptomatic colorectal cancer in patients with inflammatory bowel disease. *Gut*. 2002;51(Suppl 5):V10–2.
- Kornbluth A, Sachar DB; Practice Parameters Committee of the American College of Gastroenterology. Ulcerative colitis practice guidelines in adults: American College of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol*. 2010;105:501–23.
- Laine L, Kaltenbach T, Barkun A, et al. SCENIC international consensus statement on surveillance and management of dysplasia in inflammatory bowel disease. *Gastroenterology*. 2015;148:639–51 e28.
- Ng SC, Shi HY, Hamidi N, et al. Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet*. 2018;390:2769–78.
- Mandel MD, Miheller P, Mullner K, et al. Have biologics changed the natural history of Crohn's disease? *Dig Dis*. 2014;32:351–9.
- Astler VB, Collier FA. The prognostic significance of direct extension of carcinoma of the colon and rectum. *Ann Surg*. 1954;139:846–52.
- Bye WA, Nguyen TM, Parker CE, et al. Strategies for detecting colon cancer in patients with inflammatory bowel disease. *Cochrane Database Syst Rev*. 2017;9:CD000279.
- Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non randomised studies in meta-analyses. https://www.ohri.ca/programs/clinical_epidemiology/oxford.asp (2014). Accessed 31 Aug 2017
- Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924–6.
- Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from <https://www.handbook.cochrane.org>.
- Choi PM, Nugent FW, Schoetz DJ Jr, et al. Colonoscopic surveillance reduces mortality from colorectal cancer in ulcerative colitis. *Gastroenterology*. 1993;105:418–24.
- Karlen P, Kornfeld D, Brostrom O, et al. Is colonoscopic surveillance reducing colorectal cancer mortality in ulcerative colitis? A population based case control study. *Gut*. 1998;42:711–4.
- Lashner BA, Kane SV, Hanauer SB. Colon cancer surveillance in chronic ulcerative colitis: historical cohort study. *Am J Gastroenterol*. 1990;85:1083–7.
- Lutgens MW, Oldenburg B, Siersema PD, et al. Colonoscopic surveillance improves survival after colorectal cancer diagnosis in inflammatory bowel disease. *Br J Cancer*. 2009;101:1671–5.
- Ananthakrishnan AN, Cagan A, Cai T, et al. Colonoscopy is associated with a reduced risk for colon cancer and mortality in patients with inflammatory bowel diseases. *Clin Gastroenterol Hepatol*. 2015;13:322–9 e1.
- Goldman H. Significance and detection of dysplasia in chronic colitis. *Cancer*. 1996;78:2261–3.
- Bacchus CM, Dunfield L, Gorber SC, et al. Recommendations on screening for colorectal cancer in primary care. *CMAJ*. 2016;188:340–8.
- Leslie A, Carey FA, Pratt NR, et al. The colorectal adenoma-carcinoma sequence. *Br J Surg*. 2002;89:845–60.
- Itzkowitz SH, Harpaz N. Diagnosis and management of dysplasia in patients with inflammatory bowel diseases. *Gastroenterology*. 2004;126:1634–48.
- Watanabe T, Ajioka Y, Mitsuyama K, et al. Comparison of targeted vs random biopsies for surveillance of ulcerative colitis-associated colorectal cancer. *Gastroenterology*. 2016;151:1122–30.
- Gasia MF, Ghosh S, Panaccione R, et al. Targeted biopsies identify larger proportions of patients with colonic neoplasia undergoing high-definition

- colonoscopy, dye chromoendoscopy, or electronic virtual chromoendoscopy. *Clin Gastroenterol Hepatol*. 2016;14:704–12, e4.
34. Iacucci M, Kaplan GG, Panaccione R, et al. A randomized trial comparing high definition colonoscopy alone with high definition dye spraying and electronic virtual chromoendoscopy for detection of colonic neoplastic lesions during IBD surveillance colonoscopy. *Am J Gastroenterol*. 2018;113:225–34.
 35. Wanders LK, Dekker E, Pullens B, et al. Cancer risk after resection of polypoid dysplasia in patients with longstanding ulcerative colitis: a meta-analysis. *Clin Gastroenterol Hepatol*. 2014;12:756–64.
 36. Choi CH, Ignjatovic-Wilson A, Askari A, et al. Low-grade dysplasia in ulcerative colitis: risk factors for developing high-grade dysplasia or colorectal cancer. *Am J Gastroenterol*. 2015;110:1461–71.
 37. Beaugerie L, Itzkowitz SH. Cancers complicating inflammatory bowel disease. *N Engl J Med*. 2015;372:1441–52.
 38. Choi CR, Bakir IA, Hart AL, et al. Clonal evolution of colorectal cancer in IBD. *Nat Rev Gastroenterol Hepatol*. 2017;14:218–29.
 39. Gupta RB, Harpaz N, Itzkowitz S, et al. Histologic inflammation is a risk factor for progression to colorectal neoplasia in ulcerative colitis: a cohort study. *Gastroenterology*. 2007;133:1099–105.
 40. Rutter MD, Saunders BP, Wilkinson KH, et al. Most dysplasia in ulcerative colitis is visible at colonoscopy. *Gastrointest Endosc*. 2004;60:334–9.
 41. Lutgens MW, van Oijen MG, van der Heijden GJ, et al. Declining risk of colorectal cancer in inflammatory bowel disease: an updated meta-analysis of population-based cohort studies. *Inflamm Bowel Dis*. 2013;19:789–99.
 42. Zhao LN, Li JY, Yu T, et al. 5-Aminosalicylates reduce the risk of colorectal neoplasia in patients with ulcerative colitis: an updated meta-analysis. *PLoS ONE*. 2014;9:e94208.
 43. Askling J, Dickman PW, Karlen P, et al. Family history as a risk factor for colorectal cancer in inflammatory bowel disease. *Gastroenterology*. 2001;120:1356–62.
 44. Kisiel JB, Ahlquist DA. Stool DNA testing for cancer surveillance in inflammatory bowel disease: an early view. *Ther Adv Gastroenterol*. 2013;6:371–80.
 45. Ma C, Lumb R, Walker EV, et al. Noninvasive fecal immunochemical testing and fecal calprotectin predict mucosal healing in inflammatory bowel disease: a prospective cohort study. *Inflamm Bowel Dis*. 2017;23:1643–9.
 46. Parker B, Buchanan J, Wordsworth S, et al. Surgery versus surveillance in ulcerative colitis patients with endoscopically invisible low-grade dysplasia: a cost-effectiveness analysis. *Gastrointest Endosc*. 2017;86:1088–99 e5.
 47. Amin MB, Edge S, Greene F, et al. *AJCC Cancer Staging Manual*. 8th ed. New York: Springer International Publishing; 2017.