Potential for Aerobic Methanotrophic Metabolism on Mars

Mayumi Seto,¹ Katsuyuki Noguchi,¹ and Philippe Van Cappellen²

Abstract

Observational evidence supports the presence of methane (CH₄) in the martian atmosphere on the order of parts per billion by volume (ppbv). Here, we assess whether aerobic methanotrophy is a potentially viable metabolism in the martian upper regolith, by calculating metabolic energy gain rates under assumed conditions of martian surface temperature, pressure, and atmospheric composition. Using kinetic parameters for 19 terrestrial aerobic methanotrophic strains, we show that even under the imposed low temperature and pressure extremes (180–280 K and 6–11 hPa), methane oxidation by oxygen (O₂) should in principle be able to generate the minimum energy production rate required to support endogenous metabolism (*i.e.*, cellular maintenance). Our results further indicate that the corresponding metabolic activity would be extremely low, with cell doubling times in excess of 4000 Earth years at the present-day ppbv-level CH₄ mixing ratios in the atmosphere of Mars. Thus, while aerobic methanotrophic microorganisms similar to those found on Earth could theoretically maintain their vital functions, they are unlikely to constitute prolific members of hypothetical martian soil communities. **Key Words:** Mars—Methane—Methanotrophs—Catabolic energy production—Maintenance energy demand—Kinetic limitation. *Astrobiology* 19, 1187–1195.

1. Introduction

MARS IS CONSIDERED a promising planet in the quest for extraterrestrial life because of its resemblance to Earth and the accumulating evidence of past and potentially current surface and subsurface water activity (Pollack and Kasting, 1987; McKay *et al.*, 1996; Squyres *et al.*, 2004; Orosei *et al.*, 2018). At present, however, life would be challenged by the environmental conditions on Mars, including the scarcity of liquid water, very low atmospheric pressure and extreme temperature variations, a carbon dioxide (CO₂)-dominated atmosphere, strong UV and cosmic radiation, high concentrations of heavy metals, the lack of organic substances and nutrients, and high pH (Berry *et al.*, 2010). Nonetheless, the fact that microorganisms survive in extreme environments on Earth (Junge *et al.*, 2004; Berry *et al.*, 2010; Nicholson *et al.*, 2013; Moissl-Eichinger *et al.*, 2016) supports the possibility of microbial life under the harsh conditions found on Mars.

Based on a re-evaluation of the environmental limits for microbial life, Rummel *et al.* (2014) identified *Special*

Regions on Mars within which microorganisms similar to those found on Earth could survive and propagate. Life forms best adapted to the martian conditions are likely chemolithoautotrophic microorganisms that use inorganic carbon sources to build new biomass and generate energy from the oxidation of inorganic compounds. Any microbial cell requires a minimum energy production to support basic metabolic functions and activity in the absence of growth, the so-called maintenance energy requirement (Hoehler, 2004). Only when the catabolic energy production exceeds the maintenance energy demand, a cell becomes able to allocate the remaining energy to reproduction. To determine whether this threshold is reached for a given group of chemolithotrophs, one should thus compare the potential catabolic energy production in a given environment with the cellular maintenance requirement.

Here, we assess the minimum criteria for survival of aerobic methanotrophic microorganisms by estimating their catabolic energy production under assumed martian near-surface conditions. On Earth, aerobic methanotrophs are widely distributed in soils, sediments, and aqueous ecosystems, and some

¹Department of Chemistry, Biology, and Environmental Sciences, Faculty of Science, Nara Women's University, Nara, Japan.

²Ecohydrology Research Group, Department of Earth and Environmental Sciences, Water Institute, University of Waterloo, Waterloo, Canada.

of them have been shown to be tolerant toward various environmental stressors (Hanson and Hanson, 1996; Trotsenko and Khmelenina, 2002, 2005). These microorganisms utilize CH_4 as the electron donor and O_2 as terminal electron acceptor to generate energy, according to the overall catabolic reaction:

Many aerobic microorganisms are able to grow at relatively low O_2 concentrations, similar to those encountered in the martian atmosphere (on average 0.13% by volume, Rummel *et al.*, 2014). The extreme low abundance of CH_4 may therefore represent a more severe limitation for aerobic methanotrophs on Mars. On Earth, some methanotrophic microbes have been found to possess a “high-affinity” enzyme for CH_4 (Conrad, 1996; Hanson and Hanson, 1996), which allows them to function at ppmv levels of CH_4 in well-aerated upper soil horizons (Knief and Dunfield, 2005). Nonetheless, the current mixing ratio of CH_4 in Earth’s atmosphere (approximately 1.75 ppmv, Conrad, 1996; Bull *et al.*, 2000) is two to four orders of magnitude higher than current estimates for the martian atmosphere (Formisano *et al.*, 2004; Krasnopolsky *et al.*, 2004; Mumma *et al.*, 2009; Fonti and Marzo, 2010; Webster *et al.*, 2013, 2015, 2018). The possible sources of martian CH_4 include exogenous inputs, for example, from meteorite impacts, hydrothermal processes, for example, serpentinization reactions, and biological emissions (Atreya *et al.*, 2007).

At first sight, the very low levels of CH_4 in the martian atmosphere would seem to preclude aerobic methanotrophy. Nevertheless, cells on Mars may have much lower maintenance energy requirements than most of their terrestrial counterparts because of the very low surface temperatures. On Earth, microbial communities found at subfreezing temperatures are able to survive by greatly reducing their energy expenditures (Price and Sowers, 2004). For example, immobile or dormant nitrifiers remain viable at -40°C due to their extremely slow metabolism and, hence, very low maintenance energy requirements (Sowers, 2001; Price and Sowers, 2004).

In this article, we begin by outlining the theoretical basis for estimating cell-specific rates of catabolic energy production and maintenance energy consumption by methanotrophic microorganisms. We then derive the equations for (1) the minimum CH_4 concentration required to support cell maintenance, and (2) cell doubling times. The latter provides an indicator for the potential for growth of the microbial population. Next, we calculate the metabolic energy gain rates under near-surface martian conditions using kinetic parameters extracted from 19 known terrestrial methanotrophic strains, and discuss the uncertainties associated with the estimations and the implications for methanotrophy on Mars.

2. Methods

2.1. Catabolic energy production at given temperature, pressure, and mixing ratios

The cell-specific rate of catabolic energy production by methanotrophs depends on the Gibbs energy change of the CH_4 oxidation reaction by O_2 [Eq. (1)] (McCollom

and Shock, 1997; Amend and Shock, 2001; Shock *et al.*, 2010):

$$\Delta_r G = \Delta_r G^\circ + RT \ln \frac{f_{\text{CO}_2}}{f_{\text{O}_2}^2 f_{\text{CH}_4}}, \quad (2)$$

where $\Delta_r G$ is expressed in $\text{kJ} (\text{mol CH}_4)^{-1}$, R is the gas constant in units of $\text{kJ K}^{-1} \text{mol}^{-1}$, T is the absolute temperature in K, f_X stands for the fugacity of a chemical species X , and $\Delta_r G^\circ$ is the standard Gibbs free energy change of the reaction at a given temperature and pressure in units of $\text{kJ} (\text{mol CH}_4)^{-1}$:

$$\Delta_r G^\circ = \Delta_f G_{\text{CO}_2}^\circ + 2\Delta_f G_{\text{H}_2\text{O}}^\circ - \Delta_f G_{\text{CH}_4}^\circ - 2\Delta_f G_{\text{O}_2}^\circ, \quad (3)$$

where $\Delta_f G_X^\circ$ is the standard molal Gibbs free energy of formation of a species X in units of kJ mol^{-1} . At a given temperature and pressure, T_g and P_g , the $\Delta_f G^\circ$ values are calculated following the work of Helgeson *et al.* (1978):

$$\begin{aligned} \Delta_f G^\circ = & \Delta_f G^{\circ*} - S^{\circ*}(T_g - T^*) + \int_{T^*}^{T_g} C_P^{\circ*} dT \\ & - \int_{T^*}^{T_g} C_P^{\circ*} d \ln T + \int_{P^*}^{P_g} V^{\circ*} dP, \end{aligned} \quad (4)$$

where superscript $*$ indicates reference values at 298.15 K and/or 1 bar, while S° is the standard partial molal entropy at the reference conditions in units of kJ mol^{-1} , and C_P° and V° designate the standard partial molal isobaric heat capacity in units of $\text{kJ K}^{-1} \text{mol}^{-1}$ and the standard partial molal isobaric volume, respectively. The following empirical equation for the standard molal heat capacity in units of $\text{kJ mol}^{-1} \text{K}^{-1}$ at 1 bar is substituted into Eq. (4):

$$C_P^{\circ*}/R = A + BT - CT^2, \quad (5)$$

where A , B , and C are experimentally derived, temperature-independent coefficients for the species of interest (Supplementary Table S1). For gaseous species, $V^{\circ*} = RT/P$ in units of $\text{kJ Pa}^{-1} \text{mol}^{-1}$, and we further assume that $V^{\circ*}$ of water is constant ($18 \times 10^{-6} \text{ m}^3 \text{mol}^{-1} = 18 \times 10^{-9} \text{ kJ Pa}^{-1} \text{mol}^{-1}$), and that fugacities, f_X , obey van der Waals equation:

$$f_X = \gamma_X P_{\text{tot}} \exp \left\{ \frac{\gamma_X P_{\text{tot}}}{RT} \left(b - \frac{a}{RT} \right) \right\}, \quad (6)$$

where γ_X is the concentration of a gaseous species X in (v/v), P_{tot} denotes the total pressure (atm), and a and b are the van der Waals constants (Supplementary Table S1).

The standard formalism to describe rates of microbially catalyzed reaction processes is the Michaelis–Menten kinetic equation. For CH_4 oxidation, the cell-specific rate is as follows:

$$r = v_{\text{max}} \frac{[\text{CH}_4]}{K_{\text{CH}_4} + [\text{CH}_4]} \frac{[\text{O}_2]}{K_{\text{O}_2} + [\text{O}_2]}, \quad (7)$$

where v_{max} is the maximum cell-specific CH_4 oxidation rate ($\text{mol CH}_4 \text{ cell}^{-1} \text{h}^{-1}$), K_X are half-saturation constants (mol L^{-1}), and $[\text{CH}_4]$ and $[\text{O}_2]$ are the molar concentrations

of CH_4 and O_2 (mol L^{-1}), which are obtained from the following:

$$[X] = \frac{\gamma_X P_{\text{tot}}}{RT}, \quad (8)$$

where the gas constant, R , is expressed in units of $\text{L atm K}^{-1} \text{mol}^{-1}$. To calculate cell-specific CH_4 oxidation rates with Eq. (7), values of v_{max} and K_{CH_4} for 19 strains of aerobic methanotrophs were extracted from the literature (Knief and Dunfield, 2005). The corresponding v_{max} values range from 5.6×10^{-18} to $5.7 \times 10^{-16} \text{ mol CH}_4 \text{ cell}^{-1} \text{ h}^{-1}$, and K_{CH_4} from 1.0×10^{-6} to $2.3 \times 10^{-5} \text{ mol L}^{-1}$. For K_{O_2} , the only reported value is that for *Methylocystis* sp. strain *MOX-1* ($1.9 \times 10^{-9} \text{ mol L}^{-1}$; Van Bodegom *et al.*, 2001). This value is therefore applied in all the following calculations.

The cell-specific catabolic energy production rate, which is equivalent to the amount of power available to the microorganisms (*i.e.*, the “power supply,” LaRowe and Amend, 2015), is calculated by multiplying the Gibbs energy change of reaction of aerobic CH_4 oxidation with the cell-specific CH_4 oxidation rate:

$$[\text{Cell-specific catabolic energy production rate}] = -r \bullet \Delta_r G, \quad (9)$$

2.2. Metabolic energy requirement of endogenous metabolism

Temperature is a key environmental control on cellular energy requirements for growth, maintenance, and survival (Tijhuis *et al.*, 1993; Price and Sowers, 2004). Tijhuis *et al.* (1993) derived empirical correlations between the maintenance energy demand rate, expressed in $\text{kJ (mol C biomass)}^{-1} \text{ h}^{-1}$, and temperature for both aerobic and anaerobic bacteria in the temperature range of 2–66°C. This temperature range, however, does not encompass the subzero martian surface temperatures.

The existing, admittedly limited, data indicate that the temperature dependence of cellular metabolic energy requirements differs between mesophilic and psychrophilic or psychrotolerant microorganisms, that is, microorganisms growing and metabolically active at temperatures of 0°C and below. This is shown in Fig. 1, where the estimated metabolic energy requirement rates for maintenance and survival of nitrifiers and methanogens at temperatures between –40°C and 0°C (Price and Sowers, 2004; Tung *et al.*, 2005; Rohde *et al.*, 2008; Supplementary Table S2) are compared with the data for aerobic and anaerobic bacteria of Tijhuis *et al.* (1993). Note that the subzero maintenance energy requirements are extracted from the original article figures using the Plot Digitizer software (available at <http://plotdigitizer.sourceforge.net>). The original units of $\text{gC (gC)}^{-1} \text{ h}^{-1}$ are converted to $\text{kJ (mol C)}^{-1} \text{ h}^{-1}$ by considering the Gibbs energy needed to synthesize 1 gC of biomass, which for autotrophic nitrifiers is 291.4 kJ gC^{-1} , and for heterotrophs that utilize organic carbon source as their electron donor is estimated by using the empirical equation of Heijnen (2010):

$$Y = \left(200 + 18(6 - C)^{1.8} + \exp \left[3.8 - \gamma^{0.32} \times (3.6 + 0.4C) \right] \right) / 12.01, \quad (10)$$

where C is the number of C-atoms of a given carbon source and γ is the degree of reduction of the C atoms in the carbon source.

FIG. 1. Correlations between temperature and maintenance energy using data of aerobic and anaerobic bacteria shown as filled squares (Tijhuis *et al.*, 1993), methanogens shown as open circles (Price and Sowers, 2004) and filled triangles (Tung *et al.*, 2005), and nitrifiers shown as open squares (Rohde *et al.*, 2008) and a filled circle (Price and Sowers, 2004) (Supplementary Table S2). The dashed line is the correlation reported by Tijhuis *et al.*, and the black line is our correlation for temperatures below 10°C: $m_E = 10^{-15} \exp \left(144.9 - \frac{39,656}{T} \right)$. Light shading indicates the 95% confidence level.

As seen in Fig. 1, the trend of the maintenance energy demand at temperatures below 10°C diverges markedly from that for temperatures above 10°C, hence suggesting that the dependence of the metabolic energy requirement of psychrophiles (or psychrotolerants) is not captured by the Tijhuis *et al.* (1993) equation. The low maintenance energy requirement rates likely represent an adaptive response where cells facing severe temperature stress reduce their endogenous metabolism to the most essential functions only (Kempes *et al.*, 2017).

The temperature-dependent metabolic energy requirement rate in $\text{kJ (mol C)}^{-1} \text{ h}^{-1}$ at cold temperatures is obtained by fitting the $\leq 10^\circ\text{C}$ data in Fig. 1 to an Arrhenius-type equation. Assuming a carbon content of methanotrophic cells of $12 \text{ fg-C cell}^{-1}$ (Nihous and Masutani, 2007), the cell-specific endogenous metabolic energy requirement rate in units of $\text{kJ cell}^{-1} \text{ h}^{-1}$, m_E , is then as follows:

$$m_E = 10^{-15} \exp \left(144.9 - \frac{39,656}{T} \right), \quad (11)$$

where T is expressed in K. In the following, Eq. (11) is used to estimate the metabolic requirements of aerobic methanotrophs under martian surface temperatures. The cell-specific net energy gain rate is further obtained by subtracting m_E from the cell-specific rate of catabolic energy production given by Eq. (9):

$$[\text{Cell-specific net energy gain rate}] = -r \bullet \Delta_r G - m_E. \quad (12)$$

The cell-specific net energy gain rate is a proxy measure of the potential for a given group of microorganisms to survive in

a given environment. It is also referred to as the organisms' invasibility according to the definition of Seto (2014). When the net energy gain rate is positive, the microorganisms can, in principle, invest the energy in excess to that required for their maintenance functions into growth. By letting Eq. (12) be 0 and solving for the corresponding CH_4 concentration, we can calculate the minimum CH_4 requirement for aerobic methanotrophs to survive under the assumed martian conditions.

2.3. Cell doubling times

Assuming exponential growth, the cell doubling time is simply the inverse of the specific growth rate multiplied by $\ln 2$. The specific growth rate in turn is obtained by dividing the cell-specific net energy gain rate by the Gibbs energy required to synthesize 1 cell, Y (kJ cell^{-1}). The cell doubling time is thus given by the following:

$$[\text{Cell doubling time}] = \frac{Y \ln 2}{-r\Delta_r G - m_E}. \quad (13)$$

The value of Y is computed with Eq. (10) assuming CH_4 is the sole carbon source. This gives $Y = 1.086 \times 10^{-12} \text{ kJ cell}^{-1}$. Note that Eq. (12) does not account for the energy dissipated as heat, and therefore, Eq. (13) yields a minimum estimate of the cell doubling time.

3. Results

3.1. Thermodynamics and kinetics of aerobic methane oxidation on Mars

Values of $\Delta_r G$ and the cell-specific microbial CH_4 oxidation rate, r , are calculated for the ranges in temperature and pressure encountered at the surface of Mars (180–280 K, 6–11 hPa; Martínez *et al.*, 2017), while fixing the mixing ratios of CO_2 and O_2 to 95% and 0.145%, respectively (Haberle *et al.*, 2017). The default mixing ratio of CH_4 is set to 0.69 parts per billion by volume (ppbv), which is the average of background levels of atmospheric CH_4 reported for *in situ* measurements made with the tunable laser spectrometer on Mars' Curiosity rover (Webster *et al.*, 2015).

Terrestrial-like microbial cells could not survive exposed at the surface of Mars because of the intense UV radiation and ionizing radiation. Rather, they would be expected to reside below the surface shielded by at least 1 mm of regolith to avoid UV radiation (Rummel *et al.*, 2014) and at least 20 cm to avoid ionizing radiation (Dartnell, 2011). Methanotrophy would then rely on O_2 produced photochemically in the atmosphere and CH_4 released from source areas presumably located at depths of several km (Stevens *et al.*, 2015). In our calculations, we assume that the CH_4 soil gas concentration in the upper martian regolith equals that measured in the atmosphere (0.69 ppbv). This is reasonable in view of the low CH_4 outgassing flux of $3.6 \times 10^{-19} \text{ mol CH}_4 \text{ cm}^{-2} \text{ s}^{-1}$ estimated by Krasnopolsky *et al.* (2004), which, together with typical values for gas diffusion coefficients, yields vertical CH_4 concentration gradients of the order of $10^{-18} \text{ mol cm}^{-4}$. In other words, within the uppermost meters of regolith, the fugacity of CH_4 should be close to that in the overlying atmosphere.

For the gas mixing ratios quoted above, and at the usual reference temperature and pressure of 298.15 K and 1 atm, Equation (1) is highly exergonic, with a $\Delta_r G$ value of $-733.3 \text{ kJ (mol CH}_4)^{-1}$ [note: the corresponding $\Delta_r G^\circ$ equals $-817.9 \text{ kJ (mol CH}_4)^{-1}$]. Under martian surface conditions, the calculated Gibbs energies of reaction are of the same order of magnitude (Fig. 2), implying that the low temperatures and pressures on Mars do not significantly alter the thermodynamic energy yield of the aerobic methane oxidation reaction. Even at the very low inferred CH_4 concentrations, Equation (1) remains thermodynamically highly favorable; CH_4 concentrations two orders of magnitude higher than 0.69 ppbv only yield $\Delta_r G$ values that are around $10 \text{ kJ (mol CH}_4)^{-1}$ more negative at 230 K and 7 hPa (results not shown).

The resulting cell-specific CH_4 oxidation rates calculated with Eq. (7) fall well below the maximum rates $v_{\max}$, that is, they are strongly limited by substrate availability, in particular the low availability of CH_4 . For comparison, when the mixing ratio of CH_4 is set to the Earth's atmosphere average of 1.8 ppm, the calculated methane oxidation rates are four orders of magnitude faster than those calculated for the default martian CH_4 mixing ratio of 0.69 ppbv, all other conditions equal. The predicted limitation of martian CH_4

FIG. 2. Standard Gibbs free energy change, $\Delta_r G^\circ$, and the Gibbs free energy change, $\Delta_r G$, of methane oxidation (left and right panel, respectively) in units of $\text{kJ (mol CH}_4)^{-1}$ against temperature and pressure. The imposed mixing ratios of CO_2 , O_2 , and CH_4 are 95%, 0.145%, and 0.69 ppbv, respectively.

oxidation kinetics is a direct consequence of the relative magnitudes of the concentrations of the gaseous reactants and the corresponding half-saturation constants in the Michaelis–Menten formulation of Eq. (7). For the 19 terrestrial methanotrophic strains, values of K_{CH_4} are in the ppm range (Knief and Dunfield, 2005), that is, orders of magnitude higher than $[\text{CH}_4]$ at the surface of Mars. Because $[\text{CH}_4] \ll K_{\text{CH}_4}$, Eq. (7) yields a (near-)linear dependence of the cell-specific CH_4 oxidation rate, r , on the CH_4 concentration. In contrast, the imposed half-saturation constant for O_2 (1.9 ppm) is only slightly higher than the martian atmospheric O_2 concentration and limitation by O_2 is therefore far less severe than for CH_4 .

Based on the analysis above, we conclude that the cell-specific energy gain rates for aerobic CH_4 oxidation in the upper martian regolith [*i.e.*, $-r \cdot \Delta_r G$, Eq. (9)] are most likely kinetically, rather than thermodynamically, limited. Despite similar Gibbs energies of reaction, the cell-specific energy gain rates at the average mixing ratios of O_2 and CH_4 in Earth's atmosphere at 298.13 K and 1 atm are four orders of magnitude faster than those calculated at the default values of the mixing ratios of O_2 and CH_4 in the martian atmosphere at 230 K and 11 hPa. However, this conclusion depends on the extrapolation of the CH_4 half-saturation (or affinity) constants of terrestrial methanotrophs (K_{CH_4}) to their hypothetical martian counterparts.

3.2. Minimum CH_4 requirement for endogenous metabolism

The cell-specific net energy gain rates for all 19 strains of terrestrial aerobic methanotrophs are estimated to be positive at <250 K (shown in Fig. 3 for *Methylocystis* sp. LR1). Although the cell-specific catabolic energy production rates of methanotrophs on Mars are predicted to be much slower than those on Earth, these rates still exceed the low metabolic energy requirement rates expected at cold temperatures. Thus, terrestrial-like aerobic methanotrophy is a potentially viable metabolism under the assumed conditions for the martian upper regolith.

The minimum CH_4 levels needed to sustain endogenous metabolism of methanotrophs at 230 K and 6 hPa are shown in Fig. 4. At 230 K, Eq. (11) predicts a cell-specific maintenance energy requirement rate of 1.12×10^{-27} kJ cell $^{-1}$ h $^{-1}$. Together with the kinetic parameters extracted from the 19 terrestrial methanotrophic strains, the calculations then yield an average threshold CH_4 concentration of 2.26×10^{-6} ppbv (maximum: 9.18×10^{-6} ppbv, minimum: 6.24×10^{-8} ppbv). According to these results, ppbv levels of CH_4 should enable martian methanotrophs to survive. The high variability in the calculated threshold CH_4 concentrations in Fig. 4 primarily reflects the very steep temperature dependence of the maintenance energy requirement (m_E) imposed in the calculations (Fig. 1). For comparison, all other conditions unchanged, at 270 K, where m_E equals 1.39×10^{-16} kJ cell $^{-1}$ h $^{-1}$, the methanotrophs would require more than 10 ppmv CH_4 to fulfill their maintenance requirements.

3.3. Doubling times

For the majority of the 19 terrestrial analog strains, Eq. (13) predicts cell doubling times in excess of 10,000 Earth years at 230 K and 0.69 ppbv CH_4 (Fig. 5). *Methylosinus*

FIG. 3. Calculated cell-specific net energy gain rate of *Methylocystis* sp. LR1 under assumed martian temperature and pressure conditions. v_{max} and K_{CH_4} are set to 5×10^{-17} mol CH_4 cell $^{-1}$ h $^{-1}$ and 2.5×10^{-6} mol L $^{-1}$, respectively (Knief and Dunfield, 2005); K_{O_2} is set to 1.9×10^{-9} mol L $^{-1}$ (Van Bodgom *et al.*, 2001). The cell-specific energy gain rates are calculated using Eq. (12). The imposed mixing ratios of CO_2 , O_2 , and CH_4 are 95%, 0.145%, and 0.69 ppbv, respectively.

trichosporium OB3b yields the shortest doubling time with 4831 years. The calculated doubling times for other *M. trichosporium* species are significantly longer, however, possibly reflecting the uncertainties associated with the estimation of the kinetic parameters used in Eq. (7).

The calculated cell doubling times in Fig. 5 are significantly longer than the 160 days estimated for methanotrophic bacteria retrieved from Siberian permafrost and incubated at -20°C (Rivkina *et al.*, 2000). They are also longer than the estimated generation times of subsurface cells, which are typically on the order of hundreds of years (Biddle *et al.*, 2006; Starnawski *et al.*, 2017) and occasionally up to tens of thousands of years (Jørgensen and Boetius, 2007). Similar doubling times for martian methanotrophs would require CH_4 mixing ratios that are several orders of magnitude larger than the reported background level of 0.69 ppbv CH_4 . Even at a CH_4 mixing ratio of 1 ppmv, which is unlikely for the martian atmosphere, the average estimated cell doubling times of the methanotrophs would still be 164 Earth years. In other words, should the methanotrophs be able to generate enough energy to survive, new biomass growth would be extremely slow. As an analogue, microorganisms retrieved from the deep seafloor have been shown to maintain their vital functions, but only grow when newly supplied with energy sources (Morono *et al.*, 2011).

4. Discussion

Microbial processes account for about 69% of all CH_4 sources on Earth (Conrad, 2009) and methanogens are among the most studied microorganisms in relation to potential life on Mars (Taubner *et al.*, 2015). The potential biogenic CH_4 production on Mars has been estimated at 8.8×10^5 molecules cm^{-2} s $^{-1}$, based on inferred atmospheric production rates of CO

FIG. 4. Minimum CH_4 requirements for the 19 strains of methanotrophs at martian average temperature and pressure of 230 K and at 6 hPa; the shaded band corresponds to the values at ± 30 K. The values of v_{max} and K_{CH_4} are extracted from the rate data for 19 strains of aerobic methanotrophs compiled by Knief and Dunfield (2005); K_{O_2} is set to $1.9 \times 10^{-9} \text{ mol L}^{-1}$ (Van Bodegom *et al.*, 2001). The cell-specific energy gain rates are calculated with Eq. (12). The imposed mixing ratios of CO_2 , O_2 , and CH_4 are 95%, 0.145%, and 0.69 ppbv, respectively.

and H_2 (Krasnopolsky *et al.*, 2004). Given the reported photochemical loss rate of CH_4 in the martian atmosphere of around $2.2 \times 10^5 \text{ molecules cm}^{-2} \text{ s}^{-1}$ (Krasnopolsky *et al.*, 2004), microbial consumption of CH_4 could account for up to $6.6 \times 10^5 \text{ molecules cm}^{-2} \text{ s}^{-1}$ ($= 8.8 \times 10^5 - 2.2 \times 10^5 \text{ molecules cm}^{-2} \text{ s}^{-1}$), assuming that other CH_4 sinks are negligible. Abiotic CH_4 decomposition on Mars, however, could be catalyzed by meteorological-driven electric fields (Farrell *et al.*, 2006) and wind-driven erosion that produces highly reactive sites on mineral grain surfaces (Knak Jensen *et al.*, 2014).

Assuming that methanotrophs are the main CH_4 sink, dividing the estimated $6.6 \times 10^5 \text{ molecules cm}^{-2} \text{ s}^{-1}$ consumption rate by the cell-specific rate of CH_4 oxidation, r in Eq. (7), of the 19 strains, yields an average aerobic methanotrophic cell density per unit of regolith area of $1.6 \times 10^6 \text{ cells cm}^{-2}$ (maximum: $6.5 \times 10^6 \text{ cells cm}^{-2}$ for *Methylocapsa acidiphila* B2; minimum: $3.2 \times 10^4 \text{ cells cm}^{-2}$ for *M. trichosporium* OB3b), at 230 K and 6 hPa, and for the default mixing ratios of O_2 and CH_4 in the martian atmosphere. Further assuming a carbon content for methanotrophic cells of $12 \text{ fg-C cell}^{-1}$ (Nihous and Masutani, 2007), the average methanotrophic biomass per unit regolith surface area is then $1.9 \times 10^{-4} \text{ gC m}^{-2}$. For comparison, the average microbial biomass in the upper 0–15 cm of ocean sediment at 8 km water depth is around 0.1 gC m^{-2} (Jørgensen and Boetius, 2007). Thus, according to our results, a hypothetical methanotrophic population in martian soils would be extremely sparse.

FIG. 5. Cell doubling times for 19 strains of methanotrophs for the average of background level CH_4 reported from Webster *et al.* (2015) (0.69 ppbv, shown by solid line), and for two orders of magnitude higher than this default value (69 ppbv, dashed line) at 230 K and 6 hPa. The values of v_{max} and K_{CH_4} are extracted from the rate data for 19 strains of aerobic methanotrophs compiled by Knief and Dunfield (2005); K_{O_2} is set to $1.9 \times 10^{-9} \text{ mol L}^{-1}$ (Van Bodegom *et al.*, 2001). The cell-specific energy gain rates are calculated using Eq. (12). The imposed mixing ratios of CO_2 , O_2 , and CH_4 are set to 95%, 0.145%, and 0.69 ppbv, respectively.

The abundance of CH_4 in the martian atmosphere is still under debate. Here, we use the CH_4 mixing ratio of CH_4 of 0.69 ppbv from the work of Webster *et al.* (2015) as the default value. The latter was recently updated to 0.41 ppbv (Webster *et al.*, 2018). For this CH_4 level, our calculations still imply that methanotrophs should be able to support their endogenous metabolism, but that their cell doubling times would be longer (minimum: 8141 years). By contrast, other researchers have proposed CH_4 levels of tens of ppbv (Fonti and Marzo, 2010; Geminale *et al.*, 2011), that is, significantly higher than the default value. Nonetheless, even at these higher CH_4 mixing ratios, biomass growth of terrestrial-like methanotrophic microorganisms would remain very slow. Our calculations, however, consider a uniform CH_4 distribution in the martian regolith. More likely, methanotrophic habitats would be concentrated near subsurface sources of CH_4 or along preferential transport pathways, as these areas would exhibit local CH_4 concentrations in excess of the atmospheric concentration (Webster *et al.*, 2015, 2018).

In addition to the low production and abundance of CH_4 , the metabolic activity of methanotrophs similar to that found on Earth would likely be limited by the cold surface temperatures on Mars. The lower temperature limit for terrestrial microbial metabolic activity is still an open question. Measurable

metabolic activity of bacteria has been recorded for temperatures down to -20°C (Rivkina *et al.*, 2000; Junge *et al.*, 2004; Mykytczuk *et al.*, 2013), but convincing evidence for cell reproduction below -18°C is yet to be presented (Moissl-Eichinger *et al.*, 2016). Aerobic methanotrophs have been retrieved from tundra soils, polar lakes, and permafrost sediments (Trotsenko and Khmelenina, 2002, 2005). Type I methanotrophs are widely distributed in cold environments (0 – 10°C) (He *et al.*, 2012), and *Methylobacter*-like methanotrophs are known to be psychrophilic (Oshkin *et al.*, 2014). Viable methanotrophic bacteria from permafrost are able to oxidize CH_4 when incubated at -5°C (Khmelenina *et al.*, 2002). Thus, although it is unclear whether permafrost bacteria are metabolically active or growing *in situ*, they retain their metabolic potential at subzero temperatures. The metabolic state of these cells therefore resembles that of microorganisms deeply buried below the seafloor, which are able to reactivate their dormant metabolic functions when re-exposed to favorable conditions (Jørgensen, 2011; Morono *et al.*, 2011).

The physiological parameters of terrestrial microorganisms that are applied under assumed martian conditions have intrinsic uncertainties associated with them. For example, the cell-specific rates of CH_4 oxidation for the 19 strains are calculated by using just one reported value for the half-saturation constant of O_2 ; $K_{\text{O}_2} = 1.9$ ppm. When K_{O_2} is three orders of magnitude higher or lower than this value, the estimated average methanotrophic microbial biomass in the martian regolith becomes 8.1×10^6 or 1.6×10^6 cells cm^{-2} , respectively. Thus, in this particular case, the uncertainty on K_{O_2} would actually have a relatively small influence on the predicted abundance and activity of methanotrophs.

A much higher degree of uncertainty is associated with the Arrhenius equation describing the endogenous metabolic energy requirement at temperatures below 10°C [Eq. (11)]. This equation is based on limited data that include only one aerobic bacterial species growing on methanol. The implicit assumption that microbial maintenance energy requirements are relatively insensitive to the genetic species composition is consistent with the temperature-dependent maintenance energy relationship of Tijhuis *et al.* (1993), which holds across a variety of bacterial species (Harder, 1997). Nevertheless, assessing the validity of Eq. (11) for aerobic methanotrophs surviving at extremely cold temperatures will need further data on their endogenous metabolic energy demands. In addition, the energy conversion efficiency of CH_4 oxidation is likely temperature dependent. At extremely cold temperature, cell doubling times may become longer than the values calculated with Eq. (13), because energy dissipation as heat would be expected to increase with decreasing temperature.

The kinetic parameters for the 19 strains of terrestrial methanotrophs used to compute aerobic methane oxidation under martian conditions are based on laboratory experiments carried out under typical Earth surface temperatures and pressures. Given that for a given enzymatic reaction pathway the value of v_{max} is expected to decrease with decreasing temperature (D'Amico *et al.*, 2006), the calculated energy gains at martian surface temperatures could be overestimated, hence compromising the proposed viability of methanotrophy in the martian regolith. For example, potential rates of CH_4 oxidation measured in incubations with samples from Lake Qalluuraq, Alaska, show a progressive drop when decreasing the temperature from 21°C to 4°C (He *et al.*, 2012). However, because of long-

term acclimation to permanently cold conditions, psychrophilic microbial populations may exhibit specific metabolic rates at low temperatures that rival those of their mesotrophic counterparts at moderate temperatures (Knoblauch and Jørgensen, 1999). To our knowledge, direct evidence of such an adaptation for psychrophilic methanotrophs is currently missing. This evidence could include measurements of relatively high methane oxidation rates at low (3 – 5°C) and subzero (-5°C) temperatures (Khmelenina *et al.*, 2002; Oshkin *et al.*, 2014).

Besides cold temperatures, other environmental factors, such as low pressure, high salinity, toxins, and harmful ionizing radiation, may all impede the activity of martian methanotrophs, or for that matter microbial life in general. Further studies on extremophiles would undoubtedly strengthen our ability to better account for these additional environmental stressors on the potential survival and growth of terrestrial-like microorganisms in the martian regolith.

5. Conclusions

Theoretical Gibbs energies of reaction, empirical maintenance energy requirements, and observed kinetic parameters for 19 strains of terrestrial methanotrophs are combined to assess thermodynamic and kinetic constraints on aerobic methanotrophy in the martian regolith. Our analysis suggests that aerobic methanotrophs similar to those found on Earth could potentially support their endogenous metabolism at the cold temperatures and ppbv CH_4 levels encountered at the surface of Mars. Nonetheless, the calculations also imply that the low availability of CH_4 (rather than that of O_2) would impose a severe kinetic limitation on the cell-specific methanotrophic energy gain rates. In turn, this would result in very long cell doubling times making it unlikely that methanotrophic cells would proliferate to great numbers under current martian conditions, with the possible exception of habitats in close proximity of CH_4 source areas or along preferential efflux pathways.

Acknowledgments

This work was supported by the Canada Excellence Research Chair (CERC) Program to P.V.C. and by JSPS Grant-in-Aid for Scientific Research (C) Grant Number 19K06853 to M.S. We are thankful to the two anonymous reviewers for their constructive comments.

Author Disclosure Statement

No competing financial interests exist.

Supplementary Material

Supplementary Table S1

Supplementary Table S2

References

- Amend, J.P. and Shock, E.L. (2001) Energetics of overall metabolic reactions of thermophilic and hyperthermophilic archaea and bacteria. *FEMS Microbiol Rev* 25:175–243.
- Atreya, S.K., Mahaffy, P.R., and Wong, A.-S. (2007) Methane and related trace species on Mars: origin, implications for life, and habitability. *Planet Space Sci* 55:358–369.
- Berry, B.J., Jenkins, D.G., and Schuerger, A.C. (2010) Effects of simulated Mars conditions on the survival and growth of

- Escherichia coli* and *Serratia liquefaciens*. *Appl Environ Microbiol* 76:2386–2388.
- Biddle, J.F., Lipp, J.S., Lever, M.A., Lloyd, K.G., Sørensen, K.B., Anderson, R., Fredricks, H.F., Elvert, M., Kelly, T.J., Schrag, D.P., Sogin, M.L., Brechley, J.E., Teske, A., House, C.H., and Hinrichs, K.-U. (2006) Heterotrophic Archaea dominate sedimentary subsurface ecosystems off Peru. *Proc Natl Acad Sci U S A* 103:3846–3851.
- Bull, I.D., Parekh, N.R., Hall, G.H., Ineson, P., and Evershed, R.P. (2000) Detection and classification of atmospheric methane oxidizing bacteria in soil. *Nature* 405:175–178.
- Conrad, R. (1996) Soil microorganisms as controllers of atmospheric trace gases (H_2 , CO, CH_4 , OCS, N_2O , and NO). *Microbiol Rev* 60:609–640.
- Conrad, R. (2009) The global methane cycle: recent advances in understanding the microbial processes involved. *Environ Microbiol Rep* 1:285–292.
- D'Amico, S., Collins, T., Marx, J.C., Feller, G., and Gerday, C. (2006) Psychrophilic microorganisms: challenges for life. *EMBO Rep* 7:385–389.
- Dartnell, L.R. (2011) Ionizing radiation and life. *Astrobiology* 11:551–582.
- Farrell, W.M., Delory, G.T., and Atreya, S.K. (2006) Martian dust storms as a possible sink of atmospheric methane. *Geophys Res Lett* 33:L21203.
- Fonti, S. and Marzo, G.A. (2010) Mapping the methane on Mars. *Astron Astrophys* 512:A51.
- Formisano, V., Atreya, S., Encrenaz, T., Ignatiev, N., and Giuranna, M. (2004) Detection of methane in the atmosphere of Mars. *Science* 306:1758–1761.
- Geminale, A., Formisano, V., and Sindoni, G. (2011) Mapping methane in Martian atmosphere with PFS-MEX data. *Planet Space Sci* 59:137–148.
- Haberle, R. M., Clancy, R. T., Forget, F., Smith, M. D., and Zurek, R. W. (Eds.). (2017). *The atmosphere and climate of Mars*. Cambridge University Press, Cambridge, England.
- Hanson, R.S. and Hanson, T.E. (1996) Methanotrophic bacteria. *Microbiol Mol Biol Rev* 60:439–471.
- Harder, J. (1997) Species-independent maintenance energy and natural population sizes. *FEMS Microbiol Ecol* 23:39–44.
- He, R., Wooller, M.J., Pohlman, J.W., Quensen, J., Tiedje, J.M., and Leigh, M.B. (2012) Shifts in identity and activity of methanotrophs in Arctic Lake sediments in response to temperature changes. *Appl Environ Microbiol* 78:4715–4723.
- Heijnen, J.J. (2010) *Encyclopedia of Industrial Biotechnology, Bioprocess, Bioseparation, and Cell Technology*, 7 Volumes Set. John Wiley & Sons, pp. 267–290.
- Helgeson, H.C., Delany, J.M., Nesbitt, H.W., and Bird, D.K. (1978) Summary and critique of the thermodynamic properties of rock-forming minerals. *Am J Sci* 278:1–229.
- Hoehler, T.M. (2004) Biological energy requirements as quantitative boundary conditions for life in the subsurface. *Geobiology* 2:205–215.
- Jørgensen, B.B. (2011) Deep subseafloor microbial cells on physiological standby. *Proc Natl Acad Sci U S A* 108:18193–18194.
- Jørgensen, B.B. and Boetius, A. (2007) Feast and famine—microbial life in the deep-sea bed. *Nat Rev Microbiol* 5:770–781.
- Junge, K., Eicken, H., and Deming, J.W. (2004) Bacterial activity at -2 to -20°C in arctic wintertime sea ice. *Appl Environ Microbiol* 70:550–557.
- Kempes, C.P., van Bodegom, P.M., Wolpert, D., Libby, E., Amend, J., and Hoehler, T. (2017) Drivers of bacterial maintenance and minimal energy requirements. *Front Microbiol* 8:31.
- Khmelenina, V.N., Makutina, V.A., Kalyuzhnaya, M.G., Rivkina, E.M., Gilichinsky, D.A., and Trotsenko, Y.A. (2002) Discovery of viable methanotrophic bacteria in permafrost sediments of northeast Siberia. *Dokl. Biol. Sci [Proc USSR Biol Sci]* (translated from Russian) 384:235–237.
- Knak Jensen, S.J., Skibsted, J., Jakobsen, H.J., ten Kate, I.L., Gunnlaugsson, H.P., Merrison, J.P., Finster, K., Bak, E., Iversen, J.J., Kondrup, J.C., and Nørnberg, P. (2014) A sink for methane on Mars? The answer is blowing in the wind. *Icarus* 236:24–27.
- Knief, C. and Dunfield, P.F. (2005) Response and adaptation of different methanotrophic bacteria to low methane mixing ratios. *Environ Microbiol* 7:1307–1317.
- Knoblauch, C. and Jørgensen, B.B. (1999) Effect of temperature on sulphate reduction, growth rate and growth yield in five psychrophilic sulphate-reducing bacteria from Arctic sediments. *Environ Microbiol* 1:457–467.
- Krasnopolsky, V.A., Maillard, J.P., and Owen, T.C. (2004) Detection of methane in the martian atmosphere: evidence for life? *Icarus* 172:537–547.
- LaRowe, D.E. and Amend, J.P. (2015) Catabolic rates, population sizes and doubling/replacement times of microorganisms in natural settings. *Am J Sci* 315:167–203.
- Martínez, G.M., Newman, C.N., De Vicente-Retortillo, A., Fischer, E., Renno, N.O., Richardson, M.I., Fairén, A.G., Genzer, M., Guzewich, S.D., Haberle, R.M., Harri, A.M., Kemppinen, O., Lemmon, M.T., Smith, M.D., de la Torre-Juárez, M., and Vasavada, A.R. (2017) The modern near-surface martian climate: a review of in-situ meteorological data from Viking to Curiosity. *Space Sci Rev* 212:295–338.
- McCollom, T.M. and Shock, E.L. (1997) Geochemical constraints on chemolithoautotrophic metabolism by microorganisms in seafloor hydrothermal systems. *Geochim Cosmochim Acta* 61:4375–4391.
- McKay, D.S., Gibson Jr., E.K., Thomas-Keptra, K.L., Vail, H., Romanek, C.S., Clemett, S.J., Chillier, X.D.F., Maechling, C.R., and Zare, R.N. (1996) Search for past life on Mars: possible relic biogenic activity in Martian meteorite ALH84001. *Science* 273:924–930.
- Moissl-Eichinger, C., Cockell, C., and Rettberg, P. (2016) Venturing into new realms? Microorganisms in space. *FEMS Microbiol Rev* 40:722–737.
- Morono, Y., Terada, T., Nishizawa, M., Ito, M., Hillion, F., Takahata, N., Sano, Y., and Inagaki, F. (2011) Carbon and nitrogen assimilation in deep subseafloor microbial cells. *Proc Natl Acad Sci U S A* 20110776, 308(45):18295–18300.
- Mumma, M.J., Villanueva, G.L., Novak, R.E., Hewagama, T., Bonev, B.P., DiSanti, M.A., Mandell, A.M., and Smith, M.D. (2009) Stong release of methane on Mars in northern summer 2003. *Science* 323:1041–1045.
- Mykytczuk, N.C.S., Foote, S.J., Omelon, C.R., Southam, G., Greer, C.W., and Whyte, L.G. (2013) Bacterial growth at -15°C ; molecular insights from the permafrost bacterium *Planococcus halocryophilus* Or1. *ISME J* 7:1211–1226.
- Nicholson, W.L., Krivushin, K., Gilichinsky, D., and Schuerger, A.C. (2013) Growth of *Carnobacterium* spp. from permafrost under low pressure, temperature, and anoxic atmosphere has implications for Earth microbes on Mars. *Proc Natl Acad Sci U S A* 110:666–671.
- Nihous, G.C. and Masutani, M. (2007) Notes on the modeling of methane in aging hydrothermal plumes. *J Mar Res* 65:789–812.
- Orosei, R., Lauro, S.E., Pettinelli, E., Cicchetti, A., Coradini, M., Cosciotti, B., Di Paolo, F., Flamini, E., Mattei, E., Pajola, M., Soldovieri, F., Cartacci, M., Cassenti, F., Frigeri, A., Giuppi,

- S., Martufi, R., Masdea, A., Mitri, G., Nenna, C., Noschese, R., Restano, M., and Seu, R. (2018) Radar evidence of subglacial liquid water on Mars. *Science* 361:490–493.
- Oshkin, I.Y., Wegner, C.E., Lükec, C., Glagolev, M. V., Filippov, I. V., Pimenov, N. V., Liesack, W., and Dedysha, S.N. (2014) Gammaproteobacterial methanotrophs dominate cold methane seeps in floodplains of West Siberian rivers. *Appl Environ Microbiol* 80:5944–5954.
- Pollack, J.B. and Kasting, J.F., 1987. The case for a wet, warm climate on early Mars. *Icarus* 71:203–224.
- Price, P.B. and Sowers, T. (2004) Temperature dependence of metabolic rates for microbial growth, maintenance, and survival. *Proc Natl Acad Sci U S A* 101:4631–4636.
- Rivkina, E.M., Friedmann, E.I., McKay, C.P. and Gilichinsky, D.A. (2000) Metabolic activity of permafrost bacteria below the freezing point. *Appl Environ Microbiol* 66:3230–3233.
- Rohde, R.A., Price, P.B., Bay, R.C., and Bramall, N.E. (2008) In situ microbial metabolism as a cause of gas anomalies in ice. *Proc Natl Acad Sci U S A* 105:8667–8672.
- Rummel, J.D., Beaty, D.W., Jones, M.a, Bakermans, C., Barlow, N.G., Boston, P.J., Chevrier, V.F., Clark, B.C., de Vera, J.-P.P., Gough, R. V., Hallsworth, J.E., Head, J.W., Hipkin, V.J., Kieft, T.L., McEwen, A.S., Mellon, M.T., Mikucki, J.a, Nicholson, W.L., Omelon, C.R., Peterson, R., Roden, E.E., Sherwood Lollar, B., Tanaka, K.L., Viola, D., and Wray, J.J. (2014) A new analysis of Mars “special regions”: findings of the second MEPAG Special Regions Science Analysis Group (SR-SAG2). *Astrobiology* 14:887–968.
- Seto, M. (2014) The Gibbs free energy threshold for the invasion of a microbial population under kinetic constraints. *Geomicrobiol J* 31:645–653.
- Shock, E.L., Holland, M., Meyer-Dombard, D., Amend, J.P., Osburn, G.R., and Fischer, T.P. (2010) Quantifying inorganic sources of geochemical energy in hydrothermal ecosystems, Yellowstone National Park, USA. *Geochim Cosmochim Acta* 74:4005–4043.
- Smith, J.M. and Van Ness H.C. (1987) *Introduction to Chemical Engineering Thermodynamics*, 4th ed., McGraw-Hill, Singapore.
- Smith, M.D., Bougher, S. W., Encrenaz, T., Forget, F., and Kleinböhl, A. (2017) Thermal structure and composition. In *The Atmosphere and Climate of Mars*, edited by R.M. Haberle, R.T. Clancy, F. Forget, M.D. Smith, and R.W. Zurek, Cambridge University Press, Cambridge.
- Sowers, T. (2001) N₂O record spanning the penultimate deglaciation from the Vostok ice core. *J Geophys Res Atmos* 106:31903–31914.
- Squyres, S.W., Grotzinger, J.P., Arvidson, R.E., Bel III, J.F., Calvin, W., Christensen, P.R., Clark, B.C., Crisp, J.A., Farrand, W.H., Herkenhoff, K.E., Johnson, J.R., Klingelhöfer, G., Knoll, A.H., McLennan, S.M., McSween, Jr., H.Y., Morris, R. V, Rice Jr., J.W., Rieder, R., and Soderblom, L.A. (2004) In situ evidence for an ancient aqueous environment at Meridiani Planum, Mars. *Science* 306:1709–1714.
- Starnawski, P., Bataillon, T., Ettema, T.J.G., Jochum, L.M., Schreiber, L., Chen, X., Lever, M.A., Polz, M.F., Jørgensen, B.B., Schramm, A., and Kjeldsen, K.U. (2017) Microbial community assembly and evolution in subsurface sediment. *Proc Natl Acad Sci U S A* 201614190.
- Stevens, A.H., Patel, M.R., and Lewis, S.P. (2015) Numerical modelling of the transport of trace gases including methane in the subsurface of Mars. *Icarus* 250:587–594.
- Taubner, R.-S., Schleper, C., Firneis, M., and Rittmann, S. (2015) Assessing the ecophysiology of methanogens in the context of recent astrobiological and planetological studies. *Life* 5:1652–1686.
- Tijhuis, L., van Loosdrecht, M.C., and Heijnen, J.J. (1993) A thermodynamically based correlation for maintenance gibbs energy requirements in aerobic and anaerobic chemotrophic growth. *Biotechnol Bioeng* 42:509–519.
- Trotsenko, Y.A. and Khmelenina, V.N. (2002) Biology of extremophilic and extremotolerant methanotrophs. *Arch Microbiol* 177:123–131.
- Trotsenko, Y.A. and Khmelenina, V.N. (2005) Aerobic methanotrophic bacteria of cold ecosystems. *FEMS Microbiol Ecol* 53:15–26.
- Tung, H.C., Bramall, N.E., and Price, P.B. (2005) Microbial origin of excess methane in glacial ice and implications for life on Mars. *Proc Natl Acad Sci U S A* 102:18292–18296.
- Van Bodegom, P., Stams, F., Mollema, L., Boeke, S., and Leffelaar, P. (2001) Methane oxidation and the competition for oxygen in the rice rhizosphere. *Appl Environ Microbiol* 67:3586–3597.
- Webster, C.R., Mahaffy, P.R., Atreya, S.K., Flesch, G.J., Farley, K.A., and the MSL Science Team (2013) Low upper limit to methane abundance on Mars. *Science* 342:355–357.
- Webster, C.R., Mahaffy, P.R., Atreya, S.K., Flesch, G.J., Mischna, M.A., Meslin, P., Farley, K.A., Conrad, P.G., Christensen, L.E., Pavlov, A.A., Martín-Torres, J., Zorzano, M., McConnochie, T.H., Owen, T., Eigenbrode, J.L., Glavin, D.P., Andrew, S., Malespin, C.A., Archer Jr., D., Sutter, B., Coll, P., Freissinet, C., McKay, C.P., Moores, J.E., Schwenzer, S.P., Bridges, J.C., Navarro-Gonzalez, R., Gellert, R., Lemmon, M.T., and the MSL Science Team (2015) Mars methane detection and variability at Gale crater. *Science* 347:415–417.
- Webster, C.R., Mahaffy, P.R., Atreya, S.K., Moores, J.E., Flesch, G.J., Malespin, C., McKay, C.P., Martinez, G., Smith, C.L., Martín-Torres, J., Gomez-Elvira, J., Zorzano, M.-P., Wong, M.H., Trainer, M.G., Steele, A., Archer, D., Sutter, B., Coll, P.J., Freissinet, C., Meslin, P.-Y., Gough, R. V, House, C.H., Pavlov, A., Eigenbrode, J.L., Glavin, D.P., Pearson, J.C., Keymeulen, D., Christensen, L.E., Schwenzer, S.P., Navarro-Gonzalez, R., Pla-García, J., Rafkin, S.C.R., Vicente-Retortillo, Á., Kahanpää, H., Viudez-Moreiras, D., Smith, M.D., Harri, A.-M., Genzer, M., Hassler, D.M., Lemmon, M., Crisp, J., Sander, S.P., Zurek, R.W., and Vasavada, A.R. (2018) Background levels of methane in Mars’ atmosphere show strong seasonal variations. *Science* 360:1093 LP-1096.

Address correspondence to:

Mayumi Seto
Department of Chemistry, Biology,
and Environmental Sciences
Faculty of Science
Nara Women's University
Kita-Uoya Nishimachi
Nara 630-8506
Japan

E-mail: seto@ics.nara-wu.ac.jp

Submitted 20 August 2018

Accepted 7 May 2019

Associate Editor: Petra Rettberg

Abbreviation Used

ppbv = parts per billion by volume