Published in final edited form as:

Per Med. 2019 September ; 16(5): 365–367. doi:10.2217/pme-2019-0045.

Can Precision Medicine Help Achieve the Goal of Reducing Care When the Risks Exceed the Benefits?

Kathryn A. Phillips, PhD^{1,2}, Deborah A Marshall, PhD³, Allison W. Kurian, MD, M.Sc⁴

¹UCSF Center for Translational and Policy Research on Personalized Medicine (TRANSPERS); Department of Clinical Pharmacy, University of California San Francisco

²UCSF Helen Diller Family Comprehensive Cancer Center

³Department of Community Health Sciences, University of Calgary

⁴Departments of Medicine and of Health Research and Policy, Stanford University

Keywords

Precision Medicine; Economics; Preferences; Evidence-Based Medicine

It is widely understood that more health care does not always result in better outcomes for patients, and there is ample evidence that many health care interventions in the U.S. are unnecessary and thus drive up costs as well as cause harm. Thus there is a growing movement to develop approaches to reduce care when the risks exceed the benefits, e.g., the “Choosing Wisely” program, which aims to facilitate choosing care that is supported by evidence, not duplicative, free from harm, and truly necessary.[1]

We examine the question: **Can precision medicine help achieve the goal of reducing care when the risks exceed the benefits?** If so, precision medicine (the use of genomic information to target health care) could improve not only patient outcomes but also outcomes for health care systems and society as part of a larger agenda for U.S. health care. However, to date there has been a greater focus on the ability of precision medicine to *increase* the use of more effective treatment than its ability to *decrease* care when the risks exceed the benefits. We discuss opportunities for precision medicine to facilitate the goal of decreasing unnecessary care – but also assert that *this potential benefit is largely understudied and untapped*. We discuss how precision medicine illustrates both the opportunities and challenges to achieving the overarching goal of reducing care when the risks exceed the benefits. We focus particularly on the ubiquitous challenge that actively choosing to *forgo* treatment is typically more difficult for patients and clinicians than choosing to *obtain* treatment.[2, 3]

Corresponding Author Kathryn A. Phillips, PhD, UCSF Center for Translational and Policy Research on Personalized Medicine (TRANSPERS); Department of Clinical Pharmacy, University of California San Francisco; UCSF Philip R. Lee Institute for Health Policy; and UCSF Helen Diller Family Comprehensive Cancer Center, 3333 California St, Room 420, Box 0613, San Francisco, CA 94143, USA, Tel: (415) 502-8271, Fax: (415) 502-0792, Kathryn.Phillips@ucsf.edu.

Early studies suggested that precision medicine was *not* more likely to facilitate willingness to forgo treatment than other interventions. For example, our nationally representative 2013 study found that many individuals stated that they would *not* change decisions based on precision medicine test results: when asked if genetic testing indicated that they are not candidates for oncology therapies, most (84%) would seek a second opinion or want therapy anyway.[4]

In contrast to the early findings, there is now some evidence suggesting that precision medicine might facilitate the willingness of patients and clinicians to forgo treatment. We use the example of gene expression profiling for patients with early-stage breast cancer (GEP), which predicts whether patients are likely to benefit from chemotherapy, to illustrate both the opportunities and challenges. There is now widely accepted evidence, including results from a pivotal trial, that GEP testing can identify patients who can safely forgo chemotherapy.[5] There is also strong evidence that the actual use of chemotherapy among these patients has decreased because of GEP testing and other factors.[6] In addition to these clinical studies, there is evidence from consumer and physician surveys that individuals value the information provided by GEP and are willing to use it to make a decision to forgo treatment. For example, a quantitative preferences study found that women state that GEP scores would highly influence their willingness to forgo chemotherapy [7] and a qualitative study found that GEP would be the “deciding factor”.[8] Similar results were found in a study of physicians: 79% of physicians reported greater confidence in their treatment recommendations (including forgoing chemotherapy) as a result of GEP.[9]

Thus, GEP testing provides an example of how precision medicine might contribute to reduction of unnecessary care. However, there is not enough evidence on the extent to which GEP has been successful in reducing unnecessary care because it is a genomics test – and thus whether this example might translate to other precision medicine interventions. We assert that the example of GEP suggests missed opportunities to understand the potential impact of precision medicine and how to harness its benefits, which are relevant not only to this specific example but also more broadly.

- Importantly, **we did not find any studies directly examining whether the use of precision medicine *per se* is facilitating a greater willingness to forgo treatment and its contribution relative to other factors.** We cannot determine from existing studies whether GEP had an influence simply because it met a diagnostic gap or because GEP is based on genomic information and such tests are viewed as more accurate and useful. Studies could shed light on this question through longitudinal, time-series studies that directly examine preferences and beliefs about precision medicine and relationship to actual behaviors. This would enable analysis of changes over time and whether specific evidence such as adoption of a new precision medicine test contributed to changes in forgoing treatment.
- Few studies have **directly asked patients and clinicians about their preferences regarding forgoing treatment and the impact of precision medicine on their preferences.** Such studies could examine the hypothesis that the increasing use of precision medicine could change individual and societal

norms by increasing confidence in test results (either because results are more targeted and accurate or because patients and clinicians simply believe that they are). In addition, it is important to explore whether patients and clinicians perceive forgoing treatment as a potential loss (because they are choosing to forgo therapy that, although unlikely to benefit them, could provide benefits). There is extensive literature from the field of behavioral economics that losses are perceived to be much greater than the equivalent gains.[2, 3] Studies that examine when and how patients and clinicians perceive forgoing treatment as a loss or a gain would provide useful insights not only for precision medicine but also for other initiatives to reduce unnecessary care. This information could be used to better understand and - as relevant - shape decision-making through appropriate decision aids that enable patients and clinicians to understand the potential benefits of forgoing treatment.

- The **intersection of patient and clinician beliefs about forgoing treatment based on precision medicine and its impact on behaviors needs further exploration.** Although there is evidence that clinicians consider their patients' preferences in their decisions to use GEP,[6] there is limited evidence on how patient and clinician decision-making interact in practice to either facilitate or hinder the appropriate use of GEP testing and the implications for reducing unnecessary care. Are there characteristics of precision medicine tests that could explain discrepancies in decision-making (e.g., when physicians order GEP even when the patient is not willing to change treatment based on test results) or be used to ameliorate these discrepancies?

There are other examples of clinical areas where precision medicine potentially could facilitate a willingness to forgo treatment. One example is the use of rapid exome sequencing in neonatal intensive care units to diagnose seriously ill newborns. Emerging evidence suggests that such testing enables parents, in the case of terminally ill children, to make the difficult choice for palliative care.[10] A diagnosis thus enables the forgoing of futile care that will not benefit the patient and can cause increased parental distress and unnecessary burdens on the health care system.

In conclusion, precision medicine could potentially facilitate an increased willingness of patients and clinicians to forgo treatment. Although it might be assumed that precision medicine will, by definition, achieve this objective, there needs to be more evidence on whether, why, how, and when this effect might occur. It would also be useful to examine where precision medicine could facilitate forgoing unnecessary care but has failed to do so and the reasons, as well as to examine future opportunities for precision medicine to contribute to reduction of unnecessary care.

The need to have patients and clinicians forgo unnecessary and potentially harmful care is great. Precision medicine is not a “silver bullet” that can solve this challenge but it may offer opportunities to achieve this objective. This potential benefit needs further exploration and analysis.

Acknowledgement

The authors gratefully acknowledge input on conceptualization from Elena Elkin, PhD (Memorial Sloan Kettering Cancer Center) and input and research assistance from Michael Douglas, MS (University of California San Francisco). Dr. Elkin was not compensated for this contribution. Mr. Douglas was compensated as part of his employment. Written permission from each person acknowledged has been obtained.

Funding/Support: This study was funded by grants from the National Human Genome Research Institute (R01 HG007063; U01 HG009599) and from the National Cancer Institutes (R01 CA221870).

Disclosures:

Dr Phillips receives consulting income and received support for attending a conference where this work was discussed from Illumina, Inc. Disclosures have been reviewed by the University of California San Francisco. Dr Marshall received travel support for attending a conference where this work was discussed from Illumina Inc. Dr Kurian has received research funding from Myriad Genetics.

References

1. Choosing Wisely (2019). <https://www.choosingwisely.org/>
2. Phillips KA, Schleifer D, Hagelskamp C. Most Americans Do Not Believe That There Is An Association Between Health Care Prices And Quality Of Care. *Health Aff (Millwood)* 35(4), 647–653 (2016). [PubMed: 27044965]
3. O'Brien BJ, Gertsen K, Willan AR, Faulkner LA. Is there a kink in consumers' threshold value for cost-effectiveness in health care? *Health Econ* 11(2), 175–180 (2002). [PubMed: 11921315]
4. Garfield S, Douglas MP, Macdonald KV, Marshall DA, Phillips KA. Consumer familiarity, perspectives and expected value of personalized medicine with a focus on applications in oncology. *Per Med* 12(1), 13–22 (2015). [PubMed: 25620993]
5. Sparano JA, Gray RJ, Makower DF et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med* 379(2), 111–121 (2018). [PubMed: 29860917]
6. Kurian AW, Bondarenko I, Jagsi R et al. Recent Trends in Chemotherapy Use and Oncologists' Treatment Recommendations for Early-Stage Breast Cancer. *J Natl Cancer Inst* 110(5), 493–500 (2018). [PubMed: 29237009]
7. Marshall DA, Deal K, Bombard Y, Leighl N, Macdonald KV, Trudeau M. How do women trade-off benefits and risks in chemotherapy treatment decisions based on gene expression profiling for early-stage breast cancer? A discrete choice experiment. *BMJ Open* 6(6), e010981 (2016).
8. Bombard Y, Rozmovits L, Trudeau M, Leighl NB, Deal K, Marshall DA. Access to personalized medicine: factors influencing the use and value of gene expression profiling in breast cancer treatment. *Curr Oncol* 21(3), e426–433 (2014). [PubMed: 24940102]
9. Tsai M, Lo S, Audeh W et al. Association of 70-Gene Signature Assay Findings With Physicians' Treatment Guidance for Patients With Early Breast Cancer Classified as Intermediate Risk by the 21-Gene Assay. *JAMA Oncol* 4(1), e173470 (2018). [PubMed: 29075751]
10. Meng L, Pammi M, Saronwala A et al. Use of Exome Sequencing for Infants in Intensive Care Units: Ascertainment of Severe Single-Gene Disorders and Effect on Medical Management. *JAMA Pediatr* 171(12), e173438 (2017). [PubMed: 28973083]