Published in final edited form as:

Biosens Bioelectron. 2019 February 01; 126: 679–689. doi:10.1016/j.bios.2018.11.015.

“Emerging Neurotechnology for Antinociceptive Mechanisms and Therapeutics Discovery”

Bryan J. Black, Ph.D.¹, Rahul Atmaramani¹, Sarah Plagens¹, Zachary T. Campbell², Gregory Dussor³, Theodore J. Price³, Joseph J. Pancrazio, Ph.D.¹

¹Department of Bioengineering, University of Texas at Dallas, 800 W. Campbell Road, Richardson, TX, 75080, USA

²Department of Biological Sciences, The University of Texas at Dallas, 800 W. Campbell Road, Richardson, TX 75080, USA.

³School of Behavioral and Brain Sciences, The University of Texas at Dallas, 800 W. Campbell Road, Richardson, TX 75080, USA.

Abstract

The tolerance, abuse, and potential exacerbation associated with classical chronic pain medications such as opioids creates a need for alternative therapeutics. Phenotypic screening provides a complementary approach to traditional target-based drug discovery. Profiling cellular phenotypes enables quantification of physiologically relevant traits central to a disease pathology without prior identification of a specific drug target. For complex disorders such as chronic pain, which likely involves many molecular targets, this approach may identify novel treatments. Sensory neurons, termed nociceptors, are derived from dorsal root ganglia (DRG) and can undergo changes in membrane excitability during chronic pain. In this review, we describe phenotypic screening paradigms that make use of nociceptor electrophysiology. The purpose of this paper is to review the bioelectrical behavior of DRG neurons, signaling complexity in sensory neurons, various sensory neuron models, assays for bioelectrical behavior, and emerging efforts to leverage microfabrication and microfluidics for assay development. We discuss limitations and advantages of these various approaches and offer perspectives on opportunities for future development.

Keywords

Neurotechnology; ion imaging; neural cell biosensors; neuropathic pain; sensory neurons; hiPSC

*Corresponding author: Bryan J. Black, Ph.D., Dept. of Bioengineering, The University of Texas at Dallas, 800 W. Campbell Road, BSB 13.605, Richardson, TX 75080. Phone: 972-883-2138, Bryan.Black@utdallas.edu.

Authorship statement:

BJB and JJP organized and interpreted the core ideas and contributed the main text of the manuscript. TJP and GD contributed concepts of high content screening through electrophysiological endpoints. ZTC, TJP, and GD contributed concepts on electrophysiology measures for study of molecular mechanisms underlying pain. BJB, RA, and SP collected and analyzed data and created the manuscript figures.

Conflict of Interest Statement: None.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. INTRODUCTION

Chronic pain is the most common form of long-term disability in the world, affecting 100 million people in the United States alone (Larner, 2014). The situation becomes dire for aged populations, with more than 60% of elderly housed in U.S. care facilities suffering from chronic pain (Miu and Chan, 2014). Pharmacological treatment options for chronic pain are limited and problematic. The most prevalent approach involves prescription opioids, which are associated with a high incidence of adverse effects including long-term tolerance and abuse (Ballantyne and LaForge, 2007; Chang et al., 2015). As a result, novel therapeutics are urgently needed that offer improved efficacy and reduced side-effects.

Contemporary drug discovery strategies rely on the identification of molecular targets (i.e., druggable targets) associated with a particular pathway and subsequently probe the role of a given gene product in disease progression. Unfortunately, the yield for this type of target-directed non-opioid analgesic drug discovery has been low. Subsequent to the identification of a relevant bioactive compound, it is common to use of animal models (Munro et al., 2017) for further compound screening. As an alternative or supplemental strategy, there is increasing recognition that phenotypic screens may serve as means for discovering drugs which ameliorate chronic pain (Pruss, 2010; Zhang et al., 2014). Rather than focusing on the activity of a molecular target, modern phenotypic screens capture complex cellular level behaviors or observable traits that are physiologically relevant and central to a pathology without relying on the identification of a specific drug target or a corresponding hypothesis concerning its pathological role. Phenotypic screens are not a panacea for drug discovery. However, this approach can generate entirely novel treatments for disorders which have either complex or unknown mechanisms (Moffat et al., 2017).

To serve as a basis of a phenotypic screen, the selection of cell or tissue type should be based on pathological relevance. In addition, the assay or screening method must be capable of quantifying signals associated with pathological outcomes or cellular behavior. Sensory neurons derived from dorsal root ganglia (DRG) undergo changes in membrane excitability during chronic pain conditions and are integral to the development of chronic pain (Waxman et al., 1999; Zhang et al., 1999). Therefore, they are an excellent candidate cell type for a phenotypic screen to identify new pain therapeutics. Furthermore, it has been suggested that therapeutic strategies which target peripheral DRG neurons rather than tissues in the central nervous system may be “better and safer” (Berta et al., 2017) as peripheral input is ultimately necessary for the maintenance of neuropathic pain states (Haroutounian et al., 2014) and such strategies may not modulate reward circuitry in the brain. Here, we review the current state-of-the-art in the design and development of *in vitro* sensory neuron-based phenotypic assays for chronic peripheral neuronal hyperexcitability. Emergent technologies are described that are enabling for more advanced, higher throughput, and ultimately more physiologically relevant, phenotypic screening paradigms.

2. DISCUSSION

2.1 Bioelectrical behavior of DRG neurons

DRG are clusters of sensory neurons, each projecting a single bifurcated axon toward both the dorsal spinal cord as well as peripheral targets (skin, muscle, viscera) (Fig. 1). In the periphery, axon terminals detect both noxious and innocuous thermal, chemical, and mechanical stimuli, and transduce these signals in the form of all-or-nothing action potentials toward the spinal cord (Wall and Devor, 1981). The type and threshold of stimulus necessary to initiate afferent signals, as well as the speed and fidelity with which these signals are processed, depends on the type of sensory neuron. Small-diameter neurons with unmyelinated axons have relatively high stimulus thresholds and are called nociceptors (or C-fibers). Small to medium-diameter neurons with thinly myelinated axons, lower thresholds and higher conduction velocity are also referred to as nociceptors, but specifically as A δ -fibers. The largest-diameter sensory neurons with relatively thick myelin are low threshold and called proprioceptors since they are largely responsible for afferent feedback from muscles for position, movement, and reflex (Basbaum and Woolf, 1999). Under normal conditions, nociceptive DRG neurons exhibit relatively low levels of spontaneous spike activity (Weng et al., 2012). However, inflammation due to disease or injury may lead to increased intrinsic activity and increased sensitivity to external stimuli (Davidson et al., 2014; Djouhri, 2006; White et al., 2005). This sensitization of peripheral nociceptors significantly contributes to the manifestation of chronic pain (Krames, 2015).

2.2 Signaling complexity in sensory neurons

Whether *in vitro* or *in vivo*, sensory neuron excitability is ultimately mediated by ion channel expression and function. These include multiple classes of sodium channels, voltage-gated and two-pore potassium channels, hyperpolarization-activated cyclic nucleotide-gated channels, voltage-gated calcium channels, ligand-mediated cation channels, and gap junctions (Eduardo E Benarroch, 2015; Kim et al., 2016; Ratté and Prescott, 2016; Waxman and Zamponi, 2014). All of these independently serve as potential pharmacological targets for pain (Moldovan et al., 2013). However, the means by which ion channels may be regulated, either directly or indirectly, also involves a multitude of molecular signaling pathways (Price et al., 2016) as well as multiple cell types (Costa and Neto, 2015). Fig. 1 illustrates only some of the pathways by which ion channel expression and function are known to be regulated in an inflammatory or neuropathic pain state. These include (1) direct activation of ligand-gated ion channels by inflammatory cytokines, neurotrophins, ATP, etc., which leads to increased depolarization and activates kinase pathways. (2) Direct phosphorylation of voltage-gated sodium channel (VGSC) or transient receptor potential (TRP) ion channels by cytokine-activated kinases, including p38 MAPK (Hudmon et al., 2008), PKC (Vellani et al., 2001), cAMP-dependent PKA (Bhave et al., 2002; Fitzgerald et al., 1999), and c-Src kinases (Jin et al., 2004). (3) Kinase-activated transcription regulation via repression/activation of transcription mediators such as members of the cAMP response element binding protein (CREB) family (Cheng and Penninger, 2002) as well as activity-dependent epigenetic regulation of transcription via increased/decreased DNA methylation and histone acetylation (E. E. Benarroch, 2015). (4) Cytokine-mediated activation of axonal and/or somal pre-translational pathways, including mTOR (Kang et al., 2009; Ohannes K

Melemedjian et al., 2010; Moy et al., 2017; Obara et al., 2012; Obara and Hunt, 2014) and phosphorylation of the eIF4F complex (Barragán-Iglesias et al., 2018; Khoutorsky and Price, 2017), resulting in increased cap-dependent ion channel protein synthesis. (5) Translational repression/activation via CPEB-mediated increased/decreased polyadenylation (Richter, 2007). (6) Maintenance of the inflammatory state via increased neuronal secretion of neuropeptides, which can create an inflammatory positive-feedback loop via intraganglionic paracrine signaling between DRG soma and adjacent satellite glial cells (Huang et al., 2013; Takeda et al., 2009).

Due to this complexity, the challenge for therapeutic interventions for chronic pain is to effectively reverse a rise in excitability when there are so many distinct ion channels or molecular targets that may be sufficient to trigger underlying pathology - a situation (Ratté and Prescott, 2016) have referred to as “degeneracy”. The molecular target complexity for chronic pain suggests that assays wherein measured outcomes are more integrative and based on cellular behavior represented by phenotypic assays may have an emerging role for therapeutic discovery.

Importantly, cytokine-induced changes in sensory neuron excitability and protein synthesis appear to be conserved *in vitro*. For example, cultured adult DRG and trigeminal ganglion neurons treated with IL-6 exhibit dose-dependent increases in the phosphorylation of extracellular signal-regulated kinase (ERK), an upstream signaling protein involved in nascent protein synthesis via MnK1 and eukaryotic initiation factor activation (Ohannes K. Melemedjian et al., 2010; Tillu et al., 2012; Yan et al., 2012). Moreover, combinations of IL-6 and NGF have been shown to rapidly increase global nascent protein synthesis *in vitro* (Ohannes K. Melemedjian et al., 2010; Moy et al., 2017), and our group has recently demonstrated that adult mouse DRG neurons achieve and maintain a sensitized phenotype following IL-6 incubation when cultured on MEAs (Black et al., 2018).

2.3 Sensory neuron models

To date, the gold standard tissue model for both *in vitro* and *in vivo* pathological pain studies has been rodents. While studies with rodent sources of primary sensory neurons have greatly informed the understanding of molecular mechanisms underlying chronic pain, there are electrophysiological differences between species (Davidson et al., 2014) which may limit the interpretation of phenotypic screening results. For example, there are differences in the pharmacological sensitivity of GABA channels between human and rodent species in DRG neurons (Sheahan et al., 2018; Valeyev et al., 1996). In addition, human and rodent DRG exhibit differential expression levels of various voltage-gated sodium channels, which also exhibit different activation/inactivation kinetics (Chang et al., 2018; Dib-Hajj et al., 1999). However, even with access to primary human specimens through organizations such as the National Disease Research Interchange, the prospect of operating high-content, high-throughput screening with donated human tissue would be very challenging given the low anticipated cell yield (Valtcheva et al., 2016). Alternative sources of human tissue include human embryonic stem cells (hESCs) (Boisvert et al., 2015) and human induced pluripotent stem cells (hiPSCs) (Wainger et al., 2014) that can be guided to differentiate into sensory-like neurons. These methods potentially enable virtually unlimited sources of non-tumor-

derived human tissue, several of which are now commercially available. However, a recent large-scale study of common genetic effects in a neuronal cell type differentiated from hiPSC-derived sensory neurons was carried out by (Schwartzentruber et al., 2018). This study concluded that hiPSC-derived sensory neurons are transcriptionally distinct from human DRG, implying that current *in vitro* differentiation protocols may not be functionally or transcriptionally mature. With that in mind, we will discuss various phenotypic and functional assays which have utilized these tissue models. Our hope is that future protocols for hiPSC-derived sensory neuron differentiation and culture may yield physiologically relevant pathological pain models, and that methodologies developed using primary tissues and immortalized cell lines may be successfully extended to incorporate these tissues.

2.4 Phenotypic assays based on sensory neurons

2.4.1 Calcium and voltage imaging—Imaging intracellular ion concentrations enables either direct or indirect measures of cellular excitability from dense populations with high temporal and single cell resolution (Hempel et al., 2017). By far, the most widely used method of optically interrogating single cell or network activity is calcium imaging. The dye chemistries are well established, imaging is relatively straightforward, and calcium is broadly appreciated as integral to cellular signaling and subsequent excitability. Imaging based on either calcium-sensitive fluorescent dyes or genetically-encoded calcium indicators (GECIs) function on the basis of calcium chelation via direct ion or calmodulin protein binding (Grienberger and Konnerth, 2012). Direct or indirect calcium chelation initiates the associated fluorophore's conformational state change to either enhance the quantum yield of the fluorophore (increase/decrease brightness) or shifts its emission spectrum for quantitative ratiometric measurements. In the context of sensory neuron excitability, calcium imaging using commercial dyes has been shown compatible with primary rodent DRG (Von Banchet et al., 2005), hiPSC sensory neurons (Wainger et al., 2014), immortalized cell lines expressing relevant ion channels (De La Torre-Martínez et al., 2017), and even primary human DRG *in vitro* (Valtcheva et al., 2016) to study phases of Wallerian degeneration subsequent to injury (Vargas et al., 2015), anti-hyperalgesic drug mechanisms (Sutton et al., 2002), and functional expression of pain-relevant ion channels (Fouillet et al., 2017), among other endpoints. Recently, demonstrations of increased imaging throughput have been carried to characterize cold- and heat-sensing ion channel (TRPM8 and TRPV1) responsiveness using immortalized cell lines in a 96-well format (De La Torre-Martínez et al., 2017). Additionally, (Fouillet et al., 2017) have developed a high-throughput method based on calcium dynamics to optically interrogate Nav1.7 channel function using primary rat DRG neurons.

However, calcium imaging has limited applications due to some inherent disadvantages. (1) Calcium activity is an indirect measure of neural activity due to calcium's slow kinetics relative to cell membrane potential, and correlating fluorescent signals to action potential *in vivo* or *in vitro* patterns remains a significant challenge. (2) Since calcium is an important secondary messenger which mediates numerous intracellular processes, including maintenance of transmembrane potential, persistent synthetic calcium chelation may alter cell signaling and viability (Smith et al., 2018). Additionally, studies of Cre-based transgenic mouse models expressing GCaMP6 report aberrant cortical network activity tantamount to

seizure (Steinmetz et al., 2017) which may be due to Cre toxicity or genetic background. (3) Long-term optical monitoring of cellular excitability at visible/near-infrared wavelengths may lead to phototoxicity (Denk et al., 1994) or stimulation of photobiological processes.

Dyes associated with other intracellular ion concentrations (e.g., sodium and potassium) are also available, as are indicators of transmembrane potential changes (i.e., voltage indicator dyes). However, current sodium and potassium indicators suffer from lack of binding specificity as compared to calcium indicators (Meuwis et al., 1995; Minta and Tsien, 1989). Voltage imaging also faces inherent challenges associated with the short duration of action potentials, requiring high frame rate acquisitions from expensive detectors and leading to relatively low fluorescence photon yields (i.e., low signal-to-noise ratios) (Kulkarni and Miller, 2017). Still, moderate throughput voltage imaging experiments have been carried out in HEK-293 cells expressing human Nav1.7 alpha subunits in 96 well fluorescent imaging plate reader (FLIPR) formats using voltage dyes to screen for VGSC antagonists (Zhao et al., 2016). However, FLIPR blue dye assays do not enable sodium response times on the order of single action potentials (1–10 ms), instead offering changes in fluorescence on the order of tens of seconds. Surrogate measures of specific sodium channels have been proposed to overcome these challenges, such as thallium flux-based functional assays for the sodium channel Nav1.7 (Du et al., 2015). While the fluorescence yield and dynamics of these surrogate measures are more amenable to high-throughput screening, they are still an indirect measure of membrane excitability.

More recently, label-free methods of optically detecting single action potentials based on minute changes cell membrane deformations (electromotility) via interferometry (Batabyal et al., 2017) or changes in near-membrane protein methyl resonance peaks via vibrational spectroscopic imaging have also been reported (Lee et al., 2017). These methods offer the advantages of reduced phototoxicity, no risk of photobleaching, and high temporal resolution as compared to calcium imaging. While these outcomes are highly promising, especially for *in vitro* applications, these methods are still in their infancy and the cellular mechanisms that influence these measurements need to be further investigated. For example, in the case of vibrational spectroscopic imaging, the contributions of molecular reorientation and relative protein densities during electromotility are unknown. These methods also potentially suffer from similar inherent noise and acquisition rate challenges as label-based voltage imaging.

2.4.2 Patch-clamp electrophysiology—In terms of high-content measurements of cellular excitability and interrogation of individual membrane channel properties, manual patch-clamp (MPC) electrophysiology remains the gold standard. However, MPC is highly limited in regards to assay development due to its relative invasiveness, low-throughput, and technical difficulty. To overcome the challenges of throughput and technicality, automated patch clamp (APC) technologies have been developed. For an excellent review on 1st and 2nd generation automated patch clamp technologies for pain-related ion channel research, see (Bell and Dallas, 2017). Briefly, the most widely adopted and successful APC format involves the application of cell suspensions to planar recording chips, where a combination of gravity and negative pressure draw cells into contact with micropores (or apertures) centered on recording sites embedded within the planar chip. Once in contact, negative

pressure is maintained until the system reports a threshold impedance necessary for potential 'clamp'. To date, these formats have largely been used by pharmaceutical companies to screen for human ether-a-gogo-related gene (hERG) induced cardiac ion channel blockade (Möller and Witchel, 2011). hERG channel (Kv11.1) blockade can lead to cardiac arrhythmia, and hERG channel block has become a frequent cause for drug discovery program delays or withdrawals from clinical trial (Danker and Möller, 2014). In the same way, APC is now being used to screen for antagonists of Nav1.7, an ion channel highly implicated in nociception and chronic pain development. One such technology (SynchroPatch 768PE) successfully screened 10,000 Nav1.7 reference compounds against heterologous cell lines expressing human Nav1.7 variants with a throughput of approximately 6000 data points per day with a Z' factor of 0.72 (Li et al., 2017); highlighting the potential of APC formats for high-throughput screening targeting pain-relevant ion channels, if not nociceptor excitability.

2.4.3 Substrate-integrated microelectrode arrays—Similar to patch-clamp electrophysiology, substrate-integrated microelectrode arrays (MEAs) may be used to resolve and record individual action potentials, or spikes (Gross et al., 1985; Pancrazio et al., 1999). However, these extracellular action potentials are recorded noninvasively, enabling long-term and/or repeated monitoring of large cell populations. For excellent reviews of fundamental principles and applications of extracellular microelectrode arrays, see (Heinricher, 2004) and (Spira and Hai, 2013). Briefly, MEAs function on the principles of volume conductor theory, where regions of cell membrane serve as a current sink, or sources, both embedded within a conductive medium. As an action potential propagates, so does the region of membrane depolarization, with sink becoming source (and vice versa). Relative to stationary recording and reference electrodes, this is manifest as time-dependent potential increases/decreases, giving rise to characteristic extracellular waveforms with temporal resolution approaching or equal to patch-clamp recordings (1–10 ms). By this means, MEAs enable high-content quantitative measures of cellular excitability, such as spike rate, amplitude, bursting rates, and neuron firing synchronization (Gross et al., 1995; Selinger et al., 2004). The quality of these recordings, in terms of signal-to-noise ratio, is largely dependent on the impedance of the recording electrodes and the quality of the cell coupling. A large number of substrate materials have been successfully used to fabricate MEAs and facilitate cell adhesion, including silicon dioxide (Mastrototaro et al., 1992), silicon nitride (Connolly et al., 1990), SU-8 (Y. Zhang et al., 2010), polystyrene (Hammack et al., 2018), parylene (Charkhkar et al., 2016, 2012; Omaki et al., 2017), liquid crystal elastomer (Rihani et al., 2018), and others. Reduction in electrode impedance is dependent on several factors, but recent advances in electrodeposition, metallic 3D printing, and electrospinning have enabled nanofabrication of electrode sites; potentially enhancing long-term cell-to-electrode coupling and effectively reducing electrode impedance, resulting in increased signal-to-noise ratios (Heim et al., 2012; Jao et al., 2015; Krishnamoorthy and Zoski, 2005; Santoro et al., 2014; Weidlich et al., 2017).

A large body of previous work has demonstrated that primary neurons, particularly those that are embryonically-derived, can be reliably cultured on MEAs, and are a valuable tool for neurotoxicology (Johnstone et al., 2010) and neuropharmacology (Accardi et al., 2016;

Kulagina et al., 2004; Novellino et al., 2011; Parenti et al., 2013). Commercially available systems from Axion Biosystems, ALA Scientific, and Multichannel Systems utilize multi-well MEA plates for moderate throughput, high content screening based on extracellular action potentials (Fig. 2). These systems have been used to assess the neural effects of putative toxicants (Valdivia et al., 2014), illicit drugs (Hondebrink et al., 2016), marine neurotoxins (Nicolas et al., 2014), insecticides (Dingemans et al., 2016), as well as patient-specific genetic predispositions for epilepsy and related disorders (Tidball and Parent, 2015; Vessoni et al., 2016).

Extension of single- and multi-well MEA technology to explorations of pain-related mechanisms and therapeutics has begun. (Pearce et al., 2005) demonstrated an integrated MEA-microfluidic system for creating a precise ‘temperature clamp’ on mouse DRG neurons seeded on an MEA and examined cold sensitivity. (Yang et al., 2016) recapitulated functional effects of a sodium channel mutation associated with inherited erythromelalgia using adult rat DRG neurons on multi-well MEA plates. Cells expressing the gain-of-function mutation showed an increased level of spontaneous firing in response to temperature elevations, consistent with the warmth-induced manifestation of pain. However, the fraction of microelectrode sites showing any discernible spike activity for the cultured wild-type DRG, even at elevated temperatures, was less than 2%, a level far too sparse to support high-throughput screening. By relying on embryonically-derived tissue, Newberry et al. (2016) demonstrated cultured DRG neurons on multi-well MEAs where activity could be detected on 53% of the microelectrode sites, an observation we have largely confirmed in our laboratory. Furthermore, spontaneous spike activity could be modulated by exposure to capsaicin, a ligand for the transient receptor potential cation channel subfamily V1 (TRPV1), bradykinin, and inhibitors of voltage gated sodium and calcium channels (Newberry et al., 2016). While embryonic tissue is amenable to *in vitro* culture due to increased viability (Eide and McMurray, 2005), there are significant disadvantages due to differences in gene expression and related function as compared to later developmental stages. These include changes linked to the voltage-gated Na channel Nav 1.8/1.9 (Benn et al., 2001), mRNA associated with axonal transport, vesicle trafficking, and axonal protein synthesis (Gumy et al., 2011). Additionally, dissociated embryonic and neonatal DRG cultures show little or no apparent NGF-induced sensitization to capsaicin (Zhu, W.; Oxford, 2011), unlike adult DRG *in vivo* and *in vitro*. This raises the possible concern that embryonic DRG neurons are insensitive to cytokine mediated sensitization, which would be in stark contrast to the outcomes of studies using adult DRG neurons for patch-clamp (Von Banchet et al., 2005) and MEAs (Moy et al., 2017). With optimization of adult DRG culture conditions, it may be possible to greatly enhance the yield of active microelectrodes by a factor of 10–20 to enable cell-based assay use. Recent work in our lab indicates that higher yield of spontaneous activity may be associated with maintaining a relatively high density of non-neuronal cell types (satellite glia, Schwann cells, and fibroblasts).

Importantly, the multiwell MEA platform does not inherently exclude the use of primary human DRG or hiPSC sensory neuron models. Proof-of-concept extracellular recordings from custom MEA devices using human DRG have been demonstrated (Enright et al., 2016), as has the use of hiPSC derived sensory neurons (Odawara et al., 2017; Wainger et al., 2014), suggesting that MEAs allow maximal physiological relevance in a moderate

throughput assay platform. Our group has used commercially available hiPSC derived sensory neuron progenitors (Axol Bioscience) to monitor spontaneous and evoked activity in a multiwell MEA format. While we observed spontaneous baseline activity with excellent signal-to-noise ratios as well as electrically, chemically, and thermally evoked activity, we did not observe any culture sensitivity to human-recombinant inflammatory cytokines IL-6 or TNF- α (Fig. 3). This suggests that either culture conditions or differentiation protocols need to be further optimized to lead to cytokine-sensitive hiPSC sensory models and/or genetically modified models for studies of chronic pain mechanisms and pharmacology.

2.4.4 Compartmentalized cultures and higher-order co-cultures—As previously discussed, each DRG sensory neuron extends a single bifurcated axon toward both the dorsal spinal cord and the periphery. In the periphery, nerve terminals form intimate contact with different tissues and have differing electrophysiological and pharmacological profiles than the soma. Perhaps most importantly, it is the peripheral nerve terminals - not the cell soma - that will be directly and immediately exposed to the previously discussed ‘inflammatory soup’ following tissue injury. It is this exposure that initiates the inflammatory cascade which can lead to chronic pain conditions. Therefore, there is considerable interest in monitoring excitability changes and pharmacological profiles in physically and/or fluidically isolated soma and axons, as shown in Fig. 4. Various methods have been described to physically compartmentalize sensory neuron soma and axons, including PDMS microfluidic channels (Dworak and Wheeler, 2009; Park et al., 2006), Campenot chambers (Campenot, 1977; Pazyra-Murphy and Segal, 2008), and porous filters (Unsain et al., 2014). To date, a majority of studies using sensory neurons in compartmentalized cultures have focused on monitoring rates and extent of neurite outgrowth and myelination (Gornstein and Schwartz, 2017; Ravula et al., 2007; Wheeler et al., 2014; Yang et al., 2012), de/regeneration following axotomy (Kilinc et al., 2011; Unsain et al., 2014), and anterograde/retrograde transport or signaling (Cohen et al., 2011; Gluska et al., 2015; K. Zhang et al., 2010; Zweifel et al., 2005), among others (Gumy et al., 2011).

The use of compartmentalized cultures has been extended to studies of sensory neuron excitability in only a few cases, wherein somatic or axonal activity were monitored independently via calcium imaging (Jonas et al., 2015) and/or MPC electrophysiology (Tsantoulas et al., 2013). However, standard or commercial microfluidic devices are not designed to accommodate MPC electrodes. Moreover, MPC on single axons remains a high technical barrier. The integration of single or multi-well MEAs with PDMS microchannel devices may prove more amenable to simultaneous stimulation and recording applications. While PDMS-based microfluidics have been used previously with MEAs for drug (Enright et al., 2016) or temperature delivery (Pearce et al., 2005), to date, no study has demonstrated compartmentalized cultures on MEAs.

In addition to isolating neuronal soma from axons, compartmentalized culture platforms may also be used to isolate multiple patho-physiologically relevant cell types. For example, previous studies have incorporated DRG neurons and keratinocytes (Griscom et al., 2005; Kumamoto et al., 2014; Roggenkamp et al., 2012) or osteoblasts (Neto et al., 2014; Silva et al., 2017) in separate compartments of PDMS microsystems or Campenot chambers to mimic skin or bone innervation and to study interactions between neurons and the cell types

native to the innervated tissue. Additionally, sensory communication between DRG neurons and dorsal horn neurons has been monitored via calcium imaging in an open well format (Ohshiro et al., 2007), but not in a compartmentalized culture to date.

While compartmentalized cultures may be advantageous for separating cell types and components, careful consideration needs to be given to the microchannel design. It has been noted that standard microchannel device designs create high-shear fluid flows during seeding or medium exchange. Since relatively low fluid flow rates have been observed to drive the directionality of neurite outgrowth (Gu et al., 2014), there is concern that similar designs will suffer from microchannel crossing failure due to directionality or ‘bundling’ (van de Wijdeven et al., 2018). Additionally, both microchannel width and length need to be optimized to accommodate axonal outgrowth rates (Tsantoulas et al., 2013), mitigate the non-neuronal cell migration (Park et al., 2006), and control passive diffusion of growth factors or other pharmacological treatments. Some of these potential issues have driven the design of novel ‘open-well seeding’ platforms for generating multi-nodal compartmentalized cultures and/or maintaining custody of cells and biochemical agents (Park et al., 2009; van de Wijdeven et al., 2018) without generating high-shear stresses which may limit application or potentially modulate function.

3. PERSPECTIVE

Modern cell-based pharmacological screening strategies monitor changes in cellular phenotype as an integrative measure of a drug’s activity. Ideally, the phenotypic endpoint is highly relevant to the indication of interest and provides some insight into the underlying mechanistic pathways. However, the successful identification of a ‘hit’ ideally would not require full mechanistic understanding of the drug-tissue interaction. An important primary phenotype of neural cells is excitability. The first-order neuronal cell type involved in the transduction and transmission of peripherally-mediated pain are sensory neurons of the DRG. Therefore, future cell-based phenotypic screening paradigms targeting the identification of peripheral analgesics should enable direct measurement of DRG sensory neuron excitability. Strategies should be developed to maximize content (cellular as well as temporal resolution and intercellular communication), throughput, and ease-of-use while minimizing invasiveness. Furthermore, the mechanistic complexity of chronic pain demands a cellular model capable of recapitulating responsiveness and/or sensitization due to compounds known to be involved in human chronic pain conditions. This has driven the ongoing development of hiPSC sensory neuron models, which provide the opportunity for maximal human relevancy while also being scalable for high-throughput platforms. However, to date, no published study has demonstrated functional sensitivity of hiPSC sensory neurons to inflammatory cytokines such as IL-6, IL-1 β , TNF α , or NGF. Therefore, more work needs to be done related to expression of sensitization pathways in hiPSC-based models as well as standardization of differentiation protocols (Schwartzentruber et al., 2018). Along these lines, it may be worthwhile to investigate phenotypic contributions of either primary or hiPSC-derived non-neuronal cell types (satellite glia, Schwann cells) in heterogeneous culture models, as it has been demonstrated in other hiPSC neuron models that the presence of support cells (glia, oligodendrocytes) facilitates synaptogenesis, phenotypically relevant mRNA expression, and functional maturation (Ishii et al., 2017;

Kayama et al., 2018; Odawara et al., 2014; Schutte et al., 2018; Tang et al., 2013). In the case of chronic pain development, satellite glia and Schwann cells have been shown to facilitate sensitization and intercellular signaling via gap junctions and transmitter release (e.g., ATP) (Campana, 2007; De Logu et al., 2017; Hartlehnert et al., 2017; Huang et al., 2013; Kodama et al., 2017; Poplawski et al., 2018; Takeda et al., 2009). Therefore, developing combinatorial imaging-electrophysiological approaches may add highly relevant content without diminishing density or throughput. This may be achievable using field effect transistor-based MEAs, which have been fabricated from silicon (Fromherz et al., 1991), graphene (Blaschke et al., 2016), and organic materials (Benfenati et al., 2013) among others. In the case of graphene and organics, this offers transparent substrates and electrodes in combination with high electrode density. Furthermore, the feasibility of both high-density MEAs (Musick et al., 2009) and functional imaging (Pasca et al., 2015) has already been demonstrated using 3D culture platforms. While it is the authors' opinion that not all pharmacological screening outcomes demand the reported advantages of 3D culture, it seems to be the natural convergence for advances in both functional recording technology and tissue engineering.

Acknowledgments

Source(s) of financial support:

This work was supported by the University of Texas at Dallas. NS065926 (TJP), NS102161 (TJP), NS072204 (GD), and NS100788 (ZTC).

References

- Accardi MV, Pugsley MK, Forster R, Troncy E, Huang H, Authier S, 2016 The emerging role of in vitro electrophysiological methods in CNS safety pharmacology. *J. Pharmacol. Toxicol. Methods* 81, 47–59. 10.1016/j.vascn.2016.03.008 [PubMed: 27058269]
- Ballantyne JC, LaForge KS, 2007 Opioid dependence and addiction during opioid treatment of chronic pain. *Pain* 129, 235–255. 10.1016/j.pain.2007.03.028 [PubMed: 17482363]
- Barragán-Iglesias P, Lou TF, Bhat VD, Megat S, Burton MD, Price TJ, Campbell ZT, 2018 Inhibition of Poly(A)-binding protein with a synthetic RNA mimic reduces pain sensitization in mice. *Nat. Commun* 9 10.1038/s41467-017-02449-5
- Basbaum a I., Woolf CJ, 1999 Pain. *Curr. Biol* 9, R429–R431. [https://doi.org/S0960-9822\(99\)80273-5](https://doi.org/S0960-9822(99)80273-5) [pii] [PubMed: 10375535]
- Batabyal S, Satpathy S, Bui L, Kim Y-T, Mohanty S, Bachoo R, Davé DP, 2017 Label-free optical detection of action potential in mammalian neurons. *Biomed. Opt. Express* 8, 3700 10.1364/BOE.8.003700 [PubMed: 28856044]
- Bell DC, Dallas ML, 2017 Using automated patch clamp electrophysiology platforms in pain-related ion channel research: Insights from industry and academia. *Br. J. Pharmacol* 1–10. 10.1111/bph.13916
- Benarroch EE, 2015 Ion channels in nociceptors. *Neurology* 84, 1153–1164. 10.1212/WNL.0000000000001382 [PubMed: 25713000]
- Benarroch EE, 2015 Brain-derived neurotrophic factor: Regulation, effects, and potential clinical relevance. *Neurology* 84, 1693–1704. 10.1212/WNL.0000000000001507 [PubMed: 25817841]
- Benfenati V, Toffanin S, Bonetti S, Turatti G, Pistone A, Chiappalone M, Sagnella A, Stefani A, Generali G, Ruani G, Saguatti D, Zamboni R, Muccini M, 2013 A transparent organic transistor structure for bidirectional stimulation and recording of primary neurons. *Nat. Mater* 12, 672–80. 10.1038/nmat3630 [PubMed: 23644524]

- Benn SC, Costigan M, Tate S, Fitzgerald M, Woolf CJ, 2001 Developmental Expression of the TTX-Resistant Voltage-Gated Sodium Channels Nav 1.8 (SNS) and Nav1.9 (SNS2) in Primary Sensory Neurons. *J. Neurosci* 21, 6077–6085. <https://doi.org/21/16/6077> [pii] [PubMed: 11487631]
- Berta T, Qadri Y, Tan P-H, Ji R-R, 2017 Targeting dorsal root ganglia and primary sensory neurons for the treatment of chronic pain. *Expert Opin. Ther. Targets* 21, 695–703. 10.1080/14728222.2017.1328057 [PubMed: 28480765]
- Bhave G, Zhu W, Wang H, Brasier DJ, Oxford GS, Gereau IV RW, 2002 cAMP-dependent protein kinase regulates desensitization of the capsaicin receptor (VR1) by direct phosphorylation. *Neuron* 35, 721–731. 10.1016/S0896-6273(02)00802-4 [PubMed: 12194871]
- Black BJ, Atmaramani R, Kumaraju R, Plagens S, Romero-Ortega M, Dussor G, Price TJ, Campbell ZT, Pancrazio JJ, 2018 Adult Mouse Sensory Neurons on Microelectrode Arrays Exhibit Increased Spontaneous and Stimulus-Evoked Activity in the Presence of Interleukin-6. *J. Neurophysiol* jn.00158.2018. 10.1152/jn.00158.2018
- Blaschke BM, Lottner M, Drieschner S, Calia AB, Stoiber K, Rousseau L, Lissourges G, Garrido JA, 2016 Flexible graphene transistors for recording cell action potentials. *2D Mater.* 3 10.1088/2053-1583/3/2/025007
- Boisvert EM, Engle SJ, Hallowell SE, Liu P, Wang ZW, Li XJ, 2015 The Specification and Maturation of Nociceptive Neurons from Human Embryonic Stem Cells. *Sci. Rep* 5, 1–12. 10.1038/srep16821
- Campana WM, 2007 Schwann Cells: Activated Peripheral Glia and Their Role in Neuropathic Pain. *Brain. Behav. Immun* 21, 522–527. 10.1007/s10439-011-0452-9.Engineering [PubMed: 17321718]
- Campenot RB, 1977 Local control of neurite development by nerve growth factor. *Proc. Natl. Acad. Sci. U. S. A* 74, 4516–9. 10.1073/pnas.74.10.4516 [PubMed: 270699]
- Chang DS, Raghavan R, Christiansen S, Cohen SP, 2015 Emerging targets in treating pain. *Curr. Opin. Anaesthesiol* 28, 379–397. 10.1097/ACO.0000000000000216 [PubMed: 26087270]
- Chang W, Berta T, Kim YH, Lee S, Lee SY, Ji RR, 2018 Expression and Role of Voltage-Gated Sodium Channels in Human Dorsal Root Ganglion Neurons with Special Focus on Nav1.7, Species Differences, and Regulation by Paclitaxel. *Neurosci. Bull* 34, 4–12. 10.1007/s12264-017-0132-3 [PubMed: 28424991]
- Charkhkar H, Arreaga-salas DE, Tran T, Hammack A, Voit WE, Pancrazio JJ, Gnade BE, 2016 Novel disposable microelectrode array for cultured neuronal network recording exhibiting equivalent performance to commercially available arrays. *Sensors Actuators B. Chem* 226, 232–238. 10.1016/j.snb.2015.11.094
- Charkhkar H, Knaack GL, Gnade BE, Keefer EW, Pancrazio JJ, 2012 Development and demonstration of a disposable low-cost microelectrode array for cultured neuronal network recording. *Sensors Actuators B. Chem* 161, 655–660. 10.1016/j.snb.2011.10.086
- Cheng HYM, Penninger JM, 2002 Transcriptional mechanisms underlying neuropathic pain: DREAM, transcription factors and future pain management? *Expert Rev. Neurother* 2, 677–689. 10.1586/14737175.2.5.677 [PubMed: 19810984]
- Cohen MS, Orth CB, Kim HJ, Jeon NL, Jaffrey SR, 2011 Neurotrophin-mediated dendrite-to-nucleus signaling revealed by microfluidic compartmentalization of dendrites. *Proc. Natl. Acad. Sci* 108, 11246–11251. 10.1073/pnas.1012401108 [PubMed: 21690335]
- Connolly P, Clark P, Curtis ASG, Dow JAT, Wilkinson CDW, 1990 An Extracellular microelectrode Array for monitoring electrogenic cells in culture. *Biosens. Bioelectron* 5, 223–234. 10.1016/0956-5663(90)80011-2 [PubMed: 2206490]
- Costa FAL, Neto FLM, 2015 Satellite glial cells in sensory ganglia: its role in pain. *Brazilian J. Anesthesiol* 65, 73–81. 10.1016/j.bjane.2013.07.013
- Danker T, Möller C, 2014 Early identification of hERG liability in drug discovery programs by automated patch clamp. *Front. Pharmacol* 5 8, 1–11. 10.3389/fphar.2014.00203 [PubMed: 24478702]
- Davidson S, Copits BA, Zhang J, Page G, Ghetti A, 2014 Human sensory neurons: Membrane properties and sensitization by inflammatory mediators. *Pain* 155, 1861–1870. 10.1016/j.pain.2014.06.017.Human [PubMed: 24973718]

- De La Torre-Martínez R, Bonache MA, Llabrés-Campaner PJ, Balsera B, Fernández-Carvajal A, Fernández-Ballester G, Ferrer-Montiel A, Jesús Pérez De Vega M, González-Muñiz R, 2017 Synthesis, high-throughput screening and pharmacological characterization of β -lactam derivatives as TRPM8 antagonists. *Sci. Rep* 7, 1–13. 10.1038/s41598-017-10913-x [PubMed: 28127051]
- De Logu F, Nassini R, Materazzi S, Carvalho Gonçalves M, Nosi D, Rossi Degl'Innocenti D, Marone IM, Ferreira J, Li Puma S, Benemei S, Trevisan G, Souza Monteiro De Araújo D, Patacchini R, Bunnett NW, Geppetti P, 2017 Schwann cell TRPA1 mediates neuroinflammation that sustains macrophage-dependent neuropathic pain in mice. *Nat. Commun* 8, 1–16. 10.1038/s41467-017-01739-2 [PubMed: 28232747]
- Denk W, Delaney KR, Gelperin A, Kleinfeld D, Strowbridge BW, Tank DW, Yuste R, 1994 Anatomical and functional imaging of neurons using 2-photon laser scanning microscopy. *J. Neurosci. Methods* 54, 151–162. 10.1016/0165-0270(94)90189-9 [PubMed: 7869748]
- Dib-Hajj SD, Tyrrell L, Cummins TR, Black JA, Wood PM, Waxman SG, 1999 Two tetrodotoxin-resistant sodium channels in human dorsal root ganglion neurons. *FEBS Lett.* 462, 117–120. 10.1016/S0014-5793(99)01519-7 [PubMed: 10580103]
- Dingemans MML, Schütte MG, Wiersma DMM, de Groot A, van Kleef RGDM, Wijnolts FMJ, Westerink RHS, 2016 Chronic 14-day exposure to insecticides or methylmercury modulates neuronal activity in primary rat cortical cultures. *Neurotoxicology* 57, 194–202. 10.1016/j.neuro.2016.10.002 [PubMed: 27720795]
- Djoughri L, 2006 Spontaneous Pain, Both Neuropathic and Inflammatory, Is Related to Frequency of Spontaneous Firing in Intact C-Fiber Nociceptors. *J. Neurosci* 26, 1281–1292. 10.1523/JNEUROSCI.3388-05.2006 [PubMed: 16436616]
- Du Y, Days E, Romaine I, Abney KK, Kaufmann K, Sulikowski G, Stauffer S, Lindsley CW, Weaver CD, 2015 Development and Validation of a Thallium Flux-Based Functional Assay for the Sodium Channel NaV1.7 and Its Utility for Lead Discovery and Compound Profiling. *ACS Chem. Neurosci* 6, 871–878. 10.1021/acschemneuro.5b00004 [PubMed: 25879403]
- Dworak BJ, Wheeler BC, 2009 Novel MEA platform with PDMS microtunnels enables the detection of action potential propagation from isolated axons in culture. *Lab Chip* 9, 404–10. 10.1039/b806689b [PubMed: 19156289]
- Eide L, McMurray CT, 2005 Culture of adult mouse neurons. *Biotechniques* 38, 99–104. 10.2144/05381RR02 [PubMed: 15679091]
- Enright HA, Felix SH, Fischer NO, Mukerjee EV, Soscia D, Mcnerney M, Kulp K, Zhang J, Page G, Miller P, Ghatti A, Wheeler EK, Pannu S, 2016 Long-term noninvasive interrogation of human dorsal root ganglion neuronal cultures on an integrated microfluidic multielectrode array platform. *Analyst* 141, 5346–5357. 10.1039/C5AN01728A [PubMed: 27351032]
- Fitzgerald EM, Okuse K, Wood JN, Dolphin AC, Moss SJ, 1999 cAMP-dependent phosphorylation of the tetrodotoxin-resistant voltage-dependent sodium channel SNS. *J. Physiol* 516, 433–446. 10.1111/j.1469-7793.1999.0433v.x [PubMed: 10087343]
- Fouillet A, Watson JF, Piekars AD, Huang X, Li B, Priest B, Nisenbaum E, Sher E, Ursu D, 2017 Characterisation of Nav1.7 functional expression in rat dorsal root ganglia neurons by using an electrical field stimulation assay. *Mol. Pain* 13, 1–13. 10.1177/1744806917745179
- Fromherz P, Offenhäusser A, Vetter T, Weis J, 1991 A Neuron-Silicon Junction: A Retzius Cell of the Leech on an Insulated-Gate Field-Effect Transistor. *Science* (80-.). 252, 1290–1293.
- Gluska S, Chein M, Rotem N, Ionescu A, Perlson E, 2015 Tracking Quantum-Dot labeled neurotropic factors transport along primary neuronal axons in compartmental microfluidic chambers. *Methods Cell Biol.* 131, 1–23. 10.1016/bs.mcb.2015.06.016 [PubMed: 26794507]
- Gornstein EL, Schwartz TL, 2017 Neurotoxic mechanisms of paclitaxel are local to the distal axon and independent of transport defects. *Exp. Neurol* 288, 153–166. 10.1007/s10549-015-3663-1 [PubMed: 27894788]
- Grienberger C, Konnerth A, 2012 Imaging calcium in neurons. *Neuron* 73, 862–85. 10.1016/j.neuron.2012.02.011 [PubMed: 22405199]
- Griscom L, Chateau Y, Pennec JP, Misery L, Le Pioufle B, 2005 Co-culture of cells in PDMS microsystem for sensitized artificial skin. 2005 3rd IEEE/EMBS Spec. Top. Conf. Microtechnology Med. Biol. 2005, 184–187. 10.1109/MMB.2005.1548421

- Gross GW, Rhoades BK, Azzazy HME, Wu Ming-Chi, 1995 The use of neuronal networks on multielectrode arrays as biosensors. *Biosens. Bioelectron* 10, 553–567. 10.1016/0956-5663(95)96931-N [PubMed: 7612207]
- Gross GW, Wen WY, Lin JW, 1985 Transparent indium-tin oxide electrode patterns for extracellular, multisite recording in neuronal cultures. *J. Neurosci. Methods* 15, 243–252. 10.1016/0165-0270(85)90105-0 [PubMed: 4094480]
- Gu L, Black B, Ordonez S, Mondal A, Jain A, Mohanty S, 2014 Microfluidic control of axonal guidance. *Sci. Rep* 4 10.1038/srep06457
- Gumy LF, Yeo GSH, Tung Y-CL, Zivraj KH, Willis D, Coppola G, Lam BYH, Twiss JL, Holt CE, Fawcett JW, 2011 Transcriptome analysis of embryonic and adult sensory axons reveals changes in mRNA repertoire localization. *RNA* 17, 85–98. 10.1261/rna.2386111 [PubMed: 21098654]
- Hammack A, Rihani RT, Black BJ, Pancrazio JJ, Gnade BE, 2018 A patterned polystyrene-based microelectrode array for in vitro neuronal recordings. *Biomed. Microdevices* 20, 1–7. 10.1007/s10544-018-0295-3
- Haroutounian S, Nikolajsen L, Bendtsen TF, Finnerup NB, Kristensen AD, Hasselstrøm JB, Jensen TS, 2014 Primary afferent input critical for maintaining spontaneous pain in peripheral neuropathy. *Pain* 155, 1272–1279. 10.1016/j.pain.2014.03.022 [PubMed: 24704366]
- Hartlehnert M, Derksen A, Hagenacker T, Kindermann D, Schäfers M, Pawlak M, Kieseier BC, Meyer Zu Horste G, 2017 Schwann cells promote post-traumatic nerve inflammation and neuropathic pain through MHC class II. *Sci. Rep* 7, 1–10. 10.1038/s41598-017-12744-2 [PubMed: 28127051]
- Heim M, Yvert B, Kuhn A, 2012 Journal of Physiology - Paris Nanostructuring strategies to enhance microelectrode array (MEA) performance for neuronal recording and stimulation. *J. Physiol. - Paris* 106, 137–145. 10.1016/j.jphysparis.2011.10.001 [PubMed: 22027264]
- Heinricher MM, 2004 Principles of Extracellular Single-Unit Recording, in: *Microelectrode Recording in Movement Disorder Surgery*. pp. 8–13. 10.1055/b-0034-56092
- Hempel CM, Werley CA, Dempsey GT, Gerber DJ, 2017 Targeting neuronal function for CNS drug discovery. *Drug Discov. Today Technol* 23, 17–25. 10.1016/j.ddtec.2017.03.005 [PubMed: 28647082]
- Hondebrink L, Verboven AHA, Drega WS, Schmeink S, de Groot MWGDM, van Kleef RGDM, Wijnolts FMJ, de Groot A, Meulenbelt J, Westerink RHS, 2016 Neurotoxicity screening of (illicit) drugs using novel methods for analysis of microelectrode array (MEA) recordings. *Neurotoxicology* 55, 1–9. 10.1016/j.neuro.2016.04.020 [PubMed: 27149913]
- Huang LYM, Gu Y, Chen Y, 2013 Communication between neuronal somata and satellite glial cells in sensory ganglia. *Glia* 61, 1571–1581. 10.1002/glia.22541 [PubMed: 23918214]
- Hudmon A, Choi J-S, Tyrrell L, Black JA, Rush AM, Waxman SG, Dib-Hajj SD, 2008 Phosphorylation of Sodium Channel Nav1.8 by p38 Mitogen-Activated Protein Kinase Increases Current Density in Dorsal Root Ganglion Neurons. *J. Neurosci* 28, 3190–3201. 10.1523/JNEUROSCI.4403-07.2008 [PubMed: 18354022]
- Ishii MN, Yamamoto K, Shoji M, Asami A, Kawamata Y, 2017 Human induced pluripotent stem cell (hiPSC)-derived neurons respond to convulsant drugs when co-cultured with hiPSC-derived astrocytes. *Toxicology* 389, 130–138. 10.1016/j.tox.2017.06.010 [PubMed: 28666936]
- Jao PF, Franca EW, Fang SP, Wheeler BC, Yoon YK, 2015 Immersion lithographic patterning of electrospun nanofibers for carbon nanofibrous microelectrode arrays. *J. Microelectromechanical Syst* 24, 703–715. 10.1109/JMEMS.2014.2346503
- Jin X, Morsy N, Winston J, Pasricha PJ, Garrett K, Akbarali HI, 2004 Modulation of TRPV1 by nonreceptor tyrosine kinase, c-Src kinase. *Am. J. Physiol. Cell Physiol* 287, C558–C563. 10.1152/ajpcell.00113.2004 [PubMed: 15084474]
- Johnstone AFM, Gross GW, Weiss DG, Schroeder OHU, Gramowski A, Shafer TJ, 2010 Microelectrode arrays: A physiologically based neurotoxicity testing platform for the 21st century. *Neurotoxicology* 31, 331–350. 10.1016/j.neuro.2010.04.001 [PubMed: 20399226]
- Jonas R, Klusch A, Schmelz M, Petersen M, Carr RW, 2015 Assessment of TTX-s and TTX-r action potential conduction along neurites of NGF and GDNF cultured porcine DRG somata. *PLoS One* 10, 1–16. 10.1371/journal.pone.0139107

- Kang P, Liao M, Wester MR, Leeder JS, Pearce RE, 2009 Translating nociceptor sensitivity: the role of axonal protein synthesis in nociceptor physiology. *Eur. J. Neurosci* 29, 2253–2263. 10.1124/dmd.107.016501.CYP3A4-Mediated [PubMed: 19490023]
- Kayama T, Suzuki I, Odawara A, Sasaki T, Ikegaya Y, 2018 Temporally coordinated spiking activity of human induced pluripotent stem cell-derived neurons co-cultured with astrocytes. *Biochem. Biophys. Res. Commun* 495, 1028–1033. 10.1016/j.bbrc.2017.11.115 [PubMed: 29170135]
- Khoutorsky A, Price TJ, 2017 Translational Control Mechanisms in Persistent Pain. *Trends Neurosci.* 1–15. 10.1016/j.tins.2017.11.006
- Kilinc D, Peyrin JM, Soubeyre V, Magnifico S, Saias L, Viovy JL, Brugg B, 2011 Wallerian-like degeneration of central neurons after synchronized and geometrically registered mass axotomy in a three-compartmental microfluidic chip. *Neurotox. Res* 19, 149–161. 10.1007/s12640-010-9152-8 [PubMed: 20162389]
- Kim YS, Anderson M, Park K, Zheng Q, Agarwal A, Gong C, Saijilafu, Young LA, He S, LaVinka PC, Zhou F, Bergles D, Hanani M, Guan Y, Spray DC, Dong X, 2016 Coupled Activation of Primary Sensory Neurons Contributes to Chronic Pain. *Neuron* 91, 1085–1096. 10.1016/j.neuron.2016.07.044 [PubMed: 27568517]
- Kodama D, Hirai T, Kondo H, Hamamura K, Togari A, 2017 Bidirectional communication between sensory neurons and osteoblasts in an in vitro coculture system. *FEBS Lett.* 591, 527–539. 10.1002/1873-3468.12561 [PubMed: 28094440]
- Krames ES, 2015 The dorsal root ganglion in chronic pain and as a target for neuromodulation: A review. *Neuromodulation* 18, 24–32. 10.1111/ner.12247 [PubMed: 25354206]
- Krishnamoorthy K, Zoski CG, 2005 Fabrication of 3D gold nanoelectrode ensembles by chemical etching. *Anal. Chem* 77, 5068–5071. 10.1021/ac050604r [PubMed: 16053324]
- Kulagina NV, O'Shaughnessy TJ, Ma W, Ramsdell JS, Pancrazio JJ, 2004 Pharmacological effects of the marine toxins, brevetoxin and saxitoxin, on murine frontal cortex neuronal networks. *Toxicon* 44, 669–676. 10.1016/j.toxicon.2004.07.023 [PubMed: 15501293]
- Kulkarni RU, Miller EW, 2017 Voltage Imaging: Pitfalls and Potential. *Biochemistry* 56, 5171–5177. 10.1021/acs.biochem.7b00490 [PubMed: 28745864]
- Kumamoto J, Ichi, Nakatani M, Tsutsumi M, Goto M, Denda S, Takei K, Denda M, 2014 Coculture system of keratinocytes and dorsal-root-ganglion-derived cells for screening neurotrophic factors involved in guidance of neuronal axon growth in the skin. *Exp. Dermatol* 23, 58–60. 10.1111/exd.12288 [PubMed: 24267269]
- Larner D, 2014 Chronic pain transition: A concept analysis. *Pain Manag. Nurs* 10.1016/j.pmn.2013.04.003
- Lee HJ, Zhang D, Jiang Y, Wu X, Shih P-Y, Liao C-S, Bungart B, Xu X-M, Drenan R, Bartlett E, Cheng J-X, 2017 Label-Free Vibrational Spectroscopic Imaging of Neuronal Membrane Potential. *J. Phys. Chem. Lett* 8, 1932–1936. 10.1021/acs.jpclett.7b00575 [PubMed: 28407470]
- Li T, Lu G, Chiang EY, Chernov-Rogan T, Grogan JL, Chen J, 2017 High-throughput electrophysiological assays for voltage gated ion channels using SyncroPatch 768PE. *PLoS One* 12, 1–18. 10.1371/journal.pone.0180154
- Mastrototaro JJ, Massoud HZ, Pilkington TC, Ideker RE, 1992 Rigid and Flexible Thin-Film Multielectrode Arrays for Transmural Cardiac Recording. *IEEE Trans. Biomed. Eng* 39, 271–279. 10.1109/10.125012 [PubMed: 1555857]
- Melemedjian OK, Asiedu MN, Tillu DV, Peebles KA, Yan J, Ertz N, Dussor GO, Price TJ, 2010 IL-6- and NGF-induced rapid control of protein synthesis and nociceptive plasticity via convergent signaling to the eIF4F complex. *J. Neurosci* 30, 15113–15123. 10.1523/JNEUROSCI.3947-10.2010 [PubMed: 21068317]
- Melemedjian OK, Asiedu MN, Tillu DV, Peebles KA, Ertz N, Dussor GO, Price TJ, 2010 IL-6- and NGF-induced rapid control of protein synthesis and nociceptive plasticity via convergent signaling to the eIF4F complex. *J. Neurosci* 30, 15113–15123. 10.1523/JNEUROSCI.3947-10.2010.IL-6- [PubMed: 21068317]
- Meuwis K, Boens N, De Schryver FC, Gallay J, Vincent M, 1995 Photophysics of the fluorescent K⁺ indicator PBFI. *Biophys. J* 68, 2469–2473. 10.1016/S0006-3495(95)80428-5 [PubMed: 7647249]

- Minta A, Tsien RY, 1989 Fluorescent indicators for cytosolic sodium. *J. Biol. Chem* 264, 19449–19457. [PubMed: 2808435]
- Miu DKY, Chan KC, 2014 Under-detection of pain in elderly nursing home residents with moderate to severe dementia. *J. Clin. Gerontol. Geriatr.* 5, 23–27. 10.1016/j.jcgg.2013.09.001
- Moffat JG, Vincent F, Lee JA, Eder J, Prunotto M, 2017 Opportunities and challenges in phenotypic drug discovery: An industry perspective. *Nat. Rev. Drug Discov* 16, 531–543. 10.1038/nrd.2017.111 [PubMed: 28685762]
- Moldovan M, Alvarez S, Romer Rosberg M, Krarup C, 2013 Axonal voltage-gated ion channels as pharmacological targets for pain. *Eur. J. Pharmacol* 708, 105–12. 10.1016/j.ejphar.2013.03.001 [PubMed: 23500193]
- Möller C, Witchel H, 2011 Automated electrophysiology makes the pace for cardiac ion channel safety screening. *Front. Pharmacol* 2 11, 1–7. 10.3389/fphar.2011.00073 [PubMed: 21779246]
- Moy JK, Khoutorsky A, Asiedu MN, Black BJ, Kuhn JL, Barragán-Iglesias P, Megat S, Burton MD, Burgos-Vega CC, Melemedjian OK, Boitano S, Vagner J, Gkogkas CG, Pancrazio JJ, Mogil JS, Dussor G, Sonenberg N, Price TJ, 2017 The MNK — eIF4E signaling axis contributes to injury-induced nociceptive plasticity and the development of chronic pain. *J. Neurosci* 0220–17. 10.1523/JNEUROSCI.0220-17.2017
- Munro G, Jansen-Olesen I, Olesen J, 2017 Animal models of pain and migraine in drug discovery. *Drug Discov. Today* 22, 1103–1111. 10.1016/j.drudis.2017.04.016 [PubMed: 28476535]
- Musick K, Khatami D, Wheeler BC, 2009 Three-dimensional micro-electrode array for recording dissociated neuronal cultures. *Lab Chip* 9, 2036–2042. 10.1039/b820596e [PubMed: 19568672]
- Neto E, Alves CJ, Sousa DM, Alencastre IS, Lourenço AH, Leitão L, Ryu HR, Jeon NL, Fernandes R, Aguiar P, Almeida RD, Lamghari M, 2014 Sensory neurons and osteoblasts: close partners in a microfluidic platform. *Integr. Biol* 6, 586–595. 10.1039/C4IB00035H
- Newberry K, Wang S, Hoque N, Ahljanian MK, Kiss L, Herrington J, Graef JD, 2016 Development of a Spontaneously Active Dorsal Root Ganglia Assay using Multi-well Multielectrode Arrays. *J. Neurophysiol* jn.01122.2015. 10.1152/jn.01122.2015
- Nicolas J, Hendriksen PJM, van Kleef RGDM, de Groot A, Bovee TFH, Rietjens IMCM, Westerink RHS, 2014 Detection of marine neurotoxins in food safety testing using a multielectrode array. *Mol. Nutr. Food Res* 58, 2369–2378. 10.1002/mnfr.201400479 [PubMed: 25266399]
- Novellino a, Scelfo B, Palosaari T, Price A, Sobanski T, Shafer TJ, Johnstone a F.M., Gross GW, Gramowski A, Schroeder O, Jügel K, Chiappalone M, Benfenati F, Martinoia S, Tedesco MT, Defranchi E, D'Angelo P, Whelan M, 2011 Development of micro-electrode array based tests for neurotoxicity: assessment of interlaboratory reproducibility with neuroactive chemicals. *Front. Neuroeng* 4, 4 10.3389/fneng.2011.00004 [PubMed: 21562604]
- Obara I, Géranton SM, Hunt SP, 2012 Axonal protein synthesis: A potential target for pain relief? *Curr. Opin. Pharmacol* 12, 42–48. 10.1016/j.coph.2011.10.005 [PubMed: 22033338]
- Obara I, Hunt SP, 2014 Axonal protein synthesis and the regulation of primary afferent function. *Dev. Neurobiol* 74, 269–278. 10.1002/dneu.22133 [PubMed: 24085547]
- Odawara A, Matsuda N, Suzuki I, 2017 Pain responses in human iPSC-derived sensory neurons using MEA system, in: *Advances in Cell Based Screening*. p. 1.
- Odawara A, Saitoh Y, Alhebshi AH, Gotoh M, Suzuki I, 2014 Long-term electrophysiological activity and pharmacological response of a human induced pluripotent stem cell-derived neuron and astrocyte co-culture. *Biochem. Biophys. Res. Commun* 443, 1176–1181. 10.1016/j.bbrc.2013.12.142 [PubMed: 24406164]
- Ohshiro H, Ogawa S, Shinjo K, 2007 Visualizing sensory transmission between dorsal root ganglion and dorsal horn neurons in co-culture with calcium imaging. *J. Neurosci. Methods* 165, 49–54. 10.1016/j.jneumeth.2007.05.018 [PubMed: 17597226]
- Omaki T, Hirai Y, Kamei K, Tsuchiya T, Tabata O, 2017 Microfabrication of Embedding a Flexible Parylene-Based Microelectrode Array within Body-on-a-Chip. *Proceedings* 1, 302 10.3390/proceedings1040302
- Pancrazio JJ, Whelan JP, Borkholder DA, Ma W, Stenger DA, 1999 Development and Application of Cell-Based Biosensors. *Ann. Biomed. Eng* 27, 697–711. 10.1114/1.225 [PubMed: 10625143]

- Parenti C, Turnaturi R, Aricò G, Gramowski-Voß A, Schroeder OHU, Marrazzo A, Prezzavento O, Ronsisvalle S, Scoto GM, Ronsisvalle G, Pasquinucci L, 2013 The multitarget opioid ligand LPI's effects in persistent pain and in primary cell neuronal cultures. *Neuropharmacology* 71, 70–82. 10.1016/j.neuropharm.2013.03.008 [PubMed: 23541722]
- Park JW, Kim HJ, Byun JH, Ryu HR, Jeon NL, 2009 Novel microfluidic platform for culturing neurons: Culturing and biochemical analysis of neuronal components. *Biotechnol. J* 4, 1573–1577. 10.1002/biot.200900159 [PubMed: 19918787]
- Park JW, Vahidi B, Taylor AM, Rhee SW, Jeon NL, 2006 Microfluidic culture platform for neuroscience research. *Nat. Protoc* 1, 2128–2136. 10.1038/nprot.2006.316 [PubMed: 17487204]
- Pasca AM, Sloan SA, Clarke LE, Tian Y, Makinson CD, Huber N, Kim CH, Park JY, O'Rourke NA, Nguyen KD, Smith SJ, Huguenard JR, Geschwind DH, Barres BA, Pasca SP, 2015 Functional cortical neurons and astrocytes from human pluripotent stem cells in 3D culture. *Nat. Methods* 12, 671–678. 10.1038/nmeth.3415 [PubMed: 26005811]
- Pazyra-Murphy M, Segal R, 2008 Preparation and Maintenance of Dorsal Root Ganglia Neurons in Compartmented Cultures. *J. Vis. Exp* 1–4. 10.3791/951
- Pearce TM, Wilson JA, Oakes SG, Chiu S-Y, Williams JC, 2005 Integrated microelectrode array and microfluidics for temperature clamp of sensory neurons in culture. *Lab Chip* 5, 97–101. 10.1039/b407871c [PubMed: 15616746]
- Poplawski G, Ishikawa T, Brifault C, Lee-Kubli C, Regestam R, Henry KW, Shiga Y, Kwon H, Ohtori S, Gonias SL, Campana WM, 2018 Schwann cells regulate sensory neuron gene expression before and after peripheral nerve injury. *Glia* 1–14. 10.1002/glia.23325
- Price TJ, Das V, Dussor G, 2016 Adenosine monophosphate-activated protein kinase (AMPK) activators for the prevention, treatment and potential reversal of pathological pain. *Curr Drug Targets* 17, 908–920. 10.1007/s12671-013-0269-8.Moving [PubMed: 26521775]
- Pruss RM, 2010 Phenotypic screening strategies for neurodegenerative diseases: a pathway to discover novel drug candidates and potential disease targets or mechanisms. *CNS Neurol. Disord. Drug Targets* 9, 693–700. <https://doi.org/BSP/CDTCNSND/E-Pub/00078> [pii] [PubMed: 20942792]
- Ratté S, Prescott SA, 2016 Afferent hyperexcitability in neuropathic pain and the inconvenient truth about its degeneracy. *Curr. Opin. Neurobiol.* 36, 31–37. 10.1016/j.conb.2015.08.007 [PubMed: 26363576]
- Ravula SK, Wang MS, McClain MA, Asress SA, Frazier B, Glass JD, 2007 Spatiotemporal localization of injury potentials in DRG neurons during vincristine-induced axonal degeneration. *Neurosci. Lett* 415, 34–39. 10.1016/j.neulet.2007.01.009 [PubMed: 17267126]
- Richter JD, 2007 CPEB: a life in translation. *Trends Biochem. Sci* 32, 279–285. 10.1016/j.tibs.2007.04.004 [PubMed: 17481902]
- Rihani R, Kim H, Black B, Atmaramani R, Saed M, Pancrazio J, Ware T, 2018 Liquid Crystal Elastomer-Based Microelectrode Array for In Vitro Neuronal Recordings. *Micromachines* 9, 416 10.3390/mi9080416
- Roggenkamp D, Falkner S, Stäb F, Petersen M, Schmelz M, Neufang G, 2012 Atopic keratinocytes induce increased neurite outgrowth in a coculture model of porcine dorsal root ganglia neurons and human skin cells. *J. Invest. Dermatol* 132, 1892–1900. 10.1038/jid.2012.44 [PubMed: 22418869]
- Santoro F, Dasgupta S, Schnitker J, Auth T, Neumann E, Panaitov G, Gompper G, Offenhäusser A, 2014 Interfacing electrogenic cells with 3D nanoelectrodes: Position, shape, and size matter. *ACS Nano* 8, 6713–6723. 10.1021/nn500393p [PubMed: 24963873]
- Schutte RJ, Xie Y, Ng NN, Figueroa P, Pham AT, O'Dowd DK, 2018 Astrocyte-enriched feeder layers from cryopreserved cells support differentiation of spontaneously active networks of human iPSC-derived neurons. *J. Neurosci. Methods* 294, 91–101. 10.1016/j.jneumeth.2017.07.019 [PubMed: 28746822]
- Schwartzentruber J, Foskolou S, Kilpinen H, Rodrigues J, Alasoo K, Knights AJ, Patel M, Goncalves A, Ferreira R, Benn CL, Wilbrey A, Bictash M, Impey E, Cao L, Lainez S, Loucif AJ, Whiting PJ, Gutteridge A, Gaffney DJ, 2018 Molecular and functional variation in iPSC-derived sensory neurons. *Nat. Genet* 50, 54–61. 10.1038/s41588-017-0005-8 [PubMed: 29229984]

- Selinger JV, Pancrazio JJ, Gross GW, 2004 Measuring synchronization in neuronal networks for biosensor applications. *Biosens. Bioelectron* 19, 675–683. 10.1016/S0956-5663(03)00267-7 [PubMed: 14709385]
- Sheahan TD, Valtcheva MV, McIlvried LA, Pullen MY, Baranger DAA, Gereau RW, 2018 Metabotropic Glutamate Receptor 2/3 (mGluR2/3) Activation Suppresses TRPV1 Sensitization in Mouse, but Not Human Sensory Neurons. *Eneuro* 10, ENEURO.0412–17.2018. 10.1523/ENEURO.0412-17.2018
- Silva DI, Santos B.P. Dos, Leng J, Oliveira H, Amédée J, 2017 Dorsal root ganglion neurons regulate the transcriptional and translational programs of osteoblast differentiation in a microfluidic platform article. *Cell Death Dis.* 8 10.1038/s41419-017-0034-3
- Smith NA, Kress BT, Lu Y, Chandler-militello D, Benraiss A, Nedergaard M, 2018 Fluorescent Ca²⁺ indicators directly inhibit the Na, K-ATPase and disrupt cellular functions. *Sci. Signal* 2039 10.1126/scisignal.aal2039
- Spira ME, Hai A, 2013. Multi-electrode array technologies for neuroscience and cardiology. *Nat. Nanotechnol* 8, 83–94. 10.1038/nnano.2012.265 [PubMed: 23380931]
- Steinmetz NA, Buetfering C, Lecoq J, Lee CR, Peters AJ, Jacobs EAK, Coen P, Ollerenshaw DR, Valley MT, de Vries SEJ, Garrett M, Zhuang J, Groblewski PA, Manavi S, Miles J, White C, Lee E, Griffin F, Larkin JD, Roll K, Cross S, Nguyen TV, Larsen R, Pendergraft J, Daigle T, Tasic B, Thompson CL, Waters J, Olsen S, Margolis DJ, Zeng H, Hausser M, Carandini M, Harris KD, 2017 Aberrant Cortical Activity in Multiple GCaMP6-Expressing Transgenic Mouse Lines. *Eneuro* 4, ENEURO.0207–17.2017. 10.1523/ENEURO.0207-17.2017
- Sutton KG, Martin DJ, Pinnock RD, Lee K, Scott RH, 2002 Gabapentin inhibits high-threshold calcium channel currents in cultured rat dorsal root ganglion neurones. *Br.J.Pharmacol* 135, 257–265. 10.1038/sj.bjp.0704439 [PubMed: 11786502]
- Takeda M, Takahashi M, Matsumoto S, 2009 Contribution of the activation of satellite glia in sensory ganglia to pathological pain. *Neurosci. Biobehav. Rev* 33, 784–792. 10.1016/j.neubiorev.2008.12.005 [PubMed: 19167424]
- Tang X, Zhou L, Wagner AM, Marchetto MCN, Muotri AR, Gage FH, Chen G, 2013 Astroglial cells regulate the developmental timeline of human neurons differentiated from induced pluripotent stem cells. *Stem Cell Res.* 11, 743–757. 10.1016/j.scr.2013.05.002 [PubMed: 23759711]
- Tidball A, Parent J, 2015 Exciting cells: modeling genetic epilepsies with patient-derived induced pluripotent stem cells. *Stem Cells* 34, 27–33. 10.1002/stem.2203.Exciting [PubMed: 26373465]
- Tillu DV, Melemedjian OK, Asiedu MN, Qu N, De Felice M, Dussor G, Price TJ, 2012 Resveratrol engages AMPK to attenuate ERK and mTOR signaling in sensory neurons and inhibits incision-induced acute and chronic pain. *Mol. Pain* 8, 5 10.1186/1744-8069-8-5 [PubMed: 22269797]
- Tsantoulas C, Farmer C, Machado P, Baba K, McMahon SB, Raouf R, 2013 Probing functional properties of nociceptive axons using a microfluidic culture system. *PLoS One* 8, 1–17. 10.1371/journal.pone.0080722
- Unsain N, Heard KN, Higgins JM, Barker PA, 2014 Production and Isolation of Axons from Sensory Neurons for Biochemical Analysis Using Porous Filters. *J. Vis. Exp* 1–8. 10.3791/51795
- Valdivia P, Martin M, LeFevre WR, Ross J, Houck KA, Shafer TJ, 2014 Multi-well microelectrode array recordings detect neuroactivity of ToxCast compounds. *Neurotoxicology* 44, 204–217. 10.1016/j.neuro.2014.06.012 [PubMed: 24997244]
- Valeyev a Y., Hackman JC, Wood PM, Davidoff R. a, 1996 Pharmacologically novel GABA receptor in human dorsal root ganglion neurons. *J. Neurophysiol* 76, 3555–3558. [PubMed: 8930293]
- Valtcheva MV, Copits BA, Davidson S, Sheahan TD, Pullen MY, McCall JG, Dikranian K, Gereau RW, 2016 Surgical extraction of human dorsal root ganglia from organ donors and preparation of primary sensory neuron cultures. *Nat. Protoc* 11, 1877–1888. 10.1038/nprot.2016.111 [PubMed: 27606776]
- van de Wijdeven R, Ramstad OH, Bauer US, Halaas Ø, Sandvig A, Sandvig I, 2018 Structuring a multi-nodal neural network in vitro within a novel design microfluidic chip. *Biomed. Microdevices* 20 10.1007/s10544-017-0254-4

- Vargas ME, Yamagishi Y, Tessier-Lavigne M, Sagasti A, 2015 Live Imaging of Calcium Dynamics during Axon Degeneration Reveals Two Functionally Distinct Phases of Calcium Influx. *J. Neurosci* 35, 15026–15038. 10.1523/JNEUROSCI.2484-15.2015 [PubMed: 26558774]
- Vellani V, Mapplebeck S, Moriondo a, Davis JB, McNaughton P. a, 2001 Protein kinase C activation potentiates gating of the vanilloid receptor VR1 by capsaicin, protons, heat and anandamide. *J. Physiol* 534, 813–25. [PubMed: 11483711]
- Vessoni AT, Herai RH, Karpiak JV, Leal AMS, Trujillo CA, Quinet A, Agnez Lima LF, Menck CFM, Muotri AR, 2016 Cockayne syndrome-derived neurons display reduced synapse density and altered neural network synchrony. *Hum. Mol. Genet* 25, 1271–1280. 10.1093/hmg/ddw008 [PubMed: 26755826]
- Von Banchet GS, Kiehl M, Schaible HG, 2005 Acute and long-term effects of IL-6 on cultured dorsal root ganglion neurones from adult rat. *J. Neurochem* 94, 238–248. 10.1111/j.1471-4159.2005.03185.x [PubMed: 15953366]
- Wainger BJ, Buttermore ED, Oliveira JT, Mellin C, Lee S, Saber WA, Wang AJ, Ichida JK, Chiu IM, Barrett L, Huebner EA, Bilgin C, Tsujimoto N, Brenneis C, Kapur K, Rubin LL, Eggan K, Woolf CJ, 2014 Modeling pain in vitro using nociceptor neurons reprogrammed from fibroblasts. *Nat. Neurosci* 18, 17–24. 10.1038/nn.3886 [PubMed: 25420066]
- Wall PD, Devor M, 1981 The effect of peripheral nerve injury on dorsal root potentials and on transmission of afferent signals into the spinal cord. *Spinal Cord* 20, 95–111.
- Waxman SG, Cummins TR, Dib-Hajj S, Fjell J, Black JA, 1999 Sodium channels, excitability of primary sensory neurons, and the molecular basis of pain. *Muscle and Nerve* 22, 1177–1187. 10.1002/(SICI)1097-4598(199909)22:9<1177::AID-MUS3>3.0.CO;2-P [PubMed: 10454712]
- Waxman SG, Zamponi GW, 2014 Regulating excitability of peripheral afferents: emerging ion channel targets. *Nat. Neurosci* 17, 153–63. 10.1038/nn.3602 [PubMed: 24473263]
- Weidlich S, Krause KJ, Schnitker J, Wolfrum B, Offenhäusser A, 2017 MEAs and 3D nanoelectrodes: Electrodeposition as tool for a precisely controlled nanofabrication. *Nanotechnology* 28 10.1088/1361-6528/aa57b5
- Weng X, Smith T, Sathish J, Djouhri L, 2012 Chronic inflammatory pain is associated with increased excitability and hyperpolarization-activated current (I_h) in C- but not Aδ-nociceptors. *Pain* 153, 900–914. 10.1016/j.pain.2012.01.019 [PubMed: 22377439]
- Wheeler M. a, Heffner DL, Kim S, Espy SM, Spano AJ, Cleland CL, Deppmann CD, 2014 TNF-α/TNFR1 signaling is required for the development and function of primary nociceptors. *Neuron* 82, 587–602. 10.1016/j.neuron.2014.04.009 [PubMed: 24811380]
- White FA, Bhargoo SK, Miller RJ, 2005 Chemokines: Integrators of pain and inflammation. *Nat. Rev. Drug Discov* 4, 834–844. 10.1038/nrd1852 [PubMed: 16224455]
- Yan J, Melemedjian OK, Price TJ, Dussor G, 2012 Sensitization of dural afferents underlies migraine-related behavior following meningeal application of interleukin-6 (IL-6). *Mol. Pain* 8, 6 10.1186/1744-8069-8-6 [PubMed: 22273495]
- Yang IH, Gary D, Malone M, Dria S, Houdayer T, Belegu V, McDonald JW, Thakor N, 2012 Axon myelination and electrical stimulation in a microfluidic, compartmentalized cell culture platform. *NeuroMolecular Med.* 14, 112–118. 10.1007/s12017-012-8170-5 [PubMed: 22527791]
- Yang Y, Huang J, Mis MA, Estacion M, Macala L, Shah P, Schulman BR, Horton DB, Dib-Hajj SD, Waxman SG, 2016 Nav1.7-A1632G Mutation from a Family with Inherited Erythromelalgia: Enhanced Firing of Dorsal Root Ganglia Neurons Evoked by Thermal Stimuli. *J. Neurosci* 36, 7511–7522. 10.1523/JNEUROSCI.0462-16.2016 [PubMed: 27413160]
- Zhang JM, Song XJ, LaMotte RH, 1999 Enhanced excitability of sensory neurons in rats with cutaneous hyperalgesia produced by chronic compression of the dorsal root ganglion. *J. Neurophysiol.* 82, 3359–3366. [PubMed: 10601467]
- Zhang K, Osakada Y, Vrljic M, Chen L, Mudrakola H, Cui B, 2010 Single molecule imaging of NGF axonal transport in microfluidic devices. *Nat. Rev. Cancer* 10, 2566–2573. 10.1002/ana.22528.Toll-like
- Zhang M, Luo G, Zhou Y, Wang S, Zhong Z, 2014 Phenotypic screens targeting neurodegenerative diseases. *J. Biomol. Screen* 19, 1–16. 10.1177/1087057113499777 [PubMed: 23958650]

- Zhang Y, Zhang X, Fang J, Jiang S, Zhang Y, Gu D, Nelson RD, LaRue JC, 2010 Application of SU-8 as the insulator toward a novel planar microelectrode array for extracellular neural recording. 2010 IEEE 5th Int. Conf. Nano/Micro Eng. Mol. Syst. NEMS 2010 395–398. 10.1109/NEMS.2010.5592414
- Zhao F, Li X, Jin L, Zhang F, Inoue M, Yu B, Cao Z, 2016 Development of a rapid throughput assay for identification of hNav1.7 antagonist using unique efficacious sodium channel agonist, antillatoxin. Mar. Drugs 14, 1–12. 10.3390/md14020036
- Zhu W; Oxford GS, 2011 Differential gene expression of neonatal and adult DRG neurons correlates with the differential sensitization of TRPV1 responses to nerve growth factor. Neurosci Lett. 500, 192–196. 10.1016/j.neulet.2011.06.034.Differential [PubMed: 21741445]
- Zweifel LS, Kuruvilla R, Ginty DD, 2005 Functions and mechanisms of retrograde neurotrophin signalling. Nat. Rev. Neurosci 6, 615–625. 10.1038/nrn1727 [PubMed: 16062170]

Highlights

- Chronic pain affects 100 million people in the US and leads to long-term disability.
- New pain medications are desperately needed, as common opioid prescriptions are ineffective and/or lead to long-term tolerance and abuse.
- Identification and targeting of ‘druggable’ molecular pathways is problematic due to the signaling complexity involved in chronic pain conditions.
- As an alternative or supplemental strategy, identification of hyperexcitable phenotypes in relevant cellular models may enable higher throughput screening of analgesic compounds.
- Sensory neurons of the dorsal root ganglion are central to the development and maintenance of chronic pain conditions, are known to change phenotype in terms of spontaneous and evoked activity, relate that phenotypic change in terms of readily measureable all-or-nothing events, and may serve as a cellular sensor for hyperexcitable phenotypes.
- Promising excitability-based phenotypic assays include calcium and voltage imaging, manual and automated patch-clamp, and substrate integrated microelectrode arrays. These methods offer moderate-to-high throughput and can be incorporated with compartmentalized co-culture systems.
- Future work should focus on development of human-derived sensory neuron models (especially hiPSC) which are scalable and retain hyperexcitable phenotypes *in vitro*.

Figure 1: Schematic illustrating the signaling complexity associated with establishing and maintaining chronic neuropathic pain. Numbered actions are expounded in section 2.2. Inset summarizes the cytokine-mediated translational regulation exerted at the 5' and 3' UTR end of mature mRNA. The outcome of cytokine-mediated eIF4F cap complex formation and polyadenylation is increased translation efficiency.

Figure 2:

Screen images from the Axion Biosystems AxiS data acquisition system depicting extracellular neural activity from a 48-well microelectrode array plate cultured with primary mouse cortical neurons 19 days in vitro. A) Representative heat map showing detected spikes across the multi-well plate. The arrow identifies well F7. B) Real time recording from 16 microelectrode contacts within well F7. C) Event detection within well F7 illustrating the features of the detected spikes.

Figure 3:

Extracellular MEA-based recordings from hiPSC sensory neurons. (A) Phase contrast image of dense cellular and axonal network on MEA substrate after 48 days in vitro. Scale bar represents 20 μm . (B) Filtered continuous recording (12.5 kHz sampling rate) from a representative electrode. Vertical and horizontal scale bars represent 30 μV and 500 ms, respectively. Red inset illustrates resolution of single action potential waveforms. Vertical and horizontal scale bars represent 30 μV and 3 ms, respectively. (C) and (D) Raster plots of hiPSC sensory neuron activity in the absence and presence of 100 nM capsaicin and

temperature increase (from 37 to 42 °C). (E) hiPSC sensory neuron firing rates were insensitive to human recombinant IL-6, TNF- α , and bradykinin (Brady), but were significantly increased by addition of 100 nM capsaicin.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 4:
 (A) Schematic of PDMS-based compartmentalized culture device. (B) Phase image of DRG sensory neuron culture (day 4) extending processes from the soma into the axonal compartment. White arrows indicate examples of cell soma. (C) β -III tubulin staining of axonal compartment highlights axon density by day 8. Scale bar represents 100 μ m.