Published in final edited form as:

Methods Mol Biol. 2019 ; 1882: 309–320. doi:10.1007/978-1-4939-8879-2_27.

Orthotopic Pancreatic Tumor Mouse Models of Liver Metastasis

George Zhang¹, Yi-Chieh Nancy Du¹

¹Department of Pathology and Laboratory Medicine, Weill Cornell Medicine, New York, USA

Abstract

The survival from pancreatic cancer is poor because most patients are diagnosed after the cancer has metastasized. Liver is the most common site of pancreatic cancer metastasis. Orthotopic mouse models of liver metastasis by intrasplenically injecting the pancreatic tumor cells are useful in studying the molecular mechanisms of metastasis and evaluating therapeutic regimens.

Keywords

Metastasis; Pancreatic cancer; Liver; Intrasplenic injection; Splenectomy; Orthotopic; Mouse model

1 Introduction

Pancreatic cancer is one of the leading causes of cancer related deaths, not only for its aggressiveness but also for its difficulty of detection. Pancreatic cancers can be divided into two main groups, exocrine tumors and pancreatic neuroendocrine tumors (PanNETs). The majority of human pancreatic cancer is pancreatic ductal adenocarcinoma (PDAC), one type of pancreatic exocrine tumors. PanNETs are uncommon, but they are rapidly rising in incidence in recent years [1].

Several genetic mutations contributing to the development and progression of PDAC and PanNETs have been identified [2–4]. However, this knowledge has not improved the survival rates of patients with pancreatic cancer much. The challenge is that pancreatic cancer has often metastasized by the time of diagnosis, with the liver being the most common site of metastasis. Once a metastatic disease is diagnosed, patients can no longer be cured with surgical resection. Most cancer patients do not die from locally confined cancer, but rather from the metastatic spread, leading to diminished function of vital organs such as the liver. Thus, animal modeling of pancreatic cancer metastasis remains in huge demand in order to better understand the molecular mechanism of pancreatic cancer metastasis, establish practical approaches, and test novel therapeutic compounds.

Over the years, many mouse models of pancreatic cancer have been developed, including genetically engineered mouse models and xenograft models [5]. In genetically engineered mouse models of PDAC and PanNETs, penetrance of spontaneous metastasis is highly variable and latency is long [6–9]. Most heterotopic pancreatic cancer xenograft models are generated by subcutaneously injecting cultured human tumor cells into the skin of immunodeficient mice. However, subcutaneous xenograft tumors rarely metastasize and usually do not display the signs and symptoms that are caused by the consequences of

significant tumor burden in the liver. In order to overcome these obstacles in studying liver metastasis of pancreatic cancer, this chapter describes a method to establish an orthotopic model of liver metastasis by intrasplenically injecting the pancreatic tumor cells into mice.

The formation of detectable metastasis is the end result of a series of stochastic events that includes steps of tumor cell migration, local invasion, entry into the circulation, arrest at secondary sites, extravasation, and colonization [10]. This orthotopic model of liver metastasis starts with introducing single tumor cell suspension into mice, which bypasses early steps of metastasis.

For orthotopic human pancreatic cancer models, immunodeficient mouse hosts need to be used. The NOD.*Cg-Prkdcscid Il2rgtm1 Wjl/SzJ* (NSG) strain is recommended because it is one of the most immunodeficient mouse strains described to date [11, 12]. If mouse pancreatic tumor cell lines are used, syngeneic mouse strains could also be used to establish liver metastasis via the protocol described here, a hemispleen injection method [13], or tail vein injection [14]. A syngeneic mouse model is immunocompetent, which allows studying the mechanisms of metastasis and cancer therapies in the presence of a functional immune system. However, there can be difficulty in interpreting and predicting how a mouse immune response translates to human immunity.

2 Materials

2.1 Cell Preparation

1. Glass pipettes.
2. PBS without calcium or magnesium.
3. 0.25% trypsin solution mixed with 2.21 mM EDTA.
4. Serum-free media.
5. 50 mL conical tube.
6. Temperature controlled centrifuge that can reach at least $3000 \times g$ at 4 °C.
7. DNase I (stock 1 mg/mL in PBS without calcium or magnesium, stored at -20 °C).
8. 40 µm cell strainer.
9. 0.4% trypan blue.
10. Hemacytometer or automatic cell counter.
11. Microscope.

2.2 Tumor Cell Injection and Splenectomy

1. Immunodeficient mice or syngeneic mouse strains. The suggested age of mice is above 8 weeks old.
2. Scale.

3. Insulin syringes with 28 gauge needles.
4. Ketamine (100 mg/kg) and xylazine (10 mg/kg) cocktail with sterile water.
5. Eye ointment.
6. Electric clipper.
7. Heating pad.
8. Absorbable underpads.
9. Adhesive tape.
10. Sterile gauze.
11. 10% povidone–iodine or chlorhexidine.
12. 70% ethanol or isopropyl alcohol swabs.
13. Sterile surgical drape.
14. Sterile scalpels.
15. Sterile scissors.
16. Sterile forceps.
17. Sterile cotton swab.
18. Sterile needle holder.
19. Sterile 6–0 absorbable sutures.
20. 9 mm stainless steel wound clips.
21. Wound clip remover.
22. Analgesic. Suggestions: meloxicam (2 mg/kg) and buprenorphine (0.5 mg/kg).

2.3 Postoperative Care

1. Heating pad.
2. Heating lamp.
3. Clean cage supplied with food and water.
4. Analgesic. Suggestions: meloxicam (2 mg/kg) and buprenorphine (0.5 mg/kg).

2.4 Tissue Harvest

1. Scissors.
2. Forceps.
3. 10% buffered formalin in 50 mL conical tube.
4. Ruler.
5. 70% ethanol solution.

6. Dry ice or liquid nitrogen (optional).

3 Methods

3.1 Cell Preparation

1. In a tissue culture plate or flask, grow the number of cells desired for injection (see Note 1).
2. Check to see if the cells are healthy under the microscope. Ensure that the cells are not overconfluent or contaminated.
3. Remove the medium.
4. Wash the cells carefully with 3 mL of PBS without calcium or magnesium per 10-cm plate.
5. Wash the cells carefully with 0.4 mL of 0.25% trypsin per 10-cm plate to remove residual PBS and medium.
6. Add 0.8 mL of 0.25% trypsin per 10-cm plate to the cells.
7. Incubate at 37 °C and 5% CO₂ for about 2–3 min to detach cells. Do not keep the cells in trypsin for extended periods of time.
8. Quickly pipet up and down the cells in trypsin on the plate to detach cells and to break up cell clumps.
9. Add 1 mL serum-free media per 10-cm plate to dilute trypsin.
10. Transfer the cell suspension to a 50 mL conical tube.
11. Wash the plate(s) with 3 mL of PBS per 10-cm plate to get residual cells off the plate and transfer the cells to the same 50 mL conical tube.
12. Spin down the cells in the 50 mL conical tube for 3 min at 4 °C at 3000 × *g*. Caution: Make sure that the tubes are balanced inside the centrifuge to avoid equipment damage.
13. Carefully remove the supernatant and avoid removing the cell pellet at the bottom.
14. Wash the plate(s) with 3 mL of PBS per 10-cm plate to get more cells if necessary and add to the cell pellet in the 50 mL conical tube.
15. Resuspend the cell pellet in PBS either by vortexing quickly or pipetting up and down in PBS.
16. Spin down the cells again for 3 min at 4 °C at 3000 × *g* and carefully remove the supernatant.
17. Resuspend the cells in 0.5–1 mL of PBS. Avoid diluting the cell too much because the maximum volume for injection is 100 µL.
18. If cells are clumpy, add DNase I to the cells suspension for a final DNase I concentration of 0.1 mg/mL.

19. Pass the cells through a 40 μm cell strainer to eliminate any large clumps by holding the pipette in a vertical position with the pipette tip touching the membrane of the cell strainer, and slowly press down the cells (*see Note 2*).
20. Prepare an aliquot of cells for counting by the following.
 - a. Mix 10 μL of 0.4% trypan blue with 10 μL of cells in PBS (*see Note 3*).
 - b. Add 10 μL of the mixture to a hemacytometer and count the cells manually under a microscope or using an automatic cell counter.
21. Dilute cultured tumor cells in PBS to your desired concentration. A good starting point is 1×10^7 cells/mL.

All procedures involving mice must be approved by the Institutional Animal Care and Use Committee. The individual(s) carrying out animal experiments must be listed on the approved protocol.

3.2 Tumor Cell Injection and Splenectomy

1. Weigh the mouse on a scale. Draw up enough ketamine–xylazine cocktail (dosage: 0.1 mL/1 g body weight) into an insulin syringe.
2. Use the two-handed method to restrain the mouse. Using your dominant hand, lift the mouse by the base of the tail and place it on the wired lid of the cage. With the other hand, pick up the mouse by the scruff of the neck. Do not grasp the scruff too tightly, which may cause respiratory arrest. Hold the tail with the same hand holding the mouse.
3. Wipe the lower abdominal area of the mouse with either 70% ethanol or isopropyl alcohol swabs for injections in **steps 4 and 5**.
4. Intraperitoneally inject the ketamine–xylazine cocktail from **step 1** into the mouse.
5. Subcutaneously inject 2 mg/kg of meloxicam and 0.5 mg/kg buprenorphine as preemptive analgesia.
6. Apply eye ointment to both eyes to prevent corneal drying of the eyes.
7. Carefully remove the fur covering the left lateral abdominal wall with an electric clipper exposing an area of skin approximately 150% larger than the area of the planned incision (approximately 0.5–2 cm in length). Remove any loose fur with a moist gauze pad.
8. Perform a toe pinch withdrawal reflex test to confirm that the mouse is fully anesthetized. The duration of the anesthesia is about 30 min. Continue to monitor the depth of anesthesia throughout the entire length of the procedure. If needed, redosing 1/3–1/2 of the original dose of the cocktail can prolong anesthesia.
9. Place the mouse in right lateral recumbency and tape the mouse down onto an absorbable underpad on top of a heating pad with adhesive tape to minimize movement of the mouse.

10. Prepare the shaved area for surgery by using a gauze to scrub the area with either a 10% povidone–iodine or a chlorhexidine scrub first, and then with either 70% ethanol or isopropyl alcohol swabs.
11. Repeat the alternating scrubs two more times to ensure the area is properly sterilized.
12. Place a sterile surgical drape over the mouse. A ~2 cm² hole is cut into the drape over the intended incision site.
13. Using a sterile scalpel blade or a pair of sterile mouse surgical scissors, make a 0.5–2 cm parasagittal incision in the left sub-costal of the mouse, revealing the abdominal muscles.
14. Using sterile scissors, make a 0.5–2 cm parasagittal incision through the abdominal musculature over the spleen.
15. Exteriorize the spleen through the incision by gentle retraction with sterile forceps and hold the spleen in place with a sterile cotton-tipped applicator or placed on a sterile moist gauze pad.
16. Mix the tumor cells well using a pipette or a sterile insulin syringe (*see* Note 4). Then, draw up the desired amount of tumor cells in a maximum volume of 100 µL into an insulin syringe for injection.
17. Insert the needle, bevel up, beneath the splenic capsule (Fig. 1a). The needle should be oriented parallel to the spleen. Avoid having to puncture the spleen more than once; other-wise, the cell solution will leak out of the punctured wounds.
18. Slowly inject the cells into the splenic parenchyma, under the splenic capsule. The spleen should turn bright red during the injection (Fig. 1b).
19. Remove the needle and gently apply pressure at the site of injection with a sterile cotton swab for 3–5 min or until hemostasis.
20. 10 min after tumor injection, perform splenectomy. Identify the splenic artery and vein at the splenic hilum, and ligate both with a sterile 6–0 absorbable suture by the following: First, make a loop around the splenic artery and vein (Fig. 2a, b). Next, make two surgical knots to properly ligate the blood supply to the spleen. For each surgical knot, make two loops using the longer end around a needle holder. Using the needle holder, pull the shorter end of the suture through the double loop.
21. Trim the ends of the suture threads to approximately 0.5 cm in length. With a pair of sterile scissors, transect the ligated splenic artery and vein. Dissect the mesentery and connective tissue from the spleen (Fig. 2c). Finally, remove the spleen (Fig. 2d). This is to prevent the growth of tumors in the spleen (*see* Note 5).
22. Check the abdomen for bleeding. If bleeding is found, apply gentle pressure using a sterile cotton swab or sterile gauze until hemostasis is achieved.

23. Insert the suture needle into the abdominal musculature at a 9° angle, 1–2 mm from the wound. The needle should then exit from the opposite edge, equidistant from the wound edge, and opposite from the side of insertion.
24. Pull the thread through the abdominal musculature until 3 cm of the thread remains on the entry site.
25. Make a surgical knot. Begin by holding the needle holder parallel to the open wound. Wrap the needle end of the suture around the needle holder twice (Fig. 3a). Orient the needle holder perpendicular to the wound opening, pointed toward the entry site (Fig. 3b). Pull the 3 cm thread remaining on the entry site through the double loop around the needle holder. Pull the needle end of the suture toward the entry side (Fig. 3c).
26. Repeat **steps 23–25**, alternating the side to pull the ends of the suture.
27. Trim the sutures ends, leaving 1 cm remaining from each end. Make sure the knots are not directly on the edge of the wound.
28. Repeat **steps 23–27** down the peritoneum, lengthwise. About 0.5 cm of the suture threads can be left between each suturing site.
29. For closure of the skin, the quickest method is to apply 2–3 sterile wound clips perpendicular to the wound edge (see Note 6). If wound clips are not available, close the skin with sutures by following **steps 23–28** (Fig. 4). Make sure that the skin is everted and the tension is not too high to impede blood flow. If using sutures, ensure the knots are not directly on the wound closure.
30. Administer meloxicam or buprenorphine subcutaneously.

3.3 Postoperative Care

1. Keep the mouse warm on a heating pad. Gently rotate the mouse side to side every 15 min until it regains the ability to maintain sternal recumbency. Monitor the mouse for the return of normal breathing, blinking patterns, and the beginning of ambulation.
2. Return the mouse to standard housing after the mouse has regained the ability to achieve and maintain sternal recumbency. Place 2–4 moistened feed pellets on the bottom of cage to make the food easily accessible.
3. Place a completed surgery card on the cage.
4. Evaluate the incision site 30 min after the procedure and once more at the end of the day.
5. Administer buprenorphine (every 24 h) or meloxicam (every 4–12 h) subcutaneously within 48 h postoperation.
6. If the mouse displays any signs of distress (hunched posture, ruffled coat, lack of appetite, and/or dehydration) any time during postoperative care, warm the mouse under a heat lamp and administer 1 mL of warm sterile saline (0.9% normal saline solution) intraperitoneally.

7. The day after the surgery, continue to monitor the status of the incision wound as well as for any signs of distress. Buprenorphine or meloxicam should continue to be administered subcutaneously on this day.
8. Monitor the health of the mouse daily.
9. If wound clips are used, remove them after 7–14 days with a wound clip remover.

3.4 Tissue Harvesting

1. Depending on the cell lines, the mouse may exhibit symptoms associated with tumor development and will require euthanasia. The mouse can be euthanized by CO₂ inhalation using a flow rate of 3 L/min for 3 min at the endpoint (*see Note 1*).
2. Open the abdomen with a pair of scissors and remove the liver (Fig. 5).
3. Weigh the liver, count the number of tumor nodules, and measure the width and length of each tumor nodule. Tumor burden is calculated as volume [mm³] = $0.52 \times \text{width}^2 \times \text{length}$. The width should not be bigger than the length.
4. If necessary, tumor nodules can be separated from the liver and saved for other analysis such as reverse transcription polymerase chain reaction (RT-PCR) and Western blotting. In these cases, tumors should be frozen down in small pieces immediately in dry ice or snap-frozen in liquid nitrogen and stored at –80 °C before further processing.
5. Place the liver in 10% buffered formalin for fixation, and rock at room temperature overnight.
6. Remove the formalin into an appropriate waste container and store the liver in 70% ethanol at room temperature before histological analysis.

4 Notes

1. The number of cells for injection from different cell lines is determined based on pilot experiments. Most cells will die in the first few days after injection into the mouse. Thus, injecting too few cells will lead to zero to very few tumor nodule formations. For pilot experiments, we suggest injecting 0.5–2 million cells per mouse and euthanizing mice at different time points (3–5 weeks post injection) to optimize the number of cells for injection and endpoints.
2. If cells do not go through the cell strainer, draw the cells through from the other side of the cell strainer using a clean pipette tip.
3. The 10 µL of cells can be diluted by either 1:10 or 1:100 in PBS to decrease the number of cells to be counted manually. This will ensure a more accurate cell counting. For an automatic cell counter, the most accurate counting lies within the range of 1×10^6 – 1×10^7 cells/mL. Make sure to account for this dilution afterward when calculating the original cell concentration.

4. It is important to mix the cells well right before each injection.
5. If the spleen is not removed, tumors can grow both in the spleen and in the liver. The mouse may need to be euthanized due to the tumor burden in the spleen instead of the tumor burden in the liver. Therefore, splenectomy is performed.
6. Wound clips are preferred over sutures for the skin when performing surgery on mice due to the fact that the clips are not easily removed. Mice tend to scratch and/or bite at the surgery site during postoperative period, tearing apart the sutures in the process.

Acknowledgments

We thank Joseph Hyungjoon Na, Bi-Sen Ding, Soyoung Choi, Manasi M. Godbole, and Danny Huang. G.Z. is supported by NIH grant 1R01CA204916. Y.C.N.D. is supported by DOD grant W81XWH-16-1-0619 and NIH grant 1R01CA204916.

References

1. Kunz PL (2015) Carcinoid and neuroendocrine tumors: building on success. *J Clin Oncol* 33(16): 1855–1863. 10.1200/JCO.2014.60.2532 [PubMed: 25918282]
2. Cancer Genome Atlas Research Network. Electronic address aadhe, Cancer Genome Atlas Research N (2017) Integrated genomic characterization of pancreatic ductal adenocarcinoma. *Cancer Cell* 32(2):185–203 e113. 10.1016/j.ccell.2017.07.007 [PubMed: 28810144]
3. Scarpa A, Chang DK, Nones K, Corbo V, Patch AM, Bailey P, Lawlor RT, Johns AL, Miller DK, Mafficini A, Rusev B, Scardoni M, Antonello D, Barbi S, Sikora KO, Cingarlini S, Vicentini C, McKay S, Quinn MC, Bruxner TJ, Christ AN, Harliwong I, Idrisoglu S, McLean S, Nourse C, Nourbakhsh E, Wilson PJ, Anderson MJ, Fink JL, Newell F, Waddell N, Holmes O, Kazakoff SH, Leonard C, Wood S, Xu Q, Nagaraj SH, Amato E, Dalai I, Bersani S, Cataldo I, Dei Tos AP, Capelli P, Davi MV, Landoni L, Malpaga A, Miotto M, Whitehall VL, Leggett BA, Harris JL, Harris J, Jones MD, Humphris J, Chantrell LA, Chin V, Nagrial AM, Pajic M, Scarlett CJ, Pinho A, Rooman I, Toon C, Wu J, Pinese M, Cowley M, Barbour A, Mawson A, Humphrey ES, Colvin EK, Chou A, Lovell JA, Jamieson NB, Duthie F, Gingras MC, Fisher WE, Dagg RA, Lau LM, Lee M, Pickett HA, Reddel RR, Samra JS, Kench JG, Merrett ND, Epari K, Nguyen NQ, Zeps N, Falconi M, Simbolo M, Butturini G, Van Buren G, Partelli S, Fassan M, Australian Pancreatic Cancer Genome I, Khanna KK, Gill AJ, Wheeler DA, Gibbs RA, Musgrove EA, Bassi C, Tortora G, Pederzoli P, Pearson JV, Waddell N, Biankin AV, Grimmond SM (2017) Whole-genome landscape of pancreatic neuroendocrine tumours. *Nature* 543(7643):65–71. 10.1038/nature21063 [PubMed: 28199314]
4. Jiao Y, Shi C, Edil BH, de Wilde RF, Klimstra DS, Maitra A, Schulick RD, Tang LH, Wolfgang CL, Choti MA, Velculescu VE, Diaz LA Jr, Vogelstein B, Kinzler KW, Hruban RH, Papadopoulos N (2011) DAXX/ATRX, MEN1, and mTOR pathway genes are frequently altered in pancreatic neuroendocrine tumors. *Science* 331(6021):1199–1203. 10.1126/science.1200609 [PubMed: 21252315]
5. Qiu W, Su GH (2013) Challenges and advances in mouse modeling for human pancreatic tumorigenesis and metastasis. *Cancer Metastasis Rev* 32(1–2):83–107. 10.1007/s10555-012-9408-2 [PubMed: 23114842]
6. Bardeesy N, Aguirre AJ, Chu GC, Cheng KH, Lopez LV, Hezel AF, Feng B, Brennan C, Weissleder R, Mahmood U, Hanahan D, Redston MS, Chin L, Depinho RA (2006) Both p16(Ink4a) and the p19(Arf)-p53 pathway constrain progression of pancreatic adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A* 103(15):5947–5952. 10.1073/pnas.0601273103 [PubMed: 16585505]
7. Crabtree JS, Scacheri PC, Ward JM, Garrett-Beal L, Emmert-Buck MR, Edgemon KA, Lorang D, Libutti SK, Chandrasekharappa SC, Marx SJ, Spiegel AM, Collins FS (2001) A mouse model of multiple endocrine neoplasia, type 1, develops multiple endocrine tumors. *Proc Natl Acad Sci U S A* 98(3):1118–1123. 10.1073/pnas.98.3.1118 [PubMed: 11158604]

8. Du YC, Lewis BC, Hanahan D, Varmus H (2007) Assessing tumor progression factors by somatic gene transfer into a mouse model: Bcl-xL promotes islet tumor cell invasion. *PLoS Biol* 5(10):e276 10.1371/journal.pbio.0050276 [PubMed: 17941720]
9. Hanahan D (1985) Heritable formation of pancreatic beta-cell tumours in transgenic mice expressing recombinant insulin/simian virus 40 oncogenes. *Nature* 315(6015):115–122 [PubMed: 2986015]
10. Vanharanta S, Massague J (2013) Origins of metastatic traits. *Cancer Cell* 24(4):410–421. 10.1016/j.ccr.2013.09.007 [PubMed: 24135279]
11. Ishikawa F, Yasukawa M, Lyons B, Yoshida S, Miyamoto T, Yoshimoto G, Watanabe T, Akashi K, Shultz LD, Harada M (2005) Development of functional human blood and immune systems in NOD/SCID/IL2 receptor {gamma} chain(null) mice. *Blood* 106 (5):1565–1573. 10.1182/blood-2005-02-0516 [PubMed: 15920010]
12. Shultz LD, Lyons BL, Burzenski LM, Gott B, Chen X, Chaleff S, Kotb M, Gillies SD, King M, Mangada J, Greiner DL, Handgretinger R (2005) Human lymphoid and myeloid cell development in NOD/LtSz-scid IL2R gamma null mice engrafted with mobilized human hemopoietic stem cells. *J Immunol* 174(10):6477–6489 [PubMed: 15879151]
13. Soares KC, Foley K, Olino K, Leubner A, Mayo SC, Jain A, Jaffee E, Schulick RD, Yoshimura K, Edil B, Zheng L (2014) A preclinical murine model of hepatic metastases. *J Vis Exp* 91:51677 10.3791/51677
14. Du YC, Chou CK, Klimstra DS, Varmus H (2011) Receptor for hyaluronan-mediated motility isoform B promotes liver metastasis in a mouse model of multistep tumorigenesis and a tail vein assay for metastasis. *Proc Natl Acad Sci U S A* 108(40):16753–16758. 10.1073/pnas.1114022108 [PubMed: 21940500]

Fig. 1. Intrasplenic injection. **(a)** Insertion of the needle nearly parallel to the spleen. **(b)** As cells are being injected, the spleen will turn bright red

Fig. 2. Ligation of the splenic vessels and transection of the spleen. **(a)** Making a loop with the suture. **(b)** Securing the loop around the spleen to ligate the splenic artery and vein. **(c)** Separation of the spleen from the underlying connective tissues. **(d)** After removing the spleen, there should be minimal to no bleeding if the artery and vein were properly ligated

Fig. 3. Making a surgical knot. **(a)** Making a double loop around a needle holder using the needle end of the suture. **(b)** Rotate the needle holder 90° angle. **(c)** Pull on the end of the suture originating from the entry point of the wound

Fig. 4.
Complete closure of the incision wound

Fig. 5. Liver metastases. **(a)** Liver without metastasis. **(b)** Liver with visible macrometastases. In this example, many small metastases are located at the edge of the liver lobes