Published in final edited form as:

J Quant Spectrosc Radiat Transf. 2018 November ; 220: 119–122. doi:10.1016/j.jqsrt.2018.09.016.

Does orbital angular momentum have effect on laser's scattering by molecular atmosphere?

Wenbo Sun^{1,2,*}, Yongxiang Hu³, Carl Weimer⁴, Weilin Hou⁵, Tsengdar Lee⁶, Gorden Videen^{2,7,8}, Rosemary R. Baize³

¹Science Systems and Applications Inc., Hampton, VA 23666, USA

²Department of Astronomy and Space Science, Kyung Hee University, 1732, Deogyong-daero, Giheung-gu, Yongin-si, Gyeonggi-do 17104, Korea.

³NASA Langley Research Center, Hampton, VA 23681, USA

⁴Ball Aerospace and Technologies Corp, Boulder, CO 80301, USA

⁵United States Naval Research Laboratory, Stennis Space Center, MS 39529, USA.

⁶NASA Headquarters, Washington, DC 20546, USA

⁷Space Science Institute, 4750 Walnut Street, Boulder Suite 205, CO 80301, USA

⁸US Army Research Laboratory RDRL-CIE-S, 2800 Powder Mill Road, Adelphi Maryland 20783, USA

Abstract

Lasers with orbital angular momentum (OAM) have potential applications in communication technology, manipulation of particles, and remote sensing. Because of its unusual light-scattering properties, the OAM laser's interaction with a molecular atmosphere must be studied to ensure that it is not lossy for communication or remote-sensing applications that involve its transmission through an atmospheric environment. In this study, the finite-difference time-domain (FDTD) method [21] is applied to calculate the light scattering of the purely azimuthal (the radial mode number is assumed to be zero) Laguerre-Gaussian (LG) beams with OAM by very small dielectric particles. Not like Lorentz-Mie solutions, the FDTD method can calculate for particles off the central axis of the LG beam. It is found that when the particles are very small, and the topological charge number of the OAM of a laser is not extremely large, the laser's OAM has little effect on the scattering phase function. This suggests that Rayleigh theory can be applied directly to calculate the light scattering by atmospheric molecules. The transmission of a laser beam with OAM in a molecular atmosphere is not different from that of a regular Gaussian beam.

Keywords

Laser; orbital angular momentum; scattering; molecular atmosphere

*Mail Stop 420, NASA Langley Research Center, Hampton, VA 23681, USA wenbo.sun-1@nasa.gov.

An electromagnetic (EM) wave can carry two forms of angular momentum around the propagation direction: Light spin angular momentum (SAM) related to polarized electromagnetic field vectors spinning around the EM-beam axis and the orbital angular momentum (OAM) related to the spatial field distribution (wave-front helical shape) of light [1–9]. Light's OAM can be generated by propagating a beam through a spiral phase plate [3], diffracting on a fork-like or pitchfork hologram [4–7], applying a q-plate with a SAM sign-change [8, 9], or converting a Hermite-Gaussian beam into a Laguerre-Gaussian beam by using two cylindrical lenses [2]. Due to the nature of the spatial phase distribution, OAM beams have potential applications in manipulation of particles in optical tweezers [10], high-bandwidth information encoding in optical communications [11], high-dimensional quantum information encoding [12–14], and optical detection [15, 16]. The interaction of OAM laser beams with particles has been studied both experimentally [17] and theoretically [17–21]. Several previous studies have used a Lorenz-Mie solution to calculate the light scattering of an OAM laser beam [17–20]. With this type of analytical solution, it is difficult to incorporate an off-axis beam. Indeed, these preliminary theoretical studies only consider the on-axis case. For the OAM laser beam, the on-axis is a special situation in which we would expect the OAM effect on light scattering would be maximized. It is impossible to extrapolate meaning to the off-axis case. For instance, Rury and Freeling [18] report that angular momentum induces transparency through particles; however, they consider a small particle positioned at the central axis of an OAM laser where the intensity is at a minimum. In this case, because its size is much smaller than the laser doughnut's radius, the tiny particle has nearly no interaction with the laser. As soon as the particle is moved off axis, the situation is changed drastically [21]. In Sun et al. [21], the finite-difference time domain (FDTD) method [22–24] with a convolutional perfectly matched layer (CPML) [25] is applied to calculate the scattering of the purely azimuthal (the radial mode number p is assumed to be zero) Laguerre-Gaussian (LG) beams [21] with the OAM (i.e. topological charge number $L > 0$) by small dielectric particles. It's found that for OAM beam's interaction with dielectric particles, the forward peak in the conventional phase function (P_{11}) is reduced, and the light-scattering peak occurs at side-scattering angles, depending on particle sizes. The reduction of forward-scattering peak means that, in laser communications most of the particle-scattered noise cannot enter the receiver, thus the received light is optimally the original OAM-encoded signal. This feature of the OAM beam also implies that in lidar/radar remote sensing of atmospheric particles, most of the multiple-scattering energy will be away from the lidar/radar sensors, and this may result in an accurate profiling of particle layers in the atmosphere or in the oceans, or even in the ground when a ground penetrating radar (GPR) with OAM is applied [21]. With such unusual light-scattering properties, it is imperative to consider the effect of molecular scattering in the atmosphere. If such scattering using an OAM proves to be excessively lossy, it reduces the usefulness of the OAM laser in optical communication and remote-sensing applications. In this study, we used the same method as reported in [21] to study the scattering of the OAM laser by dielectric particles much smaller than the incident wavelength to investigate whether an OAM laser's scattering by a molecular atmosphere is different from Rayleigh scattering.

Figure 1 illustrates a particle positioned at the center of the FDTD computational domain and a laser beam with its central axis at a distance of d from the center of the computational

domain. The FDTD computational domain is bounded by the CPML [21, 24, 25], and the spatial cell size is set at $\Delta s = \lambda/200$, where λ is the incident wavelength [21–25]. The CPML is set to be $6\Delta s$ thick and a free space of $6\Delta s$ is set between the particle surface and the CPML inner surface. The time step for the FDTD simulation is set to be $\Delta t = \Delta s/(2c)$, where c is light speed in free space [21–24].

Figure 2 shows phase function P_{11} of spherical particles with a refractive index of 1.33 and size parameter $x = \pi D/\lambda = 0.48, 0.24, 0.12$, and 0.06 , where D denotes the diameter of the particle, under the incidence of Laguerre-Gaussian beams [21] with $p = 0$ and topological charge numbers $L = 0, 6, 48$, corresponding to Gaussian and Laguerre-Gaussian beams, respectively. The laser waist $w_0 = 400\Delta s$. The distance of the central axis of the beam from the particle center is $d = 200\Delta s$. We can see that the light-scattering phase function of the OAM Laguerre-Gaussian beam ($L = 6$) is nearly coincident with that from the Gaussian beam ($L = 0$) for $x < 0.48$. When $x = 0.48$, the OAM ($L = 6$) scattering is slightly weaker in the forward direction and stronger in the backward direction than the Gaussian beam ($L = 0$). When $L = 48$, the effect of the OAM on the light scattering becomes more significant for larger particles. However, when $x = 0.06$, this effect also becomes insignificant.

As illustrated in Fig. 3, the effect of the OAM on light scattering by particles is due to the phase change incurred by the OAM inside the particle, which is determined by a function $f(LD/d)$, where L is the OAM topological charge number, D is the particle size, and d is the distance between the particle center and the central axis of the OAM laser beam. LD/d is approximately the phase change in the range of the particle due to OAM. However f is also a function of particle's refractive index and shape. For beams of the same L , when d increases or D decreases, LD/d decreases and effect of the OAM on light scattering becomes smaller. This means that when particles are very small with respect to the wavelength, thus within the particle the OAM-incurred phase difference is also very small, the light scattering of the OAM laser beam is not different from that of a Gaussian beam or plane-wave incidence. As illustrated in the final panel of Fig. 2, when a particle is smaller than about $1/50$ of the wavelength ($x \sim 0.06$), its light scattering is very close to Rayleigh scattering. For a particle of $x = 0.06$, we found that when OAM $L < 480$, the scattering is not significantly different from Rayleigh scattering. When OAM $L > 480$, there is very small in the forward scattering of P_{11} . Thus we can conclude that an OAM of $L < 1000$ should have a negligible effect on the light scattering of atmospheric molecules ($x \ll 0.06$).

In this study, the FDTD method is used to calculate the scattering of a purely azimuthal Laguerre-Gaussian beams having OAM by small dielectric particles. It is found that when the particles are very small, and the order of the OAM of a laser is not extremely high, the laser's OAM has little effect on the scattering phase function. This implies that Rayleigh theory can be directly applied to calculate the light scattering of an OAM laser beam by atmospheric molecules, and the transmission of an OAM laser beam through the atmosphere is not different from that of a Gaussian beam. Note here that, although we set a specific distance of the central axis of the beam from the particle center ($d = 200\Delta s$) in this study, this does not affect the conclusion.

Acknowledgment

This work is supported by NASA ESTO ACT project “Lidar Orbital Angular Momentum Sensor”.

References

1. Hammond RT. Lorentz covariant and gauge invariant description of orbital and spin angular momentum and the non-symmetric energy momentum tensor. arXiv:1706.07665 [physics.optics] 2017.
2. Allen L, Beijersbergen MW, Spreeuw RJC, Woerdman JP. Orbital angular momentum of light and the transformation of Laguerre-Gaussian laser modes. Phys Rev A 1992; 45: 8185–9. [PubMed: 9906912]
3. Beijersbergen MW. Helical-wavefront laser beams produced with a spiral phase plate. Optics Communication 1994; 112: 321–7, doi:10.1016/0030-4018(94)90638-6.
4. Bazhenov VY, Vasnetsov MV, Soskin MS. Laser beams with screw dislocations in their wavefronts. JETP Letters 1990; 52: 429–31.
5. Bazhenov VY, Soskin MS, Vasnetsov MV. Screw dislocations in light wavefronts. J Modern Opt 1992; 39: 985–90, doi: 10.1080/09500349214551011.
6. Heckenberg NR, McDuff R, Smith CP, Rubinsztein-Dunlop H, Wegener MJ. Laser beams with phase singularities. Optical and Quantum Electronics 1992; 24: S951–62, doi:10.1007/BF01588597.
7. Soskin MS, Gorshkov V, Vasnetsov MV, Malos J, Heckenberg N. Topological charge and angular momentum of light beams carrying optical vortices. Phys Rev A 1997; 56: 4064–75, doi:10.1103/PhysRevA.56.4064.
8. Marrucci L, Manzo C, Paparo D. Optical spin-to-orbital angular momentum conversion in inhomogeneous anisotropic media. Phys Rev Lett 2006; 96: 163905, doi:10.1103/PhysRevLett.96.163905. [PubMed: 16712234]
9. Karimi E, Piccirillo B, Nagali E, Marrucci L, Santamato E. Efficient generation and sorting of orbital angular momentum eigenmodes of light by thermally tuned q-plates. Appl Phys Lett 2009; 94: 231124, doi:10.1063/1.3154549.
10. Friese MEJ, Enger J, Rubinsztein-Dunlop H, Heckenberg N. Optical angular-momentum transfer to trapped absorbing particles. Phys Rev A 1996; 54: 1593–6, doi:10.1103/PhysRevA.54.1593. [PubMed: 9913630]
11. Gibson G, Courtial J, Padgett MJ, Vasnetsov M, Pas’Ko V, Barnett SM, Franke-Arnold S. Free-space information transfer using light beams carrying orbital angular momentum. Optics Express 2004; 12: 5448–56. [PubMed: 19484105]
12. Malik M. Influence of atmospheric turbulence on optical communications using orbital angular momentum for encoding. Optics Express 2012; 20: 13195, doi:10.1364/OE.20.013195(2012). [PubMed: 22714347]
13. Boyd RW, Jha A, Malik M, O’Sullivan C, Rodenburg B, Gauthier DJ. Quantum key distribution in a high-dimensional state space: exploiting the transverse degree of freedom of the photon. Proc SPIE Advances in Photonics of Quantum Computing, Memory, and Communication IV 2011; 7948: 79480L, doi:10.1117/12.873491.
14. Barreiro JT, Wei TC, Kwiat PG. Beating the channel capacity limit for linear photonic superdense coding. Nature Physics 2008; 4: 282–6, doi:10.1038/nphys919.
15. Foo G, Palacios DM, Swartzlander GA Jr. Optical vortex coronagraph. Opt Lett 2005; 30: 3308–10. [PubMed: 16389814]
16. Sun W, Hu Y, MacDonnell DG, Weimer C, Baize RR. Technique to separate lidar signal and sunlight. Opt. Express 2016; 24: 12949–54, doi: 10.1364/OE.24.012949. [PubMed: 27410314]
17. Garbin V, Volpe G, Ferrari E, Versluis M, Cojoc D, Petrov D. Mie Scattering distinguishes the topological charge of an optical vortex. New J Phys 2009; 11: 013046.
18. Rury AS, Freeling R. Mie scattering of purely azimuthal Laguerre-Gauss beams: Angular-momentum-induced transparency. Phys Rev A 2012; 86: 053830.

19. Zambrana-Puyalto X, Molina-Terriza G. The role of the angular momentum of light in Mie scattering. Excitation of dielectric spheres with Laguerre–Gaussian modes. *J Quant Spectrosc Radiat Transfer* 2013; 126: 50–5.
20. Kiselev AD, Plutenko DO. Mie scattering of Laguerre-Gaussian beams: Photonic nanojets and near-field optical vortices. *Phys Rev A* 2014; 89: 043803
21. Sun W, Hu Y, Weimer C, Ayers K, Baize RR, Lee T. A FDTD solution of scattering of laser beam with orbital angular momentum by dielectric particles: Far-field characteristics. *J Quant Spectrosc Radiat Transfer* 2017; 188: 200–13, doi:10.1016/j.jqsrt.2016.02.006.
22. Sun W, Fu Q, Chen Z. Finite-difference time domain solution of light scattering by dielectric particles with a perfectly matched layer absorbing boundary condition. *Appl Opt* 1999; 38: 3141–51. [PubMed: 18319902]
23. Sun W, Loeb NG, Fu Q. Finite-difference time domain solution of light scattering and absorption by particles in an absorbing medium. *Appl Opt* 2002; 41: 5728–43. [PubMed: 12269573]
24. Sun W, Videen G, Fu Q, Hu Y. Scattered-field FDTD and PSTD algorithms with CPML absorbing boundary conditions for light scattering by aerosols. *J Quant Spectrosc Radiat Transfer* 2013; 131: 166–74.
25. Roden JA, Gedney SD. Convolutional PML (CPML): An efficient FDTD implementation of the CFS-PML for arbitrary media. *Microwave Opt Tech Lett* 2000; 27: 334–9.

Highlights

1. Scattering of laser with orbital angular momentum (OAM) by small particles is studied.
2. The 3D CPML FDTD technique is applied in the solution of OAM laser beam scattering.
3. The OAM has no effect on laser's scattering by molecular atmosphere.

Figure 1.

Illustration of a particle positioned at the center of the FDTD computational domain and a laser beam with its central axis at a distance d from the center of the computational domain. The FDTD computational domain is bounded by the CPML [21].

Figure 2.

Phase function P_{11} of spherical particles with a refractive index of 1.33 and size parameter $x = 0.48, 0.24, 0.12$, and 0.06 , respectively, illuminated by a Laguerre-Gaussian beams with $p = 0$. The laser waist $w_0 = 400 \text{ } s$, where $s = \lambda / 200$. The laser OAM topological charge numbers are $L = 0, 6$, and 48 . The distance of the central axis of the beam from the particle center is $d = 200 \text{ } s$.

Figure 3. Illustration of a cross cut of a particle and an incident OAM laser beam. In this figure, D denotes particle size, d is the distance from the beam's central axis to the center of the particle.