Published in final edited form as:

Semin Immunol. 2014 August ; 26(4): 315–320. doi:10.1016/j.smim.2014.04.007.

Inflammation and Wound Repair

Danny C. LeBert¹, Anna Huttenlocher^{2,3}

¹Cellular and Molecular Pathology Graduate Program, University of Wisconsin-Madison, Madison, WI 53706

²Department of Pediatrics, University of Wisconsin-Madison, Madison, WI 53706

³Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI 53706

Abstract

Wound repair requires the integration of complex cellular networks to restore tissue homeostasis. Defects in wound repair are associated with human diseases including pyoderma gangrenosum, a heterogeneous disorder that is characterized by unhealed wounds and chronic inflammation of unclear etiology. Despite its clinical importance, there remain significant gaps in understanding how different types of cells communicate to integrate inflammation and wound repair. Recent progress in wound and regenerative biology has been gained by studying genetically tractable model organisms, like zebrafish, that retain the ability to regenerate. The optical transparency and ease of genetic manipulation make zebrafish an ideal model system to dissect multi-cellular and tissue level interactions during wound repair. The focus of this review is on recent advances in understanding how inflammation and wound repair are orchestrated and integrated to achieve wound resolution and tissue regeneration using zebrafish.

Keywords

Inflammation; Chronic wounds; Wound repair; Regeneration

1. Introduction

An inability to repair wounded tissue is a major clinical problem. The response to tissue damage is different depending on the tissue type and the severity of damage (1). Some human tissues retain the ability to regenerate, like the human liver (2). Most tissues, however, have limited capacity for regeneration and the outcome of tissue damage is often scarring (1). Scarring can have detrimental effects in tissues like the heart where it leads to congestive heart failure (3–5). Tissues such as the central nervous system also have limited regeneration after injury and display poor recovery of function (6, 7). Therefore, there is considerable interest in understanding how to improve functional outcomes in human tissues that fail to regenerate or result in scar formation after tissue damage. To understand tissue repair, there has been an interest in studying vertebrate models in which wound healing

occurs with minimal scarring in an attempt to understand how to optimize healing in humans. Herein we will focus on recent studies that have used zebrafish to study wound repair and regeneration. For general reviews on wound biology, we refer the readers to many outstanding general reviews (8–12).

2. Impaired wound healing and pyoderma gangrenosum

A useful model to understand tissue repair is wound healing of the skin. In response to tissue damage, the skin is repaired through a sequence of steps that involves the interactions between different types of cells including leukocytes, blood cells, fibroblasts and epithelial cells (13, 14). The complex process of cutaneous wound repair includes distinct and often overlapping steps including the formation of a blood clot to re-establish tissue homeostasis, inflammation, re-epithelialization, granulation tissue formation and finally remodeling with the potential for scar formation (15). A disruption of these steps can lead to chronic inflammation and impaired wound healing.

There are many human diseases characterized by impaired wound healing and chronic skin ulcers including diabetes (16). In addition, inherited human diseases including immunodeficiencies like Leukocyte Adhesion Deficiency (LAD) are associated with impaired wound healing due to persistent infection and have been successfully recapitulated in the zebrafish (17, 18). There are also heterogeneous disorders characterized by non-resolving skin lesions in the absence of infection. Pyoderma gangrenosum (PG) is a rare neutrophilic dermatosis that is characterized by ulcer formation and chronic non-resolving wounds even in the absence of infection (19). PG is often associated with other underlying conditions including inflammatory bowel disease, but in many cases the etiology is unclear. There have been single gene autoinflammatory diseases associated with PG, including the autosomal dominant disorder pyogenic arthritis, pyoderma gangrenosum and acne (PAPA) syndrome (20). Other autoinflammatory disorders are also associated with unhealed wounds and pustules including early onset inflammatory bowel disease and caspase recruitment domain-containing protein 14 (CARD14) mediated psoriasis (21).

Understanding these rare autoinflammatory diseases can contribute to progress in dissecting underlying mechanisms that contribute to wound repair. For example, in the case of PAPA syndrome the prevailing view is that wounds do not heal because of persistent inflammation within the skin, due to elevated interleukin-1 beta (IL-1 β) and tumor necrosis factor alpha (TNF- α) production (20, 22–24). This is likely an important factor that contributes to the pathogenesis of PAPA syndrome. However, treatment with anti-inflammatory drugs that target IL-1 β or TNF α do not always improve disease symptoms (20). This raises the intriguing idea that other factors may contribute to chronic tissue damage in patients with PAPA syndrome. In support of this idea, a recent report using human cells suggests that macrophages may play a driving role in some forms of PAPA syndrome. In particular, a novel mutation in the SH3 domain of the adaptor protein Proline-Serine-Threonine Phosphatase-Interacting Protein 1 (PSTPIP1) in macrophages induces altered organization of the actin cytoskeleton and exaggerated release of matrix metalloproteinases (MMPs) *in vitro* (25). These changes may contribute to the tissue destruction and the development of chronic ulcers characteristic of PAPA syndrome. This raises an interesting question about

alternative drug targets like MMP inhibition or drugs that target cytoskeletal organization. Further studies will be needed to determine if MMP inhibition or alternative therapies will benefit patients with PAPA syndrome or other forms of PG. Single gene disorders such as mutations in PSTPIP1 represent a unique opportunity to dissect specific pathways involved in inflammation and wound repair. They also highlight the need for improved animal models, like zebrafish, to aid in understanding the pathogenesis and treatment of wound repair disorders.

3. Continuum from wound to resolution and repair

A fundamental question in wound biology involves an understanding of the relationship between inflammation and wound repair. It is generally believed that if inflammation does not resolve, wound healing and regeneration will not occur (26). This is particularly interesting in the context of the central nervous system (CNS), a tissue that exhibits limited recovery and regeneration after damage. In a recent review, Shechter and Schwartz propose that CNS injury may be similar to a chronic and unhealed wound (27). They go on to suggest that regeneration may be limited in the CNS because of impaired wound healing and view CNS damage as a type of chronic wound (27). Therefore, an important component of regeneration research is to understand the steps that occur during wound healing, including the links between inflammation, repair and regeneration.

Wound resolution occurs when the damaged tissue seals and generally coincides with the resolution of inflammation (28). Regeneration, on the other hand, is defined as the reconstitution of the damaged tissue in a scar-free manner that occurs after the wound resolves, representing a final stage in the repair process (15, 29). This distinction raises interesting questions: what are the signaling networks responsible for driving the healing process towards regeneration, and why are some organisms better at integrating these signals to regenerate tissues? Why do some organisms maintain the ability to regenerate tissues throughout their lifespan and how does inflammation and early signaling at damaged tissue influence the repair process? Moreover, does differential integration of inflammatory signaling drive the outcome toward chronic wound versus resolution and regeneration?

The wound repair process is highly conserved across single and multi-cellular organisms (30–32). This conservation positions simple model systems as important tools to understand how diverse cell populations integrate wound responses and to determine the kinetics of these interactions. It is important to consider that many organisms, including vertebrate genetic model systems like zebrafish, retain the ability to regenerate tissues without permanent scar formation (15, 33–36). It is possible that by understanding why some organisms perform scar-free regeneration, we will develop new insight into human wound biology. The zebrafish represents a powerful model system to dissect the wound repair process in both the larval period, where imaging is optimized, and in adult zebrafish, where many steps of the wound repair process are conserved (36, 37).

4. Zebrafish as a model to understand wound repair

The development of transgenic zebrafish lines with fluorescently labeled neutrophils, and macrophages, in addition to reporter lines that label specific tissues including epithelial tissues, has revolutionized our ability to dissect the molecular mechanisms that regulate inflammation and wound healing in zebrafish (38–48). The innate immune response to infection and tissue damage seems to be highly conserved in zebrafish, making it an ideal system to study inflammation and wound repair (41, 42, 46, 47, 49–55). Support for the conservation of innate immune functions can be found in the recapitulation of human immunodeficiency phenotypes in zebrafish models of Wiskott-Aldrich syndrome (WAS), warts-hypogammaglobulinemia-infections-myelokathexis (WHIM) syndrome, and leukocyte adhesion deficiency (LAD)-like syndrome (18, 56, 57). Moreover, zebrafish have retained the ability to regenerate many tissues, including the fins, heart, skin, photoreceptors, neurons, and the brain (36, 37, 58–63). Adult zebrafish also retain regenerative potential and have provided new insight into regeneration of cardiac tissue, kidney and the CNS (60, 61, 64–68).

Use of tools to analyze spatial and temporal changes in gene expression after injury in zebrafish has supported a central role for changes in inflammatory gene expression during zebrafish regeneration (69, 70). Moreover, transcriptome analysis during spinal cord regeneration has been used to uncover a potential role for signal transducer and activator of transcription 3 (Stat3) in orchestrating both inflammation and proliferation after injury (70). Finally, the potential of chemical tools to uncover mechanisms of regeneration in zebrafish provides an additional strength of the zebrafish model in wound biology research (71). The combination of the optical transparency, conservation of the innate immune system and genetic tractability make zebrafish an ideal model to dissect mechanisms that integrate inflammation and tissue repair.

5. Wound-induced inflammation

At the beginning of the 20th century Ilya Mechnikov first reported the response of inflammatory cells to wounds in starfish larvae. In his Nobel Prize lecture, he described the appearance of small moving “elements” that rapidly responded to wounds (72). In the years following his initial observation, significant progress has been made in characterizing the nature of these elements as innate immune cells which include neutrophils and macrophages. Neutrophils are the most abundant leukocyte, and are generally the first responders to infection and tissue damage (73). This infiltration of leukocytes is necessary to limit infection at the site of tissue damage (74). Moreover, the response of macrophages is necessary for efficient wound repair by clearing debris from sites of tissue damage (75).

There is substantial evidence to support the importance of macrophages in the wound healing process in mouse and guinea pig models, and recent studies in zebrafish also showed that depletion of macrophages results in a build-up of debris and impairs regeneration (76–80). However, inflammation can also have detrimental effects on wound repair. For example, in mouse models the presence of neutrophils can contribute to scar formation after tissue damage and slow the wound closure rate (81). In zebrafish, Li and colleagues showed using

runx1 mutants with impaired neutrophil function that larvae have improved regeneration, suggesting that neutrophils in zebrafish larvae may also have detrimental effects on regeneration and wound repair (76). However, in the CNS of adult zebrafish, a recent study suggests that acute CNS inflammation induced by injection of microparticles promotes neurogenesis and regeneration through leukotriene signaling, suggesting complex inflammatory signaling integration is necessary during wound repair (82). Taken together, a first step in understanding the role of inflammation in wound healing is to understand the signals that mediate leukocyte recruitment to wounds.

6. Signals guiding leukocyte recruitment in zebrafish

Leukocytes, both neutrophils and macrophages, are rapidly recruited to wounds in zebrafish larvae (Fig. 1) (41, 42, 73, 83, 84). This has provided a useful system to dissect mechanisms that mediate leukocyte wound attraction. The chemotaxis of leukocytes to sites of injury involves the sensing of a chemotactic gradient, polarization and migration (85). An early signal in zebrafish larvae is a wound-induced gradient of the reactive oxygen species (ROS), hydrogen peroxide (H₂O₂). Niethammer and colleagues showed that wounding induces a rapid and transient burst of H₂O₂ in the wounded epithelium of larval zebrafish, which mediates early neutrophil recruitment (86). Inhibition of epithelial ROS production impairs the initial recruitment of neutrophils to wounded tissue. To uncover mechanisms that function downstream of epithelial ROS production, it was found that the Src family kinase (SFK) Lyn becomes activated and is necessary for early neutrophil wound attraction, suggesting that wound-generated ROS can modify signaling within zebrafish neutrophils (87).

The larval zebrafish has also provided insights into other signaling pathways that mediate neutrophil recruitment to wounds, including a key role for chemokine signaling through interleukin-8 (IL-8) generated at the wound edge (88). Environmental tissue signals also mediate neutrophil recruitment, including changes in osmotic signals induced by wounding. Cellular swelling occurs after wounding and induces activation of cytosolic phospholipase A2 (cPLA2), the production of non-canonical arachidonate metabolites and leukocyte recruitment (89). Finally, wound-induced waves of calcium (Ca²⁺) in the CNS are associated with recruitment of microglial cells, supporting the idea that there are diverse environmental cues that mediate inflammation in response to damage (90). Taken together, these studies highlight the complex changes that occur within tissues that orchestrate leukocyte recruitment and inflammation in response to tissue damage and highlight the importance of the zebrafish model.

7. Resolution of inflammation at wounds

Wound resolution occurs when the wound is sealed and inflammation resolves. In part, the resolution of inflammation occurs when the inciting triggers, including damage signals or infection, are no longer present (85). However, with some diseases like pyoderma gangrenosum, there is impaired wound healing and inflammation does not resolve. This raises the intriguing question: what resolves inflammation? Is this a passive process that occurs when the positive trigger ends? Or are there active processes that mediate resolution

through the release of pro-resolving factors (91)? In mammalian models there has been substantial progress in identifying factors like resolvins and other lipid mediators, that mediate resolution of inflammation, and this has been an area of great interest in terms of its therapeutic potential (91–94). Resolution of innate immune inflammation is generally considered to be orchestrated by macrophages and is characterized by the clearance of apoptotic neutrophils by macrophages (75, 95). There is also evidence that this form of innate immune resolution occurs in zebrafish (41, 47, 76). However, studies of neutrophil behavior in the zebrafish revealed that neutrophil apoptosis at the wound site is a rare event, with less than 3% undergoing apoptosis (96).

Live imaging of transgenic zebrafish larvae has identified a novel mechanism of inflammation resolution by the reverse migration of neutrophils from wounded tissue back toward the vasculature (42). In subsequent studies, using photoconvertible Dendra2, the fate of wound-associated neutrophils was observed and it was found that neutrophils traffic repeatedly between damaged tissue and the vasculature before finally resolving this response by reverse migration away from the wound, often back into the vasculature where they traffic to diverse tissues (97). The role of these sensitized neutrophils in subsequent host immune responses has not yet been defined. Studies in mice also suggest that neutrophils can undergo reverse migration from tissues into the vasculature (98). The cues that mediate this reverse migration are just beginning to be uncovered in zebrafish but include pathways through hypoxia inducible factor-1 alpha (HIF-1 α) (96). There is also evidence that this reverse migration may be targeted using small molecules to optimize inflammation resolution (99).

An important aspect of inflammation resolution is the loss of signals that induce inflammation. An intriguing study by Pase et al., demonstrated that neutrophil myeloperoxidase (MPO) is an important factor that resolves inflammatory signaling by depleting H₂O₂ at the wound (100). The zebrafish larvae, deficient in MPO, have sustained H₂O₂ signaling at the wound and impaired inflammation resolution (100). This finding is interesting because humans who are deficient in MPO can have chronic inflammation (101). These studies highlight that the zebrafish model system has been instrumental in the identification of novel regulators of leukocyte resolution and will certainly be poised to help identify other factors which mediate resolution by attenuating pro-inflammatory signaling at the wound site or alternatively promoting neutrophil phagocytosis or reverse migration.

8. The role of early wound signaling in wound resolution and regeneration

Early signaling at wounds through ROS and calcium are necessary for leukocyte recruitment to tissue damage in epithelium and the CNS (86, 87, 90). But what impact do these early signals have on wound resolution and repair? Recent studies suggest these early wound signals may, in fact, influence the wound healing outcome. For example, tail transection in zebrafish induces early ROS signaling and activation of the SFK Fyn within the epithelium. Inhibition of early ROS or SFK signaling for just 2 hours around the time of wounding impairs late regeneration and resolution of healing 3 days later, suggesting that early signals may orchestrate the subsequent regeneration program (59). However, it is not clear if this defect in regeneration is due to impaired wound resolution or alternatively is due to a direct

link between early signaling and regeneration, independent of the effects on inflammation resolution. Early Ca^{2+} signaling waves in the epithelium and CNS in zebrafish are also associated with wound resolution and regeneration (59, 90). Moreover, a requirement for early wound signaling has been demonstrated during *Xenopus* tadpole tail limb regeneration, indicating a conserved regeneration pathway (102). Future studies will need to continue to unravel both the pathways involved in early wound signals, and their roles in specific cell types in the damaged tissue.

Among the cells that regenerate during caudal-fin regeneration are sensory neurons, which re-establish axon branching. Interestingly, regeneration of these axons occurs more rapidly when the surrounding epithelium has been wounded, supporting a key role for the wound microenvironment (103). This suggests that wounded epithelium release a factor(s) to promote axon regrowth. One such factor appears to be H_2O_2 . Addition of exogenous H_2O_2 to axon ablations in which the surrounding epithelium is not damaged improves axon regeneration. Meanwhile, blocking of H_2O_2 production in axons regenerating in the presence of epithelial injury efficiently reduces the growth-promoting effect of the skin injury (103). The study provides further evidence of inflammatory integration during damage repair, although the downstream effectors of the H_2O_2 were not identified.

There are many putative downstream targets of H_2O_2 that can mediate wound healing, including inactivation of phosphatases or activation of kinases through the oxidation of cysteine residues (104). Our recent studies have implicated Lyn and Fyn as downstream targets that mediate H_2O_2 effects during wound healing (59, 87). To further characterize the effect of H_2O_2 signaling during tissue repair, Gauron et al., performed caudal-fin amputations on adult zebrafish and studied changes to blastema function (105). The blastema is a mass of lineage-restricted progenitors that accumulate underneath the wounded epithelium. The blastema mediates the proliferation and patterning of the regenerating fin (106). Adult zebrafish produce a prolonged burst of H_2O_2 (compared to larval zebrafish) at the site of amputation. In this context, the extended burst of H_2O_2 leads to the activation of c-Jun N-terminal kinase (JNK) and a wave of epidermal apoptosis. Inhibition of the apoptosis or JNK signaling results in a smaller blastema, suggesting that H_2O_2 can influence epidermal cell proliferation through the regulation of apoptotic/JNK signaling (105).

The CNS of the zebrafish shows improved regeneration of neurons compared to mammals. However, not all nerve cell types are easily repaired in zebrafish. For instance, Mauthner cells (M axon) are myelinated neurons that regenerate poorly after spinal lesion (61, 107). The inability to regenerate is thought to be the result of inhibitory molecules in the surrounding environment or alternatively from intrinsic factors. Attempts to identify the intrinsic factors have identified cyclic adenosine monophosphate (cAMP) as a regulator of axonal sprouting *in vitro* (108–110). Acting as a second messenger, cAMP dampens inflammation and thus, inhibits the innate immune response (111). To determine if cAMP could influence the regenerative outcome of the M axon in the larval zebrafish, Bhatt et al., induced spinal lesions and pressure injected membrane-permeable dibutyl-cyclic adenosine monophosphate (db-cAMP) two days after injury. They found that addition of db-cAMP induced regeneration of the M axon, suggesting that cAMP promotes the regeneration of the injured M axon (112). Interestingly, cAMP acts to dampen the innate

immune response, suggesting the improved M axon regeneration may be mediated in part through changes in innate immune inflammation.

9. Future

Advances in imaging, genetic tools and chemical genetics have positioned zebrafish as a powerful tool to dissect the temporal and spatial relationship between inflammation and wound repair. Recent advances have unraveled key pathways that mediate early signaling at wounds and the onset of inflammation using zebrafish. The identification of reverse neutrophil migration as a mechanism that can resolve local inflammation is an example of how the zebrafish model can help to uncover new mechanisms during wound repair by exploiting live imaging to probe temporal and spatial interactions in live zebrafish. Moreover, the idea that acute inflammation can facilitate regeneration in the adult zebrafish brain raises exciting new avenues for investigations into CNS regeneration. Finally, zebrafish are poised to help unravel key pathways by which environmental cues like hypoxia regulate cell reprogramming and tissue repair (113, 114). In the long term, it is this type of information that may help to unravel mechanisms that identify new avenues for treatment of patients with chronic tissue damage.

Acknowledgements

This research was supported by the NIH National Research Service Award (NRSA) T32 GM081061 (to DC LeBert).

References

1. Stroncek JD, Reichert WM. Overview of Wound Healing in Different Tissue Types. In: Reichert WM, editor. *Indwelling Neural Implants: Strategies for Contending with the In Vivo Environment*. Boca Raton (FL); 2008.
2. Chen MF, Hwang TL, Hung CF. Human liver regeneration after major hepatectomy. A study of liver volume by computed tomography. *Annals of surgery*. 1991 3;213(3):227–9. [PubMed: 1998403]
3. Itoh G, Tamura J, Suzuki M, Suzuki Y, Ikeda H, Koike M, et al. DNA fragmentation of human infarcted myocardial cells demonstrated by the nick end labeling method and DNA agarose gel electrophoresis. *The American journal of pathology*. 1995 6;146(6):1325–31. [PubMed: 7778672]
4. Saraste A, Pulkki K, Kallajoki M, Henriksen K, Parvinen M, Voipio-Pulkki LM. Apoptosis in human acute myocardial infarction. *Circulation*. 1997 1 21;95(2):320–3. [PubMed: 9008443]
5. Ertl G, Frantz S. Healing after myocardial infarction. *Cardiovascular research*. 2005 4 1;66(1):22–32. [PubMed: 15769445]
6. Silver J, Miller JH. Regeneration beyond the glial scar. *Nature reviews Neuroscience*. 2004 2;5(2):146–56. [PubMed: 14735117]
7. Zurn AD, Bandtlow CE. Regeneration failure in the CNs: cellular and molecular mechanisms. *Advances in experimental medicine and biology*. 2006;557:54–76. [PubMed: 16955704]
8. Tal TL, Franzosa JA, Tanguay RL. Molecular signaling networks that choreograph epimorphic fin regeneration in zebrafish - a mini-review. *Gerontology*. 2010;56(2):231–40. [PubMed: 19923791]
9. Sonnemann KJ, Bement WM. Wound repair: toward understanding and integration of single-cell and multicellular wound responses. *Annual review of cell and developmental biology*. 2011;27:237–63.
10. Poss KD. Advances in understanding tissue regenerative capacity and mechanisms in animals. *Nature reviews Genetics*. 2010 10;11(10):710–22.
11. Martin P Wound healing--aiming for perfect skin regeneration. *Science*. 1997 4 4;276(5309):75–81. [PubMed: 9082989]

12. Cordeiro JV, Jacinto A. The role of transcription-independent damage signals in the initiation of epithelial wound healing. *Nature reviews Molecular cell biology*. 2013 4;14(4):249–62.
13. Singer AJ, Clark RA. Cutaneous wound healing. *The New England journal of medicine*. 1999 9 2;341(10):738–46. [PubMed: 10471461]
14. Shaw TJ, Martin P. Wound repair at a glance. *Journal of cell science*. 2009 9 15;122(Pt 18):3209–13. [PubMed: 19726630]
15. Gurtner GC, Werner S, Barrandon Y, Longaker MT. Wound repair and regeneration. *Nature*. 2008 5 15;453(7193):314–21. [PubMed: 18480812]
16. Moura LI, Cruz MT, Carvalho E. The effect of neurotensin in human keratinocytes--implication on impaired wound healing in diabetes. *Experimental biology and medicine*. 2014 1;239(1):6–12. [PubMed: 24198343]
17. Fischer A, Lisowska-Grospierre B, Anderson DC, Springer TA. Leukocyte adhesion deficiency: molecular basis and functional consequences. *Immunodeficiency reviews*. 1988;1(1):39–54. [PubMed: 3078709]
18. Deng Q, Yoo SK, Cavnar PJ, Green JM, Huttenlocher A. Dual roles for Rac2 in neutrophil motility and active retention in zebrafish hematopoietic tissue. *Developmental cell*. 2011 10 18;21(4):735–45. [PubMed: 22014524]
19. Powell FC, Schroeter AL, Su WP, Perry HO. Pyoderma gangrenosum: a review of 86 patients. *The Quarterly journal of medicine*. 1985 5;55(217):173–86. [PubMed: 3889978]
20. Smith EJ, Allantaz F, Bennett L, Zhang D, Gao X, Wood G, et al. Clinical, Molecular, and Genetic Characteristics of PAPA Syndrome: A Review. *Current genomics*. 2010 11;11(7):519–27. [PubMed: 21532836]
21. Almeida de Jesus A, Goldbach-Mansky R. Monogenic autoinflammatory diseases: concept and clinical manifestations. *Clinical immunology*. 2013 6;147(3):155–74. [PubMed: 23711932]
22. Shoham NG, Centola M, Mansfield E, Hull KM, Wood G, Wise CA, et al. Pyrin binds the PSTPIP1/CD2BP1 protein, defining familial Mediterranean fever and PAPA syndrome as disorders in the same pathway. *Proceedings of the National Academy of Sciences of the United States of America*. 2003 11 11;100(23):13501–6. [PubMed: 14595024]
23. Edrees AF, Kaplan DL, Abdou NI. Pyogenic arthritis, pyoderma gangrenosum, and acne syndrome (PAPA syndrome) associated with hypogammaglobulinemia and elevated serum tumor necrosis factor-alpha levels. *Journal of clinical rheumatology : practical reports on rheumatic & musculoskeletal diseases*. 2002 10;8(5):273–5. [PubMed: 17041385]
24. Cortis E, De Benedetti F, Insalaco A, Cioschi S, Muratori F, D'Urbano LE, et al. Abnormal production of tumor necrosis factor (TNF) -- alpha and clinical efficacy of the TNF inhibitor etanercept in a patient with PAPA syndrome [corrected]. *The Journal of pediatrics*. 2004 12;145(6):851–5. [PubMed: 15580218]
25. Starnes TW, Bennin DA, Bing X, Eickhoff JC, Grahf DC, Bellak JM, et al. The F-bar protein PSTPIP1 controls extracellular matrix degradation and filopodia formation in macrophages. *Blood*. 2014 1 13.
26. Eming SA, Krieg T, Davidson JM. Inflammation in wound repair: molecular and cellular mechanisms. *The Journal of investigative dermatology*. 2007 3;127(3):514–25. [PubMed: 17299434]
27. Shechter R, Schwartz M. Harnessing monocyte-derived macrophages to control central nervous system pathologies: no longer 'if' but 'how'. *The Journal of pathology*. 2013 1;229(2):332–46. [PubMed: 23007711]
28. Guo S, Dipietro LA. Factors affecting wound healing. *Journal of dental research*. 2010 3;89(3):219–29. [PubMed: 20139336]
29. Needham AE. Regeneration and wound-healing: Methuen; 1952.
30. Hambleton P Repair of wall damage in *Escherichia coli* recovered from an aerosol. *Journal of general microbiology*. 1971 11;69(1):81–8. [PubMed: 5004633]
31. Levin DE. Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiology and molecular biology reviews : MMBR*. 2005 6;69(2):262–91. [PubMed: 15944456]
32. Volkmann D, Baluska F. Actin cytoskeleton in plants: from transport networks to signaling networks. *Microscopy research and technique*. 1999 10 15;47(2):135–54. [PubMed: 10523792]

33. Major RJ, Poss KD. Zebrafish Heart Regeneration as a Model for Cardiac Tissue Repair. *Drug discovery today Disease models*. 2007;4(4):219–25. [PubMed: 19081827]
34. Chablais F, Jazwinska A. The regenerative capacity of the zebrafish heart is dependent on TGFβ signaling. *Development*. 2012 6;139(11):1921–30. [PubMed: 22513374]
35. Goldshmit Y, Sztal TE, Jusuf PR, Hall TE, Nguyen-Chi M, Currie PD. Fgf-dependent glial cell bridges facilitate spinal cord regeneration in zebrafish. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2012 5 30;32(22):7477–92. [PubMed: 22649227]
36. Poss KD, Keating MT, Nechiporuk A. Tales of regeneration in zebrafish. *Developmental dynamics : an official publication of the American Association of Anatomists*. 2003 2;226(2):202–10. [PubMed: 12557199]
37. Poss KD, Wilson LG, Keating MT. Heart regeneration in zebrafish. *Science*. 2002 12 13;298(5601):2188–90. [PubMed: 12481136]
38. Ward AC, McPhee DO, Condron MM, Varma S, Cody SH, Onnebo SM, et al. The zebrafish *spil* promoter drives myeloid-specific expression in stable transgenic fish. *Blood*. 2003 11 1;102(9):3238–40. [PubMed: 12869502]
39. Hsu K, Traver D, Kutok JL, Hagen A, Liu TX, Paw BH, et al. The *pu.1* promoter drives myeloid gene expression in zebrafish. *Blood*. 2004 9 1;104(5):1291–7. [PubMed: 14996705]
40. Lawson ND, Weinstein BM. In vivo imaging of embryonic vascular development using transgenic zebrafish. *Dev Biol*. 2002 8 15;248(2):307–18. [PubMed: 12167406]
41. Renshaw SA, Loynes CA, Trushell DM, Elworthy S, Ingham PW, Whyte MK. A transgenic zebrafish model of neutrophilic inflammation. *Blood*. 2006 12 15;108(13):3976–8. [PubMed: 16926288]
42. Mathias JR, Perrin BJ, Liu TX, Kanki J, Look AT, Huttenlocher A. Resolution of inflammation by retrograde chemotaxis of neutrophils in transgenic zebrafish. *Journal of leukocyte biology*. 2006 12;80(6):1281–8. [PubMed: 16963624]
43. Mathias JR, Dodd ME, Walters KB, Yoo SK, Ranheim EA, Huttenlocher A. Characterization of zebrafish larval inflammatory macrophages. *Dev Comp Immunol*. 2009 11;33(11):1212–7. [PubMed: 19619578]
44. Hall C, Flores MV, Storm T, Crosier K, Crosier P. The zebrafish lysozyme C promoter drives myeloid-specific expression in transgenic fish. *BMC Dev Biol*. 2007;7:42. [PubMed: 17477879]
45. Feng Y, Santoriello C, Mione M, Hurlstone A, Martin P. Live imaging of innate immune cell sensing of transformed cells in zebrafish larvae: parallels between tumor initiation and wound inflammation. *Plos Biol*. 2010;8(12):e1000562. [PubMed: 21179501]
46. Gray C, Loynes CA, Whyte MK, Crossman DC, Renshaw SA, Chico TJ. Simultaneous intravital imaging of macrophage and neutrophil behaviour during inflammation using a novel transgenic zebrafish. *Thrombosis and haemostasis*. 2011 5;105(5):811–9. [PubMed: 21225092]
47. Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. *mpeg1* promoter transgenes direct macrophage-lineage expression in zebrafish. *Blood*. 2011 1 27;117(4):e49–56. [PubMed: 21084707]
48. Wittamer V, Bertrand JY, Gutschow PW, Traver D. Characterization of the mononuclear phagocyte system in zebrafish. *Blood*. 2011 6 30;117(26):7126–35. [PubMed: 21406720]
49. Balla KM, Lugo-Villarino G, Spitsbergen JM, Stachura DL, Hu Y, Banuelos K, et al. Eosinophils in the zebrafish: prospective isolation, characterization, and eosinophilia induction by helminth determinants. *Blood*. 2010 11 11;116(19):3944–54. [PubMed: 20713961]
50. Bennett CM, Kanki JP, Rhodes J, Liu TX, Paw BH, Kieran MW, et al. Myelopoiesis in the zebrafish, *Danio rerio*. *Blood*. 2001 8 1;98(3):643–51. [PubMed: 11468162]
51. Danilova N, Steiner LA. B cells develop in the zebrafish pancreas. *Proceedings of the National Academy of Sciences of the United States of America*. 2002 10 15;99(21):13711–6. [PubMed: 12370418]
52. Dobson JT, Seibert J, Teh EM, Da'as S, Fraser RB, Paw BH, et al. Carboxypeptidase A5 identifies a novel mast cell lineage in the zebrafish providing new insight into mast cell fate determination. *Blood*. 2008 10 1;112(7):2969–72. [PubMed: 18635811]
53. Herbomel P, Thisse B, Thisse C. Ontogeny and behaviour of early macrophages in the zebrafish embryo. *Development*. 1999 9;126(17):3735–45. [PubMed: 10433904]

54. Langenau DM, Ferrando AA, Traver D, Kutok JL, Hezel JP, Kanki JP, et al. In vivo tracking of T cell development, ablation, and engraftment in transgenic zebrafish. *Proceedings of the National Academy of Sciences of the United States of America*. 2004 5 11;101(19):7369–74. [PubMed: 15123839]
55. Redd MJ, Cooper L, Wood W, Stramer B, Martin P. Wound healing and inflammation: embryos reveal the way to perfect repair. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2004 5 29;359(1445):777–84. [PubMed: 15293805]
56. Cvejic A, Hall C, Bak-Maier M, Flores MV, Crosier P, Redd MJ, et al. Analysis of WASp function during the wound inflammatory response--live-imaging studies in zebrafish larvae. *Journal of cell science*. 2008 10 1;121(Pt 19):3196–206. [PubMed: 18782862]
57. Walters KB, Green JM, Surfus JC, Yoo SK, Huttenlocher A. Live imaging of neutrophil motility in a zebrafish model of WHIM syndrome. *Blood*. 2010 10 14;116(15):2803–11. [PubMed: 20592249]
58. Richardson R, Slanchev K, Kraus C, Knyphausen P, Eming S, Hammerschmidt M. Adult zebrafish as a model system for cutaneous wound-healing research. *The Journal of investigative dermatology*. 2013 6;133(6):1655–65. [PubMed: 23325040]
59. Yoo SK, Freisinger CM, LeBert DC, Huttenlocher A. Early redox, Src family kinase, and calcium signaling integrate wound responses and tissue regeneration in zebrafish. *The Journal of cell biology*. 2012 10 15;199(2):225–34. [PubMed: 23045550]
60. Becker T, Bernhardt RR, Reinhard E, Wullimann MF, Tongiorgi E, Schachner M. Readiness of zebrafish brain neurons to regenerate a spinal axon correlates with differential expression of specific cell recognition molecules. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 1998 8 1;18(15):5789–803. [PubMed: 9671667]
61. Becker T, Wullimann MF, Becker CG, Bernhardt RR, Schachner M. Axonal regrowth after spinal cord transection in adult zebrafish. *The Journal of comparative neurology*. 1997 1 27;377(4):577–95. [PubMed: 9007194]
62. Fausett BV, Goldman D. A role for alpha1 tubulin-expressing Muller glia in regeneration of the injured zebrafish retina. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2006 6 7;26(23):6303–13. [PubMed: 16763038]
63. Villegas R, Martin SM, O'Donnell KC, Carrillo SA, Sagasti A, Allende ML. Dynamics of degeneration and regeneration in developing zebrafish peripheral axons reveals a requirement for extrinsic cell types. *Neural development*. 2012;7:19. [PubMed: 22681863]
64. Becker CG, Becker T. Adult zebrafish as a model for successful central nervous system regeneration. *Restorative neurology and neuroscience*. 2008;26(2–3):71–80. [PubMed: 18820403]
65. Diep CQ, Ma D, Deo RC, Holm TM, Naylor RW, Arora N, et al. Identification of adult nephron progenitors capable of kidney regeneration in zebrafish. *Nature*. 2011 2 3;470(7332):95–100. [PubMed: 21270795]
66. Fleisch VC, Fraser B, Allison WT. Investigating regeneration and functional integration of CNS neurons: lessons from zebrafish genetics and other fish species. *Biochimica et biophysica acta*. 2011 3;1812(3):364–80. [PubMed: 21044883]
67. Jopling C, Sleep E, Raya M, Marti M, Raya A, Izpisua Belmonte JC. Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature*. 2010 3 25;464(7288):606–9. [PubMed: 20336145]
68. Kroehne V, Freudenreich D, Hans S, Kaslin J, Brand M. Regeneration of the adult zebrafish brain from neurogenic radial glia-type progenitors. *Development*. 2011 11;138(22):4831–41. [PubMed: 22007133]
69. Lien CL, Schebesta M, Makino S, Weber GJ, Keating MT. Gene expression analysis of zebrafish heart regeneration. *PLoS biology*. 2006 8;4(8):e260. [PubMed: 16869712]
70. Hui SP, Sengupta D, Lee SG, Sen T, Kundu S, Mathavan S, et al. Genome wide expression profiling during spinal cord regeneration identifies comprehensive cellular responses in zebrafish. *PLoS one*. 2014;9(1):e84212. [PubMed: 24465396]
71. Mathew LK, Sengupta S, Kawakami A, Andreasen EA, Lohr CV, Loynes CA, et al. Unraveling tissue regeneration pathways using chemical genetics. *The Journal of biological chemistry*. 2007 11 30;282(48):35202–10. [PubMed: 17848559]

72. Betsuyaku T, Fukuda Y, Parks WC, Shipley JM, Senior RM. Gelatinase B is required for alveolar bronchiolization after intratracheal bleomycin. *The American journal of pathology*. 2000 8;157(2): 525–35. [PubMed: 10934155]
73. Lieschke GJ, Oates AC, Crowhurst MO, Ward AC, Layton JE. Morphologic and functional characterization of granulocytes and macrophages in embryonic and adult zebrafish. *Blood*. 2001 11 15;98(10):3087–96. [PubMed: 11698295]
74. Nathan C. Neutrophils and immunity: challenges and opportunities. *Nature reviews Immunology*. 2006 3;6(3):173–82.
75. van Furth R, Nibbering PH, van Dissel JT, Diesselhoff-den Dulk MM. The characterization, origin, and kinetics of skin macrophages during inflammation. *The Journal of investigative dermatology*. 1985 11;85(5):398–402. [PubMed: 4056454]
76. Li L, Yan B, Shi YQ, Zhang WQ, Wen ZL. Live imaging reveals differing roles of macrophages and neutrophils during zebrafish tail fin regeneration. *The Journal of biological chemistry*. 2012 7 20;287(30):25353–60. [PubMed: 22573321]
77. Leibovich SJ, Ross R. The role of the macrophage in wound repair. A study with hydrocortisone and antimacrophage serum. *The American journal of pathology*. 1975 1;78(1):71–100. [PubMed: 1109560]
78. Martin P, D'Souza D, Martin J, Grose R, Cooper L, Maki R, et al. Wound healing in the PU.1 null mouse--tissue repair is not dependent on inflammatory cells. *Current biology : CB*. 2003 7 1;13(13):1122–8. [PubMed: 12842011]
79. Shiratori Y, Hongo S, Hikiba Y, Ohmura K, Nagura T, Okano K, et al. Role of macrophages in regeneration of liver. *Digestive diseases and sciences*. 1996 10;41(10):1939–46. [PubMed: 8888704]
80. Tidball JG, Wehling-Henricks M. Macrophages promote muscle membrane repair and muscle fibre growth and regeneration during modified muscle loading in mice in vivo. *The Journal of physiology*. 2007 1 1;578(Pt 1):327–36. [PubMed: 17038433]
81. Dovi JV, He LK, DiPietro LA. Accelerated wound closure in neutrophil-depleted mice. *Journal of leukocyte biology*. 2003 4;73(4):448–55. [PubMed: 12660219]
82. Kyritsis N, Kizil C, Zocher S, Kroehne V, Kaslin J, Freudenreich D, et al. Acute inflammation initiates the regenerative response in the adult zebrafish brain. *Science*. 2012 12 7;338(6112): 1353–6. [PubMed: 23138980]
83. Redd MJ, Kelly G, Dunn G, Way M, Martin P. Imaging macrophage chemotaxis in vivo: studies of microtubule function in zebrafish wound inflammation. *Cell motility and the cytoskeleton*. 2006 7;63(7):415–22. [PubMed: 16671106]
84. Deng Q, Huttenlocher A. Leukocyte migration from a fish eye's view. *Journal of cell science*. 2012 9 1;125(Pt 17):3949–56. [PubMed: 23104739]
85. Starnes TW, Huttenlocher A. Neutrophil reverse migration becomes transparent with zebrafish. *Advances in hematology*. 2012;2012:398640. [PubMed: 22844288]
86. Niethammer P, Grabher C, Look AT, Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature*. 2009 6 18;459(7249):996–9. [PubMed: 19494811]
87. Yoo SK, Starnes TW, Deng Q, Huttenlocher A. Lyn is a redox sensor that mediates leukocyte wound attraction in vivo. *Nature*. 2011 12 1;480(7375):109–12. [PubMed: 22101434]
88. de Oliveira S, Reyes-Aldasoro CC, Candel S, Renshaw SA, Mulero V, Calado A. Cxcl8 (IL-8) mediates neutrophil recruitment and behavior in the zebrafish inflammatory response. *Journal of immunology*. 2013 4 15;190(8):4349–59.
89. Enyedi B, Kala S, Nikolich-Zugich T, Niethammer P. Tissue damage detection by osmotic surveillance. *Nat Cell Biol*. 2013 9;15(9):1123–30. [PubMed: 23934216]
90. Sieger D, Moritz C, Ziegenhals T, Prykhodzhiy S, Peri F. Long-range Ca²⁺ waves transmit brain-damage signals to microglia. *Developmental cell*. 2012 6 12;22(6):1138–48. [PubMed: 22632801]
91. Serhan CN, Chiang N, Van Dyke TE. Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nature reviews Immunology*. 2008 5;8(5):349–61.

92. Hsiao HM, Sapinoro RE, Thatcher TH, Croasdell A, Levy EP, Fulton RA, et al. A novel anti-inflammatory and pro-resolving role for resolvin D1 in acute cigarette smoke-induced lung inflammation. *PloS one*. 2013;8(3):e58258. [PubMed: 23484005]
93. Ji RR, Xu ZZ, Strichartz G, Serhan CN. Emerging roles of resolvins in the resolution of inflammation and pain. *Trends in neurosciences*. 2011 11;34(11):599–609. [PubMed: 21963090]
94. Spite M, Serhan CN. Novel lipid mediators promote resolution of acute inflammation: impact of aspirin and statins. *Circulation research*. 2010 11 12;107(10):1170–84. [PubMed: 21071715]
95. Savill J, Haslett C. Granulocyte clearance by apoptosis in the resolution of inflammation. *Seminars in cell biology*. 1995 12;6(6):385–93. [PubMed: 8748146]
96. Elks PM, van Eeden FJ, Dixon G, Wang X, Reyes-Aldasoro CC, Ingham PW, et al. Activation of hypoxia-inducible factor-1alpha (Hif-1alpha) delays inflammation resolution by reducing neutrophil apoptosis and reverse migration in a zebrafish inflammation model. *Blood*. 2011 7 21;118(3):712–22. [PubMed: 21555741]
97. Yoo SK, Huttenlocher A. Spatiotemporal photolabeling of neutrophil trafficking during inflammation in live zebrafish. *Journal of leukocyte biology*. 2011 5;89(5):661–7. [PubMed: 21248150]
98. Maletto BA, Ropolo AS, Alignani DO, Liscovsky MV, Ranocchia RP, Moron VG, et al. Presence of neutrophil-bearing antigen in lymphoid organs of immune mice. *Blood*. 2006 11 1;108(9):3094–102. [PubMed: 16835380]
99. Robertson AL, Holmes GR, Bojarczuk AN, Burgon J, Loynes CA, Chimen M, et al. A zebrafish compound screen reveals modulation of neutrophil reverse migration as an anti-inflammatory mechanism. *Science translational medicine*. 2014 2 26;6(225):225ra29.
100. Pase L, Layton JE, Wittmann C, Ellett F, Nowell CJ, Reyes-Aldasoro CC, et al. Neutrophil-delivered myeloperoxidase dampens the hydrogen peroxide burst after tissue wounding in zebrafish. *Current biology : CB*. 2012 10 9;22(19):1818–24. [PubMed: 22940471]
101. Kutter D, Devaquet P, Vanderstocken G, Paulus JM, Marchal V, Gothot A. Consequences of total and subtotal myeloperoxidase deficiency: risk or benefit? *Acta haematologica*. 2000;104(1):10–5. [PubMed: 11111115]
102. Love NR, Chen Y, Ishibashi S, Kritsiligkou P, Lea R, Koh Y, et al. Amputation-induced reactive oxygen species are required for successful *Xenopus* tadpole tail regeneration. *Nat Cell Biol*. 2013 2;15(2):222–8. [PubMed: 23314862]
103. Rieger S, Sagasti A. Hydrogen peroxide promotes injury-induced peripheral sensory axon regeneration in the zebrafish skin. *PLoS biology*. 2011 5;9(5):e1000621. [PubMed: 21629674]
104. Finkel T. Reactive oxygen species and signal transduction. *IUBMB life*. 2001 7;52(1–2):3–6. [PubMed: 11795590]
105. Gauron C, Rampon C, Bouzaffour M, Ipendey E, Teillon J, Volovitch M, et al. Sustained production of ROS triggers compensatory proliferation and is required for regeneration to proceed. *Scientific reports*. 2013;3:2084. [PubMed: 23803955]
106. Hattori N, Mochizuki S, Kishi K, Nakajima T, Takaishi H, D'Armiento J, et al. MMP-13 plays a role in keratinocyte migration, angiogenesis, and contraction in mouse skin wound healing. *The American journal of pathology*. 2009 8;175(2):533–46. [PubMed: 19590036]
107. Scott JW, Zottoli SJ, Beatty NP, Korn H. Origin and function of spiral fibers projecting to the goldfish Mauthner cell. *The Journal of comparative neurology*. 1994 1 1;339(1):76–90. [PubMed: 8106663]
108. Neumann S, Bradke F, Tessier-Lavigne M, Basbaum AI. Regeneration of sensory axons within the injured spinal cord induced by intraganglionic cAMP elevation. *Neuron*. 2002 6 13;34(6):885–93. [PubMed: 12086637]
109. Kao HT, Song HJ, Porton B, Ming GL, Hoh J, Abraham M, et al. A protein kinase A-dependent molecular switch in synapsins regulates neurite outgrowth. *Nature neuroscience*. 2002 5;5(5):431–7. [PubMed: 11976703]
110. Cai D, Qiu J, Cao Z, McAtee M, Bregman BS, Filbin MT. Neuronal cyclic AMP controls the developmental loss in ability of axons to regenerate. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2001 7 1;21(13):4731–9. [PubMed: 11425900]

111. Moore AR, Willoughby DA. The role of cAMP regulation in controlling inflammation. *Clinical and experimental immunology*. 1995 9;101(3):387–9. [PubMed: 7664483]
112. Bhatt DH, Otto SJ, Depoister B, Fetho JR. Cyclic AMP-induced repair of zebrafish spinal circuits. *Science*. 2004 7 9;305(5681):254–8. [PubMed: 15247482]
113. Foja S, Jung M, Harwardt B, Riemann D, Pelz-Ackermann O, Schroeder IS. Hypoxia supports reprogramming of mesenchymal stromal cells via induction of embryonic stem cell-specific microRNA-302 cluster and pluripotency-associated genes. *Cellular reprogramming*. 2013 Feb; 15(1):68–79.
114. Lokmic Z, Musyoka J, Hewitson TD, Darby IA. Hypoxia and hypoxia signaling in tissue repair and fibrosis. *International review of cell and molecular biology*. 2012;296:139–85. [PubMed: 22559939]

Fig. 1.

Schematic of innate immune inflammation induced by wounding in zebrafish. A) Wounding induces neutrophil recruitment through different cues including a tissue-scale gradient of hydrogen peroxide, cellular swelling-induced production of non-canonical arachadonate metabolites, and IL-8 production. B) Recruitment of macrophages occurs after neutrophils, and the cues are not well defined. After recruitment, neutrophils can reverse migrate from the damaged tissue, the function of which is unknown. C) Resolution is regulated by neutrophil reverse migration, and neutrophil apoptosis accompanied by a resolution of wound signals including hydrogen peroxide. D) Wound debridement by macrophages further contributes to wound resolution and regeneration. Macrophages phagocytose apoptotic neutrophils to mediate resolution.