Reference Distributions for the Negative Acute-Phase Proteins, Albumin, Transferrin, and Transthyretin: A Comparison of a Large Cohort to the World's Literature

Robert F. Ritchie,* Glenn E. Palomaki, Louis M. Neveux, and Olga Navolotskaia

Foundation for Blood Research, Scarborough, Maine

Clinical interpretation of the acute-phase proteins—albumin, transferrin, and transthyretin—has been hampered by the lack of accurate and precise methods for quantifying the levels and a stable and respected reference material. Now that these issues have been addressed, the community is faced with the need for credible age- and gender-specific reference values. The number of publications that address this issue, even for an analyte as familiar as albumin, is small and, in most cases, such publications lack the relevant data that would allow a combined experience to be created. We have identified 40 studies that meet our criteria: a description of the study participants' health status, of the statistical methodology, and of the laboratory technique and/or reference material used. Few of these studies

reported values stratified by gender. A summary of the published median levels by age is presented for the three analytes, along with our own age- and gender-specific medians based on a large cohort. Ten of the studies presented a 95 percent reference range, in close agreement with ours where selection was based upon reported diagnosis rather than upon determination of individual health status. This meta-analysis provides support for the reliability of our recently published methodology and reference data for the clinical interpretation of individual albumin, transferrin, and transthyretin values. As with most laboratory measurements, clinical interpretation requires that other laboratory and clinical factors be considered. *J. Clin. Lab. Anal.* 13:280–286, 1999. © 1999 Wiley-Liss, Inc.

Key words: serum proteins; acute-phase proteins; meta-analysis; CRM 470/RPPHS

INTRODUCTION

The acute-phase proteins, a complex family of about 30 different moieties, have been traditionally viewed as markers of overt inflammation, infection, tissue necrosis, or malignancy. However, their clinical application has been hampered by uncertainty in the appropriate reference ranges for a given patient. This is particularly evident in very young individuals where significant inflammation may be accompanied by elevations in the acute-phase proteins that fall well within the adult reference ranges, but would be considered high were reference ranges to be available for children and infants. The three proteins examined in this study represent the major negative acute-phase proteins (APP) whose synthesis is down-regulated by the same cytokines, IL-1 and IL-6, that drive the acute-phase response (APR). This study, like its predecessor (1), aims to identify publications from 1961 to the present that report reference data for albumin, transferrin, and transthyretin (prealbumin) and that satisfy minimal acceptable criteria. All results were standardized against CRM 470/RPPHS (2,3) before comparison.

MATERIAL AND METHODS

Identification of Published Reference Data

The methods used to search for relevant publication was the same as for the previous study (1) with the addition of searching PubMed (<http://www.ncbi.nlm.nih.gov/PubMed>) online for the same key headings as were used for MEDLINE CD-ROM (OVID Technologies, New York, NY) searches. Briefly, the literature was searched by direct review of the Cumulative Index Medicus published compendia and the computerized MEDLINE CD-ROM. Articles were considered acceptable if they: (1) identified the assay method and reference material; (2) provided details of the study population such as age; and (3) provided the information required for statistical analysis (4–102). Studies that were exclusively methodological or reported measurements only in individuals with known disease were removed from consideration.

*Correspondence to: Robert F. Ritchie, M.D., Foundation for Blood Research, Scarborough, Maine, 04074. E-mail: ritchie@fbr.org

Received 10 May 1999; Accepted 20 July 1999

Estimating the Central Estimate and Reference Range From Published Studies

For each study, either the mean, median, or geometric mean was used as the estimate of the center of the distribution. If these estimates were not directly available, they were computed using observed or smoothed centiles (e.g., the 2.5th and 97.5th centile). If none of these were available, the study was excluded from all analyses. Observed 95 percent ranges were used for the reference interval analysis. If the range was estimated from parameters or was other than the 95 percent range, the study was excluded from the reference interval analysis.

Conversion of Reported Results to a Single Reference Material

Results were standardized against CRM 470/RPPHS by conversion from package insert data from the period of the study as previously described (1). This material was released by the International Federation of Clinical Chemists (IFCC), the Community Bureau of Reference (BCR) in Brussels, and the College of American Pathologists (CAP) in the United States (the latter being labeled Reference Preparation for Proteins in Human Serum [RPPHS]).

RESULTS

Number of Studies Available for Analysis

We identified 71 publications that contained information on albumin in nondiseased individuals (4–9,12–16,18,19,21–26,28,29,32,33,35,36,45,46,54–56,58–60,62–70,72–91,94–102). Thirty-one of these studies did not include adequate information about assay methodology and/or the reference material used (9,12,14–16,22,23,25,26,29,33,35,58,60,62–64,72,76,77,79,85–90,95,99,100,102). Five studies did not include information about the age of the study subjects (8,18,19,75,94). The remaining 35 studies (4–7,13,21,24,28,32,36,45,46,54–56,59,65–70,73,74,78,80–84,91,96–98,101) were considered acceptable for the analysis of albumin measurements.

We identified 30 publications that contained information on transferrin in nondiseased individuals (4–7,9–11,18–20,26,28,37–39,47–49,53,54,56,59,62,63,69–71,76,83,84). Eleven studies did not include adequate information about assay methodology and/or the reference material used (11,26,39,47,48,53,62,63,71,76,84). Two studies did not include information about the age of study subjects (10,18). The remaining 17 studies (4–7,9,19,20,28,37,38,49,54,56,59,69,70,83) were considered acceptable for the analysis of transferrin measurements.

A total of 42 publications were identified which contained information on transthyretin in nondiseased individuals (5–8,17,18,22,27,28,30,31,34,36,40–44,46,49–53,56,57,61–63,65,69–71,73,76,78,83,92,93,98,99,101). Twenty-five

studies did not include adequate information about assay methodology and/or the reference material used (17,22,27,30,31,34,36,40,41,43,44,50,51,53,57,62,63,65,71,76,78,92,93,98,99). Two studies did not include age (8,18). The remaining 15 studies (5–7,28,42,46,49,52,56,61,69,70,73,83,101) were considered acceptable for transthyretin measurements.

For the analysis of reference range width, ten studies provided adequate information for at least one of the proteins (6,24,32,54,56,70,73,80,83,101).

Comparison of the Central Estimates

Figure 1 shows the central estimates from the published studies of albumin (a), transferrin (b), and transthyretin (c). The reported mean (or median) age of the study population is shown on the logarithmic horizontal axis versus the reported mean or median analyte level on the logarithmic vertical axis over a fourfold range in values. Each observation is shown as a circle with the smallest and darkest circles representing estimates based on over 100 observations, median sized circles representing between 50 and 99 observations, and the largest circles representing between 10 and 49 observations. If a published observation was based on less than 10 observations, it was combined with an adjacent age group from the same publication. For purposes of comparison, the solid and dashed lines represent the median levels found for males and females, respectively, in our companion study (3).

Figure 1 shows the reported albumin median values from the 35 published studies considered acceptable. Most data are for adults over age 30. Our median levels for albumin are identical to the large number of published estimates for subjects over age 60. Our estimates are somewhat lower than, but consistent with, the estimates for younger adults. The figure shows that little reliable data has been published for children under age 10, but our estimates are consistent with that which does exist. There are far fewer published median levels of transferrin than for albumin and the spread of the estimates is greater. Our median levels for adults are near the middle of the range of published estimates. The few estimates available for young children, however, are considerably lower than ours. However, five of the six published estimates under age 10 are from two studies (56,83). The estimates of our median albumin and transferrin levels (Fig. 1a,b) are remarkably consistent with published studies of single age- and gender-specific consensus values generated soon after CRM 470 was released (103,104). The published median estimates for transthyretin are similar to our estimates over the entire range of ages (Fig. 1c).

Comparison of the Reference Ranges

The reference ranges reported in the studies used in the above analyses were generally based on a small number of observations; half were based on 489 observations or less.

Fig. 1. A summary of published acute phase protein median reference values. The published median (or geometric mean) levels for albumin (A), transferrin (B) and transthyretin (C) levels are displayed on the logarithmic vertical axis (spanning a factor of four) versus the mean (or median) age on the logarithmic horizontal axis. The three symbol sizes represent the number of observations in each group (small, medium and large circles representing over 100, 50 to 99 and 10 to 49). The solid (males) and dashed (females) lines are the regressed median levels from our large cohort study (3).

Thus, it was possible to verify the health status of each individual. In contrast, our observed reference ranges were each based on over 50,000 observations and, therefore, it was not possible to verify the health status of each individual. Although the median levels derived in our study are in agreement with published ones, it is possible that our reference ranges may be wider than other smaller studies because we relied on the diagnosis provided on the laboratory slip (105). In order to determine whether this was the case, we re-analyzed the published reference ranges using our earlier finding (Fig. 1a,b,c) that the widths of the ranges are nearly completely independent of age and gender (3). Figure 2 shows the 10 previous studies for which it was possible to calculate a 2.5th to 97.5th centile range (after each study's reference limits were divided by their own population medians—conversion to multiples of the median). The reference numbers are shown on the horizontal axis, sorted by decreasing 95 percent reference-range width. The weighted average of the ranges reported in the literature (excluding those from our

study) is shown by two thin horizontal dashed lines. The 95 percent reference ranges for our study (both observed and predicted from the population parameters) are represented as thick, dashed lines. Figure 2b,c shows a similar analysis for transferrin and transthyretin reference ranges, respectively. For all three analytes, our ranges are similar to the consensus estimate from the literature. Although there are a few extreme estimates that vary greatly from the consensus, the majority of the studies are consistent. Furthermore, the reference range widths for children (56,106), when compared to an elderly group (70), also are remarkably similar even though the median levels vary widely.

study) is shown by two thin horizontal dashed lines. The 95 percent reference ranges for our study (both observed and predicted from the population parameters) are represented as thick, dashed lines. Figure 2b,c shows a similar analysis for transferrin and transthyretin reference ranges, respectively. For all three analytes, our ranges are similar to the consensus estimate from the literature. Although there are a few extreme estimates that vary greatly from the consensus, the majority of the studies are consistent. Furthermore, the reference range widths for children (56,106), when compared to an elderly group (70), also are remarkably similar even though the median levels vary widely.

Ethnic Group-Specific Reference Ranges

A total of 25 studies included in the previous analyses reported the race or ethnic heritage of the study subjects. The majority, 19, reported measurements in Caucasian or mostly Caucasian populations (4,5,7,13,24,38,45,49,65,68,69,73,78,82,83,91,96,97,101). Four studied only black Africans

Fig. 2. Reference intervals from published studies. The bars represent the interval between the 2.5th and 97.5th centile for each study with the number of subjects above the bar. The horizontal axis shows the reference number for each study. The selected centiles have been presented as multiples of the median (MoM), compensating for age, gender and reference material differences between studies. Figure 2A, 2B and 2C show the analyses for albumin, transthyretin and transferrin measurements, respectively, on identical logarithmic vertical axes. The horizontal dotted lines display the weighted consensus value at the 2.5th centile and the 97.5th centile for previously published studies. The broken bars represent the observed and predicted reference ranges from our large cohort study (3).

(42,80,81,98) and two studied only Asians (28,66). Of the 10 studies providing reference ranges displayed in Figure 2a (6,24,32,54,56,70,73,80,83,101), five did not provide information on race, four were based on Caucasians, and one was based on Blacks. Therefore, no claim for differences or similarities in reference-range widths between races can be made in this study.

DISCUSSION

With the considerable improvement in instrument performance (107) and the availability of a stable, widely accepted reference material (2), robust age- and gender-specific reference data have remained one of the last remaining barriers to improving clinical interpretation. Until now, reference ranges were based on relatively small numbers of study subjects but had the advantage of often documenting individual health sta-

min, transferrin and transthyretin measurements, respectively, on identical logarithmic vertical axes. The horizontal dotted lines display the weighted consensus value at the 2.5th centile and the 97.5th centile for previously published studies. The broken bars represent the observed and predicted reference ranges from our large cohort study (3).

tus. Our companion study (3) reports reference data for albumin, transferrin, and transthyretin based on a much larger data set with less reliance on individual health status. The current study shows our reference data, in terms of both the median and reference interval width, to be on average, consistent with these smaller studies. The advantage of our larger study is that the reference data can now be specified for any age between 1 month and 85 years of age, for both males and females.

The establishment of a reliable set of reference data for males and females throughout life facilitates future reviews comparing serum protein levels in similar populations. Laboratories are being required to design and establish local reference ranges that can account for possible ethnic and geographic differences. Protocols have been developed to assist and guide laboratories in constructing and determining serum protein reference ranges in local cohorts (103,104).

This effort has been made easier with the completion of our two studies (3,105). These reference data for presumed healthy individuals also set the stage for estimating screening or diagnostic performance by establishing the distribution of analyte values in individuals with specific diseases. The importance of this knowledge is just becoming clear.

ACKNOWLEDGMENTS

This work was supported by funds from the Foundation for Blood Research Development Program.

REFERENCES

- Ritchie RF, Palomaki GE, Neveux LM, Navolotskaia O. Reference values for immunoglobulins A, G, and M: a comparison of the world's literature to a single large cohort. *J Lab Clin Anal* 1998;12:371-377.
- Whicher JT, Ritchie RF, Johnson AM, Baudner S, Bienvenu J, Blirup-Jensen S. New international reference preparation for proteins in human serum (RPPHS). *Clin Chem* 1994;40:934-938.
- Ritchie RF, Palomaki GE, Neveux LM, Navolotskaia O, Ledue TB, Craig WY. Reference distributions for the negative acute phase proteins, albumin, transferrin and transthyretin: A practical, simple and clinically relevant approach. *Clin Chem J Clin Lab Anal* 1999;13:273-279.
- Song CS, Merkatz IR, Rifkind AB, Gillette PN, Kappas A. The influence of pregnancy and oral contraceptive steroids on the concentration of plasma proteins. *Am J Obstet Gynecol* 1970;66:227-231.
- Lyngbye J, Krøll J. Quantitative immunoelectrophoresis of proteins in serum from a normal population: season-, age-, and sex-related variations. *Clin Chem* 1971;17:495-500.
- Weeke B, Krasilnikoff PA. The concentration of 21 serum proteins in normal children and adults. *Acta Med Scand* 1972;192:149-155.
- Dotchev D, Liappis N, Hungerland H. Sex-specific differences of serum proteins in adults and influence of oral hormonal contraceptives on serum protein composition. *Clin Chem Acta* 1973;44:431-435.
- Agostoni A, Marasini B, Stabilini R, Del Ninno E, Pontello M. Multivariate analysis of serum protein assays in chronic hepatitis and postnecrotic cirrhosis. *Clin Chem* 1974;20:428-429.
- Svartman M, Potter EV, Poon-King T, Earle DP. Immunoglobulins and complement components in synovial fluid of patients with acute rheumatic fever. *J Clin Invest* 1975;56:111-117.
- Snyder S, Durham BC, Iskandrian AS, Coodley EL, Linhart JW. Serum lipids and glycoproteins in acute myocardial infarction. *Am Heart J* 1975;90:582-586.
- Statland BE, Winkel P, Killingsworth LM. Factors contributing to intra-individual variation of serum constituents: 6. Physiological day-to-day variation in concentrations of 10 specific proteins in sera of healthy subjects. *Clin Chem* 1976;22:1635-1638.
- Ramcharan S, Sponzilli EE, Wingerd JC. Serum protein fractions. Effect of oral contraceptives and pregnancy. *Obstet Gynecol* 1976;48:211-215.
- Ljunghall S, Hedstrand H, Hellsing K, Wibell L. Calcium, phosphate and albumin in serum. *Acta Med Scand* 1977;01:23-30.
- Wingerd J, Sponzilli EE. Concentrations of serum protein fractions in white women: effects of age, weight, smoking, tonsillectomy, and other factors. *Clin Chem* 1977;23:1310-1317.
- Ramsay L, Mitchell T, Shelton J. Plasma uric acid and plasma albumin in healthy subjects. *Ann Rheum Dis* 1977;36:374-377.
- Cherian AG, Hill JG. Percentile estimates of reference values for fourteen chemical constituents in sera of children and adolescents. *Am J Clin Pathol* 1978;69:24-31.
- Vahlquist A, Sjölund K, Nordén A, Peterson PA, Stigmar G, Johanson B. Plasma vitamin A transport and visual dark adaptation in diseases of the intestine and liver. *Scan J Clin Lab Invest* 1978;38:301-308.
- Milano G, Cooper EH, Goligher JC, Giles GR, Neville AM. Serum prealbumin, retinol-binding protein, transferrin, and albumin levels in patients with large bowel cancer. *J Natl Cancer Inst* 1978;61:687-691.
- Ritchie RF. Serum protein profile analysis and interpretation: some basic information. *Immunoassays in the Clinical Laboratory*. New York:AR Liss, Inc; 1979. p 227-242.
- Freer DE, Statland BE, Johnson M, Felton H. Reference values for selected enzyme activities and protein concentrations in serum and plasma derived from cord-blood specimens. *Clin Chem* 1979;25:565-569.
- Cheng MH, Lipsey AI, Blanco V, Wong HT, Spiro SH. Microchemical analysis for 13 constituents of plasma from healthy children. *Clin Chem* 1979;25:692-698.
- Baskies AM, Chretien PB, Yang C-s, Wolf GT, Makuch RW, Tu S-m, et al. Serum glycoproteins and immunoglobulins in nasopharyngeal carcinoma. *Am J Surgery* 1979;38:478-478.
- Ischei UP. Serum protein profile in sickle cell disease. *J Clin Pathol* 1979;32:117-121.
- Gardner MD, Scott R. Age- and sex-related reference ranges for eight plasma constituents derived from randomly selected adults in a Scottish new town. *J Clin Pathol* 1980;33:380-385.
- Jernigan JA, Gudat JC, Blake JL, Bowen L, Lezotte DC. Reference values for blood findings in relatively fit elderly persons. *J Am Geriatr Society* 1980;28:308-314.
- Thompson DK, Haddow JE, Smith DE, Ritchie RF. Serum protein changes in women with early breast cancer. *Cancer* 1981;48:793-798.
- Skinner M, Heller L, Rubinow A, Libbey CA, Sipe JD, Cohen AS. Depressed serum prealbumin levels in patients with familial amyloid polyneuropathy (FAP) and their non-affected relatives. *Arth Rheum* 1982;25(Suppl):S58.
- Pongpaew P, Schelp FP. Albumin, prealbumin, transferrin and serum proteinase inhibitors in asymptomatic and hospitalized Thai alcohol drinkers. *J Stud Alcohol* 1983;44:381-383.
- Haram K, Augensen K, Elsayed S. Serum protein in normal pregnancy with special reference to acute-phase reactants. *Br J Obstet Gynecol* 1983;90:139-145.
- Savaria MJM, Costa PP, Goodman DS. Studies on plasma transthyretin (prealbumin) in familial amyloidotic polyneuropathy, Portuguese type. *J Lab Clin Med* 1983;102:590-603.
- Bhatia J, Ziegler EE. Retinol-binding protein and prealbumin in cord blood of term and preterm infants. *Early Hum Dev* 1983;8:129-130.
- Ash KO, Clark SJ, Sandberg LB, Hunter E, Woodward SC. The influence of sample distribution and age on reference intervals for adult males. *Am J Clin Pathol* 1983;79:574-581.
- Pickart L. Increased ratio of plasma free fatty acids to albumin during normal aging and in patients with coronary heart disease. *Atherosclerosis* 1983;46:21-28.
- Benson MD, Dwulet FE. Prealbumin and retinol-binding protein serum concentrations in the Indiana type hereditary amyloidosis. *Arth Rheum* 1983;26: 1493-1498.
- Butler SJ, Payne RB, Gunn IR, Burns J, Paterson CR. Correlation between serum ionized calcium and serum albumin concentrations in two hospital populations. *Br Med J* 1984;289:948-950.
- Giacoia GP. Concentration of serum prealbumin and retinol-binding proteins during pregnancy. *South Med J* 1984;77:1261-1263.
- Stoddard L, Dennis W, Parvin RM, Assendelft OW van. Freeze/thaw stability of transferrin, and reference values obtained with kinetic nephelometry. *Clin Chem* 1984;30:114-115.
- Behr W, Schlimok G, Firchau V, Paul HA. Determination of reference intervals for 10 serum proteins measured by rate nephelometry, taking into consideration different sample groups and different distribution functions. *J Clin Chem Clin Biochem* 1985;23:157-166.
- DiRusso SC, Check IJ, Hunter RL. Quantitation of apo-, mono-, and

- rum and its correlation with some hematic parameters. *BioMetals* 1994;7:75–79.
83. Kanakoudi F, Drossou V, Tzimouli V, Diamanti E, Konstantinidis T, Germentis A, Kremenopolous G. Serum concentrations of 10 acute-phase proteins in healthy term and preterm infants from birth to age 6 months. *Clin Chem* 1995;41:605–608.
  84. Periquet BA, Jammes NM, Lambert WE, Tricoire J, Moussa MM, Garcia J, Ghisolfi J, Thouvenot J-P. Micronutrient levels in HIV-1-infected children. *AIDS* 1995;9:887–893.
  85. Kenny SE, Pierro A, Isherwood D, Donnell SC, Van Saene HKF, Lloyd DA. Hypoalbuminemia in surgical neonates receiving parenteral nutrition. *J Pediatr Surg* 1995;30:454–457.
  86. Donnelly CA, Seth J, Clayton RM, Phillips CI, Cuthbert J, Prescott RJ. Some blood plasma constituents correlate with human cataract. *Br J Ophthalmol* 1995;79:1036–1041.
  87. Odvina CV, Safi I, Wojtowicz CH, Barengolts EI, Lathon P, Skapars A, Desai PN, Kukreja SC. Effect of heavy alcohol intake in the absence of liver disease on bone mass in black and white men. *J Clin Endocrinol Metab* 1995;80:2499–2503.
  88. Kuroki S, Okamoto S, Naito T, Oda H, Nagase S, Sakai H, Nawata H, Yamashita H, Chijiwa K, Tanaka M. Serum 7 α -hydroxycholesterol as a new parameter of liver function in patients with chronic liver disease. *Hepatology* 1995;22:1182–1187.
  89. Maes M, Wauters A, Neels H, Scharpé S, Van Gastel A, D'Hondt P, Cosyns P, Desnyder R. Total serum protein and serum protein fractions in depression: relationship to depressive symptoms and glucocorticoid activity. *J Affect Disord* 1995;34:61–69.
  90. Shiota G, Umeki K, Okano J-I, Kawasaki H. Hepatocyte growth factor and acute phase proteins in patients with chronic liver disease. *J Med* 1995;26:295–308.
  91. Hosseine SJ, Farid R, Ghalighi MR, Feely J. Interethnic differences in drug protein binding and α_1 -acid glycoprotein concentration. *Ir J Med Sci* 1995;64:26–27.
  92. Centanni M, Maiani G, Parkes AB, N'Diaye AM, Ferro-luzzi A, Lazarus JH. Thyroid homeostasis and retinol circulating complex relationships in severe iodine-deficient area of Senegal. *J Endocrinol Invest* 1995;18:608–612.
  93. Jain SK, Rohatgi A, Raman KK, Sharma VK. Study of serum prealbumin and serum alpha-fetoprotein in cases of fulminant hepatic failure. *Journal of the Association Physicians of India* 1995;43:462–463.
  94. Prabha K, Al-Sulaiman A, Olusi SO, Sugathan TN. Reference intervals for some serum biochemical markers of bone metabolism in Kuwait. *Clin Chim Acta* 1996;249:67–75.
  95. Bacq Y, Zarka O, Bréchet J-F, Mariotte N, Vol S, Tichet J, Weill J. Liver function tests in normal pregnancy: a prospective study of 103 pregnant women and 103 matched controls. *Hepatology* 1996;23:1030–1034.
  96. Smith G, Robinson PH, Fleck A. Serum albumin distribution in early treated anorexia nervosa. *Nutrition* 1996;12:677–684.
  97. Thome J, Münch G, Müller R, Schinzel R, Kornhuber J, Blum-Degen D, Sitzmann L, Rösler M, Heidland A, Riederer P. Advanced glycation endproducts-associated parameters in the peripheral blood of patients with Alzheimer's disease. *Life Sci* 1996;59:679–685.
  98. Walter EA, Scariano JK, Easington CR, Polaco AM, Hollis BW, Dasgupta A, Pam S, Glew R. Rickets and protein malnutrition in Northern Nigeria. *J Trop Pediatr* 1997;43:98–102.
  99. Malvy D J-M, Arnaud J, Burtschy B, Sommelet D, Leverger G, Dostalova L, Amedee-Manésmé O. Antioxidant micronutrients and childhood malignancy during oncological treatment. *Med Pediatr Oncology* 1997;29:213–217.
  100. Kim KJ, Yang WS, Kim SB, Lee SK, Park JS. Fibrinogen and fibrinolytic activity in CAPD patients with atherosclerosis and its correlation with serum albumin. *Perit Dial Int* 1997;7:157–161.
  101. Fiore P, Castagnola E, Marchese N, Dufour C, Garaventa A, Mangraviti S, Cornaglia-Ferraris P. Retinol (Vitamin A) and retinol-binding protein serum levels in children with cancer at onset. *Nutrition* 1997;13:17–20.
  102. Cederholm T, Arner P, Palmblad J. Low circulating leptin levels in protein-energy malnourished chronically ill elderly patients. *J Intern Med* 1997;242:377–382.
  103. Dati F, Schumann G, Thomas L, Aguzzi F, Baudner S, Bienvenu J, et al. Consensus of a group of professional societies and diagnostic companies on guidelines for the interim reference ranges for 14 proteins in serum based on the standardization against the IFCC/BCR/CAP reference material (CRM 470). *Eur J Clin Chem Biochem* 1996;34:517–520.
  104. Itoh Y, Ichihara K, Kanno T, Sugawara T, Ohkubo A, Hirabayashi Y, et al. Serum protein standardization project in Japan: evaluation of an IFCC reference material (RPPHS/CRM470) and establishment of reference intervals. *J Clin Lab Anal* 1997;1:39–44.
  105. Ritchie RF, Palomaki GE, Neveux LM, Navolotskaia O, Ledue TB, Craig WY. Reference distributions for immunoglobulins A, G and M: A practical, simple and clinically relevant approach. *J Clin Lab Anal* 1998;12:363–370.
  106. Malvy DJM, Poveda JD, Debruyne M, Montagnon B, Burtschy B, Herbert C, et al. Laser immunonephelometry reference intervals for eight serum proteins in healthy children. *Clin Chem* 1992;38:394–399.
  107. Ledue TB, Johnson MJ, Cohen LA, Ritchie RF. Evaluation of proficiency survey results for serum immunoglobulins following the introduction of a new international reference material for human serum proteins. *Clin Chem*, 1998;44:878–879.
  108. National Committee for Clinical Laboratory Standards. How to define and determine reference intervals in the clinical laboratory; Approved Guideline. NCCLS Document C28-A. NCCLS, 771 East Lancaster Avenue, Villanova, Pennsylvania 19085, 1995.
  109. Whicher JW. Minutes of the International Federation of Clinical Chemistry Meeting of the Committee on Plasma Proteins, Scientific Division, August 20, 1997. Basel, Switzerland, and October 8, 1997, Viareggio, Italy.