Published in final edited form as:

Curr Opin Physiol. 2019 April ; 8: 188–192. doi:10.1016/j.cophys.2019.02.003.

Reconciling the functions of even-skipped interneurons during crawling, swimming, and walking

Michael Jay, David L. McLean

Department of Neurobiology Northwestern University EVANSTON, IL USA

Abstract

In all bilaterally symmetric animals, movements across the body are coordinated by interneurons that traverse the midline. Recent work is beginning to tease apart the functional complexity of interneurons labeled by the homeodomain transcription factor even-skipped, which provide a phylogenetically-conserved source of commissural excitation during locomotion in both vertebrates and invertebrates. Here we review recent studies of the roles of even-skipped neurons during locomotion in flies (EL neurons), fishes, frogs, and mice (V0v neurons). Comparisons across species reveal commonalities, which include the functional organization of even-skipped circuits based on birth order, the link between increased muscular complexity and even-skipped neuron diversity, and the hierarchical organization of even-skipped circuits based on their control of escape versus exploratory movements. We discuss how stronger links between different species enable testable predictions to further the discovery of principles of locomotor network organization.

AUTHOR DECLARATION TEMPLATE

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us.

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

We further confirm that any aspect of the work covered in this manuscript that has involved either experimental animals or human patients has been conducted with the ethical approval of all relevant bodies and that such approvals are acknowledged within the manuscript. [CAN BE DELETED IF NOT RELEVANT]

We understand that the Corresponding Author is the sole contact for the Editorial process (including Editorial Manager and direct communications with the office). He/she is responsible for communicating with the other authors about progress, submissions of revisions and final approval of proofs. We confirm that we have provided a current, correct email address which is accessible by the Corresponding Author and which has been configured to accept email from (PLEASE CONFIRM WHAT E-MAIL ADDRESS YOU WISH TO INSERT HERE)

david-mclean@northwestern.edu

Signed by all authors as follows:

[LIST AUTHORS AND DATED SIGNATURES ALONGSIDE]

David McLean 11/18/2018

Michael Jay 11/18/2018

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

V0v neurons; EL neurons; spinal cord; abdominal ganglion; neural circuits; locomotion; behavior

Introduction

Different forms of animal locomotion necessitate different patterns of muscle activation and with different levels of urgency. Whether crawling, swimming, walking or flying, commissural projection neurons that traverse the midline are required to coordinate movements across the body during escape and exploratory movements. The discovery of shared transcriptional networks regulating interneuron specification has allowed for better comparisons between locomotor-related interneurons in different species and different stages of development [1]. The hope for such comparative approaches is the identification of core operational principles that explain locomotor control in vertebrates and invertebrates alike. Here we highlight several recent investigations that are beginning to hint at evolutionarily conserved functions of a genetically-identified source of commissural excitation in the coordination of bilateral activity during various forms of locomotion, namely interneurons labeled by the homeodomain transcription factor even-skipped [2–4].

Even-skipped interneurons and crawling

During larval stages of the fly, peristaltic crawling movements using axial muscles are the primary form of exploratory locomotion [5]. Crawling is characterized by waves of bilateral muscle contractions that pass from tail-to-head (forward crawling) or head-to-tail (backward crawling). Muscles in the abdominal segments are responsible for generating these movements and neurons within isolated abdominal ganglia controlling these muscles can produce rhythmic motor patterns appropriate for crawling in the absence of sensory inputs [6]. In addition to exploratory movements, larvae also produce a forceful ‘corkscrew’ response that allows them to escape threats [5].

Relatively recent work from the Doe lab using thermogenetic and optogenetic perturbation strategies in fly larvae has revealed a key role for a subset of laterally located even-skipped (Eve) positive interneurons (known as ‘ELs’) in the coordination of bilaterally symmetric locomotor movements [7]. Each abdominal ganglion contains bilateral clusters of excitatory (acetylcholine) EL interneurons. While all ELs were commissural, their axonal trajectories were diverse. Commissural axons from cells within each cluster had either ascending (e.g., A08c, A08s) or locally bifurcating (e.g., A08e1–3) projections where they formed direct synaptic contacts with motoneurons as well as with other motor- and sensory-related interneurons. Activation or ablation of the EL population slowed crawling movements and frequently led to unilateral bends. Interestingly, the effects were limited to the amplitude of muscle contractions rather than the timing, suggesting these neurons are not part of the circuits that produce the crawling rhythm, but rather contribute to the intensity of body bends. This conclusion was supported by calcium imaging experiments in isolated preparations demonstrating EL neurons are not rhythmically active during ‘fictive’ crawling [7].

A more recent study from the Heckscher lab has revealed a more nuanced behavioral role for ELs [8]. By using a molecular birth-dating strategy, Wreden and colleagues (2017) first demonstrated that EL interneurons comprise two developmentally distinct cohorts of cells. The authors then used optogenetic techniques in freely behaving larvae to examine how these cohorts might contribute to locomotion. Optogenetic activation of late-born ELs (A08c, A08s, A08e1–3) reduced the speed of crawling and perturbed symmetric body contractions, consistent with previous findings [7]. However, stimulation of early-born ELs (A08x, A08m [9]) increased the speed of locomotion and induced corkscrew escapes. How might these two populations of ELs contribute to different behaviors? Electron microscopy revealed putative synaptic inputs to early-born ELs from mechanosensory primary and secondary afferents, whereas late-born ELs received inputs from proprioceptive primary afferents [7,8], suggesting ELs integrate different sensory information to facilitate selection of appropriate motor patterns.

In sum, these results indicate that in the developing *Drosophila* larvae, temporally arrayed cohorts of even-skipped neurons form distinct circuits to modulate the intensity and mode of locomotor output – early born neurons are active during more forceful escape movements, while later born neurons are active during weaker exploratory ones.

Even-skipped interneurons and swimming

The idea that temporal sequencing contributes to even-skipped neuron functional diversity is consistent with studies in zebrafish. Axial muscle activity during early locomotor movements in zebrafish alternates from left–right [10]. By freely swimming exploratory larval stages, rhythmic axial movements are accompanied by rhythmic movements of bilaterally paired pectoral fins [11]. By juvenile/adult stages, zebrafish also use paired pelvic fins and medial dorsal, anal and caudal fins for propulsion and stabilization [12]. As in invertebrates, even-skipped (*Evx1/2*) interneurons in the spinal cord are also commissural and excitatory (glutamate) but are referred to as V0v neurons due to their ventral origin from the spinal p0 progenitor domain in mice [13,14].

In zebrafish, V0v neurons with ascending axons (a.k.a. commissural secondary ascending interneurons or CoSAs) are the first to differentiate [15]. These neurons are integrated into zebrafish motor networks early during development and are recruited during spontaneous, whole-body bends [16], which resemble escape bends used in response to threats in older fish. Recent work from the Lewis lab has revealed that V0v neurons in frog embryos are also primarily ascending [17,18], which is a time of life when frogs engage axially-based escape movements [19,20]. These results are consistent with the idea of a conserved function for early-born *Eve/Evx* interneurons during powerful escape behaviors.

So do later born V0v neurons also participate in exploratory behaviors? In zebrafish, the next V0v neurons to differentiate have bifurcating axons that can project locally or across multiple body segments [15], followed by V0v neurons with primarily descending axon trajectories (a.k.a., multipolar commissural descending neurons or MCoDs and unipolar commissural descending neurons or UCoDs) [21]. Of these, only MCoDs have been functionally characterized in larvae and they participate exclusively in exploratory low

frequency swimming [22], which is accompanied by rhythmic activity in the pectoral fins [23]. At higher frequency escape responses, the fins are adducted against the body and MCoDs are inhibited [23]. Laser ablation of MCoDs uncouples fin movements from axial ones during exploratory swimming, but does not impact escape swimming [24]. This suggests they are not part of the rhythm generator, but instead play a role in regulating the intensity of muscle contractions in a mode-specific manner, as proposed in fly larvae.

Recent work from the El Manira lab in juvenile/adult zebrafish has begun to allow for comparisons of V0v function throughout development [25]. As zebrafish age, there is a leftward shift in the distribution of axial bending frequencies in freely swimming fish, from 20–90 Hz in embryos/larvae [21] to 1–60 Hz in juvenile/adults [26]. Larval and juvenile/adult fish spend most of their time in exploratory mode swimming at lower frequencies, which can be reliably observed in fictive preparations [27,28]. However, generating high frequency swimming in fictive preparations is more challenging. With this in mind, Björnfors & El Manira (2016) performed a series of electrophysiological recordings from V0v neurons in juvenile/adults during fictive swimming, examined their recruitment at frequencies between 1–8 Hz, and then assessed whether they had ascending, bifurcating or descending axonal trajectories [25].

As observed in larvae, a fraction of descending V0v neurons participate in low frequency exploratory swimming in juvenile/adults (< 8 Hz). There were also many ascending and bifurcating V0v neurons that were not recruited at these low frequencies, consistent with their proposed birth order. Clearly, however, morphology was not always a perfect predictor of recruitment order since a fraction of ascending and bifurcating V0v neurons were also recruited below 8 Hz, while many descending V0vs were still not. Since higher frequencies of motor activity were not generated in this study, it remains to be seen if non-recruited V0v neurons participate in faster exploratory and escape swimming or if slower V0v circuits in adults are inhibited during transitions in swimming mode as they appear to in larvae [22]. Based on previous work in adults, one would expect such switches to occur above 21 Hz, during what is referred to as ‘escape’ [26].

Regardless, the work in fish and frogs not only supports the temporal assembly of escape then exploratory even-skipped circuitry observed in flies, it also suggests that increased morphological diversity within the Eve/Evx population during development supports increased complexity of muscular coordination during locomotion.

Even-skipped interneurons and walking

Intensity and mode specific participation of morphologically diverse V0v neurons according to birth order is also supported by recent experimental and computational studies in mice. During ambulatory locomotion, increases in speed are accompanied by increases in the cyclical frequency of muscles within limbs and changes in the timing and amplitude of muscle contractions within and between them [29,30]. Earlier work from the Kiehn lab [31] genetically ablating V0v neurons during development revealed profound deficits in left–right alternation between hindlimbs at high frequencies of fictive locomotion in neonates and real

locomotion in adults. Additionally, there were more subtle deficits in forelimb–hindlimb coordination at low frequencies.

Previous tracing studies in embryonic mice from the Goulding lab revealed glutamatergic Evx1-positive V0v neurons have primarily ascending axonal projections and a small number have bifurcating axons [14]. In addition, the Arber lab has recently shown that by postnatal stages, there is a further population of descending V0v neurons that link the cervical and lumbar regions of the spinal cord that control the fore- and hindlimbs, respectively [32]. Ablation of cervico-lumbar propriospinal connections, which include V0v neurons, disrupts low intensity exploratory walking in freely behaving animals and interlimb coordination at higher speeds. In addition, recent work using c-fos expression from the Gosgnach lab has demonstrated that V0v neurons in lumbar spinal cord are activated from low (0.18 Hz) to moderately high (0.48 Hz) frequencies of fictive locomotion [33], although links between morphology and recruitment could not be performed. Nonetheless, these observations suggest that ascending and bifurcating V0v neurons are among the earliest detectable and that V0v neurons play critical roles in coordinating activity along and across the body during both exploratory and escape movements, as reported in flies and zebrafish.

The idea that V0v neurons in mice represent a morphologically diverse population with speed-dependent patterns of recruitment was recently explored computationally by Danner and colleagues [34]. One of the key predictions from this work is that locally and long-range projecting V0vs may be central to speed-dependent transitions between gaits, but are not part of the rhythm generating circuits. Under the proposed paradigm, locally, but not long-range projecting V0vs (i.e. cervical–cervical, lumbar–lumbar) receive supraspinal inhibitory drive that increases with locomotor frequency. This means transitions to the fastest gaits require inhibition of V0v neurons associated with slower gaits. Although gait-dependent inhibition needs to be confirmed physiologically, this pattern mirrors the one observed in zebrafish larvae, where V0v neurons engaged during slower exploratory swimming involving the pectoral fins are inhibited as fish transition to faster escape swimming with the fins adducted. The clear discontinuities in movements of the pectoral fins during transitions between exploratory and escape locomotion arguably reflect changes in gait despite relatively continuous variations of tail-beat frequency.

Collectively, these observations are consistent with the idea that V0v neuron differentiation in mammals follows the logic described in flies, fish, and frogs and also provides evidence that V0v neurons operating during different modes of locomotion are functionally incompatible, since switching to a faster escape mode of locomotion involves inhibition of slower exploratory circuits.

Conclusions and predictions

Despite differences in body plans and forms of locomotion, consistent patterns are emerging related to the functional organization of even-skipped interneurons for escape and exploration. From flies and fishes, there is evidence for a temporal logic to the assembly of even-skipped circuitry, where early-born neurons participate in faster escape movements and later-born neurons in slower exploratory ones. While this has yet to be tested in frogs and

mice, one prediction is that V0v neurons engaged in the most powerful modes of locomotion are among earliest born.

From flies, frogs, fishes and mice there is evidence that morphological diversity within the even-skipped population is linked to increases in muscular and behavioral complexity. Recent work from the Dasen lab has revealed remarkably conserved transcription-factor programs underlying the specification of fin motor pools and limb motor pools [35]. This likely extends to the interneuronal networks controlling their activity and so future studies of V0v neurons providing fin-control should provide insights into the evolution of limb-control circuits. One prediction is that fin/limb V0v neurons represent a more temporally delayed cohort relative to axial V0v neurons. This cohort could be equally diverse and thus provide fin/limb–axis coordination along and across the body during different modes of locomotion.

From fishes and simulations of mouse locomotion, there is evidence for a tiered organization of even-skipped circuits, where V0v neurons active at slower speeds are inhibited as V0v neurons recruited at faster speeds are engaged. Why might there be differential control of V0v populations? Even-skipped neurons do not appear to be critical for rhythm generation, but rather help establish appropriate patterns of muscle activity across and along the body. It therefore makes some sense that interneurons responsible for producing one motor pattern are inhibited as those responsible for producing another are engaged. Future work identifying the source of this inhibition, whether from the brain or the spinal cord, is now required. It would also be worth seeing whether a similar tiered organization exists in the abdominal ganglion of flies and how EL diversity may increase commensurate with changes in body plan during fly development. For example, even-skipped interneurons are found in both larval and adult locusts [36].

In flies, EL interneurons receive distinct sources of afferent input related to their participation in escapes versus crawling. In zebrafish, it is conceivable that early-born ascending V0v neurons with their more dorsally-positioned somata are targeted by primary or secondary sensory afferents, which could evoke more forceful movements. Recent work from the Wyart lab has revealed that zebrafish cerebrospinal fluid-contacting neurons are mechanosensitive [37] and synapse onto V0v-MCoD neurons [38]. Thus, it is possible that distinct types of V0v neurons receive input from different sensory modalities to select different modes of locomotion in fish as in fly larvae. V0v neurons activated during fictive locomotion in mice are distributed close to the central canal [33], so this form of afferent modulation could also apply more broadly.

A more recent study in mice from the Kiehn lab has demonstrated that descending midbrain control systems are also organized according to their role in evasion versus exploration [39]. In addition, a very recent study in zebrafish from the Koyama lab has revealed a developmental logic to the wiring of descending control systems to spinal circuits, where hindbrain neurons active during slower swimming wire up to V0v-MCoDs after hindbrain-spinal networks for fast swimming [40]. Consequently, the staggered assembly of even-skipped networks according to function and behavioral complexity could also apply to the descending control systems responsible for producing behaviors of differing urgency. While it remains to be seen how other features of organization of Eve/Evx circuitry are either

shared or specialized between flies, fishes, frogs and mice, thanks to continuing work in these animals there should be plenty of predictions to test.

Acknowledgments

The authors are supported by the National Institutes of Health (R01-NS067299) and the National Science Foundation (IOS-1456830).

References

**Of special note

*Of note

1. Goulding M: Circuits controlling vertebrate locomotion: moving in a new direction. *Nat Rev Neurosci* 2009, 10:507–518. [PubMed: 19543221]
2. Suster ML, Kania A, Liao M, Asakawa K, Charron F, Kawakami K, Drapeau P: A novel conserved *evx1* enhancer links spinal interneuron morphology and cis-regulation from fish to mammals. *Dev Biol* 2009, 325:422–433. [PubMed: 18992237]
3. Duman-Scheel M, Patel NH: Analysis of molecular marker expression reveals neuronal homology in distantly related arthropods. *Development* 1999, 126:2327–2334. [PubMed: 10225992]
4. Doe CQ, Smouse D, Goodman CS: Control of neuronal fate by the *Drosophila* segmentation gene *even-skipped*. *Nature* 1988, 333:376–378. [PubMed: 3374572]
5. Clark MQ, Zarin AA, Carreira-Rosario A, Doe CQ: Neural circuits driving larval locomotion in *Drosophila*. *Neural Dev* 2018, 13:6. [PubMed: 29673388]
6. Pulver SR, Bayley TG, Taylor AL, Berni J, Bate M, Hedwig B: Imaging fictive locomotor patterns in larval *Drosophila*. *J Neurophysiol* 2015, 114:2564–2577. [PubMed: 26311188]
7. Heckscher ES, Zarin AA, Faumont S, Clark MQ, Manning L, Fushiki A, Schneider-Mizell CM, Fetter RD, Truman JW, Zwart MF, et al.: Even-skipped(+) interneurons are core components of a sensorimotor circuit that maintains left-right symmetric muscle contraction amplitude. *Neuron* 2015, 88:314–329. [PubMed: 26439528]
8. Wreden CC, Meng JL, Feng W, Chi W, Marshall ZD, Heckscher ES: Temporal cohorts of lineage-related neurons perform analogous functions in distinct sensorimotor circuits. *Curr Biol* 2017, 27:1521–1528 e1524. [PubMed: 28502656] ** The authors establish developmentally distinct cohorts of Eve-positive interneurons contribute to distinct motor circuits. Eve-positive interneurons integrate different sensory modalities to drive slow and fast modes of motor output.
9. Ohyama T, Schneider-Mizell CM, Fetter RD, Aleman JV, Franconville R, Rivera-Alba M, Mensh BD, Branson KM, Simpson JH, Truman JW, et al.: A multilevel multimodal circuit enhances action selection in *Drosophila*. *Nature* 2015, 520:633–639. [PubMed: 25896325]
10. Brustein E, Saint-Amant L, Buss RR, Chong M, McDearmid JR, Drapeau P: Steps during the development of the zebrafish locomotor network. *J Physiol Paris* 2003, 97:77–86. [PubMed: 14706693]
11. Hale ME: Developmental change in the function of movement systems: transition of the pectoral fins between respiratory and locomotor roles in zebrafish. *Integr Comp Biol* 2014, 54:238–249. [PubMed: 24748600]
12. Siomava N, Shkil F, Voronezhskaya E, Diogo R: Development of zebrafish paired and median fin musculature: basis for comparative, developmental, and macroevolutionary studies. *Sci Rep* 2018, 8:14187. [PubMed: 30242203]
13. Lanuza GM, Gosgnach S, Pierani A, Jessell TM, Goulding M: Genetic identification of spinal interneurons that coordinate left-right locomotor activity necessary for walking movements. *Neuron* 2004, 42:375–386. [PubMed: 15134635]
14. Moran-Rivard L, Kagawa T, Saueressig H, Gross MK, Burrill J, Goulding M: *Evx1* is a postmitotic determinant of v0 interneuron identity in the spinal cord. *Neuron* 2001, 29:385–399. [PubMed: 11239430]

15. Satou C, Kimura Y, Higashijima S: Generation of multiple classes of V0 neurons in zebrafish spinal cord: progenitor heterogeneity and temporal control of neuronal diversity. *J Neurosci* 2012, 32:1771–1783. [PubMed: 22302816]
16. Knogler LD, Drapeau P: Sensory gating of an embryonic zebrafish interneuron during spontaneous motor behaviors. *Front Neural Circuits* 2014, 8:121. [PubMed: 25324729]
17. Juarez-Morales JL, Martinez-De Luna RI, Zuber ME, Roberts A, Lewis KE: Zebrafish transgenic constructs label specific neurons in *Xenopus laevis* spinal cord and identify frog V0v spinal neurons. *Dev Neurobiol* 2017, 77:1007–1020. [PubMed: 28188691]
18. Juarez-Morales JL, Schulte CJ, Pezoa SA, Vallejo GK, Hilinski WC, England SJ, de Jager S, Lewis KE: *Evx1* and *Evx2* specify excitatory neurotransmitter fates and suppress inhibitory fates through a *Pax2*-independent mechanism. *Neural Dev* 2016, 11:5. [PubMed: 26896392]
19. Roberts A, Hill NA, Hicks R: Simple mechanisms organise orientation of escape swimming in embryos and hatchling tadpoles of *Xenopus laevis*. *J Exp Biol* 2000, 203:1869–1885. [PubMed: 10821744]
20. Sillar KT, Robertson RM: Thermal activation of escape swimming in post-hatching *Xenopus laevis* frog larvae. *J Exp Biol* 2009, 212:2356–2364. [PubMed: 19617428]
21. McLean DL, Fetcho JR: Spinal interneurons differentiate sequentially from those driving the fastest swimming movements in larval zebrafish to those driving the slowest ones. *J Neurosci* 2009, 29:13566–13577. [PubMed: 19864569]
22. McLean DL, Masino MA, Koh IY, Lindquist WB, Fetcho JR: Continuous shifts in the active set of spinal interneurons during changes in locomotor speed. *Nat Neurosci* 2008, 11:1419–1429. [PubMed: 18997790]
23. Green MH, Hale ME: Activity of pectoral fin motoneurons during two swimming gaits in the larval zebrafish (*Danio rerio*) and localization of upstream circuit elements. *J Neurophysiol* 2012, 108:3393–3402. [PubMed: 23034362]
24. McLean DL, Fan J, Higashijima S, Hale ME, Fetcho JR: A topographic map of recruitment in spinal cord. *Nature* 2007, 446:71–75. [PubMed: 17330042]
25. Bjornfors ER, El Manira A: Functional diversity of excitatory commissural interneurons in adult zebrafish. *Elife* 2016, 5.** The authors provide a detailed characterization of the morphological and electrophysiological properties of V0v interneurons at juvenile/adult stages in zebrafish. The majority of V0v types are not active during low-frequency exploratory locomotion, consistent with studies at larval stages.
26. Kyriakatos A, Mahmood R, Ausborn J, Porres CP, Buschges A, El Manira A: Initiation of locomotion in adult zebrafish. *J Neurosci* 2011, 31:8422–8431. [PubMed: 21653846]
27. Gabriel JP, Mahmood R, Walter AM, Kyriakatos A, Hauptmann G, Calabrese RL, El Manira A: Locomotor pattern in the adult zebrafish spinal cord in vitro. *J Neurophysiol* 2008, 99:37–48. [PubMed: 17977928]
28. Masino MA, Fetcho JR: Fictive swimming motor patterns in wild type and mutant larval zebrafish. *J Neurophysiol* 2005, 93:3177–3188. [PubMed: 15673549]
29. McLean DL, Dougherty KJ: Peeling back the layers of locomotor control in the spinal cord. *Curr Opin Neurobiol* 2015, 33:63–70. [PubMed: 25820136]
30. Bellardita C, Kiehn O: Phenotypic characterization of speed-associated gait changes in mice reveals modular organization of locomotor networks. *Curr Biol* 2015, 25:1426–1436. [PubMed: 25959968]
31. Talpalar AE, Bouvier J, Borgius L, Fortin G, Pierani A, Kiehn O: Dual-mode operation of neuronal networks involved in left-right alternation. *Nature* 2013, 500:85–88. [PubMed: 23812590]
32. Ruder L, Takeoka A, Arber S: Long-distance descending spinal neurons ensure quadrupedal locomotor stability. *Neuron* 2016, 92:1063–1078. [PubMed: 27866798] * By retrograde labeling, the authors demonstrated the presence of long-range descending V0vs linking cervical and lumbar regions in the mouse spinal cord. Ablation of these cells revealed a role for long-range projecting V0vs during motor output which was previously not appreciated.
33. Griener A, Zhang W, Kao H, Wagner C, Gosgnach S: Probing diversity within subpopulations of locomotor-related V0 interneurons. *Dev Neurobiol* 2015, 75:1189–1203. [PubMed: 25649879]

34. Danner SM, Shevtsova NA, Frigon A, Rybak IA: Computational modeling of spinal circuits controlling limb coordination and gaits in quadrupeds. *Elife* 2017, 6.* In this study, the authors provide a computational model of quadrupedal spinal circuits. By incorporating recent results, the authors were able to successfully recapitulate the behavioural results of experimental manipulations of different cell populations. The model also suggests unique roles for local and long-range V0vs.
35. Jung H, Baek M, D'Elia KP, Boisvert C, Currie PD, Tay BH, Venkatesh B, Brown SM, Heguy A, Schoppik D, et al.: The Ancient Origins of Neural Substrates for Land Walking. *Cell* 2018, 172:667–682 e615. [PubMed: 29425489] ** The authors demonstrate that the molecular program underpinning the specification of fin and limb motor neuron pools is conserved between tetrapods and fish that use their pelvic fins to locomote. These results suggest the basic spinal networks coordinating locomotion are highly conserved.
36. Bevan S, Burrows M: Localisation of Even-skipped in the mature CNS of the locust, *Schistocerca gregaria*. *Cell Tissue Res* 2003, 313:237–244. [PubMed: 12883992]
37. Böhm UL, Prendergast A, Djenoune L, Nunes Figueiredo S, Gomez J, Stokes C, Kaiser S, Suster M, Kawakami K, Charpentier M, et al.: CSF-contacting neurons regulate locomotion by relaying mechanical stimuli to spinal circuits. *Nat Commun* 2016, 7:10866. [PubMed: 26946992]
38. Fidelin K, Djenoune L, Stokes C, Prendergast A, Gomez J, Baradel A, Del Bene F, Wyart C: State-dependent modulation of locomotion by GABAergic spinal sensory neurons. *Curr Biol* 2015, 25:3035–3047. [PubMed: 26752076] ** This study is the first to demonstrate V0v interneurons receive synaptic inputs from a unique class of cerebrospinal fluid-contacting mechanosensitive interneuron, thus further demonstrating conserved sensory integration functions across species.
39. Caggiano V, Leiras R, Goni-Errro H, Masini D, Bellardita C, Bouvier J, Caldeira V, Fisone G, Kiehn O: Midbrain circuits that set locomotor speed and gait selection. *Nature* 2018, 553:455–460. [PubMed: 29342142] * The authors identify two key midbrain structures which comprise the mesencephalic locomotor region and examine how these structures contribute to different modes of motor output.
40. Pujala A, Koyama M: Chronology-based architecture of descending circuits that underlie the development of locomotor repertoire after birth. *bioRxiv* 2018, 10.1101/425587.** The authors examined the development and connectivity between hindbrain and spinal interneurons, including the descending V0v population during zebrafish development. In doing so, they revealed that early born hindbrain-spinal circuits are recruited during escape behaviours whilst late-born hindbrain-spinal pathways are recruited during slower behaviours.

Highlights

- The transcription factor even-skipped labels interneurons in many animals
- Even-skipped interneurons are functionally organized by birth order
- Even-skipped neuron diversity is linked to muscular and behavioural complexity
- Even-skipped circuits are hierarchically arranged based on movement intensity