Published in final edited form as:

Cancer Prev Res (Phila). 2019 November ; 12(11): 771–780. doi:10.1158/1940-6207.CAPR-19-0200.

Racial/ethnic patterns of young-onset noncardia gastric cancer

Andreana N. Holowatyj, PhD, MS^{1,2,*}, Cornelia M. Ulrich, MS, PhD^{1,2}, Mark A. Lewis, MD³

¹Huntsman Cancer Institute, Salt Lake City, Utah

²Department of Population Health Sciences, University of Utah, Salt Lake City, Utah

³Department of Internal Medicine, Intermountain Healthcare, Murray, Utah

Abstract

Increasing noncardia gastric cancer incidence rates among individuals age younger than 50 years have gained much attention, particularly as causes remain unknown. Using population-based NIH/NCI's Surveillance, Epidemiology, and End Results (SEER) program data from 2007 to 2015, multivariable logistic regression was used to quantify associations between race/ethnicity and clinicodemographic features among young-onset noncardia gastric cancer patients. A total of 2,872 individuals aged 20 to 49 years were diagnosed with primary noncardia gastric cancer. Age at diagnosis, insurance status, anatomic sub-site, AJCC clinical stage, histologic type, tumor grade, surgery and county-level smoking prevalence differed by race/ethnicity (all $P < 0.003$). Compared to non-Hispanic whites, Hispanics were more likely to be diagnosed at younger ages (Odds Ratio [OR]=0.97, 95% CI 0.95-0.99), on Medicaid/uninsured (OR=3.83, 95% CI 2.89-5.08), diagnosed with higher grade tumors (OR=1.93, 95% CI 1.32-2.84), and less likely to undergo surgery (OR=0.62, 95% CI 0.44-0.88) or to reside in counties with higher smoking prevalence (OR=0.15, 95% CI 0.11-0.21) after adjustment for sex, sub-site, and histologic type. Asian/Pacific Islanders were more likely to be female (OR=1.40, 95% CI 1.04-1.88), and less likely to be diagnosed with metastatic disease (OR=0.59, 95% CI 0.37-0.95) or to reside in counties with higher smoking prevalence (OR=0.13, 95% CI 0.08-0.19). Approximately two in every five patients with young-onset noncardia gastric cancer are Hispanic. Further investigation into the molecular heterogeneity of young-onset noncardia gastric cancers by race/ethnicity to understand etiologies underlying this rising disease epidemic is warranted. This population-based cohort study sheds light that biologic and environmental factors may partly underlie race/ethnicity-related differences in young-onset noncardia gastric cancer susceptibility and outcomes.

***Corresponding Author:** Andreana N. Holowatyj, PhD, MS, Huntsman Cancer Institute, University of Utah, 2000 Circle of Hope, Room 4746, Salt Lake City, Utah 84112, Holowatyj@hci.utah.edu, Phone: 801-213-6202; Fax: 801-585-0900.
Author Contributions

A.N.H. and M.A.L. had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. A.N.H. and M.A.L. contributed to the conception and design of the study. All authors participated in the acquisition, analysis and interpretation of data; and drafting and critical revision of the manuscript for important intellectual content. A.N.H. participated in creating figures and in the statistical analysis. A.N.H., C.M.U. and M.A.L. obtained funding and provided support and supervision for this study.

Conflict of Interest

The authors declare no potential conflicts of interest with this work. C.M.U. has as cancer center director oversight over research funded by several pharmaceutical companies, but has not received funding directly herself.

Keywords

Gastric cancer; stomach; non-cardia; young-onset; race; SEER; ethnicity; Hispanic; Asian; non-Hispanic black; early-onset; gastrointestinal cancer; smoking

Introduction

As demographics, diet, excess body weight, smoking prevalence and exposures to other cancer-related risk factors have changed over time, the burden of noncardia gastric cancer has also evolved.[1-6] Population-based studies and studies from academic institutions have both reported racial/ethnic disparities in gastric cancer incidence and outcomes among all patients or among young-onset cases (aged younger than 50 years at diagnosis) relative to older-onset cases.[2, 6-8] Anderson and colleagues examined incidence rates of noncardia gastric cancer by age groups, noting that rates are rising from older to younger generations. [2] Indeed, noncardia gastric cancer incidence rates have increased among young non-Hispanic white, non-Hispanic black, and Hispanic women, as well as among young men.[6] Gupta et al. recently demonstrated that among all-comers, incident noncardia gastric cancer rate ratios were higher among non-Hispanic blacks, Hispanics, Asian or Pacific Islanders, and American Indian or Alaska Natives compared with non-Hispanic whites.[9] However, no studies to date have examined racial/ethnic differences specifically within the population of young individuals diagnosed with gastric cancer. Given the increasing burden of young-onset noncardia gastric cancers,[1, 2, 5, 6, 10] we evaluated racial/ethnic patterns among men and women diagnosed between the ages of 20 to 49 years with a first primary noncardia gastric cancer in the United States.

Patients and Methods

Young-onset noncardia gastric cancer cases were identified using the National Institutes of Health (NIH) and National Cancer Institute's (NCI) Surveillance, Epidemiology, and End Results (SEER) program.[11] The SEER program collects data on cancer incidence and mortality from 18 population-based cancer registries that cover approximately 28% of the United States population.[12] The dataset used is publicly available (<https://seer.cancer.gov/data/access.html>) and was exempt from Institutional Review Board review. This study was performed in accordance with the Declaration of Helsinki.

A case listing session in SEER*Stat was run on the SEER18 incidence dataset [11] to obtain clinicodemographic features on pathologically-confirmed first primary noncardia gastric cancer cases among individuals aged 20 to 49 years at diagnosis (young-onset noncardia gastric cancer cases). We restricted our analysis to young-onset noncardia gastric cancer cases diagnosed between 2007 and 2015 with known age and race/ethnicity, as 2007 is the first year for which individual-level insurance status data were available in SEER. Our final cohort was comprised of 2,872 patients diagnosed with a first primary young-onset noncardia gastric cancer.

Clinicodemographic characteristics examined included: patient age at diagnosis, sex, race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic/Spanish/Latino, Asian or

Pacific Islander, American Indian/Alaska Native), individual-level insurance status (privately insured, Medicaid/uninsured), anatomic sub-site (C16.1-C16.6), histologic type (recorded from pathology reports according to International Classification of Disease for Oncology [ICD-O] codes), tumor grade (well to moderately differentiated, poorly differentiated to undifferentiated), American Joint Committee on Cancer (AJCC) clinical tumor stage, cancer sequence (only 1 primary tumor, first of 2+ primaries), and surgical therapy as part of first course of therapy (yes, no). Histologic types were classified into diffuse type, intestinal type, or other type, using the Lauren classification [13] previously described in Henson et al.[14] Based upon the Lauren classification,[13] diffuse type gastric carcinoma included: diffuse carcinoma (8145); signet ring cell carcinoma (8490); and linitis plastica, a gross descriptive term referring to a signet ring cell and/or diffuse carcinoma within the stomach wall (8142). Intestinal type gastric carcinoma included: carcinoma, not otherwise specified (8010); adenocarcinoma, not otherwise specified (8140); adenocarcinoma, intestinal type (8144); and tubular adenocarcinoma (8211). All other histologic types were classified as other tumor types.

It has been previously suggested that race/ethnicity might be a proxy for socioeconomic status.[15-17] For this reason, we used individual-level insurance status as an approximation of socioeconomic status. Evidence also supports that cigarette smoking is associated with increased risk of gastric cancer.[18-21] Consequently, we used an area-based measure of adult smoking prevalence that is available in SEER for an approximation of current smokers. This measure was the percentage of current smokers age 18+ years taken from the Behavioral Risk Factor Surveillance System (BRFSS) and the National Health Interview Survey (NHIS) to model small area estimates for current smoking prevalence between 2008 to 2010 (<https://sae.cancer.gov/nhis-brfss/>). On the basis of the distribution in our cohort of patients with young-onset noncardia gastric cancer, tertiles of the percentage of current adult smokers (smoking index) were created. Tertile 1 (T1) represents the third of the cohort that resides in areas where the lowest proportion of adults are current smokers, and tertile 3 (T3) represents the proportion of the cohort that resides in areas with the higher proportion of current adult smokers, divided as follows: T1: <13% current smokers age 18+ years; T2: 13 to 17% current smokers age 18+ years; and T3: 18%+ current smokers age 18+ years.

To assess patterns of young-onset noncardia gastric cancer among individuals age 20 to 49 years at diagnosis, we compared clinicodemographic features by race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic/Spanish/Latino, Asian or Pacific Islander, and American Indian/Alaska Native) using chi-square tests and *t*-tests for categorical and continuous variables, respectively. Multivariable logistic regression was used to estimate odds ratios (OR) and 95% confidence intervals (CI) to quantify associations between clinicodemographic features and race/ethnicity as the outcome, where the reference outcome category was non-Hispanic white individuals diagnosed with young-onset gastric cancer. Associations between clinicodemographic features and race/ethnicity were assessed in adjusted models that included: age at diagnosis, sex, individual-level insurance status, anatomic sub-site, AJCC clinical stage, histologic type, tumor grade, smoking index, and the use of surgical resection, based upon patients having complete covariate information. Among patients with unknown AJCC clinical stage, reasons could include: lack of information available in the patient chart, ineligibility for pathological staging, or

classification criteria not met for assigning a valid AJCC stage value. Data were analyzed using SAS version 9.4 statistical software (SAS Institute). All tests were two-sided, and $P < 0.05$ was considered to be statistically significant.

Results

Between 2007 and 2015, a total of 2,872 individuals aged 20 to 49 years were diagnosed with a first primary young-onset noncardia gastric cancer. Baseline clinical and demographic characteristics of individuals by race/ethnicity are described in Table 1. Young-onset noncardia gastric cancers were most commonly diagnosed among Hispanics, as nearly 2 in every 5 individuals (37.8%) with young-onset noncardia gastric cancer were Hispanic. Non-Hispanic whites comprised approximately one-quarter (27.2%) of the population. Mean age of diagnosis was 41.6 years (sd 6.4 years) and differed by race/ethnicity, as Hispanics were more likely to be diagnosed at younger ages (40.8 years) compared with non-Hispanic whites, non-Hispanic blacks, Asian or Pacific Islanders, or American Indian/Alaska Natives (42.0 years, 42.4 years, 41.7 years, and 42.1 years, respectively; $P < 0.0001$). Individual-level insurance status, anatomic sub-site, AJCC clinical stage, histologic type, tumor grade, surgical resection, and county-level adult smoking prevalence also statistically significantly differed by race/ethnicity (all $P < 0.003$; Table 1 and Figure 1).

Over half (59.2%) of young-onset noncardia gastric cancers were located in the corpus and antrum (Table 1). Differences by disease stage were most pronounced between non-Hispanic whites or Asian/Pacific Islanders and Hispanic individuals—while approximately one-third (33.7%) of young-onset noncardia gastric cancers were diagnosed at stage IV among non-Hispanic whites as well as Asian or Pacific Islanders, nearly half (46.9%) of cases among Hispanics presented with metastatic disease (pairwise $P < 0.0001$; Table 1 and Figure 1). By histologic type, 41.5% of young-onset noncardia gastric cancers were of the diffuse type (which includes diffuse carcinomas, signet ring cell carcinomas, and linitis plastica) (Table 1). Non-Hispanic whites were least likely to present with poorly differentiated to undifferentiated (grade III/IV) tumors across racial/ethnic groups. Over half of young individuals (57.2%) diagnosed with noncardia gastric cancer underwent surgery as part of first course of therapy. Among young-onset noncardia gastric cancer patients, American Indian/Alaska Natives and non-Hispanic blacks were most likely to reside in areas with the highest prevalence of current adult smokers.

To evaluate associations between race/ethnicity and clinicodemographic features among individuals diagnosed with young-onset noncardia gastric cancer, we performed adjusted multivariable logistic regression analysis (Table 2). Age at diagnosis was inversely associated with Hispanic race/ethnicity (OR 0.97, 95%CI 0.95-0.99, $P = 0.008$). Compared to non-Hispanic whites with young-onset noncardia gastric cancer, Hispanics were also more likely to be on Medicaid/uninsured (OR 3.83, 95%CI 2.89-5.08, $P < 0.0001$), and were nearly twice as likely to be diagnosed with higher grade (III/IV) tumors (OR 1.93, 95%CI 1.32-2.84, $P = 0.001$). Hispanic individuals were 38% less likely to undergo surgery (OR 0.62, 95%CI 0.44-0.88, $P = 0.007$) and were 46% to 85% less likely to reside in counties with higher prevalence of adult smoking (T2: OR 0.54, 95%CI 0.40-0.74, $P = 0.0001$; T3: OR 0.15, 95%CI 0.11-0.21, $P < 0.0001$). While young non-Hispanic blacks were 82% more likely

to be diagnosed with stage II noncardia gastric cancers (OR 1.82, 95%CI 1.12-2.96, $P=0.016$), no associations with advanced stage (stage III/IV) disease were observed among young non-Hispanic blacks versus non-Hispanic whites.

Asian or Pacific Islanders with young-onset noncardia gastric cancer were statistically significantly more likely to be female (OR 1.40, 95%CI 1.04-1.88, $P=0.025$) and were 41% less likely to be diagnosed with metastatic disease (OR 0.59, 95%CI 0.37-0.95, $P=0.031$) compared to young non-Hispanic white individuals. Similar to Hispanics, Asian or Pacific Islanders with young-onset noncardia gastric cancer were also nearly twice as likely to be diagnosed with grade III/IV tumors (OR 1.91, 95%CI 1.22-2.97, $P=0.005$) and were 46% to 87% less likely to reside in counties with a higher prevalence of adult smoking (T2: OR 0.54, 95%CI 0.38-0.76, $P=0.0004$; T3: OR 0.13, 95%CI 0.08-0.19, $P<0.0001$) versus non-Hispanic whites (Table 2). Aligned with evidence that American Indians/Alaska Natives have one of the highest cigarette smoking prevalences of all racial/ethnic groups,[19-21] American Indian/Alaska Native individuals with young-onset noncardia gastric cancer were over five-fold more likely to reside in areas with the highest adult smoking prevalence compared with young non-Hispanic whites (T3: OR 5.69, 95%CI 1.29-25.02, $P=0.021$) (Table 2).

Discussion

Young-onset noncardia gastric cancer incidence rates are rising alarmingly, with underlying causes unknown.[1-3, 5-7, 10] Our investigation of 2,872 individuals diagnosed with young-onset noncardia gastric cancer in the United States from 2007 to 2015 identified racial/ethnic disparities by tumor grade, county-level adult smoking prevalence, age at diagnosis, and sex. Approximately two in every five patients with young-onset noncardia gastric cancer were Hispanic. Further, Hispanic individuals were more likely to be diagnosed at younger ages and with high grade tumors, and were less likely to undergo surgery or reside in counties with higher prevalence of adult smoking compared with non-Hispanic whites. Asian/Pacific Islanders diagnosed with young-onset noncardia gastric cancer were more likely to be female and to be diagnosed with high grade tumors, and were less likely to be diagnosed with metastatic disease or to reside in counties with higher adult smoking prevalence. American Indian/Alaska Natives diagnosed with young-onset noncardia gastric cancer were more likely to reside in areas with the highest prevalence of current adult smokers. To our knowledge, this population-based cohort study is the first to evaluate clinicodemographic characteristics, including tumor grade, county-level smoking prevalence and race/ethnicity, within the population of individuals diagnosed with young-onset noncardia gastric cancer.

The landscape of race/ethnicity is evolving across the United States, and Hispanics are expected to account for nearly one-third of the population by year 2060.[22] Aligned with these changing demographics, Islami et al. revealed gastric noncardia adenocarcinoma rates increased in younger age groups among men (across racial/ethnic groups) as well as among non-Hispanic white, non-Hispanic black, and Hispanic women using nationwide data from the North American Association of Central Cancer Registries.[6] Balakrishnan and colleagues [10] reported that gastric cancer incidence remained stable among non-Hispanic whites and non-Hispanic blacks, but noted a significant rise in incidence among Hispanic

adults in a cancer registry-based study among all patients across a single county in Texas. A recent retrospective study in Mexico examined clinicopathologic features of gastric carcinomas in Latin patients and noted an increased incidence of gastric cancer among patients age 40 years and younger compared to patients over the age of 40 years at diagnosis.[5] Using the National Cancer Database, De et al. reported significant differences in race/ethnicity between individuals age younger than 40 years versus those 40 years of age and older diagnosed with gastric adenocarcinoma in a similar sized cohort, as Hispanic individuals were significantly more likely to be diagnosed with gastric adenocarcinoma at younger ages—concordant with our results.[7] Rona and colleagues also compared clinicopathological features and outcomes among patients diagnosed with gastric carcinoma at an academic institution and noted a higher incidence of advanced stage tumors, poorly differentiated (grade III) carcinomas, and signet-cell type tumors among young patients—of which the majority were Hispanics—when compared to older counterparts.[8] Consistent with these findings, our population-based study suggests that approximately two in every five patients diagnosed with young-onset noncardia gastric cancer in the United States are Hispanic. Further, we observed that Hispanic individuals were also more likely to be diagnosed with high grade (grade III/IV) tumors. Hispanics with young-onset noncardia gastric cancer were more likely to be on Medicaid or uninsured, and were less likely to undergo surgery compared to non-Hispanic whites. Together with population-based findings that noncardia gastric cancer incidence rates are rising among young (age younger than 50 years) non-Hispanic white and Hispanic individuals,[2, 4, 23] and that higher rates of *Helicobacter pylori* seropositivity are detected among Mexican Americans compared with non-Hispanic white adults,[24] our novel findings in this population of young individuals suggest that additional research is needed to examine clinicopathologic characteristics of noncardia gastric cancers among Hispanic/Spanish/Latino subpopulations by country of origin. Genetic ancestral heritage and gene-environment interactions [25-28] may contribute to these pronounced racial/ethnic disparities in young-onset noncardia gastric cancer.

It is estimated that approximately 14% of all adults in the United States are current smokers, and that approximately four out of every ten cancer cases are linked to tobacco use.[29] Evidence also supports that cigarette smoking is associated with an increased risk of gastric cancer,[18-21] and cigarette smoking rates sharply differ by race/ethnicity. In particular, American Indian/Alaska Natives have one of the highest cigarette smoking prevalences across all racial/ethnic groups,[19-21] and the risk for gastric cancer is higher among American Indian/Alaska Native young adults compared to non-Hispanic whites.[30] Aligned with these patterns, we observed that American Indian/Alaska Natives with young-onset noncardia gastric cancer were more likely to reside in areas with the highest prevalence of current adult smokers versus young non-Hispanic whites. Together with other environmental exposures including diet, excess body weight, and microbiome composition,[2, 23, 27, 31, 32] these health behaviors may be contributing to distinct molecular phenotypes of young-onset gastrointestinal tumors by race/ethnicity. Additional studies are needed to investigate the role of these environmental exposures and health behaviors by race/ethnicity on young-onset gastrointestinal cancer outcomes.

Although our findings suggest racial/ethnic differences in noncardia gastric cancer within the population of young individuals, we acknowledge limitations of our study. SEER does

not collect individual-level data on environmental exposures and lifestyle/behavioral factors (e.g. *Helicobacter pylori* infection, history of cigarette smoking, dietary patterns) or molecular phenotypes of tumors, does not include complete information about chemotherapy and/or radiation therapy, there could exist potential reporting differences in clinical stage by race/ethnicity, as well as the small sample size in some racial/ethnic (e.g. American Indian/Alaska Native) groups. Given that race/ethnicity is classified based on self-identification in SEER, it is subject to misclassification. Our findings raise the possibility that interactions between noncardia gastric tumor biology and treatments might contribute to racial/ethnic differences among young patients, particularly by disease stage. However, our study does not provide data about treatment regimens nor the molecular phenotypes of tumors among individuals with young-onset noncardia gastric cancer.

Nevertheless, our examination of the racial/ethnic patterns of young-onset noncardia gastric cancer using a population-based cohort yields important implications for gastric cancer risk assessment and prevention. The inherent clinical heterogeneity of gastric cancer is reflected in distinct molecular subtypes of cancer cells, where integrative genomic and proteomic patterns have characterized four molecular subtypes of disease that predict survival and adjuvant chemotherapy outcomes among patients of all ages diagnosed with gastric carcinoma.[33] These molecular characteristics of gastric cancer have also revealed intriguing molecular phenotypes of gastric tumors by age at diagnosis, as patients with the genomically stable subtype—associated with the poorest prognosis, the least benefit from adjuvant chemotherapy, and less distinctive genomic alterations—were diagnosed at younger ages relative to patients with the other subtypes.[33] By race/ethnicity, a single institution study revealed differences in mutation rates among all-comers diagnosed with gastric adenocarcinoma, as non-Hispanic black patients were found to have higher rates of *TP53* mutations compared to non-Hispanic whites, Hispanic, or Asian/Pacific Islander cases.[34] Importantly, noncardia gastric cancer has a distinct disease biology from adenocarcinoma arising in the cardia, which is much more similar to esophagogastric junction and lower esophagus (Siewert I tumors) cancers in terms of natural history and risk factors for oncogenesis.[35, 36] Together, these results suggest that distinct molecular phenotypes of gastric tumors may be associated with differential response to therapies, and evaluation of these phenotypes by race/ethnicity specifically among young patients with noncardia gastric cancer could have significant implications for the clinical management of these patients.

In conclusion, these novel findings support that biologic and environmental (e.g. *Helicobacter pylori* infection, cigarette smoking, diet) factors [4, 18, 21, 23, 24, 27, 31, 32, 35] may partly underlie race-related differences in young-onset noncardia gastric cancer susceptibility and outcomes. Studies are warranted to further investigate the heterogeneity of malignancies among individuals diagnosed with young-onset noncardia gastric cancer.

Acknowledgments

Funding

A.N.H. was supported by the National Institutes of Health under Ruth L. Kirschstein National Research Service Award T32 HG008962 from the National Human Genome Research Institute. This work was also supported by grants from the National Institute of Health/National Cancer Institute (U01 CA206110, R01 CA189184, and R01 CA211705 to C.M.U.) and the Huntsman Cancer Foundation. The research reported in this publication was

supported by the National Cancer Institute of the National Institutes of Health under award number P30 CA042014. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Anderson WF, Camargo MC, Fraumeni JF Jr., et al. Age-specific trends in incidence of noncardia gastric cancer in US adults. *JAMA* 2010; 303: 1723–1728. [PubMed: 20442388]
2. Anderson WF, Rabkin CS, Turner N et al. The Changing Face of Noncardia Gastric Cancer Incidence Among US Non-Hispanic Whites. *J Natl Cancer Inst* 2018; 110: 608–615. [PubMed: 29361173]
3. Camargo MC, Anderson WF, King JB et al. Divergent trends for gastric cancer incidence by anatomical subsite in US adults. *Gut* 2011; 60: 1644–1649. [PubMed: 21613644]
4. Merchant SJ, Kim J, Choi AH et al. A rising trend in the incidence of advanced gastric cancer in young Hispanic men. *Gastric Cancer* 2017; 20: 226–234. [PubMed: 26924751]
5. Calderillo-Ruiz G, Takahashi A, Herrera M et al. P-145 Gastric cancer in young Latin women: bad prognostic factors and outcomes. *Annals of Oncology* 2019; 30.
6. Islami F, DeSantis CE, Jemal A. Incidence Trends of Esophageal and Gastric Cancer Subtypes by Race, Ethnicity, and Age in the United States, 1997–2014. *Clin Gastroenterol Hepatol* 2019; 17: 429–439. [PubMed: 29902641]
7. De B, Rhome R, Jairam V et al. Gastric adenocarcinoma in young adult patients: patterns of care and survival in the United States. *Gastric Cancer* 2018; 21: 889–899. [PubMed: 29691758]
8. Rona KA, Schwameis K, Zehetner J et al. Gastric cancer in the young: An advanced disease with poor prognostic features. *J Surg Oncol* 2017; 115: 371–375. [PubMed: 28008624]
9. Gupta S, Tao L, Murphy JD et al. Race/Ethnicity-, Socioeconomic Status-, and Anatomic Subsite-Specific Risks for Gastric Cancer. *Gastroenterology* 2019; 156: 59–62 e54. [PubMed: 30267713]
10. Balakrishnan M, George R, Sharma A et al. An Investigation into the Recent Increase in Gastric Cancer in the USA. *Dig Dis Sci* 2018; 63: 1613–1619. [PubMed: 29546646]
11. Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence - SEER 18 Regs Research Data + Hurricane Katrina Impacted Louisiana Cases, 11 2017 Sub (1973–2015 varying) - Linked To County Attributes - Total U.S., 1969–2016 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released April 2018, based on the November 2017 submission.
12. Noone AM, Howlader N, Krapcho M, Miller D, Brest A, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). *SEER Cancer Statistics Review, 1975–2015*, National Cancer Institute Bethesda, MD, https://seer.cancer.gov/csr/1975_2015/, based on November 2017 SEER data submission, posted to the SEER web site, 4 2018.
13. Lauren P The Two Histological Main Types of Gastric Carcinoma: Diffuse and So-Called Intestinal-Type Carcinoma. An Attempt at a Histo-Clinical Classification. *Acta Pathol Microbiol Scand* 1965; 64: 31–49. [PubMed: 14320675]
14. Henson DE, Dittus C, Younes M et al. Differential trends in the intestinal and diffuse types of gastric carcinoma in the United States, 1973–2000: increase in the signet ring cell type. *Arch Pathol Lab Med* 2004; 128: 765–770. [PubMed: 15214826]
15. Soneji S, Iyer SS, Armstrong K, Asch DA. Racial disparities in stage-specific colorectal cancer mortality: 1960–2005. *Am J Public Health* 2010; 100: 1912–1916. [PubMed: 20724684]
16. Holowatyj AN, Ruterbusch JJ, Rozek LS et al. Racial/Ethnic Disparities in Survival Among Patients With Young-Onset Colorectal Cancer. *J Clin Oncol* 2016; 34: 2148–2156. [PubMed: 27138583]
17. Holowatyj AN, Lewis MA, Pannier ST et al. Clinicopathologic and Racial/Ethnic Differences of Colorectal Cancer Among Adolescents and Young Adults. *Clin Transl Gastroenterol* 2019; 10: e00059. [PubMed: 31259751]
18. Sasazuki S, Sasaki S, Tsugane S, Japan Public Health Center Study G. Cigarette smoking, alcohol consumption and subsequent gastric cancer risk by subsite and histologic type. *Int J Cancer* 2002; 101: 560–566. [PubMed: 12237898]

19. Odani S, Armour BS, Graffunder CM, Garrett BE, Agaku IT. Prevalence and Disparities in Tobacco Product Use Among American Indians/Alaska Natives — United States, 2010–2015. *MMWR Morb Mortal Wkly Rep* 2017;66:1374–1378. DOI: 10.15585/mmwr.mm6650a2. [PubMed: 29267265]
20. United States Department of Health and Human Services. The health consequences of smoking—50 years of progress: a report of the Surgeon General, 2014. Atlanta, GA: US Department of Health and Human Services, CDC; 2014 <https://www.surgeongeneral.gov/library/reports/50-years-of-progress/index.html>External.
21. Nez Henderson P, Jacobsen C, Beals J, Team A-S. Correlates of cigarette smoking among selected Southwest and Northern plains tribal groups: the AI-SUPERPFP Study. *Am J Public Health* 2005; 95: 867–872. [PubMed: 15855467]
22. Siegel R, Naishadham D, Jemal A. Cancer statistics for Hispanics/Latinos, 2012. *CA Cancer J Clin* 2012; 62: 283–298. [PubMed: 22987332]
23. Chang ET, Gomez SL, Fish K et al. Gastric cancer incidence among Hispanics in California: patterns by time, nativity, and neighborhood characteristics. *Cancer Epidemiol Biomarkers Prev* 2012; 21: 709–719. [PubMed: 22374991]
24. Grad YH, Lipsitch M, Aiello AE. Secular trends in *Helicobacter pylori* seroprevalence in adults in the United States: evidence for sustained race/ethnic disparities. *Am J Epidemiol* 2012; 175: 54–59. [PubMed: 22085628]
25. Sun Y, Gu J, Ajani JA et al. Genetic and intermediate phenotypic susceptibility markers of gastric cancer in Hispanic Americans: a case-control study. *Cancer* 2014; 120: 3040–3048. [PubMed: 24962126]
26. Stern MC, Fejerman L, Das R et al. Variability in Cancer Risk and Outcomes Within US Latinos by National Origin and Genetic Ancestry. *Curr Epidemiol Rep* 2016; 3: 181–190. [PubMed: 27547694]
27. Kim J, Cho YA, Choi WJ, Jeong SH. Gene-diet interactions in gastric cancer risk: a systematic review. *World J Gastroenterol* 2014; 20: 9600–9610. [PubMed: 25071358]
28. Skierucha M, Milne AN, Offerhaus GJ et al. Molecular alterations in gastric cancer with special reference to the early-onset subtype. *World J Gastroenterol* 2016; 22: 2460–2474. [PubMed: 26937134]
29. National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health The health consequences of smoking—50 years of progress: a report of the surgeon general. Atlanta, GA: Centers for Disease Control and Prevention, 2014.
30. Weir HK, Jim MA, Marrett LD, Fairley T. Cancer in American Indian and Alaska Native young adults (ages 20–44 years): US, 1999–2004. *Cancer* 2008; 113: 1153–1167. [PubMed: 18720386]
31. Castello A, Fernandez de Larrea N, Martin V et al. High adherence to the Western, Prudent, and Mediterranean dietary patterns and risk of gastric adenocarcinoma: MCC-Spain study. *Gastric Cancer* 2018; 21: 372–382. [PubMed: 29139048]
32. Ulrich CM, Himbert C, Holowatyj AN, Hursting SD. Energy balance and gastrointestinal cancer: risk, interventions, outcomes and mechanisms. *Nat Rev Gastroenterol Hepatol* 2018; 15: 683–698. [PubMed: 30158569]
33. Sohn BH, Hwang JE, Jang HJ et al. Clinical Significance of Four Molecular Subtypes of Gastric Cancer Identified by The Cancer Genome Atlas Project. *Clin Cancer Res* 2017.
34. van Beek E, Hernandez JM, Goldman DA et al. Rates of TP53 Mutation are Significantly Elevated in African American Patients with Gastric Cancer. *Ann Surg Oncol* 2018; 25: 2027–2033. [PubMed: 29725898]
35. Kamangar F, Dawsey SM, Blaser MJ et al. Opposing risks of gastric cardia and noncardia gastric adenocarcinomas associated with *Helicobacter pylori* seropositivity. *J Natl Cancer Inst* 2006; 98: 1445–1452. [PubMed: 17047193]
36. Shao L, Li P, Ye J et al. Risk of gastric cancer among patients with gastric intestinal metaplasia. *Int J Cancer* 2018.

Characteristic		NHW	NHB	Hispanic	API	AI/AN
Age at Diagnosis	NHW		0.90	0.0007	0.34	0.56
	NHB			0.002	0.39	0.61
	Hispanic				0.06	0.43
	API					0.75
	AI/AN					
Sex	NHW		0.19	0.15	0.29	0.99
	NHB			0.87	0.03	0.65
	Hispanic				0.019	0.68
	API					0.70
	AI/AN					
Individual Insurance Status	NHW		<0.0001	<0.0001	0.09	<0.0001
	NHB			0.0003	<0.0001	0.0003
	Hispanic				<0.0001	0.01
	API					<0.0001
	AI/AN					
Anatomic Sub-site	NHW		0.45	0.008	0.0005	0.08
	NHB			0.29	0.13	0.30
	Hispanic				0.21	0.09
	API					0.18
	AI/AN					
AJCC Clinical Stage	NHW		0.003	<0.0001	0.01	0.51
	NHB			0.007	0.21	0.96
	Hispanic				<0.0001	0.72
	API					0.64
	AI/AN					
Histologic Type	NHW		<0.0001	<0.0001	<0.0001	0.0002
	NHB			<0.0001	0.0001	0.06
	Hispanic				0.03	0.03
	API					0.25
	AI/AN					
Tumor Grade	NHW		0.02	<0.0001	<0.0001	0.36
	NHB			0.0002	0.004	0.96
	Hispanic				0.79	0.19
	API					0.24
	AI/AN					
Cancer Sequence	NHW		0.68	0.01	0.27	0.51
	NHB			0.08	0.53	0.60
	Hispanic				0.34	0.98
	API					0.75
	AI/AN					
Surgery	NHW		0.05	<0.0001	0.36	0.46
	NHB			0.0006	0.01	0.98
	Hispanic				<0.0001	0.25
	API					0.28
	AI/AN					
Smoking Index	NHW		<0.0001	<0.0001	<0.0001	<0.0001
	NHB			<0.0001	<0.0001	0.009
	Hispanic				0.22	<0.0001
	API					<0.0001
	AI/AN					

Figure 1. Summary of pairwise *P*-values between race/ethnicities by clinicodemographic characteristics among patients with young-onset noncardia gastric cancer. *P*-values are presented for all pairwise comparisons. Associations with $P<0.05$, $P<0.01$, and $P<0.001$ are highlighted in light, medium, and dark grey color, respectively. NHW, non-Hispanic white; NHB, non-Hispanic black; API, Asian or Pacific Islander; AI/AN, American Indian/Alaska Native. AJCC, American Joint Committee on Cancer.

Summary of clinicodemographic characteristics by race/ethnicity among patients with young-onset noncardia gastric cancer: SEER18, 2007-2015.

Table 1.

Characteristic	Total		Non-Hispanic White		Non-Hispanic Black		Hispanic		Asian or Pacific Islander		American Indian/Alaska Native		P*
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
Total	2872		781	27.2	486	16.9	1086	37.8	478	16.6	41	1.4	
Age at Diagnosis													0.003
20-29 Years	175	6.1	43	5.5	24	4.9	83	7.6	24	5.0	1	2.4	
30-39 Years	710	24.7	170	21.8	105	21.6	303	27.9	121	25.3	11	26.8	
40-49 Years	1987	69.2	568	72.7	357	73.5	700	64.5	333	69.7	29	70.7	
Mean (sd), Years	41.6	(6.4)	42.0	(6.3)	42.4	(6.3)	40.8	(6.6)	41.7	(6.2)	42.1	(5.7)	<0.0001
Sex													0.12
Female	1434	49.9	401	51.3	231	47.5	521	48.0	260	54.4	21	51.2	
Male	1438	50.1	380	48.7	255	52.5	565	52.0	218	45.6	20	48.8	
Insurance Status													<0.0001
Privately Insured	1796	62.5	603	77.2	282	58.0	539	49.6	360	75.3	12	29.3	
Medicaid/Uninsured	983	34.2	142	18.2	182	37.4	524	48.3	108	22.6	27	65.9	
Unknown	93	3.2	36	4.6	22	4.5	23	2.1	10	2.1	2	4.9	
Anatomic Sub-site													0.0026
Fundus	296	10.3	108	13.8	57	11.7	93	8.6	34	7.1	4	9.8	
Corpus	722	25.1	203	26.0	114	23.5	294	27.1	104	21.8	7	17.1	
Antrum	979	34.1	239	30.6	167	34.4	379	34.9	183	38.3	11	26.8	
Pylorus	124	4.3	30	3.8	25	5.1	44	4.1	20	4.2	5	12.2	
Lesser Curvature	441	15.4	110	14.1	72	14.8	165	15.2	85	17.8	9	22.0	
Greater Curvature	310	10.8	91	11.7	51	10.5	111	10.2	52	10.9	5	12.2	
AJCC Clinical Stage													<0.0001
0-I	510	17.8	175	22.4	78	16.0	146	13.4	104	21.8	7	17.1	
II	371	12.9	79	10.1	74	15.2	130	12.0	81	16.9	7	17.1	
III	453	15.8	101	12.9	76	15.6	185	17.0	85	17.8	6	14.6	
IV	1128	39.3	263	33.7	177	36.4	509	46.9	161	33.7	18	43.9	
Unknown	410	14.3	163	20.9	81	16.7	116	10.7	47	9.8	3	7.3	

Characteristic	Total		Non-Hispanic White		Non-Hispanic Black		Hispanic		Asian or Pacific Islander		American Indian/Alaska Native		P*
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
Histologic Type													<0.0001
Diffuse	1192	41.5	248	31.8	169	34.8	542	49.9	217	45.4	16	39.0	
Intestinal	892	31.1	203	26.0	176	36.2	320	29.5	173	36.2	20	48.8	
Other	788	27.4	330	42.3	141	29.0	224	20.6	88	18.4	5	12.2	
Tumor Grade													<0.0001
I/II (Well to Moderately Differentiated)	476	16.6	176	22.5	88	18.1	138	12.7	66	13.8	8	19.5	
III/IV (Poorly Differentiated to Undifferentiated)	1754	61.1	377	48.3	269	55.3	743	68.4	340	71.1	25	61.0	
Unknown	642	22.4	228	29.2	129	26.5	205	18.9	72	15.1	8	19.5	
Cancer Sequence													0.14
Only One Primary	2772	96.5	745	95.4	466	95.9	1059	97.5	462	96.7	40	97.6	
First of 2+ Primaries	100	3.5	36	4.6	20	4.1	27	2.5	16	3.3	1	2.4	
Surgery													<0.0001
None	1188	41.4	275	35.2	197	40.5	543	50.0	156	32.6	17	41.5	
Yes	1642	57.2	494	63.3	281	57.8	529	48.7	314	65.7	24	58.5	
Unknown	42	1.5	12	1.5	8	1.6	14	1.3	8	1.7	0	0.0	
Smoking Index**													<0.0001
<13% current smokers	1123	39.1	194	24.8	83	17.1	587	54.1	257	53.8	2	4.9	
13-17% current smokers	675	23.5	201	25.7	104	21.4	213	19.6	153	32.0	4	9.8	
18%+ smokers	975	33.9	386	49.4	299	61.5	187	17.2	68	14.2	35	85.4	

Note. P value calculations do not include unknown values.

* Pairwise P values are presented in Figure 1.

**

Tertiles of the percentage of current smokers age 18+ years on the basis of the distribution in our cohort of patients with young-onset noncardia gastric cancer. Abbreviations: sd, standard deviation; AJCC, American Joint Committee on Cancer.

Table 2.

Multivariable logistic regression for clinicodemographic factors by race/ethnicity among patients with young-onset noncardia gastric cancer: SEER18, 2007-2015.

Characteristic	Adjusted Model**							
	Non-Hispanic Black*		Hispanic*		Asian or Pacific Islander*		American Indian/Alaska Native*	
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
Age at Diagnosis								
Years (continuous)	1.01 (0.99-1.04)	0.32	0.97 (0.95-0.99)	0.008	0.99 (0.97-1.02)	0.60	1.02 (0.96-1.10)	0.50
Sex								
Male	Ref (---)		Ref (---)		Ref (---)		Ref (---)	
Female	1.15 (0.85-1.55)	0.37	1.02 (0.79-1.32)	0.86	1.40 (1.04-1.88)	0.025	1.17 (0.53-2.60)	0.70
Insurance Status								
Privately Insured	Ref (---)		Ref (---)		Ref (---)		Ref (---)	
Medicaid/Uninsured	1.94 (1.39-2.69)	<0.0001	3.83 (2.89-5.08)	<0.0001	1.25 (0.88-1.78)	0.21	6.27 (2.75-14.26)	<0.0001
Anatomic Sub-site								
Fundus	Ref (---)		Ref (---)		Ref (---)		Ref (---)	
Corpus	1.41 (0.78-2.54)	0.26	1.11 (0.68-1.80)	0.69	1.00 (0.55-1.82)	1.00	1.01 (0.17-5.97)	0.99
Antrum	1.57 (0.89-2.78)	0.12	1.00 (0.62-1.61)	1.00	1.21 (0.68-2.16)	0.52	1.20 (0.23-6.21)	0.83
Pylorus	1.62 (0.67-3.88)	0.28	1.26 (0.60-2.67)	0.54	1.27 (0.53-3.05)	0.59	7.57 (1.20-47.81)	0.03
Lesser Curvature	1.38 (0.74-2.58)	0.31	1.07 (0.63-1.82)	0.80	1.31 (0.70-2.45)	0.40	1.58 (0.29-8.63)	0.60
Greater Curvature	1.04 (0.49-2.18)	0.93	1.17 (0.65-2.11)	0.60	1.12 (0.55-2.26)	0.76	3.20 (0.53-19.44)	0.21
AJCC Clinical Stage								
0-I	Ref (---)		Ref (---)		Ref (---)		Ref (---)	
II	1.82 (1.12-2.96)	0.016	1.36 (0.88-2.10)	0.17	1.26 (0.79-2.01)	0.34	1.39 (0.36-5.33)	0.63
III	1.18 (0.73-1.93)	0.50	1.26 (0.83-1.91)	0.28	0.86 (0.54-1.36)	0.51	0.80 (0.21-3.10)	0.75
IV	0.89 (0.54-1.44)	0.62	0.81 (0.53-1.22)	0.31	0.59 (0.37-0.95)	0.031	1.10 (0.30-3.96)	0.89
Histologic Type								
Diffuse	Ref (---)		Ref (---)		Ref (---)		Ref (---)	
Intestinal	1.35 (0.95-1.90)	0.09	0.95 (0.71-1.28)	0.74	1.28 (0.92-1.79)	0.15	1.27 (0.53-3.05)	0.60
Other	0.72 (0.44-1.20)	0.20	0.68 (0.45-1.04)	0.08	0.55 (0.33-0.91)	0.02	0.33 (0.06-1.84)	0.20
Tumor Grade								

Characteristic	Adjusted Model**					
	Non-Hispanic Black*			Hispanic*		
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
I/II	Ref (---)		Ref (---)		Ref (---)	
III/IV	1.23 (0.81-1.87)	0.34	1.93 (1.32-2.84)	0.001	1.91 (1.22-2.97)	0.005
Surgery						
None	Ref (---)		Ref (---)		Ref (---)	
Yes	0.75 (0.50-1.14)	0.17	0.62 (0.44-0.88)	0.007	1.10 (0.72-1.67)	0.66
Smoking Index***						
<13% current smokers	Ref (---)		Ref (---)		Ref (---)	
13-17% current smokers	1.13 (0.73-1.76)	0.58	0.54 (0.40-0.74)	0.0001	0.54 (0.38-0.76)	0.0004
18%+ smokers	1.79 (1.22-2.62)	0.003	0.15 (0.11-0.21)	<0.0001	0.13 (0.08-0.19)	<0.0001

Abbreviations: AJCC, American Joint Committee on Cancer; OR, odds ratio; CI, confidence interval; T, tertile.

* Referent group is non-Hispanic white patients with young-onset primary gastric cancer.

** Adjusted for age at diagnosis, sex, anatomic sub-site, insurance status, AJCC clinical stage, histologic type, tumor grade, surgery, and smoking index.

*** Tertiles of the percentage of current smokers age 18+ years on the basis of the distribution in our cohort of patients with young-onset noncardia gastric cancer.