ERRATUM

Erratum to: The cardiothoracic ratio and all-cause and cardiovascular disease mortality in patients undergoing maintenance hemodialysis: results of the MBD-5D study

Hiroaki Ogata¹ · Junji Kumasawa^{2,3,4} · Shingo Fukuma^{2,4,5} · Masahide Mizobuchi⁶ · Eriko Kinugasa¹ · Masafumi Fukagawa⁷ · Shunichi Fukuhara^{2,5} · Tadao Akizawa⁶

Published online: 4 July 2017
© Japanese Society of Nephrology 2017

Erratum to: Clin Exp Nephrol DOI 10.1007/s10157-017-1380-2

In the original publication, under Abstract, in *Results* section, the sentence “Exploratory analysis binder.” should read as:

“Exploratory analysis revealed that there were relationships between CTR and age, sex, body mass index, comorbidity of CVD, dialysis duration and intact parathyroid hormone, phosphorus, hemoglobin, and usage of phosphate binder”.

Under Results, in *Exploratory analysis investigating the relationship between CTR and covariates* section the

sentence “Multivariable regression binder (Table 4).” should read as:

“Multivariable regression analysis showed that there were relationships between CTR and age, sex, BMI, comorbidity of CVD, dialysis duration and level of iPTH, phosphorus, hemoglobin, and usage of phosphate binder (Table 4)”.

In Table 4, one row (Diabetes mellitus comorbidity) was duplicated and a row (Dialysis duration) was not included. The corrected Table 4 is given in this Erratum.

The online version of the original article can be found under doi:[10.1007/s10157-017-1380-2](https://doi.org/10.1007/s10157-017-1380-2).

✉ Hiroaki Ogata
ogatah@med.showa-u.ac.jp

¹ Department of Internal Medicine, Showa University Northern Yokohama Hospital, Chigasaki-chuo 35-1, Tsuzuki, Yokohama, Kanagawa 224-8503, Japan

² Department of Healthcare Epidemiology, Graduate School of Medicine and Public Health, Kyoto University, Kyoto, Japan

³ Department of Critical Care Medicine, Sakai City Medical Center, Osaka, Japan

⁴ Institute for Health Outcomes and Process Evaluation Research (iHope International), Kyoto, Japan

⁵ Center for Innovative Research for Communities and Clinical Excellence (CIRC2LE), Fukushima Medical University, Fukushima, Japan

⁶ Division of Nephrology, Department of Medicine, Showa University School of Medicine, Tokyo, Japan

⁷ Division of Nephrology, Endocrinology and Metabolism, Tokai University School of Medicine, Kanagawa, Japan

Variable	Unit	β coefficients	95% CI	<i>p</i> value
Age	per 1 year	0.093	(0.070 to 0.117)	<0.001
Sex, female	vs Male	1.664	(1.040 to 2.287)	<0.001
Dialysis duration	per 1 year	0.032	(0.002 to 0.064)	0.040
Cardiovascular-related disease comorbidity	vs No	1.176	(0.596 to 1.756)	<0.001
Diabetes mellitus comorbidity	vs No	−0.215	(−0.748 to 0.318)	0.424
Smoking	vs No	0.736	(−0.145 to 1.618)	0.100
Body mass index	per 1 kg/m ²	0.224	(0.079 to 0.369)	0.003
Urea clearance time/volume	per 0.1	−0.602	(−1.872 to 0.667)	0.348
Dry weight	per 1 kg	−0.096	(−0.145 to −0.046)	<0.001
Calcium	per 1 mg/dL	−0.409	(−0.341 to 0.259)	0.786
Phosphorus	per 1 mg/dL	0.211	(0.025 to 0.398)	0.027
Intact parathyroid hormone	per 100 pg/mL	0.002	(0.001 to 0.003)	<0.001
Hemoglobin	per 1 g/dL	−0.300	(−0.528 to −0.073)	0.010
Albumin	per 1 g/dL	−0.342	(−1.013 to 0.329)	0.313
C-reactive protein	per 1 mg/dL	0.046	(−0.117 to 0.209)	0.576
Vitamin D receptor activator	vs No	0.312	(−0.525 to 0.588)	0.911
Phosphate binder	vs No	0.746	(0.072 to 1.420)	0.031
Angiotensin converting enzyme inhibitor	vs No	0.145	(−0.729 to 1.019)	0.741
Beta blocker	vs No	−0.068	(−1.317 to 1.181)	0.914