ERRATUM

Erratum to: Subsequent risk of ipsilateral and contralateral invasive breast cancer after treatment for ductal carcinoma in situ: incidence and the effect of radiotherapy in a population-based cohort of 10,090 women

Lotte E. Elshof^{1,2,3} · Michael Schaapveld² · Marjanka K. Schmidt^{1,2} ·
Emiel J. Rutgers³ · Flora E. van Leeuwen² · Jelle Wesseling^{1,4}

Published online: 23 November 2016
© Springer Science+Business Media New York 2016

Erratum to: Breast Cancer Res Treat (2016)
159:553–563
DOI 10.1007/s10549-016-3973-y

In the original publication of the article, Fig. 1 was published incorrectly. The corrected Fig. 1 is given in this erratum.

The online version of the original article can be found under doi:[10.1007/s10549-016-3973-y](https://doi.org/10.1007/s10549-016-3973-y).

✉ Jelle Wesseling
j.wesseling@nki.nl

¹ Department of Molecular Pathology, The Netherlands Cancer Institute/Antoni van Leeuwenhoek, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

² Department of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute/Antoni van Leeuwenhoek, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

³ Department of Surgery, The Netherlands Cancer Institute/Antoni van Leeuwenhoek, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

⁴ Department of Pathology, The Netherlands Cancer Institute/Antoni van Leeuwenhoek, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

Fig. 1 Flow diagram for patient selection and median follow-up by initial treatment type. *iIBC* ipsilateral invasive breast cancer, *cIBC* contralateral invasive breast cancer