CORRESPONDENCE

The Interdisciplinary Management of Foot Drop

by Dr. med. Anne Elisabeth Carolus, PD Dr. med. Michael Becker, Dr. med. Jeanne Cuny, Prof. Dr. med. Rüdiger Smektala, Prof. Dr. med. Kirsten Schmieder, PD Dr. med. Christopher Brenke in issue 20/2019

Neuromuscular Electrical Stimulation (NMES) and α -Lipoic Acid

Any unilateral spontaneous foot drop as a sign of peripheral peroneal nerve palsy which is not associated with other symptoms (e.g. pain) and not explained by a preceding injury can be the first manifestation of underlying diabetes mellitus. Consequently, the patient's family history should be taken and fasting glucose levels as well as Hb1Ac should be measured. During my clinical training and the many years working in my own practice, neuromuscular electrical stimulation on weekdays and a dose of 600 mg α -lipoic acid (unfortunately no longer available on statutory health care prescription) taken in the morning have proven effective and the prognosis is good. At worst, spontaneous (potentially bilateral) foot drop can be the first sign of amyotrophic lateral sclerosis (ALS), but this is rarely the case. If ALS is suspected, the patient should be carefully examined for abnormal reflexes, abnormal muscle tone, fasciculations, and the presence of pyramidal signs. The patient should only be informed that a diagnosis of ALS is suspected, if additional signs and symptoms have clearly been identified. DOI: 10.3238/arztebl.2019.0643a

References

1. Carolus AE, Becker M, Cuny J, Smektala R, Schmieder K, Brenke C: The interdisciplinary management of foot drop. *Dtsch Arztebl Int* 2019; 116: 347–54.

Dr. med. Franz-Ulrich Beutner
Burgdorf, Germany
fub_bu@me.com

Anatomy Not Precisely Described

We welcome the article by Carolus et al. (1) on the management of foot drop, as it discusses a key interdisciplinary topic. However, it is of clinical importance to distinguish the central elements from the peripheral elements of the neuronal circuitry in a way that is anatomically correct. For the purpose of simplification, the motor system is described as consisting of an upper (central) and a lower (peripheral) neuron. The cell body of the upper neuron is located in the primary motor cortex and its axon extends in the pyramidal tract down to the anterior horn cells of the spinal cord, i.e. the lower neuron. Contrary to the description in the article, this lower neuron is not part of the pyramidal tract. Even though the cell body (perikaryon) of the motor anterior horn cell is, in fact, located in the spinal cord, i.e. in a strict sense within the central nervous system, isolated failure of this second neuron—regardless of the location—results in a typical peripheral (i.e. flaccid) paresis.

With regard to central foot drop, it should be mentioned that electronic foot drop orthoses have now been available for several years (2). The technique of functional electrical stimulation

(FES) is used to stimulate the common fibular nerve (peroneal nerve) (3), which is intact in patients with central paresis. The FES technique uses sensors to detect the need for lifting the foot while walking. Unfortunately, the costs for such a system are frequently not reimbursed by health insurances and often a major effort has to be made—e.g. by video documentation of the beneficial effect during testing of the device—to persuade the patient's insurance to cover the costs.

Finally, we would like to point out that rapid significant weight loss represents a relevant risk factor for the development of (frequently bilateral) common fibular nerve lesions with foot drop. DOI: 10.3238/arztebl.2019.0643b

References

1. Carolus AE, Becker M, Cuny J, Smektala R, Schmieder K, Brenke C: The interdisciplinary management of foot drop. *Dtsch Arztebl Int* 2019; 116: 347–54.
2. Kluding PM, Dunning K, O'Dell MW, et al.: Foot drop stimulation versus ankle foot orthosis after stroke—30-week outcomes. *Stroke* 2013; 44: 1660–9.
3. Federal Committee on Anatomical Terminology: *Terminologia Anatomica*, Stuttgart: Thieme 1998.

Prof. Dr. med. Andreas Winkelmann

Institut für Anatomie, Medizinische Hochschule Brandenburg, Brandenburg, Germany
andreas.winkelmann@mhb-fontane.de

Prof. Dr. med. Andreas Bitsch

Klinik für Neurologie, Ruppiner Kliniken GmbH, Hochschulklinikum der Medizinischen Hochschule Brandenburg, Brandenburg, Germany

In Reply:

We are grateful for the additions to our article (1). FU Beutner highlights again that polyneuropathy is a common cause of spontaneous foot drop—that is our experience as well. In addition, it should be recognized that there is a high coincidence between polyneuropathies and compression syndromes (2): Because of changes in their myelin sheaths that may result from the systemic disease, the nerves are, per se, more sensitive to local pressure. To perform HbA1c testing in patients with foot drop is a valuable suggestion for daily practice which should be added to our algorithm.

Neuromuscular electrical stimulation is briefly touched upon in our article. In animal experiments it was shown that electrical stimulation of denervated muscles can stop the development of muscle atrophy, provided muscle fiber-specific stimulation frequencies, intervals, and stimulus locations are used (3). Its effectiveness in patients remains the subject of controversy. It would miss the point to attribute a direct beneficial effect on foot drop to conventional transcutaneous electrical nerve stimulation (TENS) (4). The efficacy of TENS in the treatment of polyneuropathy-associated pain is supported by a considerable amount of evidence and a beneficial effect of TENS on spasticity

in patients after stroke has also been demonstrated (5). Furthermore, we think that the psychological benefit for patients from the perceived sensation of electrical pulses in the affected area of the foot should not be underestimated. However, one should explain to the patients that these devices are unlikely to improve the actual paralysis and thus do not represent a viable alternative to other causal treatments, if any are available in the individual case. Furthermore, TENS is no substitute for active physiotherapy which includes joint mobility and training of compensatory movements. TENS should be distinguished from direct nerve stimulation, the technique mentioned by A. Winkelmann and A. Bitsch. The method exploits the integrity of the peripheral nerve in patients with central foot drop and its beneficial effects have been demonstrated. As mentioned above, currently this treatment is hampered by its prohibitively high costs.

Given the numerous inquiries of readers—many thanks for these—here some additional information about tendon transfer techniques: The most important prerequisite for successful tendon transfer surgery is that the paresis is limited to the area innervated by the common fibular nerve, i.e. that only the tibialis anterior muscle and the peroneal muscles are affected. Patients with polyradicular lesions and advanced polyneuropathies typically do not meet this criterion. In addition, patients should not have severe pes planovalgus deformity, because in these patients

the tibialis posterior muscle, which is used for the transfer, is an important stabilizer of the inner arch of the foot. In such cases, tendon transfer surgery can be combined with subtalar arthrodesis. If these factors are taken into account when making the decision to operate, the patients' quality of life can be significantly improved by the procedure.

DOI: 10.3238/arztebl.2019.0643c

References

1. Carolus AE, Becker M, Cuny J, Smekala R, Schmieder K, Brenke C: The interdisciplinary management of foot drop. *Dtsch Arztebl Int* 2019; 116: 347–54.
2. Rajabally YA, Narasimhan M: Electrophysiological entrapment syndromes in chronic inflammatory demyelinating polyneuropathy. *Muscle Nerve* 2011; 44: 444–7.
3. Tanaka M, Nakanishi R, Murakami S, et al.: Effectiveness of daily eccentric contractions induced via kilohertz frequency transcutaneous electrical stimulation on muscle atrophy. *Acta Histochem* 2016; 118: 56–62.
4. Lin S, Sun Q, Wang H, Xie G: Influence of transcutaneous electrical nerve stimulation on spasticity, balance, and walking speed in stroke patients: A systematic review and meta-analysis. *J Rehabil Med* 2018; 50: 3–7.
5. Poulsen JB, Möller K, Jensen CV, Weisdorf S, Kehlet H, Perner A: Effect of transcutaneous electrical muscle stimulation on muscle volume in patients with septic shock. *Crit Care Med* 2011; 39: 456–61.

Corresponding author

Dr. med. Anne Elisabeth Carolus

Klinik für Neurochirurgie

Universitätsklinik Knappschafts-Krankenhaus, Bochum, Germany

AnneElisabeth.Carolus@kk-bochum.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.

CLINICAL SNAPSHOT

Maculopapular erythema, particularly pronounced over the abdomen in the area of the stretch marks (wide arrow), with slightly less reddening around the navel (narrow arrow), target-like (>) and vesicular foci (>>) on the extremities

Pronounced Polymorphic Eruption of Pregnancy in a Primipara

A 38-year-old primipara presented in the 36th week of gestation with strongly itching maculopapular erythema that was particularly pronounced over the abdomen in the area of the stretch marks, with less reddening around the navel. Target-like and vesicular foci were present on the extremities. Taken together, the skin findings, the absence of relevant medications in the patient's medical history, and the lack of potential differential diagnoses (herpes simplex virus polymerase chain reaction [PCR], varicella zoster virus PCR, histology, direct and indirect immunofluorescence) indicated the diagnosis of polymorphic eruption of pregnancy (or PUPPP, pruritic urticarial papules and plaques of pregnancy). The healthy newborn had no cutaneous abnormalities. The mother's erythema regressed completely when treated with topical antipruritics and medium-strength glucocorticoids. PUPPP is a common (incidence 1 : 200) self-limiting form of dermatosis that occurs most frequently in the final weeks of pregnancy or post partum (15 %). The individual risk factors include first pregnancy, multiple gestation, and excessive maternal weight gain. In pregnant women with pruritus the following differential diagnoses have to be ruled out: atopic gestational dermatosis, intrahepatic gestational cholestasis, gestational pemphigoid, and unspecific dermatoses of pregnancy (cutaneous drug reactions, viral exanthema).

Dr. med. Maria Isabel Schmidgen, Dr. med. Beate Weidenthaler-Barth, Hautklinik und Poliklinik, Universitätsmedizin Mainz, maria_isabel.schmidgen@unimedizin-mainz.de

Prof. Dr. med. Kerstin Steinbrink, Klinik für Dermatologie, Universitätsklinikum Münster

Conflict of interest statement: The authors declare that no conflict of interest exists.

Translated from the original German by David Roseveare.

Cite this as: Schmidgen MI, Steinbrink K, Weidenthaler-Barth B: Pronounced polymorphic eruption of pregnancy in a primipara. *Dtsch Arztebl Int* 2019; 116: 644. DOI: 10.3238/arztebl.2019.0644