Corrigendum

Journal of Dental Research
2015, Vol. 94(7) 1016–1017
© International & American Associations
for Dental Research 2015
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0022034515585305
jdr.sagepub.com

Asai K, Yamori M, Yamazaki T, Yamaguchi A, Takahashi K, Sekine A, Kosugi S, Matsuda F, Nakayama T, Bessho K; Nagahama Study Group. 2015. Tooth loss and atherosclerosis: the Nagahama Study. *J Dent Res.* 94(3 Suppl):52S–58S. (Original DOI: 10.1177/0022034514559127)

There were some errors in the Results and in Tables 1 and 2. These errors do not affect the Discussion or Conclusion of the study. The corrected text and tables are below:

Table 1. Characteristics of Study Participants.

Variable	Male (n = 2680)	Female (n = 5444)	All Participants (N = 8124)
Age, y	56.0 (28.9–83.1)	53.3 (27.2–79.5)	54.2 (27.7–80.8)
Tooth loss	4.3 (0–18.3)	3.2 (0–14.4)	3.6 (0–15.8)
CAVI	7.9 (5.4–10.3)	7.2 (5.2–9.3)	7.4 (5.2–9.7)
Hypertension	1,066 (39.8)	1,549 (23.5)	2,420 (29.8)
Diabetes	226 (8.4)	170 (2.6)	377 (4.6)
Former smoker	1,184 (44.2)	454 (8.3)	1,638 (20.2)
Current smoker	827 (30.9)	345 (6.4)	1,172 (14.4)
BMI, kg/m ²	23.4 (17.3–29.5)	21.8 (15.3–28.3)	22.3 (15.8–28.9)
SBP, mm Hg	128.6 (91.9–160.3)	119.4 (85.6–153.3)	122.4 (88.2–156.7)
DBP, mm Hg	79.9 (58.4–101.3)	73.5 (52.1–94.9)	75.6 (53.3–97.8)
HDL-C, mg/dL	57.7 (26.2–89.2)	68.7 (36.3–101.2)	65.1 (31.3–98.8)
LDL-C, mg/dL	123.0 (60.4–185.7)	123.8 (61.36–186.2)	123.5 (61.1–181.0)
HbA1c	5.2 (3.9–6.5)	5.1 (4.2–6.0)	5.1 (4.0–6.2)
Glucose, mg/dL	94.3 (51.9–136.7)	88.8 (63.2–114.4)	90.6 (58.1–123.2)
Antihypertensive medication	598 (22.3)	843 (15.5)	1,441 (17.7)
Hypoglycemic medication	141 (5.3)	88 (1.6)	229 (2.8)
Insulin	14 (0.5)	14 (0.3)	28 (0.3)

Values are presented as the mean (reference range: mean –2 SD to mean +2 SD) or n (%).

BMI, body mass index; CAVI, cardio-ankle vascular index; DBP, diastolic blood pressure; HbA1c, hemoglobin A1c; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; SBP, systolic blood pressure.

Table 2. Multivariable Linear Regression Models for CAVI and Number of Tooth Loss Adjusted for Age, Sex, BMI, Smoking Status, Hemoglobin A1c, a History of Insulin or Hypoglycemic Medication, and Interactions.

	Adjusted Model ^a					
	Unadjusted Model		Model 1		Model 2	
	β	95% CI	β	95% CI	β	95% CI
Tooth loss	0.47	(0.45 to 0.50)	0.04	(0.02 to 0.06)	0.03	(0.01 to 0.05)
Sex	–0.65	(–0.70 to –0.60)	–0.48	(–0.53 to –0.43)	–0.48	(–0.53 to –0.43)
Age	0.06	(0.06 to 0.06)	0.06	(0.05 to 0.06)	0.06	(0.05 to 0.06)
Tooth loss × sex ^b					–0.05	(–0.08 to –0.02)
Tooth loss × age category ^c					0.04	(–0.01 to 0.08)

Tooth loss was logarithmically transformed.

CI, confidence interval.

^aMultivariable linear regression analysis, adjusting for body mass index, smoking status, hemoglobin A1c, and a history of insulin or hypoglycemic medication.

^bSex (0: male, 1: female).

^cAge category (0: <60 years, 1: ≥60 years).

In the “General Characteristics” section:

The reference range for “age for women” is stated as “27.2–**79.4**” but should be “27.2–**79.5**.”

The reference range for “tooth loss for women” is stated as “0–**14.3**” but should be “0–**14.4**.”

In the “Association between Tooth Loss and CAVI” section:

The 95% CI for “CAVI and tooth loss” is stated as “0.01 to **0.6**” but should be “0.01 to **0.06**.”

In the “Interaction Effects of Sex, Age, and Tooth Loss” section:

The β (95% CI) for “ β for interaction” is stated as “**0.04 (0.02 to 0.6)**” but should be “**–0.05 (–0.08 to –0.02)**.”