Published in final edited form as:

Brain Res. 2019 January 01; 1702: 12–16. doi:10.1016/j.brainres.2018.01.001.

Complex noradrenergic dysfunction in Alzheimer's disease: Low norepinephrine input is not always to blame

Mary Gannon, Qin Wang*

Department of Cell, Developmental and Integrative Biology, University of Alabama at Birmingham, Birmingham, AL, USA

Abstract

The locus coeruleus-noradrenergic (LC-NA) system supplies the cerebral cortex with norepinephrine, a key modulator of cognition. Neurodegeneration of the LC is an early hallmark of Alzheimer's disease (AD). In this article, we analyze current literature to understand whether NA degeneration in AD simply leads to a loss of norepinephrine input to the cortex. With reported adaptive changes in the LC-NA system at the anatomical, cellular, and molecular levels in AD, existing evidence support a seemingly sustained level of extracellular NE in the cortex, at least at early stages of the long course of AD. We postulate that loss of the integrity of the NA system, rather than mere loss of NE input, is a key contributor to AD pathogenesis. A thorough understanding of NA dysfunction in AD has a large impact on both our comprehension and treatment of this devastating disease.

Keywords

Alzheimer's disease; Locus coeruleus; Noradrenergic; Norepinephrine; Dysfunction

1. Introduction to the noradrenergic system and its degeneration in Alzheimer's disease

The brain noradrenergic (NA) system plays a critical role in supporting interactions with and responses to environmental stimuli, and serves as a major contributor to normal cognitive activities in the cortex including attention, perception, and memory retrieval (Chamberlain and Robbins, 2013; Gannon et al., 2015). The majority of NA neurons are located in the locus coeruleus (LC) and, with widespread efferent projections, these neurons supply norepinephrine (NE) to the cortex, hippocampus, striatum, basal forebrain, preoptic area, and hypothalamus (Oleskevich et al., 1989; Swanson and Hartman, 1975). NE is a catecholamine that is synthesized from the amino acid tyrosine through sequential reactions catalyzed by tyrosine hydroxylase, DOPA decarboxylase and dopamine β hydroxylase (D β H). NE released from presynaptic terminals binds to different subtypes of adrenergic receptors to elicit a variety of physiological and pharmacological responses (Hein, 2006). Extracellular NE can be enzymatically degraded by enzymes such as catechol-O-

*Corresponding author at: 1918 University Boulevard, 986 MCLM, Birmingham, AL 35294, USA. qinwang@uab.edu (Q. Wang).

methyltransferases (COMT) or monoamine oxidases (MAO) (Golan et al., 2011), or be taken back up into the presynaptic terminal by the norepinephrine transporter (NET), a Na/K pump. The amount of extracellular NE at the synapse is largely dependent on how much NE is degraded and taken back to the presynaptic terminal.

Changes of the NA system in Alzheimer's disease (AD) have been long observed. As AD progresses, there is profound neuronal loss in the LC (Chan-Palay and Asan, 1989; Forstl et al., 1994; German et al., 1992; Iversen et al., 1983; Mann et al., 1980; Marcyniuk et al., 1986; Matthews et al., 2002; McMillan et al., 2011; Zarow et al., 2003). However, despite this well documented phenomenon, its impact on AD pathogenesis has not been adequately addressed to date. While a decrease in NE has often been observed in advanced AD, in some cases, NE levels in AD patients remain constant, or even elevated. In addition, alterations of different adrenergic receptors have been reported with both decreases and increases observed in the levels of receptor expression and density (Kalaria, 1989; Kalaria et al., 1989a; Ruiz et al., 1993; Shimohama et al., 1986; Szot et al., 2006, 2007). These changes at the receptor level directly influence the sensitivity and amplitude of the physiological responses elicited by NE, adding another level of complexity to the consequence of NA degeneration in AD. Furthermore, some subtypes of adrenergic receptors have been shown to regulate A β generation (Chen et al., 2014; Ni et al., 2006; Wang et al., 2013b; Yu et al., 2010) or mediate A β toxicity (Wang et al., 2013a), indicating an important role of these receptors in AD pathogenesis. In this review, we will analyze existing literature to discern, in the context of the long and progressive course of AD, NE input to the cerebral cortex and its potential effects through different adrenergic receptors. Although NE has been well appreciated as a neurotransmitter involved in cognition, it appears that NA system dysfunction in AD is much more complex than a simple loss of NE input to the cerebral cortex.

2. Compensatory mechanisms to counteract NA neuron loss occur during the long and progressive course of AD

Pathological changes in the LC, including tau accumulation in the neurons of LC, can occur as early as childhood or adolescence (Braak and Del Tredici, 2011a,b; Braak et al., 2011; Braak and Del Tredici, 2012; Braak et al., 2013). As AD progresses, degeneration of the LC neurons happens (Chan-Palay and Asan, 1989; Forstl et al., 1994; German et al., 1992; Iversen et al., 1983; Mann et al., 1980; Marcyniuk et al., 1986; Matthews et al., 2002; McMillan et al., 2011; Zarow et al., 2003). This, however, is not a rapid occurrence. Rather, the degeneration seems to be a gradual process, and there is evidence that compensatory mechanisms occur in an effort to counteract this change and maintain homeostasis.

First, at an anatomical level, NA degeneration in AD is accompanied by increased sprouting of the remaining LC neurons to different brain regions, including the prefrontal cortex (Szot et al., 2007) and hippocampus (Szot et al., 2006). NA sympathetic sprouting in the hippocampus has also been observed in late-onset AD subjects (Booze et al., 1993; Nelson et al., 2014). Secondly, changes in the enzymes responsible for NE generation have been reported in AD. Both tyrosine hydroxylase, which is the enzyme responsible for the rate limiting step in NE synthesis, (Iversen et al., 1983; Szot et al., 2006, 2007) and D β H

(Giubilei et al., 2004; Miyata et al., 1984) levels are increased in AD patients compared to control subjects. Further, D β H levels are higher in middle and late stages of AD than in early stages (Mustapic et al., 2013). Elevated levels of these enzymes can lead to an increase in production of NE. Thirdly, decreased levels of NET have been found in AD patients using both autoradiograph (Gulyas et al., 2010) and radioligand binding studies (Tejani-Butt et al., 1993) of post-mortem tissues. As mentioned earlier, NE re-uptake is mediated by NET, a Na/K pump that returns NE to the presynaptic terminal and acts as a major determinant of the synaptic NE level. A decrease in the level of this transporter leaves more NE in the synapse. Taken together, enhancement in neuronal sprouting and NE production and reduction in NE reuptake could compensate LC neuronal loss to maintain the NE levels in AD. It is also possible that these mechanisms may overcompensate for the neuronal loss, especially at the early stages of the disease, leading to increased NE levels, as seen in some AD patients (Elrod et al., 1997; Tohgi et al., 1992), and thus creating another problem for AD patients. As AD continues to progress and LC degeneration becomes increasingly profound, these compensatory mechanisms may no longer be sufficient to counteract the neuronal loss. As a result, a decrease in the tissue levels of NE can be seen.

3. Is there really a loss of norepinephrine input in AD?

Given the well-documented loss of LC neurons, it is often assumed that there is a loss of norepinephrine input to the cerebral cortex in AD. Is this really the case?

As discussed above, AD is a long and progressive process, during which degeneration of LC neurons degeneration occurs gradually along with compensation events. When remaining LC neurons cannot sufficiently compensate the overall neuronal loss, a decrease in the tissue levels of NE is observed in the temporal lobe of advanced AD patients compared to healthy controls (Adolfsson et al., 1979; D'Amato et al., 1987; Gottfries et al., 1983; Herregodts et al., 1989; Hoogendijk et al., 1999; Matthews et al., 2002; Palmer et al., 1984, 1987; Reinikainen et al., 1988; Szot et al., 2006). Hoogendijk et al. (1999) found a NE reduction in frontal medial gyrus, temporal superior gyrus, hippocampus, amygdala, thalamus, and LC with AD NE levels ranging from 34% of healthy controls in the frontal medial gyrus to 65% of healthy controls in the hippocampus. Others have seen similar results, with 12–50% decreases seen in the gyrus frontalis, putamen, frontal and temporal cortex, hypothalamus, caudate nucleus, hippocampus, and gyrus cingula (Adolfsson et al., 1979; Gottfries et al., 1983; Matthews et al., 2002; Palmer et al., 1987; Reinikainen et al., 1988, 1990). Finally, the NE metabolite MHPG was found to be decreased in the temporal cortex of older AD patients (>70 years) compared to controls, but not in younger AD patients (<70 years) (Palmer et al., 1984). However, does a 12–66% decrease in the tissue levels of NE necessarily cause a change in the extracellular levels of NE, i.e. NE input, in the cerebral cortex?

Direct measurement of the extracellular levels of NE in vivo in LC-lesioned animals suggest that this may not be not the case. In one study, the noradrenergic specific toxin N-(2-chloroethyl)-N-e thyl-2-bromobenzylamine (DSP-4) treatment led to a 75% drop in noradrenergic content. However, in microdialysis experiments, the extracellular NE levels of the DSP-4 treated rats were almost 2x greater than sham controls (Hughes and Stanford,

1998). In other studies, 50% (Kask et al., 1997), 75% (Abercrombie and Zigmond, 1989) or even 90% (Ventura et al., 2003) depletion of the tissue levels of NE content, as measured by HPLC, does not lead to a significant change in the extracellular levels of NE in the frontal cortex measured by microdialysis. It is important to note that the varied levels of noradrenergic depletion can be due to inconsistencies in the methods of drug treatment such as drug doses, numbers of injections and treatment time. Nevertheless, these well-controlled studies suggest a clear disconnect between the tissue and the extracellular levels of NE.

Several lines of evidence from AD patients support that the extracellular levels of NE are likely maintained despite a decrease in the tissue levels of NE. First, in fresh cortical samples obtained during neurosurgery from AD patients who have an average 68% reduction in cortical NE levels, as measured by HPLC following tissue homogenization, the extracellular release of endogenous NE is not different from that in cortical samples of control subjects (Palmer et al., 1987). Additionally, NE reuptake in these AD samples is reduced to 47% of the control level. This unchanged release and reduced reuptake of NE would lead to sustained extracellular NE in the cerebral cortex of AD patients.

Secondly, when measured in the cerebrospinal fluid (CSF) of living AD patients, the levels of NE are unchanged or often increased compared to age-matched control subjects (Czech et al., 2012; Elrod et al., 1997; Kaddurah-Daouk et al., 2011; Oishi et al., 1996; Raskind et al., 1984, 1999; Tohgi et al., 1992; Watson et al., 2006). Furthermore, the increase in CSF NE has been shown to correlate with the severity of AD (Elrod et al., 1997; Oishi et al., 1996) and frontal temporal dementia (FTD) (Engelborghs et al., 2008), with higher NE levels in CSF correlating with more severe cognitive deficit. Additionally, the NE metabolite MHPG has been reported to negatively correlate with cognitive function in AD (Oishi et al., 1996). Even in healthy subjects, NE levels have been reported to increase with age, and are associated with poorer cognitive performance (Wang et al., 2013c). This evidence suggests that although important for cognitive functions, a higher level of NE could be detrimental. This is consistent with studies that have shown cognitive impairment in response to stress, under which conditions NE release is elevated (Finlay et al., 1995; Goldstein et al., 1996). Exposure to uncontrollable stressors has been shown to cause rapid impairment of working memory in monkeys and rodents (Arnsten, 1998; Arnsten and Goldman-Rakic, 1998). Stress exposure in humans is also found to weaken working memory performance (Qin et al., 2009). Furthermore, blockade of α_1 AR by antagonists, such as prazosin and urapidil, has been shown to protect cognitive functions from the detrimental effects of stress exposure (Arnsten and Jentsch, 1997).

While the majority of studies have found an increase in CSF NE levels in AD patients, one study did find reduced NE levels in the CSF of AD patients with moderate cognitive decline as compared to controls (Martignoni et al., 1992). As Martignoni noted in his publication, all of his AD patients and controls were hospitalized, while only the most severe AD patients were hospitalized in some of the other published studies. It is possible that institutionalization itself or other sickness that caused hospitalization in these patients could have affected the study. Another study found decreased NE levels in ventricular CSF of postmortem AD patients (Kaddurah-Daouk et al., 2011), though postmortem CSF

measurement can be confounded by the cause of death, postmortem delay and storage period and pH of the CSF samples.

In addition to the direct measures of NE levels in CSF, studies with adrenergic antagonists point to NE levels remaining high enough in AD to exert effects through adrenergic receptors. Treatment with prazosin (Wang et al., 2009), an α_1 AR antagonist, or propranolol (Pauszek, 1991; Peskind et al., 2005; Shankle et al., 1995), a β AR antagonist, significantly improved the agitation symptoms in AD patients compared to placebo. These results indicate that NE acting on the α_1 AR or β AR is contributing to the agitation and aggressive behaviors seen in the AD patients. In addition, α_2 AR responsiveness to a yohimbine blockade is enhanced in AD, as indicated by a higher increase in CSF NE levels by yohimbine treatment in AD patients compared to that in control subjects (Peskind et al., 1995; Raskind et al., 1999). These results suggest a higher level of α_2 AR activation by the endogenous NE in AD.

Taken together, existing literature points to 12%–65% reduction in the tissue levels of NE in most AD cases, and yet, the extracellular levels of NE (i.e. NE input to the cortex) in these AD patients may very likely maintain at a level close to the normal condition. Thus, although NA degeneration occurs frequently, there may actually be a sustained NA tone in AD.

4. NE effects through the adrenergic receptors on cognition in the context of AD

NE elicits its effects through activating the adrenergic receptors, which are members of the G-protein coupled receptor (GPCR) superfamily. The α_1 subfamily of adrenergic receptors (α_{1A} , α_{1B} and α_{1D}) signals through the G_q signaling pathway and, upon activation, increase both IP3 and calcium levels (Chen and Minneman, 2005; Perez, 2007). The β ARs (β_1 , β_2 and β_3) mainly signal through the G_s protein and increase cAMP generation upon activation (Hall, 2004). The α_2 ARs (α_{2A} , α_{2B} and α_{2C}) signal through the $G_{i/o}$ signaling pathway, and activation of these receptors inhibits cAMP production and calcium influx (Knaus et al., 2007). The α_2 receptors also serve as auto-receptors on presynaptic NA terminals where they inhibit NE release. All three receptor subfamilies play important roles in normal cognitive function, which has been reviewed extensively (Chamberlain and Robbins, 2013; Gannon et al., 2015). Here we primarily focus on the effects of activation or inhibition of these receptors on cognitive functions in AD animal models or human patients.

In the non-disease state, inhibition of the β_1 and β_2 ARs with propranolol, which blocks the β_1 and β_2 ARs with equal affinity (Summers, 2006), leads to cognitive impairments, including impaired memory retrieval (Ouyang and Thomas, 2005), fear conditioning (Taherian et al., 2014), taste memory consolidation (Ruetti et al., 2014), scent recall (Veyrac et al., 2007) and spatial learning (Decker et al., 1990; Kenton et al., 2008; Saber and Cain, 2003). One might therefore assume that activating these receptors in AD patients would lead to positive outcomes. However, for the most part, blockade of the β ARs seems to be cognitively beneficial, as demonstrated in AD animal models. This may be related to the role of β AR in A β generation. Activation of β_2 AR promotes A β production through enhancing γ

secretase activity (Ni et al., 2006). Thus, a β_2 antagonist, ICI 118,551, reduces A β plaque load in AD mouse models, whereas β_2 agonists increase the A β plaque load (Ni et al., 2006). Additionally, treatment of Tg2576 mice with a β_1 antagonist, nebivolol, reduces A β generation (Wang et al., 2013b). Consistently, in both Tg2576 AD model mice and a mouse model of age-related cognitive decline (SAMP8), treatment with a non-subtype selective β blocker, propranolol, leads to an improvement in cognitive function (Dobarro et al., 2013a,b). However, in a 3xTg mouse model, ICI 118,551 actually increased A β levels and impaired cognitive function (Branca et al., 2014). This discrepancy has yet to be fully addressed, though may be related to the different models used.

In addition to β_2 AR, a number of other G protein-coupled receptors have been linked to AD by altering APP processing, and this topic is reviewed well elsewhere (Thathiah and De Strooper, 2011). Within the adrenergic receptor family, α_2 AR also promotes amyloidogenic processing of APP (Chen et al., 2014). An α_2 AR agonist, clonidine, increases A β plaque load, whereas an α_2 AR antagonist, idazoxan, decreases it and rescues the cognitive deficit observed in APP/PS1 transgenic mice (Chen et al., 2014).

Although activation of the α_1 ARs has been shown to improve spatial learning, fear learning, and working memory and attention in WT animals (Do Monte et al., 2013; Doze et al., 2011; Mishima et al., 2004; Nalepa et al., 2013; Veyrac et al., 2007), in an AD transgenic model, APP23 mice, blockade of these receptors by an antagonist, prazosin, improves the cognitive deficits (Katsouri et al., 2013). In AD patients, α_1 AR activation is thought to contribute to the observed agitation symptom (Raskind and Peskind, 1994; Sharp et al., 2007), and prazosin is effective in alleviating such symptoms (Wang et al., 2009).

Taken together, the above evidence suggest dysfunction of adrenergic receptors in regulating cognitive functions under the pathological conditions of AD. This is not totally surprising given the substantial alterations in neuroanatomy including the primary and compensatory changes in the NA system in AD. At the molecular and cellular levels, aberrant receptor densities (Kalaria et al., 1989a,b; Lemmer et al., 1993; Leverenz et al., 2001; Pascual et al., 1992; Ruiz et al., 1993; Shimohama et al., 1986; Szot et al., 2006, 2007), functional changes in G proteins (Cowburn et al., 2001; Hashimoto et al., 2004), changes in the levels of regulators, such as GRKs (Suo and Li, 2010; Suo, 2013) and β -arrestin 2 (Thathiah et al., 2013), and changes in levels and activity of down-stream effectors, such as adenylyl cyclase and cAMP (Dierssen et al., 1996; Dierssen et al., 1997; Ohm et al., 1989; Ross et al., 1993; Yamamoto-Sasaki et al., 1999; Yamamoto et al., 1996, 1997), in AD brains could all lead to dysregulation of adrenergic receptors and their malfunction in cognition.

5. Conclusions and perspectives

Pathological changes to the NA system occur early in AD. However, the underlying mechanisms driving these changes and their contributions to AD pathogenesis remain to be determined. Nonetheless, existing evidence tends to paint a more complex picture of NA dysfunction in AD than a simple, straight-forward loss of NE input to the cerebral cortex. Despite the well-documented loss of NE producing NA neurons, the extracellular levels of NE, especially at the early stages of disease, are likely unchanged or even increased due to a

number of compensatory mechanisms. Moreover, the endogenous NE target, adrenergic receptors, are dysregulated at multiple levels in AD so that activation of these receptors seem to elicit a number of deteriorating effects rather than promoting cognition. Overall, the integrity of the NA system appears to be very integral in maintaining a balance that allows for the proper operation of a variety of physiological processes that are dysregulated in AD. Alterations at the anatomical, cellular, and molecular levels of NA components in AD support that loss of the integrity of the NA system, not merely loss of the NE input, is a key contributor to AD development.

Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Abbreviations:

AR	adrenergic receptor
AD	Alzheimer's disease
APP	amyloid precursor protein
COMT	catechol-O-methyltransferases
CSF	cerebral spinal fluid
DβH	dopamine β hydroxylase
GPCR	G-protein coupled receptor
LC	locus coeruleus
MAO	monoamine oxidases
NET	norepinephrine transporter
DSP-4	N-(2-chloroethyl)-N-ethyl-2-bromobenzylamine

References

- Abercrombie ED, Zigmond MJ, 1989 Partial injury to central noradrenergic neurons: reduction of tissue norepinephrine content is greater than reduction of extracellular norepinephrine measured by microdialysis. *J. Neurosci* 9, 4062–4067. [PubMed: 2511282]
- Adolfsson R et al., 1979 Changes in the brain catecholamines in patients with dementia of Alzheimer type. *Br. J. Psychiatry* 135, 216–223. [PubMed: 486847]
- Arnsten AF, Jentsch JD, 1997 The alpha-1 adrenergic agonist, cirazoline, impairs spatial working memory performance in aged monkeys. *Pharmacol. Biochem. Behav* 58, 55–59. [PubMed: 9264070]
- Arnsten AF, 1998 The biology of being frazzled. *Science* 280, 1711–1712. [PubMed: 9660710]
- Arnsten AF, Goldman-Rakic PS, 1998 Noise stress impairs prefrontal cortical cognitive function in monkeys: evidence for a hyperdopaminergic mechanism. *Arch. Gen. Psychiatry* 55, 362–368. [PubMed: 9554432]

- Booze RM et al., 1993 Frequency analysis of catecholamine axonal morphology in human brain. II. Alzheimer's disease and hippocampal sympathetic ingrowth. *J. Neurol. Sci* 119, 110–118. [PubMed: 7902423]
- Braak H, Del Tredici K, 2011a The pathological process underlying Alzheimer's disease in individuals under thirty. *Acta Neuropathol.* 121, 171–181. [PubMed: 21170538]
- Braak H, Del Tredici K, 2011b Alzheimer's pathogenesis: is there neuron-to-neuron propagation? *Acta Neuropathol.* 121, 589–595. [PubMed: 21516512]
- Braak H et al., 2011 Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years. *J. Neuropathol. Exp. Neurol* 70, 960–969. [PubMed: 22002422]
- Braak H, Del Tredici K, 2012 Where, when, and in what form does sporadic Alzheimer's disease begin? *Curr. Opin. Neurol* 25, 708–714. [PubMed: 23160422]
- Braak H et al., 2013 Age-related appearance of dendritic inclusions in catecholaminergic brainstem neurons. *Neurobiol. Aging* 34, 286–297. [PubMed: 22503003]
- Branca C et al., 2014 Administration of a selective beta2 adrenergic receptor antagonist exacerbates neuropathology and cognitive deficits in a mouse model of Alzheimer's disease. *Neurobiol. Aging* 35, 2726–2735. [PubMed: 25034342]
- Chamberlain SR, Robbins TW, 2013 Noradrenergic modulation of cognition: therapeutic implications. *J. Psychopharmacol* 27, 694–718. [PubMed: 23518815]
- Chan-Palay V, Asan E, 1989 Alterations in catecholamine neurons of the locus coeruleus in senile dementia of the Alzheimer type and in Parkinson's disease with and without dementia and depression. *J. Comp. Neurol* 287, 373–392. [PubMed: 2570794]
- Chen Y et al., 2014 alpha(2A) adrenergic receptor promotes amyloidogenesis through disrupting APP-SorLA interaction. *Proc. Natl. Acad. Sci. USA* 111, 17296–17301. [PubMed: 25404298]
- Chen ZJ, Minneman KP, 2005 Recent progress in alpha1-adrenergic receptor research. *Acta Pharmacol Sin.* 26, 1281–1287. [PubMed: 16225747]
- Cowburn RF et al., 2001 Receptor-G-protein signalling in Alzheimer's disease. *Biochem. Soc. Symp.* 163–175 [PubMed: 11447833]
- Czech C et al., 2012 Metabolite profiling of Alzheimer's disease cerebrospinal fluid. *PLoS One* 7, e31501. [PubMed: 22359596]
- D'Amato RJ et al., 1987 Aminergeric systems in Alzheimer's disease and Parkinson's disease. *Ann. Neurol* 22, 229–236. [PubMed: 3477996]
- Decker MW, Gill TM, McGaugh JL, 1990 Concurrent muscarinic and beta-adrenergic blockade in rats impairs place-learning in a water maze and retention of inhibitory avoidance. *Brain Res.* 513, 81–85. [PubMed: 2161697]
- Dierssen M et al., 1996 Impaired cyclic AMP production in the hippocampus of a Down syndrome murine model. *Brain Res. Dev. Brain Res* 95, 122–124. [PubMed: 8873983]
- Dierssen M et al., 1997 Alterations of central noradrenergic transmission in Ts65Dn mouse, a model for Down syndrome. *Brain Res.* 749, 238–244. [PubMed: 9138724]
- Do Monte FH et al., 2013 Systemic or intra-prelimbic cortex infusion of prazosin impairs fear memory reconsolidation. *Behav. Brain Res* 244, 137–141. [PubMed: 23380678]
- Dobarro M, Gerenu G, Ramirez MJ, 2013a Propranolol reduces cognitive deficits, amyloid and tau pathology in Alzheimer's transgenic mice. *Int. J. Neuropsychopharmacol* 16, 2245–2257. [PubMed: 23768694]
- Dobarro M et al., 2013b Propranolol restores cognitive deficits and improves amyloid and Tau pathologies in a senescence-accelerated mouse model. *Neuropharmacology* 64, 137–144. [PubMed: 22824191]
- Doze VA et al., 2011 Long-term alpha1A-adrenergic receptor stimulation improves synaptic plasticity, cognitive function, mood, and longevity. *Mol. Pharmacol* 80, 747–758. [PubMed: 21791575]
- Elrod R et al., 1997 Effects of Alzheimer's disease severity on cerebrospinal fluid norepinephrine concentration. *Am. J. Psychiatry* 154, 25–30. [PubMed: 8988954]
- Engelborghs S et al., 2008 The dopaminergic neurotransmitter system is associated with aggression and agitation in frontotemporal dementia. *Neurochem. Int* 52, 1052–1060. [PubMed: 18093695]

- Finlay JM, Zigmond MJ, Abercrombie ED, 1995 Increased dopamine and norepinephrine release in medial prefrontal cortex induced by acute and chronic stress: effects of diazepam. *Neuroscience* 64, 619–628. [PubMed: 7715775]
- Forstl H et al., 1994 Disproportionate loss of noradrenergic and cholinergic neurons as cause of depression in Alzheimer's disease - a hypothesis. *Pharmacopsychiatry* 27, 11–15.
- Gannon M et al., 2015 Noradrenergic dysfunction in Alzheimer's disease. *Front. Neurosci* 9, 220. [PubMed: 26136654]
- German DC et al., 1992 Disease-specific patterns of locus coeruleus cell loss. *Ann. Neurol* 32, 667–676. [PubMed: 1449247]
- Giubilei F et al., 2004 Increased lymphocyte dopamine beta-hydroxylase immunoreactivity in Alzheimer's disease: compensatory response to cholinergic deficit? *Dement. Geriatr. Cogn. Disord* 18, 338–341. [PubMed: 15305112]
- Golan DE, Tashjian AH, Armstrong EJ, 2011 Principles of pharmacology: the pathophysiologic basis of drug therapy, Vol., Lippincott Williams & Wilkins.
- Goldstein LE et al., 1996 Role of the amygdala in the coordination of behavioral, neuroendocrine, and prefrontal cortical monoamine responses to psychological stress in the rat. *J. Neurosci* 16, 4787–4798. [PubMed: 8764665]
- Gottfries CG et al., 1983 Biochemical changes in dementia disorders of Alzheimer type (AD/SDAT). *Neurobiol. Aging* 4, 261–271. [PubMed: 6200784]
- Gulyas B et al., 2010 The norepinephrine transporter (NET) radioligand (S, S)-[18F] FMeNER-D2 shows significant decreases in NET density in the human brain in Alzheimer's disease: a post-mortem autoradiographic study. *Neurochem. Int* 56, 789–798. [PubMed: 20211213]
- Hall RA, 2004 Beta-adrenergic receptors and their interacting proteins. *Semin. Cell Dev. Biol* 15, 281–288. [PubMed: 15125891]
- Hashimoto E et al., 2004 Impairment of G(salpa) function in human brain cortex of Alzheimer's disease: comparison with normal aging. *J. Neural. Transm. (Vienna)* 111, 311–322. [PubMed: 14991457]
- Hein L, 2006 Adrenoceptors and signal transduction in neurons. *Cell Tissue Res.* 326, 541–551. [PubMed: 16896948]
- Herregodts P et al., 1989 Monoaminergic neurotransmitters in Alzheimer's disease. An HPLC study comparing presenile familial and sporadic senile cases. *J. Neurol. Sci* 92, 101–116. [PubMed: 2570132]
- Hoogendijk WJ et al., 1999 Increased activity of surviving locus ceruleus neurons in Alzheimer's disease. *Ann. Neurol* 45, 82–91. [PubMed: 9894881]
- Hughes ZA, Stanford SC, 1998 A partial noradrenergic lesion induced by DSP-4 increases extracellular noradrenaline concentration in rat frontal cortex: a microdialysis study in vivo. *Psychopharmacology (Berl)*. 136, 299–303. [PubMed: 9566816]
- Iversen LL et al., 1983 Loss of pigmented dopamine-beta-hydroxylase positive cells from locus coeruleus in senile dementia of Alzheimer's type. *Neurosci. Lett* 39, 95–100. [PubMed: 6633940]
- Kaddurah-Daouk R et al., 2011 Metabolomic changes in autopsy-confirmed Alzheimer's disease. *Alzheimer's Dementia* 7, 309–317.
- Kalaria RN, 1989 Characterization of [125I]HEAT binding to alpha 1-receptors in human brain: assessment in aging and Alzheimer's disease. *Brain Res.* 501, 287–294. [PubMed: 2555022]
- Kalaria RN et al., 1989a Adrenergic receptors in aging and Alzheimer's disease: increased beta 2-receptors in prefrontal cortex and hippocampus. *J. Neurochem* 53, 1772–1781. [PubMed: 2553864]
- Kalaria RN, Stockmeier CA, Harik SI, 1989b Brain microvessels are innervated by locus coeruleus noradrenergic neurons. *Neurosci. Lett* 97, 203–208. [PubMed: 2537480]
- Kask A et al., 1997 Overflow of noradrenaline and dopamine in frontal cortex after [N-(2-chloroethyl)-N-ethyl-2-bromobenzylamine] (DSP-4) treatment: in vivo microdialysis study in anaesthetized rats. *Naunyn Schmiedebergs Arch. Pharmacol* 355, 267–272. [PubMed: 9050022]
- Katsouri L et al., 2013 Prazosin, an alpha(1)-adrenoceptor antagonist, prevents memory deterioration in the APP23 transgenic mouse model of Alzheimer's disease. *Neurobiol. Aging* 34, 1105–1115. [PubMed: 23063647]

- Kenton L, Boon F, Cain DP, 2008 Combined but not individual administration of beta-adrenergic and serotonergic antagonists impairs water maze acquisition in the rat. *Neuropsychopharmacology* 33, 1298–1311. [PubMed: 17653108]
- Knaus AE et al., 2007 Alpha2-adrenoceptor subtypes-unexpected functions for receptors and ligands derived from gene-targeted mouse models. *Neurochem. Int* 51, 277–281. [PubMed: 17664025]
- Lemmer B et al., 1993 Beta-adrenoceptor density and subtype distribution in cerebellum and hippocampus from patients with Alzheimer's disease. *Naunyn Schmiedebergs Arch. Pharmacol* 347, 214–219. [PubMed: 8097284]
- Leverenz JB et al., 2001 Increased alpha 2-adrenergic receptor binding in locus coeruleus projection areas in dementia with Lewy bodies. *Neurobiol. Aging* 22, 555–561. [PubMed: 11445255]
- Mann DMA et al., 1980 Changes in the monoamine containing neurones of the human CNS in senile dementia. *British J. Psychiatry* 136, 533–541.
- Marcyniuk B, Mann DM, Yates PO, 1986 Loss of nerve cells from locus coeruleus in Alzheimer's disease is topographically arranged. *Neurosci. Lett* 64, 247–252. [PubMed: 3960404]
- Martignoni E et al., 1992 Cerebrospinal fluid norepinephrine, 3-methoxy-4-hydroxyphenylglycol and neuropeptide Y levels in Parkinson's disease, multiple system atrophy and dementia of the Alzheimer type. *J. Neural. Transm. Park Dis. Dement. Sect 4*, 191–205. [PubMed: 1320891]
- Matthews KL et al., 2002 Noradrenergic changes, aggressive behavior, and cognition in patients with dementia. *Biol. Psychiatry* 51, 407–416. [PubMed: 11904135]
- McMillan PJ et al., 2011 Differential response of the central noradrenergic nervous system to the loss of locus coeruleus neurons in Parkinson's disease and Alzheimer's disease. *Brain Res.* 1373, 240–252. [PubMed: 21147074]
- Mishima K et al., 2004 Characteristics of behavioral abnormalities in alpha1d-adrenoceptors deficient mice. *Behav. Brain Res* 152, 365–373. [PubMed: 15196805]
- Miyata S et al., 1984 Dopamine-beta-hydroxylase activities in serum and cerebrospinal fluid of aged and demented patients. *J. Neurol. Sci* 63, 403–409. [PubMed: 6726279]
- Mustapic M et al., 2013 Genotype-independent decrease in plasma dopamine beta-hydroxylase activity in Alzheimer's disease. *Progr. NeuroPsychopharmacol. Biol. Psychiatry* 44, 94–99.
- Nalepa I et al., 2013 alpha1-Adrenergic receptor subtypes in the central nervous system: insights from genetically engineered mouse models. *Pharmacol. Rep* 65, 1489–1497. [PubMed: 24552996]
- Nelson AR, Kolasa K, McMahon LL, 2014 Noradrenergic sympathetic sprouting and cholinergic reinnervation maintains non-amyloidogenic processing of AbetaPP. *J. Alzheimers Dis* 38, 867–879. [PubMed: 24081376]
- Ni Y et al., 2006 Activation of beta2-adrenergic receptor stimulates gamma-secretase activity and accelerates amyloid plaque formation. *Nat. Med* 12, 1390–1396. [PubMed: 17115048]
- Ohm TG, Bohl J, Lemmer B, 1989 Reduced cAMP-signal transduction in postmortem hippocampus of demented old people. *Prog. Clin. Biol. Res* 317, 501–509. [PubMed: 2557639]
- Oishi M et al., 1996 Laboratory examinations correlated with severity of dementia. *Ann. Clin. Lab. Sci* 26, 340–345. [PubMed: 8800434]
- Oleskevich S, Descarries L, Lacaille JC, 1989 Quantified distribution of the noradrenaline innervation in the hippocampus of adult rat. *J. Neurosci* 9, 3803–3815. [PubMed: 2585056]
- Ouyang M, Thomas SA, 2005 A requirement for memory retrieval during and after long-term extinction learning. *Proc. Natl. Acad. Sci. USA* 102, 9347–9352. [PubMed: 15947076]
- Palmer AM et al., 1984 Monoamine metabolite concentrations in lumbar cerebrospinal fluid of patients with histologically verified Alzheimer's dementia. *J. Neurol. Neurosurg. Psychiatry* 47, 481–484. [PubMed: 6204017]
- Palmer AM et al., 1987 Catecholaminergic neurones assessed ante-mortem in Alzheimer's disease. *Brain Res.* 414, 365–375. [PubMed: 3620938]
- Pascual J et al., 1992 Loss of high-affinity alpha 2-adrenoceptors in Alzheimer's disease: an autoradiographic study in frontal cortex and hippocampus. *Neurosci. Lett* 142, 36–40. [PubMed: 1357610]
- Pauszek ME, 1991 Propranolol for treatment of agitation in senile dementia. *Indiana Med.* 84, 16–17. [PubMed: 1997597]

- Perez DM, 2007 Structure-function of alpha1-adrenergic receptors. *Biochem. Pharmacol* 73, 1051–1062. [PubMed: 17052695]
- Peskind ER et al., 1995 Effects of Alzheimer's disease and normal aging on cerebrospinal fluid norepinephrine responses to yohimbine and clonidine. *Arch. Gen. Psychiatry* 52, 774–782. [PubMed: 7654129]
- Peskind ER et al., 2005 Propranolol for disruptive behaviors in nursing home residents with probable or possible Alzheimer disease: a placebo-controlled study. *Alzheimer Dis. Assoc. Disord* 19, 23–28. [PubMed: 15764868]
- Qin S et al., 2009 Acute psychological stress reduces working memory-related activity in the dorsolateral prefrontal cortex. *Biol. Psychiatry* 66, 25–32. [PubMed: 19403118]
- Raskind MA et al., 1984 Norepinephrine and MHPG levels in CSF and plasma in Alzheimer's disease. *Arch. Gen. Psychiatry* 41, 343–346. [PubMed: 6703854]
- Raskind MA, Peskind ER, 1994 Neurobiologic bases of noncognitive behavioral problems in Alzheimer disease. *Alzheimer Dis. Assoc. Disord* 8 (Suppl 3), 54–60. [PubMed: 7999346]
- Raskind MA et al., 1999 Patterns of cerebrospinal fluid catechols support increased central noradrenergic responsiveness in aging and Alzheimer's disease. *Biol. Psychiatry* 46, 756–765. [PubMed: 10494443]
- Reinikainen KJ et al., 1988 A post-mortem study of noradrenergic, serotonergic and GABAergic neurons in Alzheimer's disease. *J. Neurol. Sci* 84, 101–116. [PubMed: 2452858]
- Reinikainen KJ, Soininen H, Riekkinen PJ, 1990 Neurotransmitter changes in Alzheimer's disease: implications to diagnostics and therapy. *J. Neurosci. Res* 27, 576–586. [PubMed: 1981917]
- Ross BM et al., 1993 Alterations in the activity of adenylate cyclase and high affinity GTPase in Alzheimer's disease. *Brain Res.* 622, 35–42. [PubMed: 8242380]
- Ruetti E et al., 2014 Corticosterone and propranolol's role on taste recognition memory. *Pharmacol. Biochem. Behav* 127, 37–41. [PubMed: 25268313]
- Ruiz J et al., 1993 Non-adrenoceptor [3H]idazoxan binding sites (I2-imidazoline sites) are increased in postmortem brain from patients with Alzheimer's disease. *Neurosci. Lett* 160, 109–112. [PubMed: 7902542]
- Saber AJ, Cain DP, 2003 Combined beta-adrenergic and cholinergic antagonism produces behavioral and cognitive impairments in the water maze: implications for Alzheimer disease and pharmacotherapy with beta-adrenergic antagonists. *Neuropsychopharmacology* 28, 1247–1256. [PubMed: 12700678]
- Shankle WR, Nielson KA, Cotman CW, 1995 Low-dose propranolol reduces aggression and agitation resembling that associated with orbitofrontal dysfunction in elderly demented patients. *Alzheimer Dis. Assoc. Disord* 9, 233–237. [PubMed: 8749613]
- Sharp SI et al., 2007 Aggressive behavior and neuroleptic medication are associated with increased number of alpha1-adrenoceptors in patients with Alzheimer disease. *Am. J. Geriatr. Psychiatry* 15, 435–437. [PubMed: 17463193]
- Shimohama S et al., 1986 Biochemical characterization of alpha-adrenergic receptors in human brain and changes in Alzheimer-type dementia. *J. Neurochem* 47, 1295–1301. [PubMed: 3018166]
- Summers WK, 2006 The management of agitation in demented patients with propranolol. *J. Alzheimers Dis* 9, 69–75. [PubMed: 16627936]
- Suo WZ, Li L, 2010 Dysfunction of G protein-coupled receptor kinases in Alzheimer's disease. *The Scientific World Journal* 10, 1667–1678. [PubMed: 20730384]
- Suo WZ, 2013 Accelerating Alzheimer's pathogenesis by GRK5 deficiency via cholinergic dysfunction. *Adv. Alzheimer's Dis*, 148–160
- Swanson LW, Hartman BK, 1975 The central adrenergic system. An immunofluorescence study of the location of cell bodies and their efferent connections in the rat utilizing dopamine-beta-hydroxylase as a marker. *J. Comp. Neurol* 163, 467–505. [PubMed: 1100685]
- Szot P et al., 2006 Compensatory changes in the noradrenergic nervous system in the locus ceruleus and hippocampus of postmortem subjects with Alzheimer's disease and dementia with Lewy bodies. *J. Neurosci* 26, 467–478. [PubMed: 16407544]
- Szot P et al., 2007 Changes in adrenoreceptors in the prefrontal cortex of subjects with dementia: evidence of compensatory changes. *Neuroscience* 146, 471–480. [PubMed: 17324522]

- Taherian F et al., 2014 Propranolol-induced impairment of contextual fear memory reconsolidation in rats: a similar effect on weak and strong recent and remote memories. *Basic Clin. Neurosci* 5, 231–239. [PubMed: 25337385]
- Tejani-Butt SM, Yang J, Zaffar H, 1993 Norepinephrine transporter sites are decreased in the locus coeruleus in Alzheimer's disease. *Brain Res.* 631, 147–150. [PubMed: 8298987]
- Thathiah A, De Strooper B, 2011 The role of G protein-coupled receptors in the pathology of Alzheimer's disease. *Nat. Rev. Neurosci* 12, 73–87. [PubMed: 21248787]
- Thathiah A et al., 2013 [beta]-arrestin 2 regulates A[beta] generation and [gamma]-secretase activity in Alzheimer's disease. *Nat. Med* 19, 43–49. [PubMed: 23202293]
- Tohgi H et al., 1992 Concentration of monoamines and their metabolites in the cerebrospinal fluid from patients with senile dementia of the Alzheimer type and vascular dementia of the Binswanger type. *J. Neural Transmission - Parkinson's Dis. Dementia Sect* 4, 69–77.
- Ventura R et al., 2003 Norepinephrine in the prefrontal cortex is critical for amphetamine-induced reward and mesoaccumbens dopamine release. *J. Neurosci* 23, 1879–1885. [PubMed: 12629192]
- Veyrac A et al., 2007 Noradrenergic control of odor recognition in a nonassociative olfactory learning task in the mouse. *Learn Mem.* 14, 847–854. [PubMed: 18086828]
- Wang D et al., 2013a β 2 Adrenergic Receptor, Protein Kinase A (PKA) and c-Jun N-terminal Kinase (JNK) Signaling Pathways Mediate Tau Pathology in Alzheimer Disease Models. *J. Biol. Chem* 288, 10298–10307. [PubMed: 23430246]
- Wang J et al., 2013b Investigation of nebivolol as a novel therapeutic agent for the treatment of Alzheimer's disease. *J. Alzheimers Dis* 33, 1147–1156. [PubMed: 23128558]
- Wang LY et al., 2009 Prazosin for the treatment of behavioral symptoms in patients with Alzheimer disease with agitation and aggression. *Am. J. Geriatr. Psychiatry* 17, 744–751. [PubMed: 19700947]
- Wang LY et al., 2013c Cerebrospinal fluid norepinephrine and cognition in subjects across the adult age span. *Neurobiol. Aging* 34, 2287–2292. [PubMed: 23639207]
- Watson GS et al., 2006 Insulin effects on CSF norepinephrine and cognition in Alzheimer's disease. *Neurobiol. Aging* 27, 38–41. [PubMed: 16298239]
- Yamamoto-Sasaki M et al., 1999 Impaired phosphorylation of cyclic AMP response element binding protein in the hippocampus of dementia of the Alzheimer type. *Brain Res.* 824, 300–303. [PubMed: 10196463]
- Yamamoto M et al., 1996 Reduced immunoreactivity of adenylyl cyclase in dementia of the Alzheimer type. *Neuroreport* 7, 2965–2970. [PubMed: 9116220]
- Yamamoto M et al., 1997 Ca^{2+} /CaM-sensitive adenylyl cyclase activity is decreased in the Alzheimer's brain: possible relation to type I adenylyl cyclase. *J. Neural. Transm. (Vienna)* 104, 721–732. [PubMed: 9444571]
- Yu NN et al., 2010 Blocking beta2-adrenergic receptor attenuates acute stress-induced amyloid beta peptides production. *Brain Res.* 1317, 305–310. [PubMed: 20059989]
- Zarow C et al., 2003 Neuronal loss is greater in the locus coeruleus than nucleus basalis and substantia nigra in Alzheimer and Parkinson diseases. *Arch. Neurol* 60, 337–341. [PubMed: 12633144]