Published in final edited form as:

Chem Commun (Camb). 2019 March 18; 55(22): 3286–3289. doi:10.1039/c9cc01243e.

Nucleophilic addition of phosphorus(III) derivatives to squaraines: colorimetric detection of transition metal-mediated or thermal reversion

Emily P. Bacher^a, Antonio J. Lepore^a, Deisy Pena-Romero^a, Bradley D. Smith^a, Brandon L. Ashfeld^a

^aDepartment of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, United States

Abstract

Nucleophilic addition of phosphorus(III) agents to the electrophilic core of intensely colored squaraine dyes gives a bleached zwitterionic adduct in good to excellent yields (up to 99%) at room temperature. The process can be reversed by adding specific transition metal complexes with high phosphorous(III) affinity.

Graphical Abstract

While squaraine dyes are best known for their chromophoric properties and use in non-invasive imaging of biological processes, nonlinear optics, photovoltaics, and ion sensing, the structural core represents a fascinating bonding motif of relatively unexplored synthetic potential.¹ Dianiline squaraines have two *para*-substituted anilines in a 1,3-orientation around the central, four-membered ring resulting in an electron deficient core with donor-acceptor-donor resonance stabilization. This orbital arrangement produces the narrow and intense absorption and fluorescence emission maxima bands in the near infrared region of light that are characteristic of this attractive class of dyes.^{2,3} Because of its electrophilic nature, the central ring is susceptible to attack by sterically unencumbered nucleophiles (Figure 1a).^{4,5} While this nucleophilic bleaching of squaraines is problematic in fluorescence imaging,⁶ exploitation of this inherent electrophilicity is useful in the design and development of chemical sensing agents for trace analytes such as cyanide anion and thiols.

Conflicts of interest

There are no conflicts to declare.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here].
See DOI: [10.1039/x0xx00000x](https://doi.org/10.1039/x0xx00000x)

In 2008, Anslyn and coworkers demonstrated that bleached squaraine-thiol adducts are effective colorimetric dosimeters for Pd^{II} salts through the abstraction of thiolates from adduct **1** to regenerate the squaraine **2a** (Figure 1b).⁷ During the course of their study, they discovered that unlike other Pd^{II} salts (i.e., Pd(NO₃)₂, Pd(OAc)₂), a decrease in the solution absorbance was observed when >0.75 equivalents of Pd(PPh₃)₂Cl₂ were added to **1**. Additionally, they noted that the presence of PPh₃ induced loss of the squaraine blue color, but the mechanism of this bleaching process was not pursued. In conceptually related work, Zhuang⁸ and Rivera-Fuentes⁹ independently reported the reversible addition of phosphines to cyanine dyes in high-resolution cell imaging by stochastic optical reconstruction microscopy (STORM) and photoactivatable probes for induction of intracellular reductive stress, respectively.¹⁰ In contrast, addition of a phosphorus(III) reagent to squaraine **2** would result in a neutral, zwitterionic complex **3**, likely possessing distinctly different physiochemical properties (Figure 1c). Inspired by this prior art and our lab's previous explorations of phosphorus additions to electrophilic systems,¹¹ we initiated an investigation into the reactivity of dianiline squaraines with nucleophilic phosphorus(III) derivatives, and their potential as colorimetric sensing agents. Herein, we report our initial findings on the structure, stability, and reversible formation of zwitterionic squaraine-phosphine adducts.

We began our study by investigating the addition of a nucleophilic trialkyl phosphine and an *N,N*-dialkylaniline squaraine dye. Initial exposure of dianiline squaraine **2a** to P^{*n*}Bu₃ at room temperature resulted in a rapid bleaching of the dye's intense blue color (ca. 5 min) and isolation of the bench stable, pale yellow phosphine adduct **3a** in 96% yield (Scheme 1). Single crystal X-ray diffraction analysis of **3a** revealed a zwitterionic species which, to the best of our knowledge, is the first crystal structure of an adduct resulting from nucleophilic addition to a squaraine dye. The solid-state structure of **3a** has several notable features, including a distortion of the central ring geometry. As a result of the phosphine addition, the central four membered ring adopts a "kite quadrilateral" shape, rather than the square in **2a**, with two pairs of consecutive sides having unequal length. The shorter pair of bonds in the central ring are incorporated within a delocalized 3-oxopropenolate substructure that has two nearly identical C-O bond lengths of 1.230 and 1.237 Å. The associated O-C-C angles are 137° which is significantly larger than the expected angle of 120° for a linear version of this substructure.

To better understand the formation and stability of zwitterion adduct **3** as a function of the phosphorus(III) agent and dye structure, we conducted a series of experiments wherein **2a** and the more organic-soluble squaraine dyes **2b** and **2c** were exposed to either P^{*n*}Bu₃ or P(NMe₂)₃ at room temperature (Table 1). In each experiment, one equivalent of PL₃ was added to a squaraine dye **2** in CHCl₃ at room temperature, and the resulting adduct obtained via recrystallization. The addition of P^{*n*}Bu₃ to *N,N,N,N*-tetrabutyl and *N,N,N,N*-tetrabenzyl squaraines **2b** and **2c** provided the corresponding phosphine adducts **3b-c** in quantitative yield (entries 1 and 2). Similarly, the bulky, electron rich P(NMe₂)₃ gave zwitterionic adducts **3d-f** from squaraines **2a-c** respectively in excellent yields (entries 3–5). Separate samples of the zwitterionic adducts in CDCl₃ were monitored over time (>7 d) via ¹H NMR at room temperature and in each case there was <5% reversion to the parent squaraine. The adducts were also found to be permanently stable when stored as solid samples.

To evaluate the capacity of these stable dianiline squaraine-phosphorus(III) adducts to act as colorimetric metal sensors¹² we examined the chemodosimeric behavior of squaraine-phosphine complex **3b** by treating it with a series of transition metal catalysts commonly used in modern synthetic organic chemistry. Employing absorption spectroscopy to monitor the extent of dye regeneration, we observed rapid, partial reversion to the parent squaraine dye **2b** upon addition of both relatively electron rich and electron deficient Pd^{II}, Rh^I, Rh^{II}, Ir^I, Ag^I, Au^{III}, Cu^I, and Cu^{II} metal complexes. In contrast, addition of NiCl₂, Ni(cod)₂, FeCl₃, TiCl₃, or TiCl₄ did not lead to any noticeable amount of **2b**. The titration of Au(PPh₃)Cl resulted in partial reversion back to **2b** after addition of 1.2 molar equivalents of Au^I (Figure 2a). Titration of complex **3b** with either Pd(OAc)₂ or [Ir(cod)Cl]₂ showed a weaker P^{*n*}Bu₃ affinity as indicated by the greater stoichiometric amounts of transition metal complex needed to produce free **2b**. The reversion of **3b** with increasing concentration of Pd(OAc)₂ is further illustrated by the notable increase in absorbance depicted in Figure 2b. In contrast, Rh(PPh₃)₃Cl was the least effective at promoting retroversion of **3b** to dye **2b**.

Interestingly, the extent of squaraine regeneration appears to correlate with the electronics and coordination capabilities of the complexes examined. For example, titration of the coordinatively unsaturated, square planar Wilkinson's catalyst proved less effective at phosphine abstraction. This observation is consistent if one considers the electron rich nature of Rh^I in combination with the unfavorable square pyramidal geometry the complex would have to adopt upon net loss of phosphine from adduct **3b**. In contrast, the electron deficient nature of Pd(OAc)₂ and the favorable square planar geometry upon phosphine coordination results in more efficient dye regeneration. The [Ir(cod)Cl]₂ complex bearing readily exchangeable neutral cyclooctadienyl ligands proved comparable to Pd(OAc)₂, while the relative electrophilicity of Au(PPh₃)Cl results in the highest absorbance titration. It is important to note that even in the best case, full reversion back to **2b** was not observed, highlighting an important contrast with analogous squaraine-thiol adducts where the dye is more completely regenerated by Pd^{II} salts. However, an attractive feature of this squaraine-phosphine adduct is its potential to detect trace transition metal derivatives that are not sufficiently thiophilic for detection using a comparable squaraine-thiol adduct.

During our initial survey of squaraine reactivity with different phosphine agents, we observed that the addition of less nucleophilic P(*p*-MeOC₆H₄)₃ to squaraine **2a** at room temperature provided adduct **3g** in a comparatively low 51% yield. Intrigued by this comparatively low level of conversion to zwitterion **3g**, we examined the addition of P(*p*-MeOC₆H₄)₃ to squaraine **2a** in more detail. Speculating that the equilibrium between squaraine and phosphine adduct could be controlled thermally, we examined the relative ratio of **2a/3g** over a temperature range of -40 °C to 40 °C (Figure 3). As shown in Figure 2, the solution at 40 °C had the characteristic dark blue color of free **2a**, whereas cooling the solution incrementally to -40 °C indicated extensive conversion to the pale yellow-green adduct **3g**. Moreover, the thermochromic effect proved reversible as the dark blue color of the squaraine dye was restored upon warming the solution back to 40 °C.

In an effort to quantify this thermal cycling phenomenon, we used variable temperature ¹H NMR spectroscopy to monitor a single solution containing a binary mixture of P(*p*-MeOC₆H₄)₃ and the highly soluble squaraine **2b**. As illustrated in Figure 4, ¹H NMR

spectra were acquired at different temperatures for the same sample, and revealed a dynamic ratio of **3h:2b** which depended strongly on the solution temperature. Upon introduction of P(*p*-MeOC₆H₄)₃ to **2b** at 20 °C, the ¹H NMR spectrum indicated a 2.3:1 ratio of **3h:2b**. As the temperature was *lowered*, the ratio of **3h:2b** *increased* to 7.7:1 at 0 °C, 97:3 at -20 °C, and at -40 °C there was no perceptible squaraine dye remaining. Warming the sample from -40 °C to 0 °C showed that the equilibrium was regenerated without observable degradation. Indeed, three further cycles of sample thermocycling between 40 °C to -40 °C showed no indication of sample decomposition. The increase in squaraine-phosphine adduct **3** formation at lower temperature is consistent with an entropy dominated Lewis acid-base equilibrium. It should be noted that the low temperature preference for **3** is opposite to that observed for squaraine-thiol adducts where dissociation is preferred as the temperature is decreased.^{7a} As a molecular strategy, reversible squaraine thermochromism due to nucleophilic phosphorus(III) addition complements existing approaches that thermally control squaraine color by changing the degree of intermolecular packing within a squaraine-based polymer.¹³

In conclusion, the zwitterionic adducts resulting from rapid nucleophilic addition of phosphorus(III) reagents to dianiline squaraine dyes are a new class of molecular scaffolds that have the potential for further development into colorimetric sensors or thermochromic agents. While dependent on the overall nucleophilicity of the phosphorus reagent employed, the adducts from P^{*n*}Bu₃ or P(NMe₂)₃ are characterized by a stable, highly crystalline structure with an intriguing “kite quadrilateral” core architecture. The resulting equilibrium between squaraine dye and phosphorus adduct serves as the basis for colorimetric chemodosimeter detection of transition metal complexes that have a high phosphorous (III) affinity. It is particularly noteworthy that the zwitterionic adducts formed using either P^{*n*}Bu₃ or P(NMe₂)₃ are obtained in excellent yield and proved exceptionally stable at room temperature for prolonged time periods. The adduct **3b** acted as a colorimetric chemodosimeter for detection of transition metal complexes that have high phosphorous (III) affinity. The ready availability of structurally diverse phosphorus(III) reagents provides ample opportunity for the fine tuning of desired sensing or thermochromic properties. Studies toward the development of this molecular framework in synthesis are also currently underway, and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the University of Notre Dame and the National Science Foundation (CHE 1056242, CHE 1665440, and CHE 1708240) for financial support of this research and Dr. Allen Oliver for assistance with crystallography work. E.P.B. was supported by the CBBi Program and NIH training grant T32GM075762.

References

1. (a) Treibs A; Jacob K *Angew. Chem. Int. Ed* 1965, 4, 694–694. (b) Ajayaghosh A *Acc. Chem. Res* 2005, 38, 449–459. [PubMed: 15966711] (c) Avirah RR; Jyothish K; Ramaiah D *Org. Lett* 2007, 9, 121–124. [PubMed: 17192100] (d) Jyothish K; Arun KT; Ramaiah D *Org. Lett* 2004, 6, 3965–3968.

- [PubMed: 15496075] (e)Ros-Lis JV; Martínez-Máñez R; Sancenón F; Soto J; Spieles M; Rurack K Chem. Eur. J 2008, 14, 10101–10114. [PubMed: 18785676]
2. (a)Sreejith S; Joseph J; Lin M; Menon NV; Borah P; Ng HJ; Loong YX; Kang Y; Yu SW-K; Zhao Y ACS Nano 2015, 9, 5695–5704. [PubMed: 26022724] (b)Harmatys KM; Cole EL; Smith BD Mol. Pharmaceutics 2013, 10, 4263–4271.(c)Cole EL; Arunkumar E; Xiao S; Smith BA; Smith BD Org. Biomol. Chem 2012, 10, 5769–5773. [PubMed: 22159917] (d)White AG; Fu N; Leevy WM; Lee J-J; Blasco MA; Smith BD Bioconjugate Chem. 2010, 21, 1297–1304.(e)Avirah RR; Jayaram DT; Adarsh N; Ramaiah D Org. Biomol. Chem 2012, 10, 911–920. [PubMed: 22179414]
3. (a)Prabhakar C; Bhanuprakash K; Rao VJ; Balamuralikrishna M; Rao DN J. Phys. Chem. C 2010, 114, 6077–6089.(b)Wang S; Hall L; Diev VV; Haiges R; Wei G; Xiao X; Djurovich PI; Forrest SR; Thompson ME Chem. Mater 2011, 23, 4789–4798.(c)Arunkumar E; Ajayaghosh A; Daub JJ Am. Chem. Soc 2005, 127, 3156–3164.
4. Ros-Lis JV; Martinez-Manez R; Soto J Chem. Commun 2002, 2248–2249.
5. Ros-Lis JV; García B; Jiménez D; Martínez-Máñez R; Sancenón F; Soto J; Gonzalvo F; Valldcabres MC J. Am. Chem. Soc 2004, 126, 4064–4065. [PubMed: 15053569]
6. (a)Gassensmith JJ; Barr L; Baumes JM; Paek A; Nguyen A; Smith BD Org. Lett 2008, 10, 3343–3346. [PubMed: 18582079] (b)Gassensmith JJ; Baumes JM; Smith BD Chem. Commun 2009, 6329–6338.
7. (a)Houk RJT; Wallace KJ; Hewage HS; Anslyn EV Tetrahedron 2008, 64, 8271–8278. [PubMed: 19122841] (b)Hewage HS; Anslyn EV J. Am. Chem. Soc 2009, 131, 13099–13106. [PubMed: 19691345]
8. Vaughan JC; Dempsey GT; Sun E; Zhuang XJ Am. Chem. Soc 2013, 135, 1197–1200.
9. Tirla A; Rivera-Fuentes P Angew. Chem. Int. Ed 2016, 55, 14709–14712.
10. (a)Rust MJ; Bates M; Zhuang X Nat. Methods 2006, 3, 793. [PubMed: 16896339] (b)Betzig E; Patterson GH; Sougrat R; Lindwasser OW; Olenych S; Bonifacino JS; Davidson MW; Lippincott-Schwartz J; Hess HF Science 2006, 313, 1642–1645. [PubMed: 16902090] (c)Hess ST; Girirajan TPK; Mason MD Biophys. J 2006, 91, 4258–4272. [PubMed: 16980368]
11. (a)Eckert KE; Ashfeld BL Org. Lett 2018, 20, 2315–2319. [PubMed: 29634274] (b)Rodriguez KX; Vail JD; Ashfeld BL Org. Lett 2016, 18, 4514–4517. [PubMed: 27603590] (c)Wilson EE; Rodriguez KX; Ashfeld BL Tetrahedron 2015, 71, 5765–5775.(d)Haugen KC; Rodriguez KX; Chavannavar AP; Oliver AG; Ashfeld BL Tetrahedron Lett. 2015, 56, 3527–3530.(e)Chavannavar AP; Oliver AG; Ashfeld BL Chem. Commun 2014, 50, 10853–10856.
12. Kaur B; Kaur N; Kumar S Coord. Chem. Rev 2018, 358, 13–69.
13. (a)Chen G; Sasabe H; Wang X-F; Hong Z; Kido J Synth. Met 2014, 192, 10–14.(b)Yang D; Sasabe H; Jiao Y; Zhuang T; Huang Y; Pu X; Sano T; Lu Z; Kido JJ Mater. Chem. A 2016, 4, 18931–18941.(c)Galliani D; Mascheroni L; Sassi M; Turrisi R; Lorenzi R; Scaccabarozzi A; Stingelin N; Beverina L Adv. Opt. Mater 2015, 3, 1164–1168.

Figure 1. Additions to the central ring of squaraine dyes. (a) Established nucleophilic additions to squaraines. (b) Colorimetric sensors for trace metal impurities. (c) Reversible formation of bleached squaraine/phosphine adducts.

Figure 2.

A) Titration plots showing partial regeneration of squaraine **2b** absorbance at 656 nm when separate solutions of **3b** (5×10^{-5} M in DMSO) were titrated with $Rh(PPh_3)_3Cl$, $Pd(OAc)_2$, $[Ir(cod)Cl]_2$, or $Au(PPh_3)Cl$. **B)** Appearance of squaraine **2b** absorption maxima band upon titration of adduct **3b** (3×10^{-5} M in DMSO) with increasing amounts of $Pd(OAc)_2$.

Figure 3.

Photographs of a vial containing a binary mixture of $\text{P}(\text{p-MeOC}_6\text{H}_4)_3$ and **2a** in CHCl_3 at different temperatures. At 40 °C the solution exhibits a dark blue color indicating free squaraine, whereas cooling the mixture to -40 °C results in appearance of pale yellow adduct **3g**.

Figure 4. Partial ^1H NMR spectra (500 MHz, CDCl_3) for a single sample, showing how the equilibrium for reversible addition of $\text{P}(p\text{-MeOC}_6\text{H}_4)_3$ to squaraine dye **2b** to produce adduct **3h** changes with temperature. The sample temperature was cycled from 20 °C (top spectrum) through seven different temperatures to end up at -40 °C (bottom). The table provides the **3h:2b** ratio at each temperature, as determined by ^1H NMR peak integration.

Scheme 1.
Observed product **3a** for addition of P^nBu_3 to squaraine **2a**.

Table 1.

Addition of phosphine derivatives to squaraine **2** to form adduct **3**.^a

Entry	PL ₃	R	Product	Yield (%)
1	P ⁿ Bu ₃	Bu (2b)	3b	>99
2	P ⁿ Bu ₃	Bn (2c)	3c	>99
3	P(NMe ₂) ₃	Me (2a)	3d	91
4	P(NMe ₂) ₃	Bu (2b)	3e	>99
5	P(NMe ₂) ₃	Bn (2c)	3f	86

^aConditions: **2** (0.1 mmol) and PL₃ (0.11 mmol) in room temperature CHCl₃ (0.1 M).