Published in final edited form as:

Chem Res Toxicol. 2019 March 18; 32(3): 370–396. doi:10.1021/acs.chemrestox.9b00028.

RBOH-Dependent ROS Synthesis and ROS Scavenging by Plant Specialized Metabolites To Modulate Plant Development and Stress Responses

Jordan M. Chapman[†], Joëlle K. Muhlemann[†], Sheena R. Gayomba[†], Gloria K. Muday^{*}

Department of Biology and Centers for Molecular Signaling and Redox Biology and Medicine, Wake Forest University, Winston-Salem, North Carolina 27109, United States

Abstract

Reactive oxygen species (ROS) regulate plant growth and development. ROS are kept at low levels in cells to prevent oxidative damage, allowing them to be effective signaling molecules upon increased synthesis. In plants and animals, NADPH oxidase/respiratory burst oxidase homolog (RBOH) proteins provide localized ROS bursts to regulate growth, developmental processes, and stress responses. This review details ROS production via RBOH enzymes in the context of plant development and stress responses and defines the locations and tissues in which members of this family function in the model plant *Arabidopsis thaliana*. To ensure that these ROS signals do not reach damaging levels, plants use an array of antioxidant strategies. In addition to antioxidant machineries similar to those found in animals, plants also have a variety of specialized metabolites that scavenge ROS. These plant specialized metabolites exhibit immense structural diversity and have highly localized accumulation. This makes them important players in plant developmental processes and stress responses that use ROS-dependent signaling mechanisms. This review summarizes the unique properties of plant specialized metabolites, including carotenoids, ascorbate, tocopherols (vitamin E), and flavonoids, in modulating ROS homeostasis. Flavonols, a subclass of flavonoids with potent antioxidant activity, are induced during stress and development, suggesting that they have a role in maintaining ROS homeostasis. Recent results using genetic approaches have shown how flavonols regulate development and stress responses through their action as antioxidants.

Graphical Abstract

^{*}Corresponding Author: muday@wfu.edu.

[†]J.M.C., J.K.M., and S.R.G. contributed equally to this work.

The authors declare no competing financial interest.

1. INTRODUCTION

Reactive oxygen species (ROS) function as signaling molecules in both plant and mammalian cells. ROS homeostasis is an important feature of productive ROS signaling. High levels are toxic to the cell, but keeping ROS at low levels allows small increases in ROS to initiate signaling events in plant growth, development, and stress responses.^{1,2} ROS are a group of highly reactive oxygen-containing molecules that can readily accept electrons from more stable compounds. Each reactive oxygen species has unique reactivity, signaling properties, and sites of production.³ Nicotinamide adenine dinucleotide phosphate (NADPH) oxidases (NOXs), also called respiratory burst oxidase homologues (RBOHs) in plants, are critical enzymes that produce ROS in response to hormone and environmental signals.⁴ Furthermore, plants have class III peroxidases, which are able to produce superoxide compounds.⁵ Superoxide can be dismutated into H_2O_2 by superoxide dismutase (SOD),⁶ and in the presence of iron, H_2O_2 can be made into hydroxyl radicals by the Fenton reaction.⁶ In this review, we focus on highlighting how distinct members of the RBOH family display tissue-specific localization and regulation of plant development and stress responses.

Plants have evolved diverse enzymatic and non-enzymatic machineries to prevent ROS from reaching damaging levels.⁷ This review summarizes the mechanisms that plants share with animal cells but also highlights plant-derived specialized metabolites that are used to scavenge ROS, with a focus on carotenoids, ascorbate, tocopherols, and flavonoids. These metabolites exhibit large structural diversity and function. We highlight recent advances in using genetic approaches to dissect the *in vivo* antioxidant role of one of these metabolite classes, termed flavonols, in regulating a number of ROS-dependent processes.

2. NADPH OXIDASES/RBOHS AS ROS SIGNALING HUBS

NOX enzymes were first identified in human phagocytic cells for their role in generating ROS during pathogen defense.^{8,9} The human NADPH oxidase family has seven members, NOX1–NOX7.¹⁰ Most NADPH oxidase proteins catalyze the reduction of oxygen to a superoxide anion ($O_2^{\bullet-}$) using NADPH as an electron donor. The superoxide anion can subsequently produce H_2O_2 through enzymatic or non-enzymatic dismutation. NOX4 is an exception, producing H_2O_2 directly.^{11,12} Mammalian NADPH oxidases are membrane protein complexes composed of the catalytic membrane-bound flavocytochrome b_{558} and additional subunits. NOX2, the enzyme most similar to the plant RBOH family, has a catalytic subunit gp91^{phox} and a p22^{phox} subunit along with three additional cytosolic subunits: p40^{phox}, p47^{phox}, and p67^{phox}.¹³

RBOH enzymes were identified in plants on the basis of their sequence similarity to the gp91^{phox} subunit of NOX2.¹⁴ Plant RBOH proteins are also integral plasma membrane proteins with six transmembrane domains. The third and fifth transmembrane domains contain two pairs of histidine residues, which are attachment sites for the heme groups that facilitate electron transport.¹⁵ Although plant RBOH proteins have NADPH and flavin adenine dinucleotide binding domains on the C-terminus, similar to the mammalian phagocytic gp91^{phox},^{16,17} they also contain an extension of approximately 300 amino acids at the N-terminus. This region contains two calcium-binding EF hand motifs and two phosphorylation sites that participate in regulation of enzyme activity.¹⁴ The role of RBOH proteins in ROS signaling was previously reviewed,⁴ but new findings, summarized below, provide more information on the distinct role of these enzymes in developmental and stress contexts.

The *RBOH* gene family is best characterized in the model plant *Arabidopsis thaliana* because of the availability of molecular genetic tools. The *Arabidopsis* *RBOH* family has 10 members, *RBOHA* to *RBOHJ*,^{18,19} with distinct expression patterns and functional roles in development and stress response. Functions of the different RBOHs in plant growth and development are illustrated in Figure 1. The tissue-specific expression of transcripts encoding all of the isoenzymes has been examined using genome-wide transcriptional profiling and is summarized in Figure 2. The distinct RBOH isoenzyme expression patterns across plant tissues are generally linked to their reported functions.

Substantial information about RBOH function has been gathered using null mutants or plants with other genetic manipulations, such as site-directed mutations to alter protein activity. For two isoenzymes, *RBOHA* and *RBOHG*, neither mutant phenotypes nor function *in planta* have been reported. The following sections are organized by tissue and integrate current knowledge about each isoenzyme's function, expression, and regulation. These findings are also summarized by isoenzyme in Table 1.

2.1. Roots.

The *Arabidopsis* root has been well-characterized because of its simple organization, with genetically defined development regulated by hormonal and environmental stimuli; ROS have emerged as signals that modulate root development. There are four root tissue layers

organized radially: the epidermis, endodermis, cortex, and central vascular cylinder.²⁰ In addition, there are four distinct longitudinal zones arranged by maturity (Figure 3).²⁰

The meristematic zone is at the root tip and contains the stem cell niche, a pool of actively dividing stem cells and newly differentiated cells.²¹ These root meristem cells divide asymmetrically, where one daughter cell differentiates to make up one of the four tissue layers.²² The final cell identity of these daughter cells is largely dependent on positional cues of neighboring cells.^{22,23} Above the meristematic zone is the transition zone, where newly differentiated cells leave the meristem to undergo slow growth and are defined by isodiametric shape (equal length and width).^{24,25} Rapid expansion of the root occurs in the elongation zone, where cells increase in length in a matter of hours.^{26,27} Finally, the maturation zone, which encompasses the rest of the root, is delineated from the elongation zone by the initiation of root hair bulges, which extend from epidermal cells and elongate into root hairs.^{28,29} In addition to producing root hairs, the maturation zone is also the site of lateral root initiation. Lateral root primordia form from pericycle cells in the center of the primary root and develop into lateral roots after emergence through the epidermis.²⁸ The intricate higher-order root architecture is formed through continuous branching of lateral roots. The lateral root formation zone can be loosely defined as the length of the primary root encompassing the first emerged lateral root to last lateral root primordium.^{30,31}

Reactive oxygen species have been implicated in both developmental and environmental responses in roots, with established roles for RBOHC, RBOHD, and RBOHF. The function of these specific RBOH in roots is consistent with the high levels of accumulation of the transcripts encoding these enzymes in roots (Figure 2).

2.1.1. Primary Root Development and Elongation.—Plant root growth and elongation enables exploration of soil/growth media for nutrients.³² RBOHF-mediated primary root elongation is dependent on nutrient and sucrose levels.^{33,34} In complete nutrient media with 3% sucrose, primary root elongation in *rboh*d and *rboh*f single mutants and the *rboh*f/*d* double mutant is comparable to the wild-type.³⁴ Under these conditions, treatment with the plant hormone abscisic acid (ABA) inhibits primary root growth in all lines, with a smaller effect seen in mutant plants.³⁴ Another study using sucrose-free and low-nutrient media showed that *rboh*f and the double mutants had reduced primary root length in the absence of ABA compared with *rboh*d and the wild-type.³³ When ABA was applied to these lines, primary root length was inhibited in *rboh*d and the wild-type, but *rboh*f and *rboh*d/*f* showed no change compared to control conditions, suggesting that *rboh*f mutants are insensitive to ABA.³³ These studies point to a specific role of RBOHF in ABA-mediated signaling under low-nutrient conditions.

In addition to adapting their growth to nutrient levels, plant roots also alter their architecture to maximize water uptake from the soil/growth medium. Hydrotropism, the growth of roots toward water (or away from low water potential), is negatively regulated by RBOHC.³⁵ H₂O₂ accumulation in the elongation zone is presumably a signal for hydrotropic bending, as the sites of bending and H₂O₂ accumulation colocalize in this area in wild-type plants.³⁵ Assays exposing the *rboh*c mutant (which will be denoted by its more common name, *rh*d2) to low water potential showed that *rh*d2 roots are more effective in bending away from low

water potential than wild-type roots.³⁵ Enhanced bending in *rhd2* is attributed to decreased H₂O₂ content in the elongation zone, as measured by Amplex Red in extracts from root apices. The *rboh*d mutant, which also had lower H₂O₂ levels in whole seedlings but wild-type H₂O₂ levels in root apices, had a similar hydrotropic bending response to wild-type plants.³⁵ These results suggest that RBOHC provides superoxide that is then converted to H₂O₂ and that this elevated H₂O₂ reduces hydrotropic bending.

Roots contain an endodermal barrier called the Casparian strip that aids in selective uptake of nutrients and water.^{36,37} Surrounding the vascular tissue of the root, the Casparian strip is an impermeable layer adjacent to the endodermis in the maturation zone of the root and is composed of lignin and suberin, which are phenolic and lipophilic polymers, respectively.^{36,37} Radial diffusion into the root system through the cell wall is prevented by the Casparian strip, which forces water and nutrients to enter the vascular system through plasma membrane-localized transporters.^{36,37} In *Arabidopsis*, formation of the Casparian strip requires controlled deposition of lignin formed by oxidative coupling of monolignol compounds by laccases and peroxidases.³⁸ Lignification of the Casparian strip is regulated by RBOHF, which was the only NADPH oxidase found in a large-scale forward genetic screen of mutants with an impaired endodermal barrier.³⁹ Consistent with this function, *RBOHF* transcripts accumulate in the maturation zone in the stele, cortex, and endodermis (Figure 2), and *RBOHF*–mCHERRY protein fusions under control of the native promoter shows distinct fluorescence at the Casparian strip domain.³⁹ The *rboh*f mutant has delayed Casparian strip formation, consistent with the idea that superoxide produced by RBOHF is necessary to form lignin.³⁹ RBOHF activity coordinates with peroxidase 64 (PER64) to increase efficiency of Casparian strip lignification.³⁹ Collectively, these observations show distinct roles of RBOH proteins in modulation of primary root architecture and development.

2.1.2. Lateral Roots.—The initiation and elongation of lateral roots, which are root branches that develop from the primary root, lead to a robust root system that anchors plants and maximizes moisture and nutrient uptake. Recent evidence implicates ROS as a developmental signal in this process. Lateral root primordia initiate in the pericycle, a cell layer within the primary root, and must emerge through several other cell layers, including the epidermis.⁴⁰ The inhibitory effect of RBOHD and RBOHF on lateral root development has been reported. Single mutants of *rboh*d and *rboh*f revealed no difference in lateral root development, while *rboh*d/*f* double mutants had increased lateral root density in comparison with the single mutants and the wild-type.^{41,42} The homeostasis of different ROS compounds is pertinent to lateral root development. The double mutant had reduced levels of H₂O₂ throughout the root relative to the wild-type, as demonstrated by 3,3-diaminobenzidine (DAB) staining, and increased levels of superoxide in the mature region of the root compared with the wild-type, as measured using dihydroethidium (DHE) and nitro blue tetrazolium (NBT).⁴¹ The increased levels of superoxide in the root maturation zone of the double mutant are inhibited by treatment with peroxidase inhibitors, reducing superoxide levels and lateral root density to wild-type levels.⁴¹ This phenotype was shown to be independent of auxin, the plant hormone that stimulates lateral root initiation.⁴¹ These data suggest that superoxide produced by RBOHD and RBOHF is directly channeled toward

H₂O₂, which then inhibits lateral root development. The increased superoxide in the *rbohdf* mutant is attributed to increased activity of class III peroxidases, a unique class of plant-specific enzymes.⁴¹ Consistent with the role of class III peroxidases as positive regulators in lateral root development, overexpression of *PER7* and *PER57* results in increased lateral root emergence, while mutants have the opposite effect.⁴³ The balance of superoxide/H₂O₂ or H₂O₂ itself produced from RBOHD and RBOHF likely inhibits these peroxidases. Taken together, these results suggest that specific reactive oxygen species have opposing effects on lateral root number: H₂O₂ produced downstream of RBOHD and RBOHF enzymes has an inhibitory effect on lateral root number, while superoxide produced by class III peroxidases promotes lateral root number.

A second study reported impaired lateral root emergence in roots treated with diphenylene iodonium (DPI), an RBOH inhibitor, and in higher-order RBOH mutants lacking RBOHD and/or RBOHE in particular.⁴⁴ In roots, RBOHE is expressed in the three layers overlying the lateral root primordium: the endodermis, cortex, and epidermis.⁴⁴ A loss-of-function mutant of RBOHE has delayed lateral root emergence. *RBOHE* transcripts accumulate across the cell types in the maturation zone (Figure 2), and its transcript abundance increases after treatment of roots with auxin, as judged by quantitative reverse transcription polymerase chain reaction (qRT-PCR) analysis of whole roots.⁴⁴ However, the effect of auxin did not depend on the auxin influx carriers AUX1 and LAX3, as RBOHE was expressed at the same level as in the wild-type in *aux1* and *lax3* mutants in the absence of auxin treatment.⁴⁴ A *RBOHE* promoter-driven β -glucuronidase (GUS) reporter accumulated above the site of lateral root initiation, suggesting that localized ROS synthesis may facilitate emergence of lateral roots. Targeted overexpression of RBOHD in lateral root primordia and overlying tissue induced lateral root emergence.⁴⁴ The authors suggest that this localized ROS may drive changes in cell wall structure to facilitate the emergence of roots through surrounding tissues.⁴⁴

2.1.3. Root Hair Elongation.—Root hairs are single-cell extensions of epidermal cells that increase root surface area to maximize water and nutrient uptake (Figure 1).^{29,32} A screen for root hair developmental defects identified the *root hair defective 2* (*rhd2*) mutant, which has shorter root hairs than the wild-type (Figure 3).^{45,46} This mutation is mapped to the *RBOHC* gene, suggesting a role for ROS in regulation of this process.⁴⁵ *RBOHC* transcripts accumulate in epidermal cells and elongating root hairs, and RBOHC proteins fused to green fluorescent protein (GFP) localize to root hair tips,⁴⁷ consistent with ROS accumulation promoting polar tip growth.⁴⁵ Further studies showed that hydroxyl radicals ($\cdot$ OH), formed downstream of superoxide, are the specific ROS that stimulate root hair elongation.⁴⁵ Localized synthesis of ROS at the tips of root hairs via RBOHC is essential for proper root hair morphology, as global application of $\cdot$ OH did not complement the short root hair phenotype of *rhd2* but rather, led to spherical root hairs as a result of depolarized cell expansion.⁴⁵ Hydroxyl radicals facilitate cell wall loosening⁴⁸ and are required to form a tip-focused Ca²⁺ gradient in the root hair, both of which are necessary for elongation.^{45,49} Staining the wild-type and the *rhd2* mutant with calcium green dextran showed that, unlike the wild-type, the *rhd2* mutant did not form a Ca²⁺ gradient.⁴⁵ The Ca²⁺ gradient was restored in *rhd2* after exposure to $\cdot$ OH via the Fenton reaction (by addition of Cu²⁺, H₂O₂,

and ascorbate).^{45,50} Detailed below are the processes that control localization of *RBOHC* transcripts and proteins and regulation of RBOHC activity.

Chromatin immunoprecipitation (ChIP) was used to identify proteins that bind to the *RBOHC* promoter.⁵¹ This assay identified the root hair cell-localized transcription factor root hair defective six-like 4 (RSL4),⁵¹ which was previously found to promote root hair length.⁵² The *rs14-1* mutant has decreased ROS levels and shorter root hairs compared to wild-type plants, although transcription levels of *RBOHC* were not measured in the mutant.^{51,52} *RSL4* expression was found to be controlled by several auxin response factors (ARFs), including ARF5, ARF7, ARF8, and ARF19.^{51,52} As auxin has been shown to induce root hair formation,^{53,54} this ARF–RSL network may induce localized ROS synthesis that drives root hair formation in response to auxin.^{53,54}

Root hairs emerge outward from existing epidermal cells through polarized cell growth. Polarized cell growth requires cell wall expansion, which is a turgor-driven process where pressure exerted by the plasma membrane is met with resistance at the cell wall, stimulating cell wall elongation to accommodate the increase in volume.⁵⁵ RBOHC trafficking and localization is a highly regulated process that is controlled by the FERONIA (FER) receptor-like kinase⁵⁶ and members of the Rho-related GTPases of plants (ROP) family.^{57–60} The extracellular domain of FER binds to pectin components of the cell wall and activates Ca^{2+} channels that maintain cell wall integrity during cell elongation.^{56,61,62} Accordingly, *fer* and *rhd2* single mutants have similar impairments in root hair morphology and decreased ROS accumulation in root hairs, suggesting that they function in the same pathway.⁵⁶ Pull-down assays suggest that FER interacts with ROP2,⁵⁶ a protein that promotes root hair tip growth upstream of RBOHC.⁵⁸ Plants that have impaired ROP2 function or altered expression of *ROP2* have deformed root hairs.^{58,59} These phenotypes are RBOHC-dependent, as crosses between the *rhd2* mutant and constitutively active ROP2 lines fail to produce superoxide and have short root hairs.⁵⁹ Finally, endosomal cycling of RBOHC at the plasma membrane may occur through fine-tuning of ROP2 localization and stability through GDP dissociation inhibitor (GDI) proteins that participate in RAC/ROP recycling and post-translational modification.^{47,60,63}

RBOHC activity is also regulated by binding of Ca^{2+} at its N-terminal EF hand motifs and by serine phosphorylation. Site-directed mutagenesis to block the function of the EF hand motifs reduced ROS production by RBOHC in HEK293T cells,⁴⁷ a mammalian cell line that lacks endogenous NOX2 and NOX5 activity.^{64,65} Serine phosphorylation can occur through the interaction of RBOHC and the calcineurin B-like-interacting protein kinase 26 (CIPK26)–calcineurin B-like protein 1 (CBL1) sensor complex.^{47,66} Heterologous expression of wild-type or serine-to-alanine mutated versions of RBOHC showed that ROS production was dampened in serine-mutated RBOHC proteins, suggesting that phosphorylation at S318 and S322 amplifies ROS production in response to Ca^{2+} influx.⁶⁶ The authors suggested that phosphorylation of RBOHC may act in concert with Ca^{2+} oscillation to modulate ROS production in developing root hairs.^{49,66} Interestingly, root hair development in *cbl1/cipk26* double mutants and single mutants and in *rhd2* mutant lines harboring the serine-mutated RBOHC protein displayed wild-type root hair morphology,

suggesting that the two conserved serine residues are not required *in planta* or that other phosphorylation sites/kinases are involved in RBOHC activity.⁶⁶

Maize (*Zea mays*) and rice (*Oryza sativa*) genomes encode RBOH proteins named *Zea mays* roothairless5 (*Zmrth5*), and *Oryza sativa* NADPH oxidase 3 (*OsNOX3*) with high sequence similarity to *Arabidopsis* RBOHH and RBOHJ, respectively.^{67,68} Although these proteins have greater sequence similarity to pollen tube-localized *Arabidopsis* RBOHH and RBOHJ (described below), *Zmrth5* and *OsNOX3* localize to root epidermal cells and root hairs. These monocot proteins have four transmembrane domains,^{67,68} suggesting that RBOH topology differs among monocots and dicots.¹⁹ Superoxide and hydrogen peroxide levels, as visualized by NBT and DAB, were reduced in root hairs of *Zmrth5* and *Osnox3* mutants relative to parental lines. Mutations in *ZmRTH5* and *OxNOX3* result in reduced root hair initiation and elongation.^{67,68} This phenotype differs from the phenotype of the *Arabidopsis* *rhd2* mutant that forms near wild-type numbers of root hairs, but with reduced elongation.^{45,67,68} These results support the role of RBOH-dependent ROS synthesis in root hair formation across the plant kingdom, despite structural differences in these particular RBOH proteins.

2.2. Guard Cell Signaling.

Stomata are pores found on leaves and other aerial organs that enable gas exchange between plants and the atmosphere. Stomata are surrounded by two guard cells that are able to expand, causing the stomatal pore to open, or to become flaccid, closing the pore,^{69,70} in a process requiring a ROS burst. Stomatal closure and opening is highly regulated, as open stomata allow exchange of gases for photosynthesis, but also allow water loss. Stomatal closure is stimulated by either the hormone ABA or environmental signals, which trigger an influx of Ca^{2+} into the guard cells.^{69,70} This Ca^{2+} influx causes an efflux of potassium and anions through K^{+} channels and slow anion channel 1 (SLAC1), respectively.⁷¹ Water then exits the guard cells, and the loss of turgor pressure causes the cells to shrink, closing the pore. The opposite is true for opening of stomata: K^{+} ions move into the guard cell vacuole, causing water influx into the vacuole and expanding the cells around the stomatal pore.⁷²

ABA activates RBOH enzymes, creating a ROS burst that induces stomatal closure^{33,73–76} and reduces water loss. ABA-induced increases in ROS, as observed with chloromethyl-2',7'-dichlorofluorescein diacetate (CM-H₂DCFDA) and peroxy orange 1 (PO1), precede stomatal closure, as shown in Figure 4 for tomato.⁷⁶ The increase in ROS and the resulting changes in stomatal closure are blocked by treatment with DPI, which blocks RBOH enzymes (Figure 4).⁷⁶

The antioxidant glutathione has also been implicated in modulating ROS-induced stomatal closure, as mutations that impair its synthesis have elevated ROS and guard cell closure.^{77,78} In *Arabidopsis*, the RBOH enzymes that regulate stomatal opening are RBOHD and RBOHF.^{73–75} The single mutant *rbohF* shows partial reduction of stomatal closure after ABA treatment, while *rbohD* is comparable to the wild-type.³³ In the presence of DPI⁷⁶ or in the *rbohD/f* double mutant,³³ the ABA-mediated ROS burst and subsequent stomatal closure is impaired. The *rbohD/f* double mutant has impaired Ca^{2+} influx and stomatal closure that are rescued by exogenous H_2O_2 treatment³³ via activation of Ca^{2+} ion channels.⁷⁹ Mutation

of guard cell hydrogen peroxide resistant 1 (GHR1), a receptor kinase-like protein, results in impaired ABA- and H₂O₂-induced stomatal closure, indicating that GHR1 is upstream of RBOH but downstream of ABA.^{80,81} GHR1 has been shown to regulate SLAC1 through CPK3,⁸² highlighting its importance in activating key players in stomatal closure.

In addition to ABA, stomatal closure is also regulated by phosphatidic acid (PA)-induced synthesis of reactive nitrogen species (RNS), including nitric oxide (NO).⁸³ Induction of the phospholipase D α 1 (PLD α 1) by H₂S increased PA levels and elevated ROS compared with untreated plants.⁸⁴ Disruption of PLD α 1 reduced PA, impairing ROS and NO production in response to ABA.⁸⁵ Furthermore, PA binding sites on RBOHD and RBOHF were identified in the 300 amino acid N-terminal extension; disruption of these binding sites resulted in impaired ROS synthesis and reduced ABA-dependent stomatal closure.⁸⁵

RBOH-dependent ROS bursts modulate calcium channels, but Ca²⁺ signals also regulate RBOHF activity through CBL, Ca²⁺ sensor proteins and CIPKs.^{86,87} CIPK26 has been identified as interacting with RBOHF by a yeast two-hybrid screen and confirmed by co-immunoprecipitation.⁸⁶ In HEK293T cells, coexpression of CIPK26 and RBOHF resulted in a decrease in ROS levels, suggesting that CIPK26 directly binds to RBOHF to negatively regulate ROS production.⁸⁶ However, expression of CIPK26, RBOHF, and CBL1 or CBL9 strongly enhances ROS production.⁸⁷ These results reinforce the positive regulation of calcium channels by RBOH-produced ROS bursts.⁸⁷

2.3. Reproduction.

Several RBOHs have been linked to plant sexual reproduction, highlighting a role for ROS in this process. Plant reproduction occurs in flowers when pollen, the male gametophyte, fertilizes the female gametophyte located within the ovule to initiate seed and fruit development.⁸⁸ During pollination, pollen grains are transferred from the anther to the stigma (female reproductive tissue) of the flower, where they germinate and generate a pollen tube that navigates long distances through female reproductive tissues.⁸⁹ Fertilization is completed when the pollen tube reaches the ovule and releases sperm cells to fertilize the female gametes.

A role for ROS has been identified during several stages of pollen development and fertilization,^{90–95} with localized ROS synthesis needed for fertilization, but with stress-induced ROS inhibiting this process.⁹⁶ Multiple RBOH enzymes have been implicated in this process in *Arabidopsis* and other plants, with specific roles defined for RBOHB, –E, –H, and –J (as illustrated in Figure 1 and summarized in Table 1). Transcripts encoding these proteins have been detected in plant reproductive structures, and the phenotypes of these mutants suggest distinct developmental roles for each RBOH enzyme.

2.3.1. Anther and Pollen Development.—RBOHE has expression patterns and functions tied to plant reproduction, with a role in the formation and development of pollen grains (Figures 1 and 2).⁹⁰ *RBOHE* transcripts are detected in mature pollen (Figure 2) and in the tapetum, a layer of cells in the anther that surrounds the developing pollen.⁹⁰ During pollen development, programmed cell death of tapetal cells releases lipids, flavonoids (specialized plant metabolites with antioxidant activity) and proteins that are utilized by

developing pollen.⁹⁷ In *Arabidopsis*, programmed cell death of the tapetum coincides with the stages of flower development when pollen mitotic division occurs. The *rbohe* mutant had reduced ROS accumulation in the anther, as judged by NBT staining, and delayed programmed cell death of the tapetum.⁹⁰ The *rbohe* mutant has impaired pollen viability, with mutant pollen displaying an abnormal outer cell wall layer and a collapsed shape.⁹⁰ *RBOHE* was found to be transcriptionally regulated by the transcription factors MYB80 and ABORTED MICROSPORES,⁹⁰ both of which control tapetum development and pollen formation.^{98–100}

2.3.2. Pollen Tube Growth and Fertilization.—Fully developed, mature pollen grains are released from the anther and transferred to the stigma of the flower, where they germinate to generate a pollen tube that will deliver sperm cells to the ovules for fertilization. The involvement of RBOHs in pollen germination and tube growth has been demonstrated using genetic approaches in tobacco and *Arabidopsis*. Potocky et al.⁹¹ showed that in tobacco a pollen-specific RBOH is necessary for pollen tube growth and production of superoxide. In *Arabidopsis*, RBOHH and RBOHJ, whose transcript abundances are at the highest levels in mature and germinating pollen (Figure 2), are involved in apical growth of the pollen tube (Figure 1) and thus plant sexual reproduction.^{92–94} While single mutants in RBOHH and RBOHJ have phenotypes comparable to the wild-type, double mutants display reduced pollen tube growth and integrity and seed set, consistent with redundant roles of these two pollen tube-expressed RBOHs.

Tip growth of the pollen tube occurs in an oscillatory manner and requires precise coordination between cell expansion and deposition of new cell wall materials through exocytosis.¹⁰¹ Compromised coordination between cell expansion and deposition of new cell wall materials leads to reduced tube growth and cell wall integrity, resulting in premature tube bursting. In *Arabidopsis*, ROS generated by RBOHH and RBOHJ dampen growth rate oscillations.⁹³ In the wild-type, the oscillatory growth dynamics do not change over time, leading to ordered tip growth. In the double mutant, however, two types of oscillation patterns are observed: high-amplitude oscillations with short growth pulses occur initially, and slower and dampened growth oscillations are observed prior to tube bursting.⁹³ The growth behavior in the *rboh/hj* double mutant indicates that ROS reduce maximal growth rates, thereby allowing timely deposition of cell wall materials through exocytosis. Mutations in the Ca²⁺-binding EF hand motifs of RBOHH and RBOHJ resulted in impaired Ca²⁺-activated ROS production.⁹² In addition to Ca²⁺, ROS production by RBOHH and RBOHJ was shown to be activated by phosphorylation, with the two signals operating synergistically.⁹²

A critical feature of the function of RBOHH and RBOHJ in tip growth is targeting of these proteins to the plasma membrane at the tip of pollen tubes. These two RBOHs are localized to distinct but overlapping domains,⁹³ with RBOHH being targeted to the subapical region of the plasma membrane and RBOHJ accumulating in more distal regions of the pollen tube plasma membrane. Similarly to RBOHC, RBOHH and RBOHJ are targeted to the plasma membrane by exocytosis and undergo endosomal recycling.⁹³ Treatment with brefeldin A, which blocks endosomal transport, revealed that RBOHH and RBOHJ are inserted into the

plasma membrane in the region behind the tip and then internalized at the pollen tube tip, with lower internalization rates for RBOHH than for RBOHJ.⁹³

The pollen-specific RBOHH and RBOHJ are regulated by the receptor-like kinases (RLKs) ANXUR1 and ANXUR2, which are colocalized in the same plasma membrane domain at the pollen tube tip.⁹⁴ ANXUR1 and ANXUR2 were proposed to detect changes in cell wall architecture and initiate responses if alterations to the cell wall architecture are too profound. Their loss results in decreased pollen tube integrity, while their overexpression leads to increased deposition of cell wall materials, exocytosis, and arrested tube growth.⁹⁴ By regulating RBOHH and RBOHJ, they coordinate cell wall/pollen tube integrity with the internal pollen tube growth machinery.

ROS also function in later stages of the plant reproductive process. RBOH-derived ROS play a role in pollen tube rupture and sperm cell discharge during fertilization of the female gametophyte.⁹⁵ Pharmacological inhibition of RBOH enzymes with DPI prevented RBOH-dependent ROS production in the synergid cells located within the ovule.⁹⁵ These cells are essential in pollen tube guidance toward the ovule¹⁰² and for the release of sperm cells from the pollen tube, with ROS generated in the synergid cells inducing rupture of pollen tubes to discharge sperm cells to the ovule.⁹⁵

2.3.3. Seed Development.—ROS also function during seed development. Seeds undergo a process called “after-ripening” when they are stored under dry conditions at room temperature. During this process, seeds are released from their dormant state, and seed germination rates are enhanced.¹⁰³ Seed after-ripening is modulated by ROS.¹⁰⁴ *RBOHB* transcripts accumulate in embryos¹⁰⁴ and whole seeds with a greater than 5-fold increase when seeds are imbibed (Figure 2). The RBOHB protein functions to generate superoxide and regulate the germination kinetics of seeds (Figure 1).¹⁰⁴ Fresh seeds of an *Arabidopsis rboh* mutant germinated faster than wild-type seeds, and this difference was abolished in after-ripened seeds.¹⁰⁴ This highlights a role for RBOHB-generated ROS in slowing germination in fresh seeds.

RBOHB is post-transcriptionally regulated to control seed development and hormone-regulated dormancy.¹⁰⁴ *RBOHB* mRNA in embryos has two splicing variants: the smaller *RBOHB-α* variant has intron 1 spliced out, while this intron is retained in the larger *RBOHB-β* splice variant leading to premature termination and a nonfunctional protein.¹⁰⁴ Expression of the two variants was developmentally and hormonally regulated in seeds, with after-ripened seeds containing the functional *RBOHB-α*.¹⁰⁴ Additionally, imbibition, which is treatment of seeds with moisture, to initiate germination increases the abundance of the *RBOHB-α* variant and ROS levels (as judged by NBT staining), while accumulation of the *RBOHB-α* transcript variant was blocked by ABA, a hormone that induces seed dormancy.¹⁰⁴ These results suggest a mechanism by which ROS can be generated specifically in after-ripened seeds and its synthesis can be blocked by ABA to maintain seed dormancy.

2.4. Stress Responses.

Elevations in ROS in response to biotic and abiotic stress are some of the best-characterized redox responses in plants.² ROS increases have been reported in response to bacterial

infection and shown to protect the plant from bacterial invasion.¹⁰⁵ These studies have identified pertinent RBOH enzymes and dissected the molecular mechanisms by which these enzymes are activated. ROS is elevated through RBOH enzymes in response to multiple abiotic stresses, including ozone, salt, and moisture and iron deficiency.^{42,106–109} In some cases these ROS bursts are protective, and in other cases they appear to trigger programmed cell death. ROS, along with Ca²⁺ and electrical signals, orchestrate long-distance signal transduction in plants.¹²² Insight into cell-to-cell signal transmission via ROS was previously reviewed in detail.¹¹⁰ This section summarizes the roles of specific RBOH isoenzymes in biotic and abiotic stress.

2.4.1. Biotic Stress Response.—Plants are constantly exposed to bacterial and fungal pathogens and mount elaborate defense mechanisms against them. RBOHD has been implicated in ROS synthesis as a defense against pathogenic bacteria and fungi, including *Pseudomonas syringae* and *Peronospora parasitica*.¹¹¹ In *rboh*d and *rboh*d/f mutants, pathogen-induced ROS synthesis is reduced in guard cells, and plants are more susceptible to both pathogens.¹⁰⁵ The role of RBOHs in biotic stress response is tied to pathogen-associated molecular pattern (PAMP)-triggered immune responses.^{112,113} There is an enhanced ROS burst during PAMP in the absence of mitogen-activated protein kinase 4 (MAPK4), suggesting that MAPK4 works to downregulate a RBOHD-mediated ROS burst.¹¹⁴ Major PAMP-triggered immune responses are activated by bacterial flagellin protein 22 (flg22) and elongation factor TU (EF-Tu), which activate pattern recognition receptors (PRRs), including flagellin sensitive 2 (FLS2) and elongation factor receptor TU (EFR). Upon recognition of flg22 and EF-Tu, PRRs recruit a coreceptor, brassinosteroid insensitive1 (BRI1)-associated kinase 1 (BAK1), which initiates a pathogen defense signaling cascade¹¹⁵ including phosphorylation of botrytis-induced kinase 1 (BIK1) and AVRPPHB susceptible-like 1 (PBL1). The N-terminal domain of RBOHD directly interacts with BIK1,^{112,113} PBL1,¹¹³ FLS2,¹¹² and EFR in a ligand-independent manner.¹¹³ BIK1 is able to directly transphosphorylate RBOHD.¹¹³ BIK1 is negatively regulated by calcium-dependent protein kinase 28 (CPK28), ensuring an optimal level of BIK1 accumulation.¹¹⁶ FLS2, BAK1, BIK1, CPK28, RBOHD, and cysteine-rich receptor-like kinase 36 (CRK36), which enhances the ROS response,¹¹⁷ work together to create a feedback loop regulating pathogen defense signaling.

The RBOHD protein has multiple phosphorylation sites that modulate the enzyme activity during pathogen response. The N-terminal domain of RBOHD has six serine phosphorylation sites.^{112,113} Three serines are phosphorylated solely by BIK1, while one serine can be phosphorylated by BIK1 or CPKs and two other serines are phosphorylated by CPKs.¹¹³ To determine the impact of BIK1 phosphorylation of RBOHD, serine-to-alanine mutants were made for BIK1-specific phosphorylation sites. While the single mutations did not have a significant effect on the flg22-induced ROS burst, double and triple site mutations diminished this response.¹¹³ Similarly, in the *rboh*d null mutant, the ROS burst is abrogated upon treatment with flg22 or EF-Tu.¹¹⁸ The triple site mutant showed no differences in ABA-induced stomatal closure, indicating that ABA-dependent activation of RBOHD and RBOHF is not mediated by BIK1 phosphorylation.¹¹³

Pathogen response elicits both Ca^{2+} and ROS induction, where ROS is required to maintain prolonged Ca^{2+} signaling.¹¹⁹ HEK293T cells transfected with *FLAG:RBOHD* show that ROS levels increase after ionomycin-induced Ca^{2+} accumulation, suggesting that RBOHD is regulated by Ca^{2+} binding.¹¹⁹ Furthermore, mutations in Ca^{2+} binding sites (EF-hand motifs) in *FLAG:RBOHD*-transfected cells abolished ionomycin-induced ROS production.¹¹⁹ Calcium measurements in wild-type guard cells treated with flg22 show an initial Ca^{2+} peak followed by a second peak or a plateau.¹¹⁸ In *rboh*d, the initial peak is comparable to that for the wild-type while the second peak/plateau is reduced, suggesting that an initial ROS burst is required to maintain Ca^{2+} signaling.¹¹⁸ These results indicate a feedback loop between Ca^{2+} and RBOHD. BIK1 is able to phosphorylate RBOHD upon treatment with Ca^{2+} channel inhibitors,^{112,113} suggesting Ca^{2+} -independent BIK1 phosphorylation within guard cells during pathogen attack.

Ethylene, a gaseous plant hormone, is involved in biotic stress signaling pathways upstream of RBOHD activation. Ethylene treatment increases transcript accumulation of *FLS2*, the gene encoding the flg22 receptor.^{120,121} FLS2 protein accumulation was reduced in two mutants with defects in ethylene signaling: ethylene response 1 (ETR1), an ethylene receptor, and ethylene insensitive 2 (EIN2), an essential signaling protein.¹²⁰ Similarly, flg22-induced ROS detected using luminol and horseradish peroxidase is decreased in *etr1-1* and *ein2-1* mutants relative to the wild-type.¹²⁰

2.4.2. Abiotic Stress Response.—Plants experience constant fluctuations in their environment, such as light, water, salt, nutrients, and temperature, and are subject to abiotic stress when these factors are not at optimal levels. Reactive oxygen species produced in plants in response to abiotic stress can lead to two outcomes: acclimation to stress through ROS signaling events or ROS-induced cell death. Abiotic stress can elicit rapid responses, where local and systemic changes occur in seconds to minutes, or slow responses, where changes take place in tens of minutes to hours and affect the whole plant.¹²² A recent review integrates data suggesting that metabolic and physiological changes induced by abiotic stress are mediated through both rapid and slow responses, where pulses of transcript abundance changes are regulated by ROS and Ca^{2+} waves to prime and/or acclimate plants to fluctuations in their environment.¹²² RBOHD and RBOHF play major roles in ROS production in response to multiple abiotic stresses,¹²³ while RBOHI plays a role in root abiotic responses.¹⁰⁶

An example of detrimental ROS accumulation occurs in plants exposed to ozone, an atmospheric pollutant.¹⁰⁷ Ozone treatment of wild-type plants elevated ROS levels in guard cells, which then spread to neighboring mesophyll cells.¹⁰⁷ ROS levels in mesophyll cells, measured by DCF fluorescence, were decreased in the *rboh*d, *rboh*f, and *rboh*d/*f* mutants, suggesting that RBOHD and RBOHF propagate the ROS signal to nearby cells.¹⁰⁷ Ozone treatment caused similar levels of cell death and necrotic lesions in leaves of wild-type and *rboh*d plants, while *rboh*f single and *rboh*d/*f* double mutants exhibited smaller lesions.¹⁰⁷ Resistance to ozone stress in *rboh*f and *rboh*d/*f* suggest that the ROS accumulation in response to ozone is not an acclamatory signal.¹⁰⁷ In other cases, such as hypoxia and salt stress, ROS is a necessary signal to initiate expression of acclimation genes.^{42,108}

While ROS produced by RBOHD and RBOHF accelerate cell death in response to ozone,¹⁰⁷ these same RBOHs facilitate plant survival in saline-rich⁴² and hypoxic environments.¹⁰⁸ In wild-type plants, salt stress increases ROS levels in roots^{42,108} and hypoxia increases ROS in leaves,⁴² leading to higher levels of Ca^{2+} .^{42,108} In contrast, these stresses result in lower ROS accumulation in *rbohdf* double and single mutants, as measured by DAB or DCF.^{42,108} Leaf damage in response to salt stress (chlorosis and browning) or hypoxia stress (pale leaves with necrotic lesions) was accentuated in *rbohdf* double mutants compared with wild-type or single mutant plants.^{42,108} RBOHI also contributes to hypoxia tolerance.¹²⁴ Although ROS levels were not measured, *rbohi* mutants had paler leaves and lower seedling survival than the wild-type under hypoxic conditions.¹²⁴

RBOHD and RBOHF function in response to multiple stresses, but there are different physiological outcomes for each stress. Salt-stressed *rbohdf* mutants had elevated sodium and reduced potassium contents compared with wild-type plants, suggesting that ROS regulates Na^+ and K^+ homeostasis.⁴² Treatment with DPI to inhibit RBOH activity in the wild-type increased sodium to similar levels as *rbohdf* while exogenous H_2O_2 decreased the sodium content in *rbohdf* mutants, consistent with ROS mediating salt stress response. Hypoxia-induced genes involved in anaerobic central metabolism were downregulated in *rbohi*¹²⁴ and *rbohdf* relative to the wild-type, suggesting that the RBOHD- and RBOHF-dependent ROS burst initiates transcriptional cascades to shift aerobic metabolism to anaerobic respiration.¹⁰⁸

A specific role of RBOHD was found in response to iron deficiency.¹⁰⁹ Iron deficiency increases transcript accumulation of root-specific genes that orchestrate iron uptake to decrease leaf chlorosis.^{125,126} Like other stresses, ROS signaling is central in the iron deficiency response.¹²⁷ Iron-deficient growth conditions increase transcript accumulation of *RBOHD* and *RBOHF* in wild-type plants.¹⁰⁹ Similar to *RBOHC*, *RBOHD* interacts with a member of the ROP family, *ROP6*.¹⁰⁹ *RBOHD* transcript accumulation is dependent on *ROP6*, and leaves of *rop6* and *rbohdf* mutants were more chlorotic compared with wild-type plants and *rbohf* mutants under iron deficiency.¹⁰⁹ Consistent with a decrease in *RBOHD* transcript levels, ROS production is decreased in *rop6*, and the mutant is unable to induce expression of iron uptake genes,¹⁰⁹ consistent with a ROS-dependent transcriptional response cascade.

Ethylene increases in response to stress¹²⁸ and modulates growth and development through transcriptional networks.^{129,130} Responses to multiple stresses, such as heat, high aluminum levels, flooding, high light, and drought, are linked to the transcription factor ethylene response factor 74 (*ERF74*), which was shown to increase expression of a luciferase reporter driven by a *RBOHD* promoter sequence.¹²⁸ The *erf 74* mutant had decreased ROS production, while the *35S::ERF74* overexpression line had increased ROS-dependent DCF fluorescence during response to these stresses. Wild-type plants treated with DPI or in *rbohdf* and *erf 74* mutants had decreased transcript abundance of genes known to respond to the different stresses, suggesting that RBOHD-synthesized ROS is necessary for the activation of these genes, presumably through a ROS-regulated transcription factor.¹²⁸ While *ERF74* regulates *RBOHD* expression, *ERF6* expression is induced downstream of RBOHD-mediated ROS signals.¹³¹ The *rbohdf* mutant has reduced levels of *ERF6* transcripts, and

these transcripts increased after treatment with H_2O_2 .¹³¹ The *erf6* had decreased accumulation of transcripts encoding antioxidant proteins, suggesting that ERF6 may play a role in protecting plants from excess ROS.¹³¹

RBOHI, which is also expressed in seeds and roots (Figure 2), was recently shown to play a role in seed germination and plant survival under drought conditions.¹⁰⁶ Germination of *rboh1* mutant seeds was delayed relative to the wild-type upon exposure to mannitol, which mimics drought stress conditions. When subjected to water withholding, leaves of soil-grown *RBOHI*-overexpressing plants were more resistant to wilting and exhibited better recovery upon rewatering compared with wild-type plants.¹⁰⁶ The *rboh1* mutants and *RBOHI*-overexpressing plants were indistinguishable from the wild-type under control conditions, indicating a stress-specific function for RBOHI.¹⁰⁶ Collectively, the expression patterns of *RBOHs* show specific (in the case of RBOHI) or broad (as seen in RBOHD and RBOHF) functions in response to abiotic stress.

3. MECHANISMS THAT SCAVENGE ROS

3.1. Enzymatic Machineries That Scavenge ROS.

Plant cells utilize many of the same enzymatic pathways to reduce or neutralize ROS that have been described for animal cells. Multiple enzyme classes are involved in direct or indirect scavenging of ROS.^{132–135} Catalase in particular catalyzes the conversion of H_2O_2 to H_2O and O_2 .¹³² In *Arabidopsis*, there are three catalase proteins: CAT1, CAT2, and CAT3.^{136,137} Genetic studies have shown that deletion of *CAT2* reduces catalase activity by 90% in *Arabidopsis* leaves, while there is only a slight reduction in catalase activity with the deletion of *CAT1* or *CAT3*.¹³² *CAT2* and *CAT3* transcript abundances in seedlings are regulated by the circadian clock, with *CAT2* transcript abundance peaking in the morning¹³⁸ and *CAT3* in the evening.¹³⁹ Catalase enzymes are highly involved in the stress response and also have developmental implications, which have been previously reviewed.^{132,140}

Localization of catalase proteins was determined using GUS reporters and qRT-PCR, with *CAT1* localizing mainly in the cotyledon, rosette leaves, roots, and the siliques. In contrast, *CAT2* is expressed in germinating seeds, cotyledons, flowers, bolts, and siliques,¹⁴¹ while *CAT3* is found in roots and young leaves.¹⁴¹ Subcellular localization of catalase enzymes using biochemical fractionation has demonstrated that all three catalase enzymes are in the mitochondria¹⁴² and in the peroxisome, consistent with the presence of a peroxisomal targeting sequences.¹⁴³ Although one study found catalase activity in chloroplast fractions,¹⁴⁴ other authors have suggested that there may have been peroxisomal contamination of the chloroplast fraction.¹⁴³

Peroxidases are enzymes that function to reduce peroxide and have been shown to modulate ROS regulation and redox signaling in *Arabidopsis*.^{140,145} In many of these peroxidases, the catalytic activity depends on a peroxidatic cysteine and a resolving cysteine.¹⁴⁶ These enzymes are able to detect the levels of ROS on the basis of their oxidation status.¹³⁵ There are many families of peroxidases, including peroxiredoxins, ascorbate peroxidase, and glutathione peroxidase, each of which functions as an enzymatic antioxidant. After reduction of H_2O_2 , each peroxidase is reduced back to the active form by a pathway of enzymes or

small molecules. For example, glutathione peroxidases are oxidized after reducing H₂O₂. They are reduced by glutathione (GSH), which is thereby converted to its oxidized form, glutathione disulfide (GSSG). GSSG is then reduced by glutathione reductase (GR), with reduced GR regenerated by electrons from NADPH. In *Arabidopsis*, glutathione peroxidases¹⁴⁷ and GSH¹⁴⁸ have been reviewed with respect to their antioxidant ability and role as redox sensors.^{149,150}

Another peroxidase in *Arabidopsis*, ascorbate peroxidase, utilizes L-ascorbic acid to reduce H₂O₂, converting the ascorbate to monodehydroascorbate, which can either be reduced by monodehydroascorbate reductase (MDAR) back to ascorbate or become dihydroascorbate.¹⁵¹ Dihydroascorbate is recycled back to L-ascorbic acid by dehydroascorbate reductase (DHAR) using GSH.¹⁵¹ The role of ascorbate peroxidase in maintaining ROS homeostasis in the chloroplast^{152,153} and in response to abiotic stress¹⁵⁴ has recently been reviewed.

Thioredoxin and glutaredoxin are found at multiple subcellular localizations and are reduced by NADPH and glutathione, respectively. Thioredoxin has been implicated in the ability to reduce peroxiredoxins. Peroxiredoxins are able to become sequentially oxidized, resulting in a cysteine sulfinic acid (-SOOH) that can be reduced to a cysteine sulfenic acid (-SOH) by sulfiredoxin. The localization of peroxiredoxins as well as their catalytic function and physiological significance in plants was recently reviewed in detail,^{133,146} as was the function of both glutaredoxin and thioredoxins.^{134,149,150,155,156} The detailed mechanisms of action and regulation of these pathways are beyond the scope of this review and have been previously reviewed.^{133,134,146,149,150,155,156}

3.2. Plant-Specific Non-enzymatic ROS Scavengers.

To assume their signaling roles, ROS must reach certain threshold levels to oxidize their downstream target proteins. These protein oxidation events will then propagate the ROS signal to initiate the specific growth and developmental programs and stress responses described above.^{157,158} However, to prevent ROS from reaching concentrations that cause oxidative damage, ROS levels must be tightly regulated by antioxidants.^{159–161} Below we highlight plant specialized metabolites with ROS scavenging activity and the diversity of their known roles in plant growth, development, and stress responses. These metabolites are often beneficial to human health, making their biosynthetic pathways targets of metabolic engineering for biofortification. Two well-known examples are golden rice, which has increased carotenoid levels,¹⁶² and purple tomatoes, with high abundance of anthocyanins.¹⁶³

3.2.1. Carotenoids.—Carotenoids are lipid-soluble pigment molecules of the terpenoid family that produce orange and red colors in plants.^{164,165} Carotenoids in the chloroplast increase photosynthetic efficiency by acting as accessory pigments, protein stabilizers, and antioxidants.^{164–166} Carotenoids are essential, as homozygous mutations in early pathway enzymes cause seedling lethality.^{167,168} There are two classes of carotenoids: carotenes (α -carotene and β -carotene) are pure hydrocarbon molecules, as opposed to their oxygen-containing derivatives, xanthophylls (lutein, zeaxanthin, antheraxanthin, violaxanthin, and

neoxanthin).^{164,165} The biosynthetic pathway of carotenoids is well-established and was recently reviewed elsewhere.^{164,165}

Carotenoids accumulate in the light-harvesting centers of Photosystems I and II (PSI and PSII), which are two complexes that mediate photosynthesis, where they scavenge ROS byproducts of photosynthesis.¹⁶⁹ High light intensity increases production of singlet oxygen ($^1\text{O}_2$),¹⁶⁹ a short-lived, highly damaging ROS responsible for the majority of photooxidative damage.¹⁷⁰ High light intensity can also excite chlorophyll to the triplet state (^3Chl), which interacts with O_2 to form $^1\text{O}_2$.¹⁶⁹ Xanthophylls directly quench 95% of ^3Chl , leaving a small percentage available for $^1\text{O}_2$ formation.^{169,171} A small fraction of xanthophylls are found in the lipid bilayer, where they also reduce lipid peroxidation by $^1\text{O}_2$.^{171–174}

Non-photochemical quenching (NPQ), a process that allows excess light energy absorbed by chlorophyll to be dissipated as heat, is crucial in preventing $^1\text{O}_2$ formation.^{169,175,176} In wild-type plants, NPQ depends on zeaxanthin, which is produced in response to excess light.^{173,177–179} NPQ is repressed in the *non-photochemical quenching 1* (*npq1*) mutant, which does not synthesize zeaxanthin because of a defect in violaxanthin de-epoxidase.^{173,178} As a consequence, $^1\text{O}_2$ levels are higher in the *npq1* mutant, as measured by the fluorescent probe Singlet Oxygen Sensor Green (SOSG).¹⁸⁰ Constitutive production of zeaxanthin in the zeaxanthin epoxidase-defective *npq2* mutant leads to faster NPQ induction compared with wild-type plants and *npq1*.¹⁷⁸ This is presumably due to higher antioxidant activity from increased zeaxanthin levels, although $^1\text{O}_2$ levels have not been reported for *npq2*.

Normal xanthophyll composition in the light harvesting centers is altered in mutants with defective carotenoid enzymes.¹⁸¹ Mutation of lycopene ϵ -cyclase in *lut2* mutants abolishes lutein production and increases flux into the β -carotene branch.^{182–184} As a consequence, lutein is replaced by violaxanthin, leading to destabilization of the light harvesting complex and increased ^3Chl in the *lut2-1* mutant.^{171,183,184} Compared with wild-type plants, decreased NPQ or unstable light-harvesting complexes have been reported for the majority of carotenoid mutants,^{181,185} and increased $^1\text{O}_2$ production has been reported in *lut2*, *lut5*, and *chy1/chy2* mutants, which have defects in the xanthophyll-producing enzymes cytochrome P450 enzyme CYP973A/LUT5 and carotenoid hydroxylase 1 and 2 (CHY1 and CHY2).^{186,187} Higher-order mutants tend to have more compromised NPQ and photoprotection, with increased $^1\text{O}_2$ levels reported in more recent articles.^{187–190} A quadruple mutant that does not produce any xanthophyll compounds, *no xanthophyll* (*nox*, defective in CHY1, CHY2, LUT1, and LUT5) has extreme sensitivity to light and decreased photosynthetic efficiency.¹⁹¹ Although *nox* does synthesize α - and β -carotenes, $^1\text{O}_2$ levels were dramatically higher in the mutant compared with the wild-type, suggesting that xanthophylls play a major role in ROS scavenging.¹⁹¹ Moreover, *nox* mutants exhibited extreme sensitivity to light, as photoinhibition studies and $^1\text{O}_2$ measurements were performed at a light intensity of $200 \mu\text{mol photons s}^{-1} \text{ m}^{-2}$, a level 5-fold lower than commonly used for “excess light”.¹⁹¹ As a consequence, *nox* mutants could survive only under very low light ($20 \mu\text{mol photons s}^{-1} \text{ m}^{-2}$).¹⁹¹ Collectively, these studies suggest that optimal $^1\text{O}_2$ scavenging requires a mixture of xanthophyll compounds.

Binding of β -carotene affects the stability of the photosystem core complex.¹⁹² The *suppressor of zeaxanthin-less 1 (szl1)* mutant has decreased lycopene β -cyclase (LycB) activity, increasing flux into the α -carotene branch.^{186,189} Replacement of β -carotene with α -carotene in PSI increases photosensitivity and $^1\text{O}_2$ production in *szl1* compared with the wild-type.¹⁸⁶ Additionally, an oxidation product of β -carotene, β -cyclocitral (β -CC), is involved in acclimation to excess light. Changes in transcription levels in response to β -CC are specific to $^1\text{O}_2$ stress and participate in retrograde signaling to promote acclimation to excess light.^{193,194} Lipid peroxidation and decreased photosynthetic efficiency caused by high light stress are mitigated in plants exposed to β -CC prior to high light exposure compared with untreated plants.¹⁹⁴ Collectively, carotenoids assume a number of distinct and important roles to mitigate high light stress by direct scavenging of $^1\text{O}_2$ and promotion of acclimation.

3.2.2. Ascorbate.—L-Ascorbic acid (vitamin C) is a water-soluble, plant-derived antioxidant that is essential to humans and serves diverse functions in plants, especially during abiotic and biotic stresses. Besides acting as a cofactor in plant enzymatic reactions, it can donate electrons and thereby act as a ROS scavenger.^{195,196} However, the iron-reducing activity of L-ascorbic acid can lead to the formation of Fe^{2+} from Fe^{3+} and thus to increased $^{\bullet}\text{OH}$ production by the Fenton reaction.¹⁹⁷ In humans, dietary vitamin C is required for collagen and carnitine biosynthesis,^{198,199} and its deficiency in humans, better known as scurvy, results in weakness, joint pain, gum bleeding, and skin hemorrhages.

The L-ascorbic acid biosynthetic pathway in plants converts L-galactose to L-ascorbic acid. The final reaction occurs on the inner mitochondrial membrane. From there, L-ascorbic acid is transported to different subcellular compartments, including the cell wall, to assume its different functions.²⁰⁰ L-Ascorbic acid is present at high concentrations in the different subcellular compartments (mM range in the chloroplast and cytosol)^{201,202} and thus serves as a major antioxidant in plants.

L-Ascorbic acid reacts with free radicals and Fe^{3+} to produce monodehydroascorbate or can be converted to monodehydroascorbate by ascorbate peroxidase, which concomitantly catalyzes the conversion of H_2O_2 to water.²⁰³ Monodehydroascorbate can either be recycled to L-ascorbic acid through MDAR^{204,205} or be non-enzymatically converted to dehydroascorbate. Dehydroascorbate is converted to L-ascorbic acid by DHAR, which uses glutathione a reducing agent.²⁰⁵

Several mutants with altered L-ascorbic acid levels, termed *vtc* mutants, were isolated in *Arabidopsis*, enabling characterization of L-ascorbic acid functions in plants.^{206,207} A 5-fold decrease in L-ascorbic acid levels in the *vtc2* mutant, which has a defect in the synthesis of the L-ascorbic acid precursor l-galactose, did not result in reduced plant growth.²⁰⁶ This indicates that in the absence of stress, only minimal levels of l-ascorbic acid are necessary to support plant growth. However, the *vtc* mutants, as well as mutants defective in ascorbate peroxidase or ascorbate recycling, are more sensitive to environmental stresses than the wild-type, in particular to photooxidative stress.^{160,208} During high light stress, removal of H_2O_2 by ascorbate peroxidases and 2-cysteine peroxidases reduced superoxide levels, as shown by NBT staining of leaves, to protect the photosynthetic machinery.²⁰⁹ In heat-

stressed leaves with an inactivated oxygen-evolving complex, L-ascorbic acid acted as an electron donor to PSII, thereby sustaining electron transport.²¹⁰ The roles of L-ascorbic acid against biotic stresses are more complex, as protective effects from certain pathogens were observed at reduced L-ascorbic acid levels,^{211,212} while in other cases elevated ascorbate increased pathogen resistance.²¹³

3.2.3. Tocochromanols.—Tocochromanols, such as vitamin E, include tocopherols and tocotrienols, which are lipid-soluble compounds produced mainly by photosynthetic organisms. They are synthesized by condensation of tyrosine-derived homogentisate and a polyprenyl side chain; the side chain is saturated in tocopherols and contains three unsaturated double bonds in tocotrienols.²¹⁴ The homogentisate moiety forms a chromanol ring and has antioxidant capacities,^{215,216} while the polyprenyl side chain anchors tocochromanols into membranes. Tocochromanols accumulate in photosynthetic membranes and plastoglobules inside the chloroplast.²¹⁷ Because of their presence in photosynthetic membranes, the main antioxidant activity of tocochromanols is to scavenge lipid peroxyl radicals generated by lipid peroxidation²¹⁸ through $^1\text{O}_2$. Additionally, tocochromanols have excellent *in vitro* $^1\text{O}_2$ quenching and scavenging capacities.^{219,220} These compounds are also important in the human diet, as tocochromanols are essential for reproduction.²²¹

Arabidopsis mutants with altered tocochromanol biosynthesis, termed *vte* mutants, have provided insight into the roles of tocochromanols. The *vte1* mutant does not exhibit any differences in growth, photosynthetic quantum yield, or chlorophyll content under normal light conditions compared to the wild-type, and only minimal differences were observed under high light stress.²²² Similar results were observed in *Synechocystis*, a unicellular photosynthetic cyanobacteria that synthesizes tocopherols.²²³ In a tocopherol-deficient *Synechocystis* strain with a mutation in the homogentisate phytyltransferase, growth under high light conditions did not change from the wild-type to the mutant.²²³ Treatments that induce superoxide or $^1\text{O}_2$ production, combined with high light stress, slowed the growth of the wild-type and the tocopherol-deficient *Synechocystis* mutant,²²³ suggesting that tocopherols are not essential in scavenging these ROS. However, treatment with polyunsaturated fatty acids, which generate lipid peroxyl radicals by auto-oxidation in the presence of oxygen, slowed the growth of the tocopherol-deficient *Synechocystis* mutant,²²³ indicating that tocopherols protect cells from lipid peroxidation. A similar function for tocopherols in protecting lipids from oxidation during seed germination and early seedling development was uncovered in *Arabidopsis thaliana* using the tocopherol-deficient mutants *vte1* and *vte2*.^{224,225}

Beyond their role in protecting plant cells against lipid peroxidation, tocopherols have also been shown to scavenge $^1\text{O}_2$ formed by PSII.²²⁶ Singlet oxygen formation can induce degradation of the D1 protein, a protein that forms the reaction center of PSII. The role of tocopherols as $^1\text{O}_2$ scavengers was shown to be particularly important when the turnover rate of the D1 protein is high.²²⁶

3.2.4. Flavonoids.—Flavonoids are a large class of plant metabolites that act as antioxidants to lower ROS levels in plants. Flavonoid antioxidants have been studied to have positive effects on human health including in cancer and age-related diseases.^{227,228}

Flavonoids regulate a wide array of plant developmental programs and stress responses, prompting intense study of their biosynthesis and regulation.^{229–232} The flavonoid biosynthetic pathway, which is shown in Figure 5, begins with the condensation of *p*-coumaroyl-CoA with three malonyl-CoAs, resulting in synthesis of multiple flavonoid subgroups, including flavonols (described in detail in the next section), flavones, flavan-3-ols, flavanones, anthocyanins, and isoflavonoids.^{228,233} Specific pathway details have been reviewed elsewhere.^{229,232}

The first committed step of flavonoid biosynthesis is catalyzed by chalcone synthase (CHS) to produce naringenin chalcone. Naringenin, isomerized from naringenin chalcone, is a precursor for the flavone and isoflavonoid branches. Flavonols are produced from dihydrokaempferol, an intermediate downstream of naringenin. The flavan-3-ol branch is produced from leucoanthocyanidins downstream of dihydrokaempferol, as is the first committed step in the production of anthocyanins, the final subclass produced in the flavonoid pathway. The flavonoid compounds can be further modified by glycosylation of hydroxyl groups with different carbohydrates, which further increases the chemodiversity of flavonoid compounds.

Flavonoid composition is species-specific, with some plant species being able to produce all of the flavonoid subclasses while in others only some subclasses accumulate.²³⁴ For instance, flavones²³⁵ such as apigenin, luteolin, baicalein, and chrysin are found in herbs such as parsley and thyme, while the flavanones eriodictyol, hesperetin, and naringenin accumulate to high levels in citrus fruits.²³⁶ Anthocyanins are red, blue, and purple pigments found in most plant species, with high concentrations found in many berries, fruits, and flowers.^{237,238} Flavan-3-ols, which include catechins and epicatechins, are most commonly associated with tea leaves,²²⁸ but are also found in berries, apples, and cacao. Finally, isoflavonoids are most commonly found in soybeans and other legumes,²³⁴ although they are present in other plant species. The differences in concentrations and types of flavonoids found across the plant kingdom are consistent with species-specific regulation and modulation of the biochemical pathway.

What is particularly striking is the difference in the antioxidant activities of these subclasses of flavonoids, suggesting that some flavonoid subclasses with higher antioxidant capacity have a more significant role in redox-regulated processes. Flavonoids share a similar chemical backbone with two benzene rings linked by a heterocyclic pyran ring, enabling them to donate one or more electrons to less stable molecules such as ROS.²³⁹ Many of the flavonoid backbones are decorated by hydroxylation, methylation, glycosylation, and/or acylation, which alters their ability to scavenge ROS, thus leading to a range of antioxidant activities *in vivo* that reflect the immense flavonoid chemodiversity. Flavan-3-ols, flavonols, and anthocyanins are able to act as potent antioxidants for lipid peroxidation *in vitro*.^{240,241} Studies investigating the flavonoid scavenging ability of H₂O₂ and ¹O₂ found that anthocyanins and flavonols are the most effective in reducing these ROS,²⁴¹ while flavones are the least effective, with no antioxidant capacity for H₂O₂ and very little capacity for ¹O₂. Other classes show selectivity: flavanones show no antioxidant capacity for ¹O₂ and very little capacity for H₂O₂, whereas flavan-3-ols have high scavenging ability for H₂O₂ and

very low ability to scavenge $^1\text{O}_2$.²⁴¹ Because of their high antioxidant activity, flavonols and anthocyanins are some of the most interesting and best-studied members of this family.

Flavonoid accumulation is induced by a variety of ROS-inducing abiotic stresses, hormones, and developmental signals.^{229,231,232,242,243} Expression of genes encoding flavonoid metabolic enzymes is highly regulated by transcription factors belonging to the R2R3-MYB, basic helix–loop–helix (bHLH), and WD40-type families of transcription factors. These factors associate in a ternary complex (termed the MYB–bHLH–WD40 complex)²⁴⁴ and mediate the changes in flavonoid synthesis in response to environmental factors^{229,231} and hormones.^{242,243}

Anthocyanins, the most downstream products of the flavonoid pathway, are the best-studied flavonoid subclass because their pigmentation allows easy visualization of their stress- or developmentally-induced synthesis. The transcriptional machinery that drives stress-induced anthocyanin synthesis includes a number of MYB proteins, including MYB75, also called production of anthocyanin pigment 1 (PAP1), PAP2,²⁴⁵ MYB113, and MYB114.²⁴⁶ A variety of abiotic stresses, including high light, drought, high sucrose, and oxidative stress, have been shown to cause increased anthocyanin synthesis via PAP1.^{247–249} An *Arabidopsis* line that overaccumulates *PAP1* (*pap1-D*) has constitutively high anthocyanin accumulation driven by increased transcription of dihydroflavonol-4-reductase (DFR), the metabolite branch point channeling dihydroflavonols into anthocyanins.^{250,251} The presence of elevated anthocyanins offers protection from the damage caused by these stresses.^{247,252} To demonstrate the protective role of flavonoids, flavonoid-deficient *transparent testa 4* (*tt4*), which has a defect in CHS, was shown to be more sensitive to drought and oxidative stress and to lose the protective effect of *pap1-D* in a double mutant.²⁵² Consistent with flavonoids acting as antioxidants to reduce ROS overaccumulation, *pap1-D* has reduced H_2O_2 levels in leaf tissue during drought stress compared with the wild-type, while *tt4* has elevated levels, as detected by DAB.²⁵² H_2O_2 treatment was shown to induce transcription levels of zinc finger of *Arabidopsis thaliana* 6 (ZAT6), which is a positive regulator of flavonoid synthesis, suggesting that ROS can also stimulate antioxidant production.²⁵¹ Knockdown of ZAT6 showed decreased anthocyanin accumulation, while ZAT6 overexpression resulted in increased anthocyanin levels.²⁵¹ However, these studies did not look at changes in endogenous ROS levels after anthocyanin accumulation/reduction in these mutant lines.

Anthocyanin accumulation is a typical phenotype associated with high light stress. One regulatory process in facilitating the response to high light is to deactivate negative regulators of anthocyanin biosynthesis. The transcription factor TCP15 negatively regulates anthocyanin biosynthesis under normal conditions.^{253,254} However, during prolonged exposure to high light, TCP15 is rendered inactive.²⁵³ This deactivation is facilitated by increased ROS levels, as oxidation of a cysteine residue in the protein prevents binding of TCP15 to DNA.²⁵³ Once TCP15 is inactivated, anthocyanin accumulation can occur, which may be a protective mechanism to prevent damage from high light stress.²⁵³ In contrast, two NAC transcription factors, ANAC032 and NAC019, are induced in response to high light/increased sucrose²⁵⁴ and drought,²⁵⁵ respectively. These transcription factors have also been shown to be involved in repression of anthocyanin biosynthesis. Treatments of ANAC032 overexpression lines with the catalase inhibitor 3-amino-1,2,4-triazole (3-AT), with the goal

of increasing ROS accumulation, reduced the anthocyanin accumulation compared with 3-AT-treated wild-type and ANAC032 repressor plants. These results suggest that ANAC032 represses anthocyanin synthesis under high light and in the presence of elevated ROS levels.²⁵⁴

Despite visible changes in anthocyanin levels upon environmental stresses or during development, changes in anthocyanins could reflect altered flux through the entire pathway and affect levels of metabolites belonging to the flavonoid subclasses outlined above. In particular, the dihydroflavonols are precursors of both flavonols (through the action of flavonol synthase (FLS)) and anthocyanins (through the action of DFR) (Figure 5), so these two steps represent important branch points. Flavonol compounds synthesized from dihydrokaempferol include kaempferol, quercetin, isorhamnetin, and myricetin, which have distinct antioxidant activities.^{256–260} In *Arabidopsis*, 32 different flavonol derivatives with distinct glycosylation patterns have been identified by liquid chromatography-mass spectrometry (LC-MS),²⁶¹ but the antioxidant capacities of only a few of these compounds have been measured. *In vitro* data suggest that quercetin may be the best ROS scavenger.²⁴⁰ Quercetin 3-*O*-Glc has a lower IC₅₀ value for superoxide ions than kaempferol 3-*O*-Glc, indicating that quercetin is a better superoxide scavenger *in vitro*. Similarly, quercetin is able to scavenge ¹O₂ better than kaempferol, although myricetin and isorhamnetin have been reported to be better at scavenging ¹O₂ than quercetin.²⁴¹ Among the flavonols, quercetin has the highest antioxidant capacity for H₂O₂, followed by kaempferol, myricetin, and isorhamnetin.²⁴¹ These insights into regulation of flavonol synthesis, flavonol localization, and their range of antioxidant activity set the stage for understanding the specific functions of these molecules.

4. GENETIC APPROACHES THAT UNCOVERED THE ROLE OF FLAVONOLS AND ROS IN PLANT DEVELOPMENT AND STRESS RESPONSES

Multiple stress responses increase anthocyanin levels,²⁵⁰ which are readily detected because of changes in the concentrations of these pigments, but few studies have investigated how this impacts accumulation of earlier pathway products like flavonols. Additionally, in many publications the effect of changes in anthocyanin accumulation on ROS homeostasis or flavonol levels has not been examined. Recent experimental advances have made it possible to study flavonols directly. LC-MS can be used to quantify changes in accumulation of specific flavonols, while the identification of diphenylboric acid 2-aminoethyl ester (DPBA), which binds flavonols and becomes fluorescent, allows their visualization *in planta*.²⁶³ Both methods have revealed developmental, hormonal, and stress-induced synthesis of these compounds. Finally, the power of genetic systems like those found in *Arabidopsis* is that we can begin to elucidate the specific functions of individual flavonoid molecules through examination of genetic mutants or tissues in which these flavonols and anthocyanins are uncoupled.

Flavonols are particularly interesting because of their cellular localization. In plant cells, ROS accumulates in the cytoplasm, nucleus, and cell wall, as determined using confocal

imaging. This localization overlaps with flavonol accumulation patterns and subcellular localization of enzymes involved in flavonol synthesis.^{96,242,264,265} Since anthocyanins localize to the vacuole,^{266,267} flavonols are more likely to be the active compounds in modulating cytoplasmic and nuclear ROS.^{76,242,264,268} In support of the hypothesis that flavonols act as an important class of ROS scavengers, recent studies have shown that flavonols alter growth and development by locally regulating ROS levels. Using well-established genetic systems, this section examines the function of flavonols as ROS scavengers in a variety of stress and developmental contexts.

4.1. Roots.

Root tissues synthesize and accumulate flavonols, but do not synthesize later flavonoid pathway intermediates such as anthocyanins because of the absence of transcripts encoding DFR, a branch-point enzyme controlling the conversion of dihydroflavonol precursors into anthocyanins.²⁴² The role of flavonols in modulating root architecture was initially tied to their ability to negatively regulate transport of the hormone auxin. Mutants that do not make flavonols have elevated auxin transport compared with the wild-type,^{269–272} and these studies revealed that quercetin derivatives are the active flavonols that negatively regulate auxin transport in roots.²⁴² Additionally, auxin treatment of wild-type plants increased accumulation of transcripts encoding flavonoid transcription factors and enzymes of the early flavonol biosynthetic pathway, thereby increasing flavonol biosynthesis, perhaps as part of a negative feedback loop.²⁴² Recent reports have uncovered a role of flavonols in modulating root architecture via antioxidant activity, as summarized below. Additionally, several lines of evidence suggest that auxin transport is regulated by the levels of ROS and RNS,²⁷³ suggesting that the regulation of auxin transport by flavonols might be linked to flavonol modulation of ROS status.

ROS gradients define the root meristematic and elongation zones, which are areas of cell division and elongation, respectively.²⁷⁴ The ROS gradient is characterized by elevated superoxide accumulation in the meristem, which decreases toward the elongation zone, while H₂O₂ accumulates in the elongation zone and is reduced in the meristem.²⁷⁴ The spatial balance of superoxide and H₂O₂ is maintained by the transcription factor UPBEAT1 (UBP1), which represses accumulation of transcripts encoding peroxidases in the elongation zone.²⁷⁴ The *tt4* mutant had increased superoxide levels in the meristem compared with wild-type roots, as measured by NBT staining, which was suggested to override the superoxide/H₂O₂ gradients in the mutant.²⁷⁵ Treatment of wild-type roots with H₂O₂ induced flavonol accumulation, which is hypothesized to reduce meristem size.²⁷⁵

Light induces accumulation of anthocyanins as described above and also increases flavonol accumulation in roots through rapid induction of transcripts encoding enzymes of the early flavonol biosynthetic pathway.^{233,276} Visualization of flavonols using DPBA and flavonol metabolite profiling by LC-MS show that in plants where shoots are exposed to light and roots are either shielded (dark-grown) or exposed to light (light-grown), flavonol levels are higher in light-grown roots.^{275,277} Wild-type roots exposed to light have reduced cell number in the meristem and shorter primary roots compared with roots shielded from light.²⁷⁵ In the *tt4* mutant, the effect of light on root growth is lost, with light-grown *tt4* seedlings

resembling the dark-grown wild-type.²⁷⁵ The longer root length and increased cell number in *tt4* are reversed by application of the flavonol quercetin.²⁷⁵ In the wild-type, exposure of the roots to light leads to lower superoxide levels compared with dark-grown roots, which may be due to increased flavonol synthesis.²⁷⁵ Consistent with this finding, superoxide levels in the meristem of *tt4* are comparable to those of dark-grown wild-type roots, suggesting that light-dependent flavonol antioxidant activity regulates root growth.²⁷⁵ The authors suggest that regulation of flavonol synthesis may occur through cytokinin- and UPB1-H₂O₂-mediated pathways, as *UPB1*-overexpressing plants had increased flavonol levels and H₂O₂.²⁷⁵ We note, however, that the meristem cell number in *UPB1*-overexpressing seedlings was not quantified in response to light or darkness and that overexpression of *UPB1* was previously reported to reduce meristem cell number and primary root length compared with the wild-type.^{274,275} It would be interesting to measure meristem cell number and ROS in crosses between *tt4* and *UPB1*.

The relationship between flavonols, ROS, and root hair development has been explored in tomato (*Solanum lycopersicum*).²⁷² Roots of the *anthocyanin reduced (are)* mutant, which contains a point mutation in flavonoid-3-hydroxylase (F3H),^{259,260} exhibit reduced flavonol accumulation as measured by LC-MS and increased ROS levels as measured by DCF fluorescence.²⁷² ROS accumulates in epidermal cells and root hairs, with higher ROS levels in the *are* mutant linked to greater root hair density compared with the wild-type (Figure 6).²⁷² Treatment with a general antioxidant, L-ascorbic acid, reduced ROS levels and root hair number in the *are* mutant, while H₂O₂ treatment of wild-type plants increased ROS and root hair number.²⁷² Genetic complementation of the *are* mutant restored flavonol synthesis, reduced ROS accumulation, and returned root development to wild-type patterns, consistent with the role of flavonols as antioxidants to modulate root development.²⁷²

Bending of roots away from light (negative phototropism) may also be modulated by localized synthesis of flavonol antioxidants.²⁷⁵ Wild-type roots grown in the dark and then exposed to directional light show increased flavonol accumulation on the light-exposed side, which also has a higher elongation rate to allow bending away from the light.²⁷⁵ Accumulation of GUS activity driven by the *CHS* promoter in transgenic lines and flavonol products overlapped with the convex (outer) side of bent roots.²⁷⁵ Localized flavonol production is suggested to fine-tune bending, as a reduced bending angle was observed in wild-type roots treated uniformly with quercetin and in *UPB1*-overexpressing lines.²⁷⁵ Additionally, the absence of flavonols in the *tt4* mutant also resulted in reduced bending angle.²⁷⁵ Consistent with this result is the finding that ROS are an important signal in asymmetric growth during root gravitropism.²⁷⁸ Despite these lines of evidence, we note that ROS localization or quantification was not reported in bent roots, which is an important question for future studies on root tropism.

4.2. Guard Cells.

The role of an ABA-dependent ROS burst in guard cells to initiate stomatal closure is clearly established.^{73,279,280} Elevated ABA triggers H₂O₂ production through activation of RBOH enzymes as described above.^{33,76,281} ROS levels then must reach a certain threshold to oxidize their downstream targets, which may include ion channels that initiate stomatal

closure.^{79,282} The localized accumulation of flavonols in guard cells suggested the testable hypothesis that these molecules may act to modulate this ROS to fine-tune guard cell closure and prevent oxidative damage.

Flavonol accumulation in guard cells suggests a role in regulation of ROS homeostasis in a diversity of species, including fava bean (*Vicia faba*), pea (*Pisum sativa*), onion (*Allium cepa*), tomato (*Solanum lycopersicum*), and *Arabidopsis*.^{76,264,283–286} Transcripts of flavonoid biosynthetic enzymes CHS and FLS were shown to accumulate in *Arabidopsis* guard cells using a GUS promoter and GFP construct, respectively.^{264,271} The accumulation of flavonols in guard cells but not in the surrounding pavement cells of *Arabidopsis* and tomato has been shown using DPBA visualized through confocal microscopy.^{76,264,287} DPBA signal accumulates in the nucleus, consistent with prior reports,^{76,242,264,268} where it may regulate redox-sensitive transcription factors and/or prevent ROS-induced DNA damage. The *Arabidopsis tt4–2* mutant did not exhibit a DPBA signal and showed increased ROS in guard cells, as visualized by DCF fluorescence.²⁶⁴ Consistent with enhanced ROS signaling in *tt4–2*, ABA-induced stomatal closure was greater in this mutant.²⁶⁴ Similarly in tomato, flavonoid production is reduced in the *are* mutant, resulting in decreased DPBA signal in guard cells (Figure 7). In contrast, the tomato *anthocyanin without (aw)* mutant, which has elevated levels of flavonols, has enhanced DPBA signal in guard cells.⁷⁶ These results are consistent with flavonol synthesis and accumulation in guard cells and their function to reduce ROS in these cells.²⁸⁸

The relationship between ROS, flavonols, and ABA-dependent stomatal closure was studied in greater detail in tomato.⁷⁶ Levels of total ROS, as detected by DCF fluorescence, and H₂O₂, as detected by PO1, were shown to increase in response to ABA treatment preceding stomatal closure.⁷⁶ The levels of DCF and PO1 were shown to be inversely proportional to the flavonol levels, with increased DCF and PO1 signals in the *are* mutant and decreased DCF signal in the *aw* mutant compared with the wild-type (Figure 7).⁷⁶ The ABA-induced DCF signal increases are dependent on RBOH enzymes, as they are not detected after treatment with DPI (Figure 4). Time course experiments showed that the rates of ABA-dependent stomatal closure were proportional to the rate of change in H₂O₂ in these flavonol mutants (Figure 7). Additionally, this study also examined the process of light-dependent stomatal opening. In wild-type plants, ROS decreases substantially after exposure to light. Elevated flavonols in the *aw* mutant facilitated this decrease, and reduced flavonols in the *are* mutant impaired this process.⁷⁶ Finally, consistent with a direct relationship between flavonol-modulated ROS and stomatal aperture, the water loss from *are* mutant leaves was greater compared with the wild-type, while the *aw* mutant had less water loss.⁷⁶

The accumulation of flavonols in guard cells to control ROS-induced stomatal closure is regulated by the hormone ethylene. Increasing ethylene levels by treating wild-type plants with ethylene gas, or comparing untreated plants to the *ethylene overproducing 1 (eto1)* mutant, which has endogenous overproduction of ethylene,²⁸⁹ increased flavonol accumulation in guard cells.^{76,264} Coincident with elevated flavonols, there was a substantial reduction in ROS accumulation, as shown with DCF in guard cells of *eto1*²⁶⁴ or ethylene-treated wild-type *Arabidopsis* or tomato plants.^{76,264} Consequently, ethylene-elevated flavonols and decreased ROS resulted in decreased rates of ABA-induced stomatal closure

compared with untreated plants.^{76,264} Similarly, induction of flavonol accumulation in *Arabidopsis* guard cells was observed after treatment with the plant growth regulator 5-aminolevulinic acid, which decreased ROS levels and ABA-induced stomatal closure.²⁸⁷

In addition to their function in guard cell signaling, flavonols may also modulate the development of stomata and surrounding pavement cells. The altered flavonol glycosylation profiles in the *Arabidopsis roll* mutant, which has a defect in a rhamnose synthase gene, reducing flavonol rhamnosylation and increasing flavonol glycosylation, are responsible for multiple phenotypes, including pavement and guard cell development, resulting in larger stomata and decreased stomatal density.²⁶⁵ These phenotypes were rescued when the *roll* mutation was crossed into the flavonoid-deficient *tt4-2* mutant, verifying that these phenotypes are linked to elevated glycosylated flavonols.²⁶⁵ Additionally, soybean isolines containing elevated concentrations of kaempferol-tri-*O*-glucoside showed decreased stomatal density in response to blue light.^{290,291} However, these studies did not examine the changes in ROS status to ask whether these effects were mediated via flavonol action as antioxidants. Taken together, these experiments implicate flavonols in control of stomatal closure and development, with many of these developmental changes mediated by alterations in ROS signaling

4.3. Pollen.

During plant sexual reproduction, pollen grains germinate on the stigma of flowers to generate a pollen tube that grows through female reproductive tissues to fertilize ovules. As mentioned above, many stages of plant reproduction—from pollen development to fertilization of ovules—are modulated by ROS. Flavonols and their glycosylated derivatives accumulate in plant reproductive structures, specifically in pollen, anthers, and stigmas,^{96,292–294} raising the intriguing possibility that flavonols in reproductive structures serve as regulators of ROS-dependent processes. Flavonols were found to regulate pollen viability, germination, and tube growth and integrity.^{96,292,295–299} *CHS-RNAi* tomato plants had reduced pollen tube growth, which led to reduced or no production of seeds.²⁹⁷ Pollen germination and tube growth also were severely inhibited in a petunia *CHS* mutant, leading to reduced seed set.^{292,295} Pharmacological complementation of petunia *CHS* mutant pollen with flavonols or an extract from wild-type anthers rescued germination and tube growth *in vitro*,²⁹⁵ while treatment with other flavonoids such as flavones, flavanones, and dihydroflavanols did not,²⁹⁹ pointing to a specific role for flavonols. Addition of flavonols to the germination medium also promoted germination and tube growth of wild-type tobacco pollen.²⁹⁹ *In situ* application of kaempferol or quercetin onto stigmas during self-crosses of the petunia *CHS* mutant or a maize *CHS* mutant resulted in a complete seed set, while self-crosses without flavonol application failed to do so.²⁹⁶

Paradoxically, flavonols were not found to be required for fertilization in *Arabidopsis*,^{300,301} despite the presence of significant flavonol metabolite pools in *Arabidopsis* pollen.^{293,294} Indeed, the *Arabidopsis tt4* mutant had undetectable flavonol levels in reproductive organs yet exhibited pollen tube growth and a seed set similar to the wild-type.^{300,301} Surprisingly, the flower flavonoid transporter (FFT), which is expressed in anthers, is required for pollen development, viability, and release and thus for the formation of a complete seed set.³⁰² It is

still unclear whether this transporter has specificity toward flavonols or promiscuous activity toward other compounds necessary for pollen development. The latter could explain discrepancies between the dispensability of flavonols for pollen development in *Arabidopsis* and the reliance of reproduction on functional transport of flavonols.

While the studies mentioned above hypothesized that flavonols act as signaling molecules in pollen, their function to reduce the levels of ROS in pollen was only recently reported.⁹⁶ Muhlemann et al.⁹⁶ showed that the tomato *are* mutant had reduced levels of flavonols in pollen grains and tubes. This mutation does not affect anthocyanin accumulation in pollen, as this metabolite is absent in this tissue in both the wild-type and the *are* mutant.⁹⁶ The *are* mutant had impaired pollen viability, germination, and tube growth and integrity (Figure 8). ROS levels in *are* pollen grains and tubes, as measured using DCF and PO1 fluorescence, were inversely proportional to the flavonol levels, pointing to a role of flavonols as ROS scavengers in pollen (Figure 8).⁹⁶ Treatment with the antioxidant L-ascorbic acid or inhibition of ROS synthesis with the RBOH inhibitor DPI rescued pollen tube growth and integrity, respectively, in the *are* mutant, which confirmed that elevated ROS in the mutant caused inhibition of pollen tube growth and integrity.⁹⁶

Heat stress has profound inhibitory effects on pollen viability^{303–305} and pollen tube growth,^{306–308} and a recent study by Muhlemann et al.⁹⁶ demonstrated that this inhibitory effect is linked to elevated ROS. These authors found that elevated temperatures increased ROS in tomato pollen tubes and resulted in reduced pollen viability and tube growth and integrity.⁹⁶ The detrimental effects of elevated temperature on pollen tube growth and integrity were alleviated by supplementing the pollen germination medium with the antioxidant L-ascorbic acid, demonstrating that elevated ROS generated during high-temperature stress underlie the inhibition of pollen tube growth and integrity.⁹⁶ In the flavonol-deficient *are* mutant, the high-temperature effects on pollen viability and tube growth and integrity were exaggerated,⁹⁶ suggesting that flavonols protect pollen from heat-induced ROS.

5. CONCLUSIONS

ROS are potent signals that can oxidize proteins, thereby changing their activity and leading to growth and developmental changes. Plants have tissue-specific expression of the RBOH family of enzymes that can be activated by hormones or other environmental signals to mediate a range of developmental responses. Equally important to productive ROS signaling is maintenance of ROS homeostasis to prevent its concentration from reaching damaging levels. Regulation of ROS levels in plants relies on a large network of specialized metabolites with antioxidant activities, such as carotenoids and flavonoids. These metabolites originate from different biosynthetic pathways and exhibit distinct subcellular localizations and functions in plant growth, development, and stress responses. Because of their human health-promoting antioxidant activities and visually attractive appearance, metabolites of the flavonoid class have received much attention. Recent experiments using genetic approaches have revealed the *in planta* functions of flavonols as ROS scavengers and their impact on ROS-dependent developmental and stress response programs.

ACKNOWLEDGMENTS

The authors appreciate the editorial input by Leslie Poole, Anthony Postiglione, and Andria Harkey.

Funding

The authors were supported by NSF IOS-1558046 (G.K.M., S.R.G., and J.M.C.), USDA 2016-67013-24746 (G.K.M. and J.K.M.), Swiss NSF P2SKP3_161684 (J.K.M.), and NIH T32 GM127261 (J.M.C.).

Biography

Jordan M. Chapman is a Ph.D. candidate at Wake Forest University. She received her B.S. in Chemistry with a concentration in Biochemistry from Wake Forest University. Her current work focuses on identification of plant proteins that undergo reversible oxidation, the role of flavonols in maintaining ROS homeostasis, and how flavonols and oxidized proteins modulate root architecture.

Joëlle K. Mühlemann is currently a research assistant professor in the Department of Biology at Wake Forest University. She received her M.Sc. in Environmental Sciences from ETH Zürich in Switzerland and her Ph.D. from Purdue University. Her research focuses on flavonols and how they protect pollen from temperature stress, and on hormonal signaling in plant development.

Sheena R. Gayomba is a postdoctoral researcher at Wake Forest University. She received her M.S. and Ph.D. from Cornell University. Her research uses a genetic approach to uncover the role of flavonol antioxidants in root development in the model plant *Arabidopsis thaliana*. She is also interested in understanding the role of flavonols and reactive oxygen species in modulating plant responses to environmental stresses.

Gloria K. Muday is a professor in the Department of Biology at Wake Forest University and Director of the Center for Molecular Signaling. She received her Ph.D. in Biochemistry from Purdue University. The research in her laboratory examines plant hormone signaling networks, flavonoid biosynthesis, and reactive oxygen species signaling and the role of these processes in plant development and response to abiotic stress.

REFERENCES

- (1). Willems P, Mhamdi A, Stael S, Storme V, Kerchev P, Noctor G, Gevaert K, and Van Breusegem F (2016) The ROS Wheel: Refining ROS Transcriptional Footprints. *Plant Physiol* 171, 1720–1733. [PubMed: 27246095]
- (2). Baxter A, Mittler R, and Suzuki N (2014) ROS as Key Players in Plant Stress Signalling. *J. Exp. Bot* 65, 1229–1240. [PubMed: 24253197]
- (3). Mittler R (2017) ROS Are Good. *Trends Plant Sci* 22, 11–19. [PubMed: 27666517]
- (4). Suzuki N, Miller G, Morales J, Shulaev V, Torres MA, and Mittler R (2011) Respiratory Burst Oxidases: The Engines of ROS Signaling. *Curr. Opin. Plant Biol* 14, 691–699. [PubMed: 21862390]
- (5). Luthje S, and Martinez-Cortes T (2018) Membrane-Bound Class III Peroxidases: Unexpected Enzymes with Exciting Functions. *Int. J. Mol. Sci* 19, 2876.
- (6). Mittler R (2002) Oxidative Stress, Antioxidants and Stress Tolerance. *Trends Plant Sci* 7, 405–410. [PubMed: 12234732]

- (7). Foyer CH, and Noctor G (2005) Redox Homeostasis and Antioxidant Signaling: A Metabolic Interface between Stress Perception and Physiological Responses. *Plant Cell* 17, 1866–1875. [PubMed: 15987996]
- (8). Sbarra AJ, and Karnovsky ML (1959) The Biochemical Basis of Phagocytosis. I. Metabolic Changes During the Ingestion of Particles by Polymorphonuclear Leukocytes. *J. Biol. Chem* 234, 1355–1362. [PubMed: 13654378]
- (9). Segal AW, and Abo A (1993) The Biochemical Basis of the NADPH Oxidase of Phagocytes. *Trends Biochem. Sci* 18, 43–47. [PubMed: 8488557]
- (10). Nauseef WM (2008) Biological Roles for the Nox Family NADPH Oxidases. *J. Biol. Chem* 283, 16961–16965. [PubMed: 18420576]
- (11). Martyn KD, Frederick LM, von Loehneysen K, Dinanier MC, and Knaus UG (2006) Functional Analysis of Nox4 Reveals Unique Characteristics Compared to Other NADPH Oxidases. *Cell. Signalling* 18, 69–82. [PubMed: 15927447]
- (12). Takac I, Schroder K, Zhang L, Lardy B, Anilkumar N, Lambeth JD, Shah AM, Morel F, and Brandes RP (2011) The E-Loop Is Involved in Hydrogen Peroxide Formation by the NADPH Oxidase Nox4. *J. Biol. Chem* 286, 13304–13313. [PubMed: 21343298]
- (13). Clark RA, Volpp BD, Leidal KG, and Nauseef WM (1990) Two Cytosolic Components of the Human Neutrophil Respiratory Burst Oxidase Translocate to the Plasma Membrane during Cell Activation. *J. Clin. Invest* 85, 714–721. [PubMed: 2155923]
- (14). Keller T, Damude HG, Werner D, Doerner P, Dixon RA, and Lamb C (1998) A Plant Homolog of the Neutrophil NADPH Oxidase gp91^{phox} Subunit Gene Encodes a Plasma Membrane Protein with Ca²⁺ Binding Motifs. *Plant Cell* 10, 255–266. [PubMed: 9490748]
- (15). Finegold AA, Shatwell KP, Segal AW, Klausner RD, and Dancis A (1996) Intramembrane Bis-Heme Motif for Transmembrane Electron Transport Conserved in a Yeast Iron Reductase and the Human NADPH Oxidase. *J. Biol. Chem* 271, 31021–31024. [PubMed: 8940093]
- (16). Segal AW, West I, Wientjes F, Nugent JH, Chavan AJ, Haley B, Garcia RC, Rosen H, and Scrase G (1992) Cytochrome B-245 Is a Flavocytochrome Containing FAD and the NADPH-Binding Site of the Microbicidal Oxidase of Phagocytes. *Biochem. J* 284, 781–788. [PubMed: 1320378]
- (17). Taylor WR, Jones DT, and Segal AW (1993) A Structural Model for the Nucleotide Binding Domains of the Flavocytochrome b₂₄₅ β -Chain. *Protein Sci* 2, 1675–1685. [PubMed: 8251942]
- (18). Torres MA, Onouchi H, Hamada S, Machida C, Hammond-Kosack KE, and Jones JD (1998) Six *Arabidopsis thaliana* Homologues of the Human Respiratory Burst Oxidase (gp91^{phox}). *Plant J* 14, 365–370. [PubMed: 9628030]
- (19). Sagi M, and Fluhr R (2006) Production of Reactive Oxygen Species by Plant NADPH Oxidases. *Plant Physiol* 141, 336–340. [PubMed: 16760484]
- (20). Pierre-Jerome E, Drapek C, and Benfey PN (2018) Regulation of Division and Differentiation of Plant Stem Cells. *Annu. Rev. Cell Dev. Biol* 34, 289–310. [PubMed: 30134119]
- (21). Scheres B (2007) Stem-Cell Niches: Nursery Rhymes across Kingdoms. *Nat. Rev. Mol. Cell Biol* 8, 345–354. [PubMed: 17450175]
- (22). Pillitteri LJ, Guo X, and Dong J (2016) Asymmetric Cell Division in Plants: Mechanisms of Symmetry Breaking and Cell Fate Determination. *Cell. Mol. Life Sci* 73, 4213–4229. [PubMed: 27286799]
- (23). van den Berg C, Willemsen V, Hage W, Weisbeek P, and Scheres B (1995) Cell Fate in the *Arabidopsis* Root Meristem Determined by Directional Signalling. *Nature* 378, 62–65. [PubMed: 7477287]
- (24). Baluska F, Kubica S, and Hauskrecht M (1990) Postmitotic 'Isodiametric' Cell Growth in the Maize Root Apex. *Planta* 181, 269–274. [PubMed: 24196802]
- (25). Baluska F, Barlow PW, and Kubica S (1994) Importance of the Post-Mitotic Isodiametric Growth (Pig) Region for Growth and Development of Roots. *Plant Soil* 167, 31–41.
- (26). van der Weele CM, Jiang HS, Palaniappan KK, Ivanov VB, Palaniappan K, and Baskin TI (2003) A New Algorithm for Computational Image Analysis of Deformable Motion at High Spatial and Temporal Resolution Applied to Root Growth. Roughly Uniform Elongation in the Meristem and Also, after an Abrupt Acceleration, in the Elongation Zone. *Plant Physiol* 132, 1138–1148. [PubMed: 12857796]

- (27). Baluska F, Mancuso S, Volkmann D, and Barlow PW (2010) Root Apex Transition Zone: A Signalling-Response Nexus in the Root. *Trends Plant Sci* 15, 402–408. [PubMed: 20621671]
- (28). Grierson C, Nielsen E, Ketelaarc T, and Schiefelbein J (2014) Root Hairs. *Arabidopsis Book* 12, e0172. [PubMed: 24982600]
- (29). Muller M, and Schmidt W (2004) Environmentally Induced Plasticity of Root Hair Development in *Arabidopsis*. *Plant Physiol* 134, 409–419. [PubMed: 14730071]
- (30). Dolan L, Janmaat K, Willemsen V, Linstead P, Poethig S, Roberts K, and Scheres B (1993) Cellular Organisation of the *Arabidopsis thaliana* Root. *Development* 119, 71–84. [PubMed: 8275865]
- (31). Dubrovsky JG, Soukup A, Napsucially-Mendivil S, Jeknic Z, and Ivanchenko MG (2009) The Lateral Root Initiation Index: An Integrative Measure of Primordium Formation. *Ann. Bot* 103, 807–817. [PubMed: 19151042]
- (32). Giehl RF, and von Wiren N (2014) Root Nutrient Foraging. *Plant Physiol* 166, 509–517. [PubMed: 25082891]
- (33). Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE, Bodde S, Jones JD, and Schroeder JI (2003) NADPH Oxidase *AtrbohD* and *AtrbohF* Genes Function in ROS-Dependent ABA Signaling in *Arabidopsis*. *EMBO J* 22, 2623–2633. [PubMed: 12773379]
- (34). Jiao Y, Sun L, Song Y, Wang L, Liu L, Zhang L, Liu B, Li N, Miao C, and Hao F (2013) *AtrbohD* and *AtrbohF* Positively Regulate Abscisic Acid-Inhibited Primary Root Growth by Affecting Ca^{2+} Signalling and Auxin Response of Roots in *Arabidopsis*. *J. Exp. Bot* 64, 4183–4192. [PubMed: 23963673]
- (35). Krieger G, Shkolnik D, Miller G, and Fromm H (2016) Reactive Oxygen Species Tune Root Tropic Responses. *Plant Physiol* 172, 1209–1220. [PubMed: 27535793]
- (36). Barbosa ICR, Rojas-Murcia N, and Geldner N (2019) The Casparian Strip-One Ring to Bring Cell Biology to Lignification? *Curr. Opin. Biotechnol* 56, 121–129. [PubMed: 30502636]
- (37). Doblas VG, Geldner N, and Barberon M (2017) The Endodermis, a Tightly Controlled Barrier for Nutrients. *Curr. Opin. Plant Biol* 39, 136–143. [PubMed: 28750257]
- (38). Naseer S, Lee Y, Lapierre C, Franke R, Nawrath C, and Geldner N (2012) Casparian Strip Diffusion Barrier in *Arabidopsis* Is Made of a Lignin Polymer without Suberin. *Proc. Natl. Acad. Sci. U. S. A* 109, 10101–10106. [PubMed: 22665765]
- (39). Lee Y, Rubio MC, Alassimone J, and Geldner N (2013) A Mechanism for Localized Lignin Deposition in the Endodermis. *Cell* 153, 402–412. [PubMed: 23541512]
- (40). Malamy JE, and Benfey PN (1997) Organization and Cell Differentiation in Lateral Roots of *Arabidopsis thaliana*. *Development* 124, 33–44. [PubMed: 9006065]
- (41). Li N, Sun L, Zhang L, Song Y, Hu P, Li C, and Hao FS (2015) *AtrbohD* and *AtrbohF* Negatively Regulate Lateral Root Development by Changing the Localized Accumulation of Superoxide in Primary Roots of *Arabidopsis*. *Planta* 241, 591–602. [PubMed: 25399352]
- (42). Ma L, Zhang H, Sun L, Jiao Y, Zhang G, Miao C, and Hao F (2012) NADPH Oxidase *AtrbohD* and *AtrbohF* Function in ROS-Dependent Regulation of Na^+/K^+ Homeostasis in *Arabidopsis* under Salt Stress. *J. Exp. Bot* 63, 305–317. [PubMed: 21984648]
- (43). Manzano C, Pallero-Baena M, Casimiro I, De Rybel B, Orman-Ligeza B, Van Isterdael G, Beeckman T, Draye X, Casero P, and Del Pozo JC (2014) The Emerging Role of Reactive Oxygen Species Signaling during Lateral Root Development. *Plant Physiol* 165, 1105–1119. [PubMed: 24879433]
- (44). Orman-Ligeza B, Parizot B, de Rycke R, Fernandez A, Himschoot E, Van Breusegem F, Bennett MJ, Perilleux C, Beeckman T, and Draye X (2016) RboH-Mediated ROS Production Facilitates Lateral Root Emergence in *Arabidopsis*. *Development* 143, 3328–3339. [PubMed: 27402709]
- (45). Foreman J, Demidchik V, Bothwell JH, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JD, Davies JM, and Dolan L (2003) Reactive Oxygen Species Produced by NADPH Oxidase Regulate Plant Cell Growth. *Nature* 422, 442–446. [PubMed: 12660786]
- (46). Schiefelbein JW, and Somerville C (1990) Genetic Control of Root Hair Development in *Arabidopsis thaliana*. *Plant Cell* 2, 235–243. [PubMed: 12354956]

- (47). Takeda S, Gapper C, Kaya H, Bell E, Kuchitsu K, and Dolan L (2008) Local Positive Feedback Regulation Determines Cell Shape in Root Hair Cells. *Science* 319, 1241–1244. [PubMed: 18309082]
- (48). Schöpfer P (2001) Hydroxyl Radical-Induced Cell-Wall Loosening in Vitro and in Vivo: Implications for the Control of Elongation Growth. *Plant J* 28, 679–688. [PubMed: 11851914]
- (49). Wymer CL, Bibikova TN, and Gilroy S (1997) Cytoplasmic Free Calcium Distributions During the Development of Root Hairs of *Arabidopsis thaliana*. *Plant J* 12, 427–439. [PubMed: 9301093]
- (50). Morgan AR, Cone RL, and Terry ME (1976) The Mechanism of DNA Strand Breakage by Vitamin C and Superoxide and the Protective Roles of Catalase and Superoxide Dismutase. *Nucleic Acids Res* 3, 1139–1150. [PubMed: 181730]
- (51). Mangano S, Denita-Juarez SP, Choi HS, Marzol E, Hwang Y, Ranocha P, Velasquez SM, Borassi C, Barberini ML, Aptekmann AA, Muschietti JP, Nadra AD, Dunand C, Cho HT, and Estevez JM (2017) Molecular Link between Auxin and ROS-Mediated Polar Growth. *Proc. Natl. Acad. Sci. U. S. A* 114, 5289–5294. [PubMed: 28461488]
- (52). Bhosale R, Giri J, Pandey BK, Giehl RFH, Hartmann A, Traini R, Truskina J, Leftley N, Hanlon M, Swarup K, Rashed A, Voss U, Alonso J, Stepanova A, Yun J, Ljung K, Brown KM, Lynch JP, Dolan L, Vernoux T, Bishopp A, Wells D, von Wiren N, Bennett MJ, and Swarup R (2018) A Mechanistic Framework for Auxin Dependent Arabidopsis Root Hair Elongation to Low External Phosphate. *Nat. Commun* 9, 1409. [PubMed: 29651114]
- (53). Rahman A, Hosokawa S, Oono Y, Amakawa T, Goto N, and Tsurumi S (2002) Auxin and Ethylene Response Interactions during Arabidopsis Root Hair Development Dissected by Auxin Influx Modulators. *Plant Physiol* 130, 1908–1917. [PubMed: 12481073]
- (54). Pitts RJ, Cernac A, and Estelle M (1998) Auxin and Ethylene Promote Root Hair Elongation in *Arabidopsis*. *Plant J* 16, 553–560. [PubMed: 10036773]
- (55). Cosgrove DJ (2018) Diffuse Growth of Plant Cell Walls. *Plant Physiol* 176, 16–27. [PubMed: 29138349]
- (56). Duan Q, Kita D, Li C, Cheung AY, and Wu HM (2010) FERONIA Receptor-like Kinase Regulates RHO GTPase Signaling of Root Hair Development. *Proc. Natl. Acad. Sci. U. S. A* 107, 17821–17826. [PubMed: 20876100]
- (57). Yang Z (2002) Small GTPases: Versatile Signaling Switches in Plants. *Plant Cell* 14, S375–S388. [PubMed: 12045289]
- (58). Jones MA, Shen JJ, Fu Y, Li H, Yang Z, and Grierson CS (2002) The Arabidopsis Rop2 GTPase Is a Positive Regulator of Both Root Hair Initiation and Tip Growth. *Plant Cell* 14, 763–776. [PubMed: 11971133]
- (59). Jones MA, Raymond MJ, Yang Z, and Smirnov N (2007) NADPH Oxidase-Dependent Reactive Oxygen Species Formation Required for Root Hair Growth Depends on ROP GTPase. *J. Exp. Bot* 58, 1261–1270. [PubMed: 17301029]
- (60). Carol RJ, Takeda S, Linstead P, Durrant MC, Kakesova H, Derbyshire P, Drea S, Zarsky V, and Dolan L (2005) A RhoGDP Dissociation Inhibitor Spatially Regulates Growth in Root Hair Cells. *Nature* 438, 1013–1016. [PubMed: 16355224]
- (61). Feng W, Kita D, Peaucelle A, Cartwright HN, Doan V, Duan Q, Liu MC, Maman J, Steinhorst L, Schmitz-Thom I, Yvon R, Kudla J, Wu HM, Cheung AY, and Dinneny JR (2018) The FERONIA Receptor Kinase Maintains Cell-Wall Integrity During Salt Stress through Ca²⁺ Signaling. *Curr. Biol* 28, 666–675. [PubMed: 29456142]
- (62). Shih HW, Miller ND, Dai C, Spalding EP, and Monshausen GB (2014) The Receptor-like Kinase FERONIA Is Required for Mechanical Signal Transduction in *Arabidopsis* Seedlings. *Curr. Biol* 24, 1887–1892. [PubMed: 25127214]
- (63). Chai S, Ge FR, Feng QN, Li S, and Zhang Y (2016) *PLURIPETALA* Mediates ROP2 Localization and Stability in Parallel to *SCN1* but Synergistically with *TIP1* in Root Hairs. *Plant J* 86, 413–425. [PubMed: 27037800]
- (64). Banfi B, Molnar G, Maturana A, Steger K, Hegedus B, Demareux N, and Krause KH (2001) A Ca²⁺-Activated NADPH Oxidase in Testis, Spleen, and Lymph Nodes. *J. Biol. Chem* 276, 37594–37601. [PubMed: 11483596]

- (65). Shiose A, and Sumimoto H (2000) Arachidonic Acid and Phosphorylation Synergistically Induce a Conformational Change of *P47^{phox}* to Activate the Phagocyte NADPH Oxidase. *J. Biol. Chem* 275, 13793–13801. [PubMed: 10788501]
- (66). Zhang X, Koster P, Schlucking K, Balcerowicz D, Hashimoto K, Kuchitsu K, Vissenberg K, and Kudla J (2018) CBL1-CIPK26-Mediated Phosphorylation Enhances Activity of the NADPH Oxidase RBOHC, but Is Dispensable for Root Hair Growth. *FEBS Lett* 592, 2582–2593. [PubMed: 29992537]
- (67). Nestler J, Liu S, Wen TJ, Paschold A, Marcon C, Tang HM, Li D, Li L, Meeley RB, Sakai H, Bruce W, Schnable PS, and Hochholdinger F (2014) *Roothairless5*, Which Functions in Maize (*Zea mays* L.) Root Hair Initiation and Elongation Encodes a Monocot-Specific NADPH Oxidase. *Plant J* 79, 729–740. [PubMed: 24902980]
- (68). Wang GF, Li WQ, Li WY, Wu GL, Zhou CY, and Chen KM (2013) Characterization of Rice NADPH Oxidase Genes and Their Expression under Various Environmental Conditions. *Int. J. Mol. Sci* 14, 9440–9458. [PubMed: 23629674]
- (69). Qi J, Song CP, Wang B, Zhou J, Kangasjarvi J, Zhu JK, and Gong Z (2018) Reactive Oxygen Species Signaling and Stomatal Movement in Plant Responses to Drought Stress and Pathogen Attack. *J. Integr. Plant Biol* 60, 805–826. [PubMed: 29660240]
- (70). Zhang J, De-Oliveira-Ceciliato P, Takahashi Y, Schulze S, Dubeaux G, Hauser F, Azoulay-Shemer T, Toldsepp K, Kollist H, Rappel WJ, and Schroeder JI (2018) Insights into the Molecular Mechanisms of CO₂-Mediated Regulation of Stomatal Movements. *Curr. Biol* 28, R1356–R1363. [PubMed: 30513335]
- (71). Hedrich R, and Geiger D (2017) Biology of SLAC1-Type Anion Channels—From Nutrient Uptake to Stomatal Closure. *New Phytol* 216, 46–61. [PubMed: 28722226]
- (72). Daszkowska-Golec A, and Szarejko I (2013) Open or Close the Gate—Stomata Action under the Control of Phytohormones in Drought Stress Conditions. *Front. Plant Sci* 4, 138. [PubMed: 23717320]
- (73). Singh R, Parihar P, Singh S, Mishra RK, Singh VP, and Prasad SM (2017) Reactive Oxygen Species Signaling and Stomatal Movement: Current Updates and Future Perspectives. *Redox Biol* 11, 213–218. [PubMed: 28012436]
- (74). Munemasa S, Hauser F, Park J, Waadt R, Brandt B, and Schroeder JI (2015) Mechanisms of Absciscic Acid-Mediated Control of Stomatal Aperture. *Curr. Opin. Plant Biol* 28, 154–162. [PubMed: 26599955]
- (75). Roelfsema MR, and Hedrich R (2005) In the Light of Stomatal Opening: New Insights into ‘the Watergate’. *New Phytol* 167, 665–691. [PubMed: 16101906]
- (76). Watkins JM, Chapman JM, and Muday GK (2017) Absciscic Acid-Induced Reactive Oxygen Species Are Modulated by Flavonols to Control Stomata Aperture. *Plant Physiol* 175, 1807–1825. [PubMed: 29051198]
- (77). Okuma E, Jahan MS, Munemasa S, Hossain MA, Muroyama D, Islam MM, Ogawa K, Watanabe-Sugimoto M, Nakamura Y, Shimoishi Y, Mori IC, and Murata Y (2011) Negative Regulation of Absciscic Acid-Induced Stomatal Closure by Glutathione in *Arabidopsis*. *J. Plant Physiol* 168, 2048–2055. [PubMed: 21764168]
- (78). Munemasa S, Muroyama D, Nagahashi H, Nakamura Y, Mori IC, and Murata Y (2013) Regulation of Reactive Oxygen Species-Mediated Absciscic Acid Signaling in Guard Cells and Drought Tolerance by Glutathione. *Front. Plant Sci* 4, 472. [PubMed: 24312112]
- (79). Pei ZM, Murata Y, Benning G, Thomine S, Klusener B, Allen GJ, Grill E, and Schroeder JI (2000) Calcium Channels Activated by Hydrogen Peroxide Mediate Absciscic Acid Signalling in Guard Cells. *Nature* 406, 731–734. [PubMed: 10963598]
- (80). Devireddy AR, Zandalinas SI, Gomez-Cadenas A, Blumwald E, and Mittler R (2018) Coordinating the Overall Stomatal Response of Plants: Rapid Leaf-to-Leaf Communication During Light Stress. *Sci. Signaling* 11, eaam9514.
- (81). Hua D, Wang C, He J, Liao H, Duan Y, Zhu Z, Guo Y, Chen Z, and Gong Z (2012) A Plasma Membrane Receptor Kinase, GHR1, Mediates Absciscic Acid- and Hydrogen Peroxide-Regulated Stomatal Movement in *Arabidopsis*. *Plant Cell* 24, 2546–2561. [PubMed: 22730405]

- (82). Sierla M, Horak H, Overmyer K, Waszczak C, Yarmolinsky D, Maierhofer T, Vainonen JP, Denessiouk K, Salojärvi J, Laanemets K, Toldsepp K, Vahisalu T, Gauthier A, Puukko T, Paulin L, Auvinen P, Geiger D, Hedrich R, Kollist H, and Kangasjarvi J (2018) The Receptor-like Pseudokinase GHR1 Is Required for Stomatal Closure. *Plant Cell* 30, 2813–2837. [PubMed: 30361234]
- (83). Bright J, Desikan R, Hancock JT, Weir IS, and Neill SJ (2006) ABA-Induced NO Generation and Stomatal Closure in Arabidopsis Are Dependent on H₂O₂ Synthesis. *Plant J* 45, 113–122. [PubMed: 16367958]
- (84). Scuffi D, Nietzel T, Di Fino LM, Meyer AJ, Lamattina L, Schwarzlander M, Laxalt AM, and Garcia-Mata C (2018) Hydrogen Sulfide Increases Production of NADPH Oxidase-Dependent Hydrogen Peroxide and Phospholipase D-Derived Phosphatidic Acid in Guard Cell Signaling. *Plant Physiol* 176, 2532–2542. [PubMed: 29438048]
- (85). Zhang Y, Zhu H, Zhang Q, Li M, Yan M, Wang R, Wang L, Welti R, Zhang W, and Wang X (2009) Phospholipase Dα1 and Phosphatidic Acid Regulate NADPH Oxidase Activity and Production of Reactive Oxygen Species in ABA-Mediated Stomatal Closure in *Arabidopsis*. *Plant Cell* 21, 2357–2377. [PubMed: 19690149]
- (86). Kimura S, Kawarazaki T, Nibori H, Michikawa M, Imai A, Kaya H, and Kuchitsu K (2013) The CBL-Interacting Protein Kinase CIPK26 Is a Novel Interactor of Arabidopsis NADPH Oxidase AtRbohF That Negatively Modulates Its ROS-Producing Activity in a Heterologous Expression System. *J. Biochem* 153, 191–195. [PubMed: 23162070]
- (87). Drerup MM, Schlucking K, Hashimoto K, Manishankar P, Steinhorst L, Kuchitsu K, and Kudla J (2013) The Calcineurin B-Like Calcium Sensors CBL1 and CBL9 Together with Their Interacting Protein Kinase CIPK26 Regulate the Arabidopsis NADPH Oxidase RBOHF. *Mol. Plant* 6, 559–569. [PubMed: 23335733]
- (88). Drews GN, and Koltunow AMG (2011) The Female Gametophyte. *Arabidopsis Book* 9, e0155. [PubMed: 22303279]
- (89). Cheung AY, Boavida LC, Aggarwal M, Wu HM, and Feijo JA (2010) The Pollen Tube Journey in the Pistil and Imaging the *in Vivo* Process by Two-Photon Microscopy. *J. Exp. Bot* 61, 1907–1915. [PubMed: 20363865]
- (90). Xie HT, Wan ZY, Li S, and Zhang Y (2014) Spatiotemporal Production of Reactive Oxygen Species by NADPH Oxidase Is Critical for Tapetal Programmed Cell Death and Pollen Development in Arabidopsis. *Plant Cell* 26, 2007–2023. [PubMed: 24808050]
- (91). Potocky M, Jones MA, Bezvoda R, Smirnov N, and Zarsky V (2007) Reactive Oxygen Species Produced by NADPH Oxidase Are Involved in Pollen Tube Growth. *New Phytol* 174, 742–751. [PubMed: 17504458]
- (92). Kaya H, Nakajima R, Iwano M, Kanaoka MM, Kimura S, Takeda S, Kawarazaki T, Senzaki E, Hamamura Y, Higashiyama T, Takayama S, Abe M, and Kuchitsu K (2014) Ca²⁺-Activated Reactive Oxygen Species Production by Arabidopsis RbohH and RbohJ Is Essential for Proper Pollen Tube Tip Growth. *Plant Cell* 26, 1069–1080. [PubMed: 24610725]
- (93). Lassig R, Gutermuth T, Bey TD, Konrad KR, and Romeis T (2014) Pollen Tube NAD(P)H Oxidases Act as a Speed Control to Dampen Growth Rate Oscillations During Polarized Cell Growth. *Plant J* 78, 94–106. [PubMed: 24506280]
- (94). Boisson-Dernier A, Lituiev DS, Nestorova A, Franck CM, Thirugnanarajah S, and Grossniklaus U (2013) Anxur Receptor-Like Kinases Coordinate Cell Wall Integrity with Growth at the Pollen Tube Tip Via NADPH Oxidases. *PLoS Biol* 11, e1001719. [PubMed: 24302886]
- (95). Duan Q, Kita D, Johnson EA, Aggarwal M, Gates L, Wu HM, and Cheung AY (2014) Reactive Oxygen Species Mediate Pollen Tube Rupture to Release Sperm for Fertilization in *Arabidopsis*. *Nat. Commun* 5, 3129. [PubMed: 24451849]
- (96). Muhlemann JK, Younts TLB, and Muday GK (2018) Flavonols Control Pollen Tube Growth and Integrity by Regulating ROS Homeostasis During High Temperature Stress. *Proc. Natl. Acad. Sci. U. S. A* 115, E11188–E11197. [PubMed: 30413622]
- (97). Hsieh K, and Huang AH (2007) Tapetosomes in *Brassica tapetum* Accumulate Endoplasmic Reticulum-Derived Flavonoids and Alkanes for Delivery to the Pollen Surface. *Plant Cell* 19, 582–596. [PubMed: 17307923]

- (98). Sorensen AM, Krober S, Unte US, Huijser P, Dekker K, and Saedler H (2003) The *Arabidopsis* *ABORTED MICROSPORES (AMS)* Gene Encodes a MYC Class Transcription Factor. *Plant J* 33, 413–423. [PubMed: 12535353]
- (99). Xu J, Ding Z, Vizcay-Barrena G, Shi J, Liang W, Yuan Z, Werck-Reichhart D, Schreiber L, Wilson ZA, and Zhang D (2014) Aborted Microspores Acts as a Master Regulator of Pollen Wall Formation in *Arabidopsis*. *Plant Cell* 26, 1544–1556. [PubMed: 24781116]
- (100). Phan HA, Iacuone S, Li SF, and Parish RW (2011) The MYB80 Transcription Factor Is Required for Pollen Development and the Regulation of Tapetal Programmed Cell Death in *Arabidopsis thaliana*. *Plant Cell* 23, 2209–2224. [PubMed: 21673079]
- (101). Chebli Y, Kaneda M, Zerkour R, and Geitmann A (2012) The Cell Wall of the *Arabidopsis* Pollen Tube-Spatial Distribution, Recycling, and Network Formation of Polysaccharides. *Plant Physiol* 160, 1940–1955. [PubMed: 23037507]
- (102). Higashiyama T, Yabe S, Sasaki N, Nishimura Y, Miyagishima S, Kuroiwa H, and Kuroiwa T (2001) Pollen Tube Attraction by the Synergid Cell. *Science* 293, 1480–1483. [PubMed: 11520985]
- (103). Finch-Savage WE, and Leubner-Metzger G (2006) Seed Dormancy and the Control of Germination. *New Phytol* 171, 501–523. [PubMed: 16866955]
- (104). Muller K, Carstens AC, Linkies A, Torres MA, and Leubner-Metzger G (2009) The NADPH-Oxidase *AtrbohB* Plays a Role in *Arabidopsis* Seed after-Ripening. *New Phytol* 184, 885–897. [PubMed: 19761445]
- (105). Torres MA, Dangl JL, and Jones JD (2002) *Arabidopsis* Gp91phox Homologues *AtrbohD* and *AtrbohF* Are Required for Accumulation of Reactive Oxygen Intermediates in the Plant Defense Response. *Proc. Natl. Acad. Sci. U. S. A* 99, 517–522. [PubMed: 11756663]
- (106). He H, Yan J, Yu X, Liang Y, Fang L, Scheller HV, and Zhang A (2017) The NADPH-Oxidase *AtrbohI* Plays a Positive Role in Drought-Stress Response in *Arabidopsis thaliana*. *Biochem. Biophys. Res. Commun* 491, 834–839. [PubMed: 28559135]
- (107). Joo JH, Wang S, Chen JG, Jones AM, and Fedoroff NV (2005) Different Signaling and Cell Death Roles of Heterotrimeric G Protein α and β Subunits in the *Arabidopsis* Oxidative Stress Response to Ozone. *Plant Cell* 17, 957–970. [PubMed: 15705948]
- (108). Liu B, Sun L, Ma L, and Hao FS (2017) Both *AtrbohD* and *AtrbohF* Are Essential for Mediating Responses to Oxygen Deficiency in *Arabidopsis*. *Plant Cell Rep* 36, 947–957. [PubMed: 28337518]
- (109). Zhai L, Sun C, Feng Y, Li D, Chai X, Wang L, Sun Q, Zhang G, Li Y, Wu T, Zhang X, Xu X, Wang Y, and Han Z (2018) *Atrop6* Is Involved in Reactive Oxygen Species Signaling in Response to Iron-Deficiency Stress in *Arabidopsis thaliana*. *FEBS Lett* 592, 3446–3459. [PubMed: 30238451]
- (110). Choi WG, Miller G, Wallace I, Harper J, Mittler R, and Gilroy S (2017) Orchestrating Rapid Long-Distance Signaling in Plants with Ca^{2+} , ROS and Electrical Signals. *Plant J* 90, 698–707. [PubMed: 28112437]
- (111). Morales J, Kadota Y, Zipfel C, Molina A, and Torres MA (2016) The *Arabidopsis* NADPH Oxidases *RbohD* and *RbohF* Display Differential Expression Patterns and Contributions During Plant Immunity. *J. Exp. Bot* 67, 1663–1676. [PubMed: 26798024]
- (112). Li L, Li M, Yu L, Zhou Z, Liang X, Liu Z, Cai G, Gao L, Zhang X, Wang Y, Chen S, and Zhou JM (2014) The FLS2-Associated Kinase BIK1 Directly Phosphorylates the NADPH Oxidase *RbohD* to Control Plant Immunity. *Cell Host Microbe* 15, 329–338. [PubMed: 24629339]
- (113). Kadota Y, Sklenar J, Derbyshire P, Strassfeld L, Asai S, Ntoukakis V, Jones JD, Shirasu K, Menke F, Jones A, and Zipfel C (2014) Direct Regulation of the NADPH Oxidase *RBOHD* by the PRR-Associated Kinase BIK1 during Plant Immunity. *Mol. Cell* 54, 43–55. [PubMed: 24630626]
- (114). Xu J, Xie J, Yan C, Zou X, Ren D, and Zhang S (2014) A Chemical Genetic Approach Demonstrates That MPK3/MPK6 Activation and NADPH Oxidase-Mediated Oxidative Burst Are Two Independent Signaling Events in Plant Immunity. *Plant J* 77, 222–234. [PubMed: 24245741]

- (115). Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nurnberger T, Jones JD, Felix G, and Boller T (2007) A Flagellin-Induced Complex of the Receptor FLS2 and BAK1 Initiates Plant Defence. *Nature* 448, 497–500. [PubMed: 17625569]
- (116). Monaghan J, Matschi S, Romeis T, and Zipfel C (2015) The Calcium-Dependent Protein Kinase CPK28 Negatively Regulates the BIK1-Mediated PAMP-Induced Calcium Burst. *Plant Signaling Behav* 10, e1018497.
- (117). Lee DS, Kim YC, Kwon SJ, Ryu CM, and Park OK (2017) The Arabidopsis Cysteine-Rich Receptor-Like Kinase CRK36 Regulates Immunity through Interaction with the Cytoplasmic Kinase BIK1. *Front. Plant Sci* 8, 1856. [PubMed: 29163585]
- (118). Ranf S, Eschen-Lippold L, Pecher P, Lee J, and Scheel D (2011) Interplay between Calcium Signalling and Early Signalling Elements During Defence Responses to Microbe- or Damage-Associated Molecular Patterns. *Plant J* 68, 100–113. [PubMed: 21668535]
- (119). Ogasawara Y, Kaya H, Hiraoka G, Yumoto F, Kimura S, Kadota Y, Hishinuma H, Senzaki E, Yamagoe S, Nagata K, Nara M, Suzuki K, Tanokura M, and Kuchitsu K (2008) Synergistic Activation of the *Arabidopsis* NADPH Oxidase AtrbohD by Ca^{2+} and Phosphorylation. *J. Biol. Chem* 283, 8885–8892. [PubMed: 18218618]
- (120). Mersmann S, Bourdais G, Rietz S, and Robatzek S (2010) Ethylene Signaling Regulates Accumulation of the Fls2 Receptor and Is Required for the Oxidative Burst Contributing to Plant Immunity. *Plant Physiol* 154, 391–400. [PubMed: 20592040]
- (121). Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, and Zimmermann P (2008) Genevestigator V3: A Reference Expression Database for the Meta-Analysis of Transcriptomes. *Adv. Bioinf* 2008, 420747.
- (122). Kollist H, Zandalinas SI, Sengupta S, Nuhkat M, Kangasjarvi J, and Mittler R (2019) Rapid Responses to Abiotic Stress: Priming the Landscape for the Signal Transduction Network. *Trends Plant Sci* 24, 25–37. [PubMed: 30401516]
- (123). Miller G, Schlauch K, Tam R, Cortes D, Torres MA, Shulaev V, Dangl JL, and Mittler R (2009) The Plant NADPH Oxidase RbohD Mediates Rapid Systemic Signaling in Response to Diverse Stimuli. *Sci. Signaling* 2, ra45.
- (124). Lin IS, Wu YS, Chen CT, Chen GH, Hwang SG, Jauh GY, Tzen JTC, and Yang CY (2017) Atrbohi Confers Submergence Tolerance and Is Involved in Auxin-Mediated Signaling Pathways under Hypoxic Stress. *Plant Growth Regul* 83, 277–285.
- (125). Gayomba SR, Zhai Z, Jung HI, and Vatamaniuk OK (2015) Local and Systemic Signaling of Iron Status and Its Interactions with Homeostasis of Other Essential Elements. *Front. Plant Sci* 6, 716. [PubMed: 26442030]
- (126). Jeong J, Merkovich A, Clyne M, and Connolly EL (2017) Directing Iron Transport in Dicots: Regulation of Iron Acquisition and Translocation. *Curr. Opin. Plant Biol* 39, 106–113. [PubMed: 28689052]
- (127). Le CT, Brumbarova T, Ivanov R, Stoof C, Weber E, Mohrbacher J, Fink-Straube C, and Bauer P (2016) Zinc Finger of *Arabidopsis thaliana* 12 (Zat12) Interacts with Fer-Like Iron Deficiency-Induced Transcription Factor (FIT) Linking Iron Deficiency and Oxidative Stress Responses. *Plant Physiol* 170, 540–557. [PubMed: 26556796]
- (128). Yao Y, He RJ, Xie QL, Zhao XH, Deng XM, He JB, Song L, He J, Marchant A, Chen XY, and Wu AM (2017) Ethylene Response Factor 74 (Erf74) Plays an Essential Role in Controlling a Respiratory Burst Oxidase Homolog D (RbohD)-Dependent Mechanism in Response to Different Stresses in Arabidopsis. *New Phytol* 213, 1667–1681. [PubMed: 28164334]
- (129). Chang KN, Zhong S, Weirauch MT, Hon G, Pelizzola M, Li H, Huang SS, Schmitz RJ, Urlich MA, Kuo D, Nery JR, Qiao H, Yang A, Jamali A, Chen H, Ideker T, Ren B, Bar-Joseph Z, Hughes TR, and Ecker JR (2013) Temporal Transcriptional Response to Ethylene Gas Drives Growth Hormone Cross-Regulation in Arabidopsis. *eLife* 2, e00675. [PubMed: 23795294]
- (130). Harkey AF, Watkins JM, Olex AL, DiNapoli KT, Lewis DR, Fetrow JS, Binder BM, and Muday GK (2018) Identification of Transcriptional and Receptor Networks That Control Root Responses to Ethylene. *Plant Physiol* 176, 2095–2118. [PubMed: 29259106]

- (131). Sewelam N, Kazan K, Thomas-Hall SR, Kidd BN, Manners JM, and Schenk PM (2013) Ethylene Response Factor 6 Is a Regulator of Reactive Oxygen Species Signaling in *Arabidopsis*. PLoS One 8, e70289. [PubMed: 23940555]
- (132). Mhamdi A, Queval G, Chaouch S, Vanderauwera S, Van Breusegem F, and Noctor G (2010) Catalase Function in Plants: A Focus on *Arabidopsis* Mutants as Stress-Mimic Models. J. Exp. Bot 61, 4197–4220. [PubMed: 20876333]
- (133). Liebthal M, Maynard D, and Dietz KJ (2018) Peroxiredoxins and Redox Signaling in Plants. Antioxid. Redox Signaling 28, 609–624.
- (134). Meyer Y, Buchanan BB, Vignols F, and Reichheld JP (2009) Thioredoxins and Glutaredoxins: Unifying Elements in Redox Biology. Annu. Rev. Genet 43, 335–367. [PubMed: 19691428]
- (135). Bindoli A, Fukuto JM, and Forman HJ (2008) Thiol Chemistry in Peroxidase Catalysis and Redox Signaling. Antioxid. Redox Signaling 10, 1549–1564.
- (136). Frugoli JA, Zhong HH, Nuccio ML, McCourt P, McPeck MA, Thomas TL, and McClung CR (1996) Catalase Is Encoded by a Multigene Family in *Arabidopsis thaliana* (L.) Heynh. Plant Physiol 112, 327–336. [PubMed: 8819328]
- (137). Zhong HH, Young JC, Pease EA, Hangarter RP, and McClung CR (1994) Interactions between Light and the Circadian Clock in the Regulation of Cat2 Expression in *Arabidopsis*. Plant Physiol 104, 889–898. [PubMed: 12232134]
- (138). Sharma YK, and Davis KR (1994) Ozone-Induced Expression of Stress-Related Genes in *Arabidopsis thaliana*. Plant Physiol 105, 1089–1096. [PubMed: 12232267]
- (139). Zhong HH, and McClung CR (1996) The Circadian Clock Gates Expression of Two *Arabidopsis* Catalase Genes to Distinct and Opposite Circadian Phases. Mol. Gen. Genet 251, 196–203. [PubMed: 8668130]
- (140). Del Rio LA, and Lopez-Huertas E (2016) ROS Generation in Peroxisomes and Its Role in Cell Signaling. Plant Cell Physiol 57, 1364–1376. [PubMed: 27081099]
- (141). Du YY, Wang PC, Chen J, and Song CP (2008) Comprehensive Functional Analysis of the Catalase Gene Family in *Arabidopsis thaliana*. J. Integr. Plant Biol 50, 1318–1326. [PubMed: 19017119]
- (142). Heazlewood JL, Tonti-Filippini JS, Gout AM, Day DA, Whelan J, and Millar AH (2004) Experimental Analysis of the Arabidopsis Mitochondrial Proteome Highlights Signaling and Regulatory Components, Provides Assessment of Targeting Prediction Programs, and Indicates Plant-Specific Mitochondrial Proteins. Plant Cell 16, 241–256. [PubMed: 14671022]
- (143). Reumann S, Babujee L, Ma C, Wienkoop S, Siemsen T, Antonicelli GE, Rasche N, Luder F, Weckwerth W, and Jahn O (2007) Proteome Analysis of Arabidopsis Leaf Peroxisomes Reveals Novel Targeting Peptides, Metabolic Pathways, and Defense Mechanisms. Plant Cell 19, 3170–3193. [PubMed: 17951448]
- (144). Allen JF, and Whatley FR (1978) Effects of Inhibitors of Catalase on Photosynthesis and on Catalase Activity in Unwashed Preparations of Intact Chloroplasts. Plant Physiol 61, 957–960. [PubMed: 16660434]
- (145). Noctor G, Reichheld JP, and Foyer CH (2018) ROS-Related Redox Regulation and Signaling in Plants. Semin. Cell Dev. Biol 80, 3–12. [PubMed: 28733165]
- (146). Lee ES, Kang CH, Park JH, and Lee SY (2018) Physiological Significance of Plant Peroxiredoxins and the Structure-Related and Multifunctional Biochemistry of Peroxiredoxin 1. Antioxid. Redox Signaling 28, 625–639.
- (147). Passaia G, and Margis-Pinheiro M (2015) Glutathione Peroxidases as Redox Sensor Proteins in Plant Cells. Plant Sci 234, 22–26. [PubMed: 25804806]
- (148). Hasanuzzaman M, Nahar K, Anee TI, and Fujita M (2017) Glutathione in Plants: Biosynthesis and Physiological Role in Environmental Stress Tolerance. Physiol. Mol. Biol. Plants 23, 249–268. [PubMed: 28461715]
- (149). Martins L, Trujillo-Hernandez JA, and Reichheld JP (2018) Thiol Based Redox Signaling in Plant Nucleus. Front. Plant Sci 9, 705. [PubMed: 29892308]
- (150). Rouhier N, Cerveau D, Couturier J, Reichheld JP, and Rey P (2015) Involvement of Thiol-Based Mechanisms in Plant Development. Biochim. Biophys. Acta, Gen. Subj 1850, 1479–1496.

- (151). Smirnov N (2018) Ascorbic Acid Metabolism and Functions: A Comparison of Plants and Mammals. *Free Radical Biol. Med* 122, 116–129. [PubMed: 29567393]
- (152). Dietz K-J (2016) Thiol-Based Peroxidases and Ascorbate Peroxidases: Why Plants Rely on Multiple Peroxidase Systems in the Photosynthesizing Chloroplast? *Mol. Cells* 39, 20–25. [PubMed: 26810073]
- (153). Maruta T, Sawa Y, Shigeoka S, and Ishikawa T (2016) Diversity and Evolution of Ascorbate Peroxidase Functions in Chloroplasts: More Than Just a Classical Antioxidant Enzyme? *Plant Cell Physiol* 57, 1377–1386. [PubMed: 26738546]
- (154). Pandey S, Fartyal D, Agarwal A, Shukla T, James D, Kaul T, Negi YK, Arora S, and Reddy MK (2017) Abiotic Stress Tolerance in Plants: Myriad Roles of Ascorbate Peroxidase. *Front. Plant Sci* 8, 581. [PubMed: 28473838]
- (155). Wu Q, Yang J, Cheng N, Hirschi KD, White FF, and Park S (2017) Glutaredoxins in Plant Development, Abiotic Stress Response, and Iron Homeostasis: From Model Organisms to Crops. *Environ. Exp. Bot* 139, 91–98.
- (156). Nikkanen L, Toivola J, Diaz MG, and Rintamaki E (2017) Chloroplast Thioredoxin Systems: Prospects for Improving Photosynthesis. *Philos. Trans. R. Soc.,B* 372, 20160474.
- (157). Liu P, Zhang H, Wang H, and Xia Y (2014) Identification of Redox-Sensitive Cysteines in the Arabidopsis Proteome Using OxiTRAQ, a Quantitative Redox Proteomics Method. *Proteomics* 14, 750–762. [PubMed: 24376095]
- (158). Wang H, Wang S, Lu Y, Alvarez S, Hicks LM, Ge X, and Xia Y (2012) Proteomic Analysis of Early-Responsive Redox-Sensitive Proteins in *Arabidopsis*. *J. Proteome Res* 11, 412–424. [PubMed: 22050424]
- (159). Munne-Bosch S, Queval G, and Foyer CH (2013) The Impact of Global Change Factors on Redox Signaling Underpinning Stress Tolerance. *Plant Physiol* 161, 5–19. [PubMed: 23151347]
- (160). Conklin PL, Williams EH, and Last RL (1996) Environmental Stress Sensitivity of an Ascorbic Acid-Deficient Arabidopsis Mutant. *Proc. Natl. Acad. Sci. U. S. A* 93, 9970–9974. [PubMed: 8790441]
- (161). Apel K, and Hirt H (2004) Reactive Oxygen Species: Metabolism, Oxidative Stress, and Signal Transduction. *Annu. Rev. Plant Biol* 55, 373–399. [PubMed: 15377225]
- (162). Ye X, Al-Babili S, Kloti A, Zhang J, Lucca P, Beyer P, and Potrykus I (2000) Engineering the Provitamin A (β -Carotene) Biosynthetic Pathway into (Carotenoid-Free) Rice Endosperm. *Science* 287, 303–305. [PubMed: 10634784]
- (163). Butelli E, Titta L, Giorgio M, Mock HP, Matros A, Peterek S, Schijlen EG, Hall RD, Bovy AG, Luo J, and Martin C (2008) Enrichment of Tomato Fruit with Health-Promoting Anthocyanins by Expression of Select Transcription Factors. *Nat. Biotechnol* 26, 1301–1308. [PubMed: 18953354]
- (164). Llorente B, Martinez-Garcia JF, Stange C, and Rodriguez-Concepcion M (2017) Illuminating Colors: Regulation of Carotenoid Biosynthesis and Accumulation by Light. *Curr. Opin. Plant Biol* 37, 49–55. [PubMed: 28411584]
- (165). Sun T, Yuan H, Cao H, Yazdani M, Tadmor Y, and Li L (2018) Carotenoid Metabolism in Plants: The Role of Plastids. *Mol. Plant* 11, 58–74. [PubMed: 28958604]
- (166). Santabarbara S, Agostini A, Casazza AP, Zucchelli G, and Carbonera D (2015) Carotenoid Triplet States in Photosystem II: Coupling with Low-Energy States of the Core Complex. *Biochim. Biophys. Acta, Bioenerg* 1847, 262–275.
- (167). Qin G, Gu H, Ma L, Peng Y, Deng XW, Chen Z, and Qu LJ (2007) Disruption of Phytoene Desaturase Gene Results in Albino and Dwarf Phenotypes in *Arabidopsis* by Impairing Chlorophyll, Carotenoid, and Gibberellin Biosynthesis. *Cell Res* 17, 471–482. [PubMed: 17486124]
- (168). Dong H, Deng Y, Mu J, Lu Q, Wang Y, Xu Y, Chu C, Chong K, Lu C, and Zuo J (2007) The *Arabidopsis Spontaneous Cell Death1* Gene, Encoding a ζ -Carotene Desaturase Essential for Carotenoid Biosynthesis, Is Involved in Chloroplast Development, Photoprotection and Retrograde Signalling. *Cell Res* 17, 458–470. [PubMed: 17468780]
- (169). Pospisil P (2012) Molecular Mechanisms of Production and Scavenging of Reactive Oxygen Species by Photosystem II. *Biochim. Biophys. Acta, Bioenerg* 1817, 218–231.

- (170). Triantaphylides C, Krischke M, Hoeberichts FA, Ksas B, Gresser G, Havaux M, Van Breusegem F, and Mueller MJ (2008) Singlet Oxygen Is the Major Reactive Oxygen Species Involved in Photooxidative Damage to Plants. *Plant Physiol* 148, 960–968. [PubMed: 18676660]
- (171). Mozzo M, Dall'Osto L, Hienerwadel R, Bassi R, and Croce R (2008) Photoprotection in the Antenna Complexes of Photosystem II: Role of Individual Xanthophylls in Chlorophyll Triplet Quenching. *J. Biol. Chem* 283, 6184–6192. [PubMed: 18079125]
- (172). Gruszecki WI, and Strzalka K (2005) Carotenoids as Modulators of Lipid Membrane Physical Properties. *Biochim. Biophys. Acta, Mol. Basis Dis* 1740, 108–115.
- (173). Havaux M, Bonfils JP, Lutz C, and Niyogi KK (2000) Photodamage of the Photosynthetic Apparatus and Its Dependence on the Leaf Developmental Stage in the *npq1* Arabidopsis Mutant Deficient in the Xanthophyll Cycle Enzyme Violaxanthin De-Epoxidase. *Plant Physiol* 124, 273–284. [PubMed: 10982442]
- (174). Havaux M, Dall'Osto L, and Bassi R (2007) Zeaxanthin Has Enhanced Antioxidant Capacity with Respect to All Other Xanthophylls in Arabidopsis Leaves and Functions Independent of Binding to Psii Antennae. *Plant Physiol* 145, 1506–1520. [PubMed: 17932304]
- (175). Liguori N, Xu P, van Stokkum IHM, van Oort B, Lu Y, Karcher D, Bock R, and Croce R (2017) Different Carotenoid Conformations Have Distinct Functions in Light-Harvesting Regulation in Plants. *Nat. Commun* 8, 1994. [PubMed: 29222488]
- (176). Rutherford AW, Paterson DR, and Mullet JE (1981) A Light-Induced Spin-Polarized Triplet Detected by EPR in Photosystem II Reaction Centers. *Biochim. Biophys. Acta, Bioenerg* 635, 205–214.
- (177). Latowski D, Kuczynska P, and Strzalka K (2011) Xanthophyll Cycle—A Mechanism Protecting Plants against Oxidative Stress. *Redox Rep* 16, 78–90. [PubMed: 21722416]
- (178). Niyogi KK, Grossman AR, and Bjorkman O (1998) Arabidopsis Mutants Define a Central Role for the Xanthophyll Cycle in the Regulation of Photosynthetic Energy Conversion. *Plant Cell* 10, 1121–1134. [PubMed: 9668132]
- (179). Adams WW, Demmig-Adams B, and Winter K (1990) Relative Contributions of Zeaxanthin-Related and Zeaxanthin-Unrelated Types of 'High-Energy-State' Quenching of Chlorophyll Fluorescence in Spinach Leaves Exposed to Various Environmental Conditions. *Plant Physiol* 92, 302–309. [PubMed: 16667275]
- (180). Dall'Osto L, Holt NE, Kaligotla S, Fuciman M, Cazzaniga S, Carbonera D, Frank HA, Alric J, and Bassi R (2012) Zeaxanthin Protects Plant Photosynthesis by Modulating Chlorophyll Triplet Yield in Specific Light-Harvesting Antenna Subunits. *J. Biol Chem* 287, 41820–41834. [PubMed: 23066020]
- (181). Lokstein H, Tian L, Polle JE, and DellaPenna D (2002) Xanthophyll Biosynthetic Mutants of *Arabidopsis thaliana*: Altered Nonphotochemical Quenching of Chlorophyll Fluorescence Is Due to Changes in Photosystem II Antenna Size and Stability. *Biochim. Biophys. Acta, Bioenerg* 1553, 309–319.
- (182). Fiore A, Dall'Osto L, Fraser PD, Bassi R, and Giuliano G (2006) Elucidation of the β -Carotene Hydroxylation Pathway in *Arabidopsis thaliana*. *FEBS Lett* 580, 4718–4722. [PubMed: 16890225]
- (183). Dall'Osto L, Lico C, Alric J, Giuliano G, Havaux M, and Bassi R (2006) Lutein Is Needed for Efficient Chlorophyll Triplet Quenching in the Major LHCII Antenna Complex of Higher Plants and Effective Photoprotection in Vivo under Strong Light. *BMC Plant Biol* 6, 32. [PubMed: 17192177]
- (184). Pogson BJ, Niyogi KK, Bjorkman O, and DellaPenna D (1998) Altered Xanthophyll Compositions Adversely Affect Chlorophyll Accumulation and Nonphotochemical Quenching in *Arabidopsis* Mutants. *Proc. Natl Acad. Sci. U. S. A* 95, 13324–13329. [PubMed: 9789087]
- (185). Tian L, Magallanes-Lundback M, Musetti V, and DellaPenna D (2003) Functional Analysis of β - and *e*-Ring Carotenoid Hydroxylases in Arabidopsis. *Plant Cell* 15, 1320–1332. [PubMed: 12782726]
- (186). Cazzaniga S, Li Z, Niyogi KK, Bassi R, and Dall'Osto L (2012) The Arabidopsis Szl1 Mutant Reveals a Critical Role of Beta-Carotene in Photosystem I Photoprotection. *Plant Physiol* 159, 1745–1758. [PubMed: 23029671]

- (187). Dall'Osto L, Fiore A, Cazzaniga S, Giuliano G, and Bassi R (2007) Different Roles of Alpha- and Beta-Branch Xanthophylls in Photosystem Assembly and Photoprotection. *J. Biol Chem* 282, 35056–35068. [PubMed: 17913714]
- (188). Kalituho L, Rech J, and Jahns P (2006) The Roles of Specific Xanthophylls in Light Utilization. *Planta* 225, 423–439. [PubMed: 16896791]
- (189). Li Z, Ahn TK, Avenson TJ, Ballottari M, Cruz JA, Kramer DM, Bassi R, Fleming GR, Keasling JD, and Niyogi KK (2009) Lutein Accumulation in the Absence of Zeaxanthin Restores Nonphotochemical Quenching in the *Arabidopsis thaliana* Npq1 Mutant. *Plant Cell* 21, 1798–1812. [PubMed: 19549928]
- (190). Niyogi KK, Shih C, Soon Chow W, Pogson BJ, Dellapenna D, and Bjorkman O (2001) Photoprotection in a Zeaxanthin- and Lutein-Deficient Double Mutant of Arabidopsis. *Photosynth. Res* 67, 139–145. [PubMed: 16228323]
- (191). Dall'Osto L, Piques M, Ronzani M, Molesini B, Alborese A, Cazzaniga S, and Bassi R (2013) The *Arabidopsis nox* Mutant Lacking Carotene Hydroxylase Activity Reveals a Critical Role for Xanthophylls in Photosystem I Biogenesis. *Plant Cell* 25, 591–608. [PubMed: 23396829]
- (192). Caffarri S, Tibiletti T, Jennings RC, and Santabarbara S (2014) A Comparison between Plant Photosystem I and Photosystem II Architecture and Functioning. *Curr. Protein Pept. Sci* 15, 296–331. [PubMed: 24678674]
- (193). Ramel F, Ksas B, Akkari E, Mialoundama AS, Monnet F, Krieger-Liszka A, Ravanat JL, Mueller MJ, Bouvier F, and Havaux M (2013) Light-Induced Acclimation of the Arabidopsis Chlorina1 Mutant to Singlet Oxygen. *Plant Cell* 25, 1445–1462. [PubMed: 23590883]
- (194). Ramel F, Birtic S, Ginies C, Soubigou-Taconnat L, Triantaphylides C, and Havaux M (2012) Carotenoid Oxidation Products Are Stress Signals That Mediate Gene Responses to Singlet Oxygen in Plants. *Proc. Natl Acad. Sci. U. S. A* 109, 5535–5540. [PubMed: 22431637]
- (195). Smirnoff N, and Wheeler GL (2000) Ascorbic Acid in Plants: Biosynthesis and Function. *Crit. Rev. Plant Sci* 19, 267–290.
- (196). Rose RC, and Bode AM (1993) Biology of Free Radical Scavengers: An Evaluation of Ascorbate. *FASEB J* 7, 1135–1142. [PubMed: 8375611]
- (197). Buettner GR, and Jurkiewicz BA (1996) Catalytic Metals, Ascorbate and Free Radicals: Combinations to Avoid. *Radiat. Res* 145, 532–541. [PubMed: 8619018]
- (198). Murad S, Grove D, Lindberg KA, Reynolds G, Sivarajah A, and Pinnell SR (1981) Regulation of Collagen Synthesis by Ascorbic Acid. *Proc. Natl Acad. Sci. U. S. A* 78, 2879–2882. [PubMed: 6265920]
- (199). Rebouche CJ (1991) Ascorbic Acid and Carnitine Biosynthesis. *Am. J. Clin. Nutr* 54, 1147S–1152S. [PubMed: 1962562]
- (200). Wheeler GL, Jones MA, and Smirnoff N (1998) The Biosynthetic Pathway of Vitamin C in Higher Plants. *Nature* 393, 365–369. [PubMed: 9620799]
- (201). Foyer C, Rowell J, and Walker D (1983) Measurement of the Ascorbate Content of Spinach Leaf Protoplasts and Chloroplasts during Illumination. *Planta* 157, 239–244. [PubMed: 24264153]
- (202). Zechmann B (2011) Subcellular Distribution of Ascorbate in Plants. *Plant Signaling Behav* 6, 360–363.
- (203). Noctor G, and Foyer CH (1998) Ascorbate and Glutathione: Keeping Active Oxygen under Control. *Annu. Rev. Plant Physiol. Plant Mol Biol* 49, 249–279. [PubMed: 15012235]
- (204). Lisenbee CS, Lingard MJ, and Trelease RN (2005) Arabidopsis Peroxisomes Possess Functionally Redundant Membrane and Matrix Isoforms of Monodehydroascorbate Reductase. *Plant J* 43, 900–914. [PubMed: 16146528]
- (205). Chew O, Whelan J, and Millar AH (2003) Molecular Definition of the Ascorbate–Glutathione Cycle in *Arabidopsis* Mitochondria Reveals Dual Targeting of Antioxidant Defenses in Plants. *J. Biol. Chem* 278, 46869–46877. [PubMed: 12954611]
- (206). Conklin PL, Saracco SA, Norris SR, and Last RL (2000) Identification of Ascorbic Acid-Deficient *Arabidopsis thaliana* Mutants. *Genetics* 154, 847–856. [PubMed: 10655235]

- (207). Veljovic-Jovanovic SD, Pignocchi C, Noctor G, and Foyer CH (2001) Low Ascorbic Acid in the *vtc-1* Mutant of *Arabidopsis* Is Associated with Decreased Growth and Intracellular Redistribution of the Antioxidant System. *Plant Physiol* 127, 426–435. [PubMed: 11598218]
- (208). Filkowski J, Kovalchuk O, and Kovalchuk I (2004) Genome Stability of *vtc1*, *tt4*, and *tt5* *Arabidopsis thaliana* Mutants Impaired in Protection against Oxidative Stress. *Plant J* 38, 60–69. [PubMed: 15053760]
- (209). Awad J, Stotz HU, Fekete A, Krischke M, Engert C, Havaux M, Berger S, and Mueller MJ (2015) 2-Cysteine Peroxiredoxins and Thylakoid Ascorbate Peroxidase Create a Water-Water Cycle That Is Essential to Protect the Photosynthetic Apparatus under High Light Stress Conditions. *Plant Physiol* 167, 1592–1603. [PubMed: 25667319]
- (210). Toth SZ, Nagy V, Puthur JT, Kovacs L, and Garab G (2011) The Physiological Role of Ascorbate as Photosystem II Electron Donor: Protection against Photoinactivation in Heat-Stressed Leaves. *Plant Physiol* 156, 382–392. [PubMed: 21357184]
- (211). Pavet V, Olmos E, Kiddle G, Mowla S, Kumar S, Antoniw J, Alvarez ME, and Foyer CH (2005) Ascorbic Acid Deficiency Activates Cell Death and Disease Resistance Responses in *Arabidopsis*. *Plant Physiol* 139, 1291–1303. [PubMed: 16244149]
- (212). Mukherjee M, Larrimore KE, Ahmed NJ, Bedick TS, Barghouthi NT, Traw MB, and Barth C (2010) Ascorbic Acid Deficiency in *Arabidopsis* Induces Constitutive Priming That Is Dependent on Hydrogen Peroxide, Salicylic Acid, and the Npr1 Gene. *Mol. Plant-Microbe Interact* 23, 340–351. [PubMed: 20121455]
- (213). Botanga CJ, Bethke G, Chen Z, Gallie DR, Fiehn O, and Glazebrook J (2012) Metabolite Profiling of *Arabidopsis* Inoculated with *Alternaria brassicicola* Reveals That Ascorbate Reduces Disease Severity. *Mol Plant-Microbe Interact* 25, 1628–1638. [PubMed: 23134520]
- (214). Pellaud S, and Mene-Saffrane L (2017) Metabolic Origins and Transport of Vitamin E Biosynthetic Precursors. *Front. Plant Sci* 8, 1959. [PubMed: 29184568]
- (215). Yamauchi R (1997) Vitamin E: Mechanism of Its Antioxidant Activity. *Food Sci. Technol. Int.*, Tokyo 3, 301–309.
- (216). Muller L, Theile K, and Bohm V (2010) In Vitro Antioxidant Activity of Tocopherols and Tocotrienols and Comparison of Vitamin E Concentration and Lipophilic Antioxidant Capacity in Human Plasma. *Mol. Nutr. Food Res* 54, 731–742. [PubMed: 20333724]
- (217). Vidi PA, Kanwischer M, Baginsky S, Austin JR, Csucs G, Dormann P, Kessler F, and Brehelin C (2006) Tocopherol Cyclase (Vte1) Localization and Vitamin E Accumulation in Chloroplast Plastoglobule Lipoprotein Particles. *J. Biol. Chem* 281, 11225–11234. [PubMed: 16414959]
- (218). Fukuzawa K, Tokumura A, Ouchi S, and Tsukatani H (1982) Antioxidant Activities of Tocopherols on Fe²⁺-Ascorbate-Induced Lipid Peroxidation in Lecithin Liposomes. *Lipids* 17, 511–513. [PubMed: 7121213]
- (219). Fahrenholtz SR, Doleiden FH, Trozzolo AM, and Lamola AA (1974) On the Quenching of Singlet Oxygen by α -Tocopherol. *Photochem. Photobiol* 20, 505–509. [PubMed: 4457927]
- (220). Fukuzawa K, Matsuura K, Tokumura A, Suzuki A, and Terao J (1997) Kinetics and Dynamics of Singlet Oxygen Scavenging by α -Tocopherol in Phospholipid Model Membranes. *Free Radical Biol. Med* 22, 923–930. [PubMed: 9119263]
- (221). Galli F, Azzi A, Birringer M, Cook-Mills JM, Eggersdorfer M, Frank J, Cruciani G, Lorkowski S, and Ozer NK (2017) Vitamin E: Emerging Aspects and New Directions. *Free Radical Biol. Med* 102, 16–36. [PubMed: 27816611]
- (222). Porfirova S, Bergmuller E, Tropf S, Lemke R, and Dormann P (2002) Isolation of an *Arabidopsis* Mutant Lacking Vitamin E and Identification of a Cyclase Essential for All Tocopherol Biosynthesis. *Proc. Natl. Acad. Sci. U. S. A* 99, 12495–12500. [PubMed: 12213958]
- (223). Maeda H, Sakuragi Y, Bryant DA, and Dellapenna D (2005) Tocopherols Protect *Synechocystis* sp. Strain PCC 6803 from Lipid Peroxidation. *Plant Physiol* 138, 1422–1435. [PubMed: 15965015]
- (224). Sattler SE, Gilliland LU, Magallanes-Lundback M, Pollard M, and DellaPenna D (2004) Vitamin E Is Essential for Seed Longevity and for Preventing Lipid Peroxidation during Germination. *Plant Cell* 16, 1419–1432. [PubMed: 15155886]

- (225). Sattler SE, Mene-Saffrane L, Farmer EE, Krishke M, Mueller MJ, and DellaPenna D (2006) Nonenzymatic Lipid Peroxidation Reprograms Gene Expression and Activates Defense Markers in Arabidopsis Tocopherol-Deficient Mutants. *Plant Cell* 18, 3706–3720. [PubMed: 17194769]
- (226). Kruk J, Hollander-Czytko H, Oettmeier W, and Trebst A (2005) Tocopherol as Singlet Oxygen Scavenger in Photosystem II. *J. Plant Physiol* 162, 749–757. [PubMed: 16008099]
- (227). Ververidis F, Trantas E, Douglas C, Vollmer G, Kretzschmar G, and Panopoulos N (2007) Biotechnology of Flavonoids and Other Phenylpropanoid-Derived Natural Products. Part I: Chemical Diversity, Impacts on Plant Biology and Human Health. *Biotechnol. J* 2, 1214–1234. [PubMed: 17935117]
- (228). Panche AN, Diwan AD, and Chandra SR (2016) Flavonoids: An Overview. *J. Nutr. Sci* 5, e47. [PubMed: 28620474]
- (229). Winkel-Shirley B (2002) Biosynthesis of Flavonoids and Effects of Stress. *Curr. Opin. Plant Biol* 5, 218–223. [PubMed: 11960739]
- (230). Taylor LP, and Grotewold E (2005) Flavonoids as Developmental Regulators. *Curr. Opin. Plant Biol* 8, 317–323. [PubMed: 15860429]
- (231). Fini A, Brunetti C, Di Ferdinando M, Ferrini F, and Tattini M (2011) Stress-Induced Flavonoid Biosynthesis and the Antioxidant Machinery of Plants. *Plant Signaling Behav* 6, 709–711.
- (232). Gayomba SR, Watkins JM, and Muday GK (2016) Flavonols Regulate Plant Growth and Development through Regulation of Auxin Transport and Cellular Redox Status In Recent Advances in Polyphenol Research (Yoshida K, Cheynier V, and Quideau S, Eds.) John Wiley & Sons: Chichester, U.K.
- (233). Winkel-Shirley B (2001) Flavonoid Biosynthesis. A Colorful Model for Genetics, Biochemistry, Cell Biology, and Biotechnology. *Plant Physiol* 126, 485–493. [PubMed: 11402179]
- (234). Tohge T, Perez de Souza L, and Fernie AR (2018) On the Natural Diversity of Phenylacetylated-Flavonoid and Their in Planta Function under Conditions of Stress. *Phytochem. Rev* 17, 279–290. [PubMed: 29755304]
- (235). Jiang N, Doseff AI, and Grotewold E (2016) Flavones: From Biosynthesis to Health Benefits. *Plants* 5, 27.
- (236). Khan MK, Zill-E-Huma, and Dangles O (2014) A Comprehensive Review on Flavanones, the Major Citrus Polyphenols. *J. Food Compos. Anal* 33, 85–104.
- (237). Liu Y, Tikunov Y, Schouten RE, Marcelis LFM, Visser RGF, and Bovy A (2018) Anthocyanin Biosynthesis and Degradation Mechanisms in Solanaceous Vegetables: A Review. *Front. Chem* 6, 52. [PubMed: 29594099]
- (238). Holton TA, and Cornish EC (1995) Genetics and Biochemistry of Anthocyanin Biosynthesis. *Plant Cell* 7, 1071–1083. [PubMed: 12242398]
- (239). Heim KE, Tagliaferro AR, and Bobilya DJ (2002) Flavonoid Antioxidants: Chemistry, Metabolism and Structure–Activity Relationships. *J. Nutr. Biochem* 13, 572–584. [PubMed: 12550068]
- (240). Pietta PG (2000) Flavonoids as Antioxidants. *J. Nat. Prod* 63, 1035–1042. [PubMed: 10924197]
- (241). Csepregi K, and Hideg E (2018) Phenolic Compound Diversity Explored in the Context of Photo-Oxidative Stress Protection. *Phytochem. Anal* 29, 129–136. [PubMed: 28895264]
- (242). Lewis DR, Ramirez MV, Miller ND, Vallabhaneni P, Ray WK, Helm RF, Winkel BS, and Muday GK (2011) Auxin and Ethylene Induce Flavonol Accumulation through Distinct Transcriptional Networks. *Plant Physiol* 156, 144–164. [PubMed: 21427279]
- (243). Grünwald W, De Smet I, Lewis DR, Lofke C, Jansen L, Goeminne G, Vanden Bossche R, Karimi M, De Rybel B, Vanholme B, Teichmann T, Boerjan W, Van Montagu MC, Gheysen G, Muday GK, Friml J, and Beeckman T (2012) Transcription Factor Wrky23 Assists Auxin Distribution Patterns During Arabidopsis Root Development through Local Control on Flavonol Biosynthesis. *Proc. Natl. Acad. Sci. U. S. A* 109, 1554–1559. [PubMed: 22307611]
- (244). Hichri I, Barrieu F, Bogs J, Kappel C, Delrot S, and Lauvergeat V (2011) Recent Advances in the Transcriptional Regulation of the Flavonoid Biosynthetic Pathway. *J. Exp. Bot* 62, 2465–2483. [PubMed: 21278228]

- (245). Borevitz JO, Xia Y, Blount J, Dixon RA, and Lamb C (2000) Activation Tagging Identifies a Conserved Myb Regulator of Phenylpropanoid Biosynthesis. *Plant Cell* 12, 2383–2394. [PubMed: 11148285]
- (246). Gonzalez A, Zhao M, Leavitt JM, and Lloyd AM (2008) Regulation of the Anthocyanin Biosynthetic Pathway by the Ttg1/Bhlh/Myb Transcriptional Complex in Arabidopsis Seedlings. *Plant J* 53, 814–827. [PubMed: 18036197]
- (247). Teng S, Keurentjes J, Bentsink L, Koornneef M, and Smeekens S (2005) Sucrose-Specific Induction of Anthocyanin Biosynthesis in Arabidopsis Requires the Myb75/Pap1 Gene. *Plant Physiol* 139, 1840–1852. [PubMed: 16299184]
- (248). Solfanelli C, Poggi A, Loreti E, Alpi A, and Perata P (2006) Sucrose-Specific Induction of the Anthocyanin Biosynthetic Pathway in Arabidopsis. *Plant Physiol* 140, 637–646. [PubMed: 16384906]
- (249). Cominelli E, Gusmaroli G, Allegra D, Galbiati M, Wade HK, Jenkins GI, and Tonelli C (2008) Expression Analysis of Anthocyanin Regulatory Genes in Response to Different Light Qualities in *Arabidopsis thaliana*. *J. Plant Physiol* 165, 886–894. [PubMed: 17766004]
- (250). Xu Z, Mahmood K, and Rothstein SJ (2017) ROS Induces Anthocyanin Production Via Late Biosynthetic Genes and Anthocyanin Deficiency Confers the Hypersensitivity to ROS-Generating Stresses in Arabidopsis. *Plant Cell Physiol* 58, 1364–1377. [PubMed: 28586465]
- (251). Shi H, Liu G, Wei Y, and Chan Z (2018) The Zinc-Finger Transcription Factor ZAT6 Is Essential for Hydrogen Peroxide Induction of Anthocyanin Synthesis in *Arabidopsis*. *Plant Mol. Biol* 97, 165–176. [PubMed: 29675814]
- (252). Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, Michael AJ, Tohge T, Yamazaki M, and Saito K (2014) Enhancement of Oxidative and Drought Tolerance in Arabidopsis by Overaccumulation of Antioxidant Flavonoids. *Plant J* 77, 367–379. [PubMed: 24274116]
- (253). Viola IL, Camoirano A, and Gonzalez DH (2016) Redox-Dependent Modulation of Anthocyanin Biosynthesis by the Tcpl Transcription Factor Tcpl5 During Exposure to High Light Intensity Conditions in Arabidopsis. *Plant Physiol* 170, 74–85. [PubMed: 26574599]
- (254). Mahmood K, Xu Z, El-Kereamy A, Casaretto JA, and Rothstein SJ (2016) *The Arabidopsis* Transcription Factor Anac032 Represses Anthocyanin Biosynthesis in Response to High Sucrose and Oxidative and Abiotic Stresses. *Front Plant Sci* 7, 1548. [PubMed: 27790239]
- (255). Wang J, Lian W, Cao Y, Wang X, Wang G, Qi C, Liu L, Qin S, Yuan X, Li X, Ren S, and Guo YD (2018) Overexpression of Bonac019, a Nac Transcription Factor from *Brassica oleracea*, Negatively Regulates the Dehydration Response and Anthocyanin Biosynthesis in Arabidopsis. *Sci. Rep* 8, 13349. [PubMed: 30190519]
- (256). Schoenbohm C, Martens S, Eder C, Forkmann G, and Weisshaar B (2000) Identification of the *Arabidopsis thaliana* Flavonoid 3'-Hydroxylase Gene and Functional Expression of the Encoded P450 Enzyme. *Biol. Chem* 381, 749–753. [PubMed: 11030432]
- (257). Tohge T, Yonekura-Sakakibara K, Niida R, Watanabe-Takahashi A, and Saito K (2007) Phytochemical Genomics in *Arabidopsis thaliana*: A Case Study for Functional Identification of Flavonoid Biosynthesis Genes. *Pure Appl. Chem* 79, 811–823.
- (258). Muzac I, Wang J, Anzellotti D, Zhang H, and Ibrahim RK (2000) Functional Expression of an *Arabidopsis* Cdna Clone Encoding a Flavonol 3'-O-Methyltransferase and Characterization of the Gene Product. *Arch. Biochem. Biophys* 375, 385–388. [PubMed: 10700397]
- (259). Yoder JJ, Belzile F, Tong Y, and Goldsbrough A (1994) Visual Markers for Tomato Derived from the Anthocyanin Biosynthetic Pathway. *Euphytica* 79, 163–167.
- (260). De Jong WS, Eannetta NT, De Jong DM, and Bodis M (2004) Candidate Gene Analysis of Anthocyanin Pigmentation Loci in the Solanaceae. *Theor. Appl. Genet* 108, 423–432. [PubMed: 14523517]
- (261). Yonekura-Sakakibara K, Tohge T, Matsuda F, Nakabayashi R, Takayama H, Niida R, Watanabe-Takahashi A, Inoue E, and Saito K (2008) Comprehensive Flavonol Profiling and Transcriptome Coexpression Analysis Leading to Decoding Gene-Metabolite Correlations in Arabidopsis. *Plant Cell* 20, 2160–2176. [PubMed: 18757557]

- (262). Agati G, Azzarello E, Pollastri S, and Tattini M (2012) Flavonoids as Antioxidants in Plants: Location and Functional Significance. *Plant Sci* 196, 67–76. [PubMed: 23017900]
- (263). Sheahan JJ, and Rechnitz GA (1992) Flavonoid-Specific Staining of *Arabidopsis thaliana*. *BioTechniques* 13, 880–883. [PubMed: 1282347]
- (264). Watkins JM, Hechler PJ, and Muday GK (2014) Ethylene-Induced Flavonol Accumulation in Guard Cells Suppresses Reactive Oxygen Species and Moderates Stomatal Aperture. *Plant Physiol* 164, 1707–1717. [PubMed: 24596331]
- (265). Ringli C, Bigler L, Kuhn BM, Leiber RM, Diet A, Santelia D, Frey B, Pollmann S, and Klein M (2008) The Modified Flavonol Glycosylation Profile in the *Arabidopsis Ro//* Mutants Results in Alterations in Plant Growth and Cell Shape Formation. *Plant Cell* 20, 1470–1481. [PubMed: 18567791]
- (266). Debeaujon I, Peeters AJ, Leon-Kloosterziel KM, and Koornneef M (2001) The Transparent Testa12 Gene of *Arabidopsis* Encodes a Multidrug Secondary Transporter-Like Protein Required for Flavonoid Sequestration in Vacuoles of the Seed Coat Endothelium. *Plant Cell* 13, 853–871. [PubMed: 11283341]
- (267). Shitan N (2016) Secondary Metabolites in Plants: Transport and Self-Tolerance Mechanisms. *Biosci., Biotechnol., Biochem* 80, 1283–1293. [PubMed: 26940949]
- (268). Saslowsky DE, Warek U, and Winkel BS (2005) Nuclear Localization of Flavonoid Enzymes in *Arabidopsis*. *J. Biol. Chem* 280, 23735–23740. [PubMed: 15817473]
- (269). Brown DE, Rashotte AM, Murphy AS, Normanly J, Tague BW, Peer WA, Taiz L, and Muday GK (2001) Flavonoids Act as Negative Regulators of Auxin Transport in Vivo in *Arabidopsis*. *Plant Physiol* 126, 524–535. [PubMed: 11402184]
- (270). Peer WA, and Murphy AS (2007) Flavonoids and Auxin Transport: Modulators or Regulators? *Trends Plant Sci* 12, 556–563. [PubMed: 18198522]
- (271). Kuhn BM, Geisler M, Bigler L, and Ringli C (2011) Flavonols Accumulate Asymmetrically and Affect Auxin Transport in *Arabidopsis*. *Plant Physiol* 156, 585–595. [PubMed: 21502189]
- (272). Maloney GS, DiNapoli KT, and Muday GK (2014) The *anthocyanin reduced* Tomato Mutant Demonstrates the Role of Flavonols in Tomato Lateral Root and Root Hair Development. *Plant Physiol* 166, 614–631. [PubMed: 25006027]
- (273). Fernández-Marcos M, Sanz L, Lewis DR, Muday GK, and Lorenzo O (2013) Control of Auxin Transport by Reactive Oxygen and Nitrogen Species. *Signaling Commun. Plants* 17, 103–117.
- (274). Tsukagoshi H, Busch W, and Benfey PN (2010) Transcriptional Regulation of ROS Controls Transition from Proliferation to Differentiation in the Root. *Cell* 143, 606–616. [PubMed: 21074051]
- (275). Silva-Navas J, Moreno-Risueno MA, Manzano C, Tellez-Robledo B, Navarro-Neila S, Carrasco V, Pollmann S, Gallego FJ, and Del Pozo JC (2016) Flavonols Mediate Root Phototropism and Growth through Regulation of Proliferation-to-Differentiation Transition. *Plant Cell* 28, 1372–1387. [PubMed: 26628743]
- (276). Fuglevand G, Jackson JA, and Jenkins GI (1996) UV-B, UV-A, and Blue Light Signal Transduction Pathways Interact Synergistically to Regulate Chalcone Synthase Gene Expression in *Arabidopsis*. *Plant Cell* 8, 2347–2357. [PubMed: 8989887]
- (277). Buer CS, and Muday GK (2004) The *Transparent Testa4* Mutation Prevents Flavonoid Synthesis and Alters Auxin Transport and the Response of *Arabidopsis* Roots to Gravity and Light. *Plant Cell* 16, 1191–1205. [PubMed: 15100399]
- (278). Joo JH, Bae YS, and Lee JS (2001) Role of Auxin-Induced Reactive Oxygen Species in Root Gravitropism. *Plant Physiol* 126, 1055–1060. [PubMed: 11457956]
- (279). Sierla M, Waszczak C, Vahisalu T, and Kangasjarvi J (2016) Reactive Oxygen Species in the Regulation of Stomatal Movements. *Plant Physiol* 171, 1569–1580. [PubMed: 27208297]
- (280). Mittler R, and Blumwald E (2015) The Roles of ROS and ABA in Systemic Acquired Acclimation. *Plant Cell* 27, 64–70. [PubMed: 25604442]
- (281). Murata Y, Mori IC, and Munemasa S (2015) Diverse Stomatal Signaling and the Signal Integration Mechanism. *Annu. Rev. Plant Biol* 66, 369–392. [PubMed: 25665132]
- (282). Song Y, Miao Y, and Song CP (2014) Behind the Scenes: The Roles of Reactive Oxygen Species in Guard Cells. *New Phytol* 201, 1121–1140. [PubMed: 24188383]

- (283). Takahama U (1988) Oxidation of Flavonols by Hydrogen Peroxide in Epidermal and Guard Cells of *Vicia faba* L. *Plant Cell Physiol* 29, 433–438.
- (284). Schnabl H, Weissenböck D, and Scharf H (1986) *In Vivo* Microspectrophotometric Characterization of Flavonol Glycosides in *Vicia faba* Guard and Epidermal Cells. *J. Exp. Bot* 37, 61–72.
- (285). Weissenböck D, Hedrich R, and Sachs G (1986) Secondary Phenolic Products in Isolated Guard Cell, Epidermal Cell and Mesophyll Cell Protoplasts from Pea (*Pisum sativum* L.) Leaves: Distribution and Determination. *Protoplasma* 134, 141–148.
- (286). Weissenböck D, Schnabl H, Scharf H, and Sachs G (1987) On the Properties of Fluorescing Compounds in Guard and Epidermal Cells of *Allium cepa* L. *Planta* 171, 88–95. [PubMed: 24227274]
- (287). An Y, Feng X, Liu L, Xiong L, and Wang L (2016) ALA-Induced Flavonols Accumulation in Guard Cells Is Involved in Scavenging H₂O₂ and Inhibiting Stomatal Closure in *Arabidopsis* Cotyledons. *Front. Plant Sci* 7, 1713. [PubMed: 27895660]
- (288). Brunetti C, Sebastiani F, and Tattini M (2018) Aba, Flavonols, and the Evolvability of Land Plants. *Plant Sci*, DOI: 10.1016/j.plantsci.2018.12.010.
- (289). Chae HS, Faure F, and Kieber JJ (2003) The *eto1*, *eto2*, and *eto3* Mutations and Cytokinin Treatment Increase Ethylene Biosynthesis in *Arabidopsis* by Increasing the Stability of ACS Protein. *Plant Cell* 15, 545–559. [PubMed: 12566591]
- (290). Liu-Gitz L, Britz SJ, and Wergin WP (2000) Blue Light Inhibits Stomatal Development in Soybean Isolines Containing Kaempferol-3-O-2g-Glycosylgentiobioside (K9), a Unique Flavonoid Glycoside. *Plant, Cell Environ* 23, 883–891.
- (291). Buttery BR, and Buzzell RI (1987) Leaf Traits Associated with Flavonol Glycoside Genes in Soybean. *Plant Physiol* 85, 20–21. [PubMed: 16665658]
- (292). Pollak PE, Vogt T, Mo Y, and Taylor LP (1993) Chalcone Synthase and Flavonol Accumulation in Stigmas and Anthers of *Petunia hybrida*. *Plant Physiol* 102, 925–932. [PubMed: 12231878]
- (293). Stracke R, Jahns O, Keck M, Tohge T, Niehaus K, Fernie AR, and Weissshaar B (2010) Analysis of Production of Flavonol Glycosides-Dependent Flavonol Glycoside Accumulation in *Arabidopsis thaliana* Plants Reveals Myb11-, Myb12- and Myb111-Independent Flavonol Glycoside Accumulation. *New Phytol* 188, 985–1000. [PubMed: 20731781]
- (294). Yonekura-Sakakibara K, Nakabayashi R, Sugawara S, Tohge T, Ito T, Koyanagi M, Kitajima M, Takayama H, and Saito K (2014) A Flavonoid 3-O-Glucoside:2'-O-Glucosyltransferase Responsible for Terminal Modification of Pollen-Specific Flavonols in *Arabidopsis thaliana*. *Plant J* 79, 769–782. [PubMed: 24916675]
- (295). Napoli CA, Fahy D, Wang HY, and Taylor LP (1999) White Anther: A *Petunia* Mutant That Abolishes Pollen Flavonol Accumulation, Induces Male Sterility, and Is Complemented by a Chalcone Synthase Transgene. *Plant Physiol* 120, 615–622. [PubMed: 10364414]
- (296). Mo Y, Nagel C, and Taylor LP (1992) Biochemical Complementation of Chalcone Synthase Mutants Defines a Role for Flavonols in Functional Pollen. *Proc. Natl. Acad. Sci. U. S. A* 89, 7213–7217. [PubMed: 11607312]
- (297). Schijlen EG, de Vos CH, Martens S, Jonker HH, Rosin FM, Molthoff JW, Tikunov YM, Angenent GC, van Tunen AJ, and Bovy AG (2007) RNA Interference Silencing of Chalcone Synthase, the First Step in the Flavonoid Biosynthesis Pathway, Leads to Parthenocarpic Tomato Fruits. *Plant Physiol* 144, 1520–1530. [PubMed: 17478633]
- (298). Taylor LP, and Jorgensen R (1992) Conditional Male Fertility in Chalcone Synthase-Deficient *Petunia*. *J. Hered* 83, 11–17.
- (299). Ylstra B, Touraev A, Moreno RM, Stoger E, van Tunen AJ, Vicente O, Mol JN, and Heberle-Bors E (1992) Flavonols Stimulate Development, Germination, and Tube Growth of Tobacco Pollen. *Plant Physiol* 100, 902–907. [PubMed: 16653074]
- (300). Burbulis IE, Iacobucci M, and Shirley BW (1996) A Null Mutation in the First Enzyme of Flavonoid Biosynthesis Does Not Affect Male Fertility in *Arabidopsis*. *Plant Cell* 8, 1013–1025. [PubMed: 8672888]
- (301). Ylstra B, Muskens M, and Van Tunen AJ (1996) Flavonols Are Not Essential for Fertilization in *Arabidopsis thaliana*. *Plant Mol. Biol* 32, 1155–1158. [PubMed: 9002614]

- (302). Thompson EP, Wilkins C, Demidchik V, Davies JM, and Glover BJ (2010) An Arabidopsis Flavonoid Transporter Is Required for Anther Dehiscence and Pollen Development. *J. Exp. Bot* 61, 439–451. [PubMed: 19995827]
- (303). Pressman E, Peet MM, and Pharr DM (2002) The Effect of Heat Stress on Tomato Pollen Characteristics Is Associated with Changes in Carbohydrate Concentration in the Developing Anthers. *Ann. Bot* 90, 631–636. [PubMed: 12466104]
- (304). Sato S, Kamiyama M, Iwata T, Makita N, Furukawa H, and Ikeda H (2006) Moderate Increase of Mean Daily Temperature Adversely Affects Fruit Set of *Lycopersicon esculentum* by Disrupting Specific Physiological Processes in Male Reproductive Development. *Ann. Bot* 97, 731–738. [PubMed: 16497700]
- (305). Sato S, Peet MM, and Thomas JF (2002) Determining Critical Pre- and Post-Anthesis Periods and Physiological Processes in *Lycopersicon esculentum* Mill. Exposed to Moderately Elevated Temperatures. *J. Exp. Bot* 53, 1187–1195. [PubMed: 11971929]
- (306). Dupuis I, and Dumas C (1990) Influence of Temperature Stress on *in Vitro* Fertilization and Heat Shock Protein Synthesis in Maize (*Zea mays* L.) Reproductive Tissues. *Plant Physiol* 94, 665–670. [PubMed: 16667763]
- (307). Snider JL, and Oosterhuis DM (2011) How Does Timing, Duration and Severity of Heat Stress Influence Pollen-Pistil Interactions in Angiosperms? *Plant Signaling Behav* 6, 930–933.
- (308). Snider JL, Oosterhuis DM, Loka DA, and Kawakami EM (2011) High Temperature Limits *in Vivo* Pollen Tube Growth Rates by Altering Diurnal Carbohydrate Balance in Field-Grown *Gossypium Hirsutum* Pistils. *J. Plant Physiol* 168, 1168–1175. [PubMed: 21256621]
- (309). Wang SS, Zhu XN, Lin JX, Zheng WJ, Zhang BT, Zhou JQ, Ni J, Pan ZC, Zhu SH, and Ding WN (2018) *OsNOX3*, Encoding a NADPH Oxidase, Regulates Root Hair Initiation and Elongation in Rice. *Biol. Plant*, DOI: 10.1007/s10535-018-0814-3.
- (310). Kobayashi M, Ohura I, Kawakita K, Yokota N, Fujiwara M, Shimamoto K, Doke N, and Yoshioka H (2007) Calcium-Dependent Protein Kinases Regulate the Production of Reactive Oxygen Species by Potato NADPH Oxidase. *Plant Cell* 19, 1065–1080. [PubMed: 17400895]
- (311). Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, and Provart NJ (2007) An “Electronic Fluorescent Pictograph” Browser for Exploring and Analyzing Large-Scale Biological Data Sets. *PLoS One* 2, e718. [PubMed: 17684564]

Figure 1. Respiratory burst oxidase homologues (RBOHs) have developmental roles in distinct plant organs and tissues in *Arabidopsis thaliana*. This figure illustrates the plant model species *Arabidopsis thaliana* (left) and highlights plant tissues and developmental responses, in which the different RBOHs are implicated in controlling signaling or development (right). The image of the *Arabidopsis* plant was reprinted with permission from ref 311.

Figure 2.

Tissue and developmental patterns of *Arabidopsis* *RBOH* transcript abundance. The abundance of the transcripts encoding all the *RBOH* enzymes were quantified by microarray and were extracted from the eFP Browser using the developmental map and tissue-specific data sets.³¹¹ The values were normalized relative to the median transcript abundance within the eFP Browser, and the fold changes from that value are reported. Conditional formatting was used within each transcript column to highlight their abundances across tissues and cell types, with values in the lowest 10% shown in dark blue, those in the top 10% in red, and those at the 50th percentile in white. The font color was changed to white for the values in the highest 20%. The locus identifiers for the individual genes are listed at the bottom of the figure.

Figure 3.

The *rhb2-6/rbohC* mutant has impaired root hair elongation compared with the wild-type (Col-0). (A) An *Arabidopsis* root stained with propidium iodide, showing cell outlines found in the meristematic, transition, and elongation zones. Scale bar = 100 μm . (B) Comparison of young *Arabidopsis* root tips for the wild-type (Col-0) and the root-hair-defective (*rhb2*) mutant, which has a mutation in the *RBOHC* gene. The area above the dashed lines indicates the maturation zone, where root hairs form. Scale bar = 200 μm .

Figure 4.

ABA-induced ROS burst in guard cells precedes stomatal closure in wild-type tomato leaves and requires RBOH activity. (A) Increases in DCF fluorescence were visualized in guard cells across a 45 min time course of ABA treatment in wild-type tomato leaves. Scale bar = 5 μm . (B) DCF fluorescence in entire guard cells and stomatal aperture shown as functions of time after treatment with 20 μM ABA. Asterisk and number signs represent significant differences in DCF fluorescence ($P < 0.001$) and stomatal aperture ($P < 0.009$) between time 0 and indicated times or treatment. (C) DCF fluorescence were quantified with and without 100 μM DPI at 0 and 45 min of treatment with ABA. Asterisks and number signs represents significant differences ($P < 0.001$) in DCF fluorescence between time 0 and the indicated times or between untreated and treated, respectively. Statistics were determined using two-way analysis of variance followed by Tukey's posthoc test with $N = 70$. Reproduced with permission from ref 76. Copyright 2017 American Society of Plant Biologists.

Figure 5.

Flavonoid biosynthesis pathway highlighting the six major subclasses of flavonoids. Mutants are noted in parentheses under the enzyme abbreviation, with *Arabidopsis* mutants in italics and tomato mutants in bold italics. The flavonol biosynthesis branch is further detailed in the red box. General flavonoid structures are presented for the six subclasses of flavonoids. The chemical bonds and/or substituents highlighted in red indicate sites of structural variation within each subclass. Abbreviations: CHS, chalcone synthase; CHI, chalcone isomerase; FNS, flavone synthase; F3H, flavanone 3-hydroxylase; IFS, isoflavone synthase; DFR, dihydroflavonol 4-reductase; LAR, leucoanthocyanidin reductase; ANS/LDOX, anthocyanidin synthase/leucoanthocyanidin dioxygenase; OMT, *O*-methyltransferase; (RT), rhamnosyl transferase; UFGT, UDP-glucose flavonoid 3-*O*-glucotransferase; F3'5'H, flavonoid 3'5'-hydroxylase; F3'H, flavonoid 3'-hydroxylase; (FLS), flavonol synthase; OMT-1, *O*-methyltransferase-1.

Figure 6.

ROS levels and root hair numbers are greater in roots of the *are* mutant, which is impaired in flavonol synthesis, compared with the wild-type (VF36). (A) ROS levels are concentrated in the epidermal cells and root hairs of the maturation zone of roots, where root hairs initiate. ROS is elevated in the mutant, as shown in tile-scan confocal images of the *are* mutant and wild-type (VF36) roots stained with CM-H₂DCFDA. Scale bar = 300 μ m. (B) The higher number of root hairs in the *are* mutant than the wild-type is evident in bright-field images of primary roots. Scale bars = 150 μ m. (C) Quantification of root hair numbers in mature regions of primary roots grown on control media. Reproduced with permission from ref 272. Copyright 2014 American Society of Plant Biologists.

Figure 7.

Levels of flavonols are inversely proportional to the levels of ROS in tomato guard cells, with more rapid closure and higher ROS. (A) Confocal micrographs show DPBA-bound flavonols in yellow and chlorophyll autofluorescence in magenta in the *are* mutant with reduced flavonols and the *aw* mutant with increased flavonols compared with the parental lines (VF36 and AC, respectively). Scale bar = 5 μ m. (B) DCF fluorescence is shown in green, and the levels are inversely proportional to flavonol levels. (C) ABA closes guard cells with closure proportional to levels of ROS. (D) Changes in DCF fluorescence upon ABA addition were quantified and are greater in the *are* mutant with low flavonols than the wild-type, while elevated flavonols in the *aw* mutant show the lowest rate of DCF changes. Reproduced with permission from ref 76. Copyright 2017 American Society of Plant Biologists.

Figure 8.

Flavonols regulate pollen development by scavenging ROS. (A) Confocal micrographs of the wild-type tomato (VF36) and the flavonol-deficient *anthocyanin reduced* (*are*) mutant show reduced pollen viability in the mutant. Pollen grains were stained with FDA (live pollen grains, shown in green) and PI (dead pollen grains, shown in magenta). Scale bar = 200 μm . (B) Bright-field images of pollen tube tips of VF36 and the *are* mutant show that flavonols reduce swelling of the pollen tube tip. Scale bar = 10 μm . (C) Confocal micrographs of VF36 and *are* mutant pollen grains and tubes stained with the general ROS sensor CM-H₂DCFDA show that flavonols reduce ROS levels (as evidenced by DCF fluorescence) in pollen grains and tubes. Scale bars = 50 μm . (D) Confocal micrographs of VF36 and *are* mutant pollen grains stained with the hydrogen peroxide-specific sensor peroxy orange 1 (PO1) show elevated hydrogen peroxide in the flavonol-deficient *are* mutant. Scale bar = 50 μm . These images were reprinted from ref 96.

Table 1.

Arabidopsis Respiratory Burst Oxidase Members^a

Gene name Locus ID	Localization	Mutant phenotype	Regulation	Orthologues	References
AtRBOHA AT5G07390	Primary Root: Maturation zone (endodermis, stele) Lateral root primordia Basal meristem of primary root	Unknown	Unknown		39,44
AtRBOHB AT1G09090	Maturation zone in primary root Seed embryo	Faster germination of fresh seeds	ABA		39,104
AtRBOHC/RHD2 AT5G51060	Root epidermal cells Root hairs Columella, elongation, and differentiation zone in primary root	Short root hairs Reduced superoxide production at root hair apoplast	Transcription: RSL4 Subcellular localization: FER, ROP2, SCN1 Protein modification: CBL1, CIPK26	OsNOX3 (rice) Zmrtb5 (maize)	44,45,47,51,52,56,59,60,66,67,309
AtRBOHD AT5G47910	Root maturation zone Lateral root primordia	Reduced primary root elongation Increased lateral root number* Reduced ABA-mediated stomatal closure* Reduced pathogen response (flg22) Resistance to ozone stress* Reduced survival under salt and hypoxia stresses Increased survival under salt or hypoxia stress* Response to iron deficiency, heat, high aluminum, flooding, high light, and drought	Protein modification: BIK1, PBL1, FLS2, ERF74 Alternate binding: PA, Ca ²⁺	SlRBOHB (potato) NbrBOHB	33,34,41,42,44,48,105,107–109,112–114,118,119,128,310
AtRBOHE AT1G19230	Tapetum Root: Basal meristem Cortex Endodermis Epidermal cells overlying lateral root primordia	Delayed degeneration of tapetum Reduced pollen viability Abnormal pollen grain shape and exine layer Delayed emergence of lateral roots	Transcription: MYB80 and AMS (tapetum), NAA (roots)		44,90
AtRBOHF/SGN4 AT1G64060	Periphery of lateral root primordia Elongation to lateral root formation zone in the primary root Vascular tissue in leaves Guard cells	Delayed Casparian strip formation Reduced primary root elongation Increased lateral root number* Reduced ABA mediated guard cell closure* Resistance to ozone stress* Increased survival under salt or hypoxia stress*	Protein modification: CIPK26 Alternate binding: PA		33,34,39,41,42,44,79,85–87,107,108,111
AtRBOHG AT4G25090	Mature roots	Unknown	Unknown		311
AtRBOHH AT5G60010	Pollen	Reduced pollen tube growth and integrity*	ANXUR1/ANXUR2 Ca ²⁺ Phosphorylation		92–94

Gene name Locus ID	Localization	Mutant phenotype	Regulation	Orthologues	References
AIRBOHI AT4G11230	Seeds Leaves Roots	Delayed germination under drought stress conditions Decreased survival under hypoxia stress	Unknown		311
AIRBOHJ AT3G45810	Pollen	Reduced pollen tube growth and integrity*	ANXUR1/ANXUR2 and Ca ²⁺ Phosphorylation		92–94

^aCommonly used gene names and locus numbers are shown along with expression patterns and phenotypes of mutants. Known regulatory events are listed. Orthologues in other species are included. Asterisks indicate phenotypes in *rbohD* for *rbohH/j* double mutants.