Published in final edited form as:

Trends Parasitol. 2019 May ; 35(5): 336–338. doi:10.1016/j.pt.2019.03.006.

Robbing Host Phosphatidic Acid to Survive: A Strategy of a Fly Parasite

Isabelle Coppens^{1,*}

¹Department of Molecular Microbiology and Immunology, Johns Hopkins University Bloomberg School of Public Health, 615 North Wolfe Street, Baltimore, MD 21205, USA

Abstract

Drosophila flies can be infected by an obligate fungal intracellular parasite, *Tubulinosema ratisbonensis*, resulting in a swollen abdomen and often death. Within the fly, the parasite multiplies in the cytoplasm of adipocytes of the fat body, feeds on host lipid droplets, and has a specific requirement for dietary phosphatidic acid.

Parasites love fat. Parasites deviously scavenge and consume lipids from their hosts. And sometimes, it costs the infected host its life. A prime example is illustrated by the obligate intracellular parasite *T. ratisbonensis* that infects and proliferates in *Drosophila melanogaster* – our major model system in modern biology, resulting in a shortened lifespan and reduced fertility for the infested fly. If kept at 25°C and left uncured, infested *Drosophila* stocks in laboratories can be devastated within a week. To explore strategies to cure or protect *Drosophila* cultures from *Tubulinosema* infection, a better understanding of the systemic effects of an infection on the fly and the identification of host factors exploited by the parasite allowing its successful proliferation, is crucial. By dissecting *Drosophila* metabolic networks through selective feedings, or the silencing of specific metabolic genes, Franchet and colleagues [1] have identified a major contributor to the rapid growth of *Tubulinosema* in the lipid metabolism of the fly.

T. ratisbonensis belongs to the phylum Microspora that comprises spore-forming, intracellular protist parasites that infect vertebrates and invertebrates, even those that are themselves parasites. Microsporidia produce environmentally resistant spores that are equipped with a sophisticated invasion machinery, consisting of an extrusion apparatus that contains a coiled polar tube, that is, a long, flexible, syringe-like structure [2]. During host cell invasion, the polar tube explosively extrudes from the spore, pierces the targeted host cell plasma membrane, and inoculates the host cell with infective sporoplasm, that is, the central mass of cytoplasm within the spore. Then, intracellular sporoplasm matures into a meront that undergoes several rounds of asexual reproduction, leading to the generation of up to several hundred meronts in a single host cell.

As with any microsporidian, *T. ratisbonensis* has lost many essential genes in response to its adaptation to the lifestyle of intracellular parasitism [3]. While it possesses a nucleus and

*Correspondence: icoppen1@jhu.edu (I. Coppens).

intracytoplasmic membranous system, it lacks many typical eukaryotic characteristics. For example, the parasite has ribosomes that resemble those of prokaryotes, a primitive mitochondrion (mitosome), an atypical Golgi apparatus, and no peroxisomes. Its genome lacks many genes involved in fatty acid anabolism (e.g., acetyl-CoA carboxylase, fatty acid synthase) and fatty acid β -oxidation (e.g., acyl-CoA oxidase). Reduced to host dependency and scavenging, *Tubulinosema* relies on its own transporter activities to steal host metabolites during its intracellular life stage.

A hallmark of *Drosophila* infection with *Tubulinosema* is the impressive swelling of the fly's abdomen (Figure 1A,B). Within the fly, *Tubulinosema* preferentially proliferates in cells of the fat body, a large organ distributed throughout the fly body (Figure 1C). The fat body plays a major role in intermediate metabolism in insects and is the central storage depot of nutrients and energy reserves [4]. The predominant cell of the fat body is the adipocyte, characterized by the accumulation of numerous lipid droplets (LDs) containing triacylglycerols (TAGs). By targeting a fat body adipocyte (the storage sanctuary) as a host cell to establish infection, the parasite secures for itself a steady and copious supply of nutrients. The choice of adipocytes as sites of multiplication for *Tubulinosema* in the fly gives us a first clue of the parasite's need from the host: likely a biosynthetic or stored molecule, for example, glucose, free fatty acid, glycogen, and/or TAGs. Indeed, as *T. ratisbonensis* meronts expand in the fat body, the LDs of infected adipocytes progressively dissolve, as a consequence of their consumption by the parasite. This 'cellular vampirism' results in a dramatic collapse of the entire fat body lobule structure followed by the fly's death.

Which nutrients, present in adipocytes, does *Tubulinosema* covet? As a first approach to solving this enigma, the authors gave a super-rich meal (yeast) to the infected flies and observed an acceleration of the parasite's proliferation. This confirms a relationship between *Tubulinosema* virulence and a high nutrient density diet of the fly. As a next approach to decipher which particular biochemical compounds coming from the food are responsible for this phenomenon, the authors selectively added amino acids, sterols, fatty acids, or sugars to the standard sucrose solution on which flies feed. Only supplementation with fatty acids, for example, oleate and palmitate, led to a doubling of parasite load. This information leads to a second clue on the parasite's requirement for growth by pointing to polar lipids as the desired nutrients for meronts.

As a third approach to refine which host lipid metabolic processes are required for parasite proliferation, the authors silenced specific genes involved in fatty acid synthesis, transport or β -oxidation, lipid droplet biogenesis or degradation, and regulation of lipid metabolism. Reduced expression of a host intestinal TAG lipase involved in the digestion of dietary TAG impairs parasite replication, establishing that dietary intake of lipids contributes to the virulence of the infection. Markedly, silencing key enzymes involved in the synthesis of phosphatidic acid from fatty acids to its direct glycerol phosphate precursors negatively impacts the meront's growth. Conversely, preventing the use of phosphatidic acid by the fly through blockade of phosphatidylethanolamine, phosphatidylcholine, or TAG synthesis favors the proliferation of *Tubulinosema*. When phosphatidic acid is directly injected into the fly, the parasite load even quadruples. These assays offer the final clue to the parasite's

nutritional need from the host by narrowing down the list of dietary lipids hijacked by the parasite to phosphatidic acid.

Why and how phosphatidic acid magically boosts the proliferation of *T. ratisbonensis* in *Drosophila* needs to be investigated. Phosphatidic acid is a peculiar lipid, being both a key building block of phospholipid synthesis and a major second messenger conveying signaling information [5]. With its coneshaped geometry, phosphatidic acid locally changes membrane topology and plays a role in membrane trafficking and fusion events. It interacts with numerous proteins, including kinases and phosphatases, leading to pleiotropic functions. Many protozoan parasites require phosphatidic acid for their growth. To this point, phosphatidic acid is both synthesized and salvaged from mammalian cells by the intravacuolar parasite *Toxoplasma gondii*, and this lipid plays a central role in the parasite's cell cycle and infectivity, controlling invasion and egress from the host cell [6,7].

How *Tubulinosema* accesses host phosphatidic acid or other lipids from the LDs of adipocytes needs further investigation. Like *T. ratisbonensis*, several pathogens that infect vertebrates find a home in adipose tissues, multiply in adipocytes, feeding on or exploiting host LDs as part of anti-immunity strategies [8,9]. Mammalian adipocytes subsist for up to 10 years, providing thus a long-term environment for a pathogen, and they offer a copious source of nutrients, thereby presenting many advantages to pathogens. Adipose tissues are targeted by *Trypanosoma cruzi*, *Trypanosoma brucei*, and *Plasmodium falciparum* and constitute a sequestration site during their chronic phase of infection in humans. Many intracellular pathogens such as *T. cruzi*, *Leishmania amazonensis*, and *T. gondii* internalize host LDs into their parasitophorous vacuole, likely to benefit from their neutral lipid content. *T. gondii* retrieves fatty acids stored in host LDs, and the depletion of LDs from the host cell is detrimental for the parasite's growth [10]. Based on these examples, interference with host LD-pathogen interactions could be harmful for the intruder, leading to its death by starvation.

Acknowledgments

The author is grateful to Julia Romano from Johns Hopkins University for her critical reading of the manuscript. The images of the figure have been kindly provided by Dr D. Ferrandon and Adrien Franchet from the Université of Strasbourg, France.

References

1. Franchet A (2019) Phosphatidic acid as a limiting host metabolite for the proliferation of the microsporidium *Tubulinosema ratisbonensis* in *Drosophila* flies. *Nat. Microbiol* 4, 645–655 [PubMed: 30692666]
2. Franzen C (2004) Microsporidia: how can they invade other cells? *Trends Parasitol.* 20, 275–279 [PubMed: 15147678]
3. Nakjang S et al. (2013) Reduction and expansion in microsporidian genome evolution: new insights from comparative genomics. *Genome Biol. Evol* 5, 2285–2303 [PubMed: 24259309]
4. Arrese EL and Soulages JL (2010) Insect fat body: energy, metabolism, and regulation. *Annu. Rev. Entomol* 55, 207–225 [PubMed: 19725772]
5. Tanguy E et al. (2019) Phosphatidic acid: from pleiotropic functions to neuronal pathology. *Front. Cell. Neurosci* 13, 2 [PubMed: 30728767]

6. Bullen HE et al. (2016) Phosphatidic acid-mediated signaling regulates microneme secretion in *Toxoplasma*. *Cell Host Microbe*, 19, 349–360 [PubMed: 26962945]
7. Bisio H et al. (2019) Phosphatidic acid governs natural egress in *Toxoplasma gondii* via a guanylate cyclase receptor platform. *Nat. Microbiol* 4, 420–428 [PubMed: 30742070]
8. Tanowitz HB et al. (2017) A safe haven for parasites? *Trends Parasitol.* 33, 276–284 [PubMed: 28007406]
9. Saka HA and Valdivia R (2012) Emerging roles for lipid droplets in immunity and host–pathogen interactions. *Annu. Rev. Cell Dev. Biol* 28, 411–437 [PubMed: 22578141]
10. Nolan SJ et al. (2017) Host lipid droplets: An important source of lipids salvaged by the intracellular parasite *Toxoplasma gondii*. *PLoS Pathog.* 13, e1006362 [PubMed: 28570716]

Figure 1. Infection and Colonization of *Drosophila* by *Tubulinosema ratisbonensis*.

(A) Illustration of the normal morphology of an uninfected *Drosophila*. By contrast, upon infection of the fly with *T. ratisbonensis* for 9 days as shown in (B), the fitness and body corpulence of the fly is dramatically altered, with a distended, swollen abdomen. *T. ratisbonensis* meronts preferentially target host adipocytes wherein they proliferate and progressively fill the fat body lobule of the fly. (C) Illustration of the area of fat body adipocytes, enriched in lipid droplets (red), that is being invaded. Most of the DAPI signal (blue) originates from the meronts. The yellow staining corresponds to mature spores of the parasites. FITC-actin is shown in green. The figure is courtesy of the D. Ferrandon group from the Université of Strasbourg, France.