Original Article

Screening for dietary fat intake of grade six children: self-assessment vs. maternal assessment

Friede Wenhold*, Una MacIntyre* and Paul Rheeder†

*Department of Human Nutrition, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa, and †Division Clinical Epidemiology, School of Health Systems and Public Health, University of Pretoria, Pretoria, South Africa

Abstract

As part of justifiable nutrition promotion, this study aimed to determine internal consistency of a dietary fat screener and to compare self-assessment to maternal assessment of fat intake of grade six (about 12 years old) learners in a South African public primary school. The children completed in school a pictorial, quantitative food frequency-type screener consisting of 10 high-fat food categories; mothers individually completed a text version. Internal consistency was measured with item-total correlations, Cronbach's alpha and the split-half method. Child-mother comparison was based on kappa (κ) statistics, McNemar's tests, Wilcoxon signed-rank test and the Bland-Altman method. In total, 101 (93.5%) children and 78 (72.2%) mothers responded. The screener was internally consistent, regardless of data source and statistical technique. For portion sizes and frequency of intake, children consistently reported higher intake than mothers. This resulted in systematic error, also evidenced by a significant difference from zero for the difference between child's and mother's final test scores for the whole group, and for boys and girls separately (always $P < 0.001$). In 76% of the pairs, classification into high fat or prudent intake was identical, yet the chance-corrected agreement was poor ($\kappa = 0.16$) and non-agreement was non-symmetrical ($P = 0.001$). Children and mothers reported high fat intakes (93% and 75%, respectively). It was concluded that the dietary fat screener was internally consistent, yet children and mothers did not agree in their assessment. The high fat intakes reported by children and mothers warrant measurement refinement and implementation of primary prevention programmes.

Keywords: child, dietary assessment, internal consistency, fat intake, mother, agreement.

Correspondence: Friede Wenhold, Department of Human Nutrition, Faculty of Health Sciences, University of Pretoria, Private Bag x323, Arcadia, 0007 Pretoria, South Africa. E-mail: friede.wenhold@up.ac.za

Introduction

South Africa has a high burden of non-communicable diseases (Mayosi *et al.* 2009), most of which can be attributed to high blood cholesterol (Norman *et al.* 2007). High dietary fat intakes are a major risk factor for hypercholesterolaemia (Van Horn *et al.* 2008). Dietary habits originate in childhood and may track into adulthood [American Dietetic Association (ADA) 2004], pointing to the importance of early

prevention, for example, in primary school children. Furthermore, the existing relatively high prevalences of overweight and obesity, especially among white South African school children (Armstrong *et al.* 2006), appear to be increasing (Armstrong *et al.* 2011).

Focused dietary prevention programmes should be planned around current intakes (ADA 2004). This requires food-based dietary assessment. Usually, children between the ages of 8 and 12 can

perform their own dietary assessment (McPherson *et al.* 2000; Burrows *et al.* 2010) because their cognitive abilities are adequately developed and because this age group makes many own and out-of-home food choices. In the case of the assessment of dietary fat intake specifically, it is, however, possible that such an assessment may not capture the full picture. While some sources of fat in the diet may be obvious, others require at least some specialised knowledge of foods, food preparation and food composition (in terms of brands and food labels) in order to differentiate between, for example, different types of dairy products, table fats and cooking methods, all of which children may not have. Also, children rely on their caregivers to provide much of their food intake (Livingstone & Robson 2000; McPherson *et al.* 2000).

The school setting offers numerous opportunities for nutrition promotion (Wenhold *et al.* 2008) – including the assessment of dietary intakes – but then this should be quick and easy, suggesting a screening approach. There are no validated dietary fat screeners available in South Africa, especially not for children. The MEDFICTS dietary assessment questionnaire, referring to the intake of meat, eggs, dairy, fried foods, fats in baked goods, convenience foods, table fats and snacks – the main contributors to fat intake – is widely used and has been validated in various US adult populations (Schwartz *et al.* 2000; Kris-Etherton *et al.* 2001; Taylor *et al.* 2003; Holmes *et al.* 2005; Teal *et al.* 2007; Mochari *et al.* 2008). Its performance in children and other contexts is however unknown. As a starting point for determining this, the question arises whether the information obtained from children and their caregivers in a South African urban setting is comparable.

This paper has two objectives with respect to a modified ('South Africanised') MEDFICTS dietary fat screener for grade six children: first, to determine the internal consistency of the screener when completed by the children and by their mothers and, second, to compare a self-assessment to a maternal assessment in relation to reported portion sizes and frequency of intake and the resultant screener scores and classifications and implied fat intake.

Internal consistency reflects the extent to which individual items in a test (in our case, the food categories in the screener) measure similar characteristics (in our case, dietary fat intake) (based on Higgins & Straub 2006). Hence, should the screener be internally consistent, it would contribute to the reliability and validity thereof. Reliability is understood to be the extent to which the screener yields the same results on repeated trials, and validity addresses the question how well the screener measures what it is intended to measure. Valid measurements require consistency of observation, with reliability being a necessary but not a sufficient condition for validity. Random errors show up as different results for ostensibly the same measurement and are associated with lack of reliability. Systematic errors (bias) push the result in the same direction, thereby negatively affecting validity (based on Higgins & Straub 2006).

Materials and methods

The study was an observational investigation. The study population comprised all 108 learners of three grade six classes of a middle-class, Afrikaans medium public primary school in Pretoria, the capital city of South Africa. The racial/ethnic composition of the study population does not reflect the South African

Key messages

- The dietary fat screener was internally consistent, when completed by grade six children and by their mothers.
- Mothers and their 12-year-old children did not agree about the fat intake of the child; there was variability and systematic error. Mothers reported lower frequencies of intake and, in particular, smaller portion sizes of high-fat food categories.
- Assessments by children and their mothers suggest high fat intakes. Implementation of nutrition promotion programmes for school children and refined measurement techniques for fat intake are needed.

demographics, but refers to white children. Reporting accuracy of energy intake among children may be related to race/ethnicity (Forrestal 2011), and white South African children, a minority group, appear to be more affected by overweight and obesity (Armstrong *et al.* 2006). All learners who returned signed informed consent and assent forms were included in the sample.

The dietary fat screener

A 'South Africanised' MEDFICTS was developed. For each food category, example foods typically consumed by the target group were identified by a team of dietitians often working with school children; the fat content of these foods was checked against the South African food composition database (Wolmarans *et al.* 2010) and food wrappers. These example foods were listed as part of the tool. For the meat category, only the high fat group was retained from the original MEDFICTS because of local availability and assumed recognisability by the target group. A composite image containing photographs of all food items included in the category was created for each food category. Based on a number of unpublished developmental evaluation studies in a school similar to the one in this study, the reference portion size was age-adjusted and appropriate portion size estimation aids were identified.

Data collection

For the *self-assessment*, each grade six class was divided into three groups of about 12 children, resulting in a total of nine groups. Data collection was done per group by one researcher (FW), in the mother tongue of the children, in the conference room of the school. A U-shaped seating arrangement was used, with separators between adjacent participants to ensure privacy of response. Administration was standardised following pre-testing. It was researcher-paced and supported by visual aids. In an attempt to assess usual intake, children were requested to record their typical ('normal') eating habits since the beginning of the year (i.e. 'since you were in grade six'), making the reference period about 9 months. It was assumed that

this explicit time frame would be meaningful for South African school children as it coincided with a cognitive reality, that is, the school year.

For each food category, an explanatory sentence was made and the full-colour composite photograph of all the relevant foods was projected on a screen. This was to enhance visual appeal and accuracy (Wallen *et al.* 2011). Every food was mentioned and distinguishing features of the food category (e.g. *full fat/whole* milk, *'paper-wrapped'* margarines and *'lite/diet/fat-free'* labels) were highlighted. A filter question 'Do you eat foods such as those on the picture?' started the questioning per food category. Participants who responded 'no' to the filter question were requested to put down their pens and only proceed when instructed to do so. Correct answering technique was demonstrated on a poster replica of the answer sheet.

Typical frequency of intake was requested next: 'If you eat foods like those on the picture everyday, write down the number of times you usually eat it during one day on the line saying "per day". If you do not eat such food everyday, move to the "per week" line, and fill in there how often in the course of a week you usually eat such foods. Only write in the "per day" or the "per week" line.' A practical example of how to answer this was given, including a child-friendly 'case study' where more than one food item from a food category was eaten in a day.

The last question for each food category dealt with usual amount consumed. For this purpose, a combination of two-dimensional portion size estimation aids (geometric shapes, e.g. a 90-mm-diameter circle for meat), household measures (cups and spoons, e.g. a 250-mL measuring cup for milk) and photographs (e.g. chocolate, nuts, chips and high fat crackers), was used. Children were instructed to mark '2' if the amount usually eaten was similar, '1' if it was about half as much and '3' if it was one and a half times as much as the reference amount, respectively, corresponding to medium, small and large portion sizes. Total contact time per group was about 45 min.

Weight and height of the children were measured with a digital scale (Tanita, Tokyo, Japan; precision 100 g) and portable height gauge (to the nearest

0.1 cm) using standard techniques (Lee & Nieman 2010) by one measurer, in a private room.

For the *maternal assessment*, the text version of the dietary fat screener was sent to the child's home together with the information letter and the informed consent form. A direct caregiver was requested to complete the screener with respect to the grade six children; hence, 'maternal' in this study refers to a primary caregiver.

Data analysis

The *scoring* of the dietary fat screener was as follows. If frequency of intake of a food category was reported as daily consumption, this was converted to weekly consumption by multiplying by 7. Weekly intake was categorised and scored as specified in the original MEDFICTS tool (Kris-Etherton *et al.* 2001). Non-consumption and intakes less than once per week were scored zero; once or more (up to three times per week) scored three points; more than three times per week was scored seven. When ranges (e.g. 1–2 times) were reported, this was coded as the midpoint (e.g. 1.5 times). Small, medium or large usual portions were, respectively, scored one, two or three. Category scores were the product of the frequency and the portion score. All 10 category scores were added to create a final score. The final score, which could range from 0 to 210, was categorised as 'high fat' if it was more than 68. Otherwise, it was classified as 'prudent' as healthy children (as opposed to, e.g. people in cardiac rehabilitation) constituted our population and because we did not include a low-fat meat food category. Description of central tendency included the mode (the most frequently reported value) and the median (the value above and below which half of the values fell).

Internal consistency was determined by item-total correlations (Pearson) between all 10 category scores and the final scores, Cronbach's coefficient alpha and the split-half method, whereby the 10 food categories were randomly assigned to two groups, and Pearson's correlation coefficient was calculated between the mean category scores of the groups.

The comparison of the *self-assessment to the maternal assessment* involved determining the percentage

of pairs with perfect (= identical) agreement, the kappa statistic (κ) to estimate chance-corrected proportional agreement of categorical variables (portion size, categorised frequency of intake, categorised final score) interpreted according to the guidelines suggested by Altman (1991), McNemar's test for symmetry, Spearman's correlation coefficients, Wilcoxon signed-rank test for assessing the significance of the difference between children's and mothers' final scores, and the Bland–Altman method for assessing agreement (Altman 1991).

Data were analysed using SAS (mainframe version 8.2, SAS Institute Inc., Cary, NC, USA) and BMDP statistical software release 7.1 (BMDP Statistical Software Inc., Los Angeles, CA, USA). For the anthropometric description of the participants, the Centers for Disease Control and Prevention (CDC) 2000 growth data files for boys and girls aged 2–20 years (http://www.cdc.gov/growthcharts/cdc_charts.htm) and the accompanying SAS software were used for describing mean age, weight, height and body mass index (BMI, in kg m^{-2}) z-scores (<http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm>). Age was calculated in months based on the actual date of assessment and date of birth as obtained from school records.

Ethics

The Research Ethics Committee of the Faculty of Health Sciences (University of Pretoria) approved the project. Permission to do the study was obtained from the relevant school authorities, and informed maternal consent and child assent were obtained. No incentives, apart from a (unannounced) pen and a snack for the children following the data collection, were offered.

Results

Sample

Maternal consent, assent and complete screener data were obtained from 101 (93.5%) children. A total of 78 (72.2%) caregivers returned a completed dietary fat screener for their child. Of these, six contained at

Table 1. Anthropometric description of sample (mean $\pm$ standard deviation) ($n=78$)

	Age (months)	Weight (kg)	Weight-for-age z-score*	Height (m)	Height-for-age z-score*	BMI (kg m ⁻²)	BMI-for-age z-score*
Boys ($n=36$)	148.7 $\pm$ 4.3	49.1 $\pm$ 15.3	0.36 $\pm$ 1.3	1.52 $\pm$ 0.08	-0.01 $\pm$ 0.9	21.1 $\pm$ 5.3	0.51 $\pm$ 1.2
Girls ($n=42$)	147.7 $\pm$ 3.5	48.8 $\pm$ 10.2	0.43 $\pm$ 0.9	1.55 $\pm$ 0.06	0.19 $\pm$ 0.8	20.3 $\pm$ 3.4	0.42 $\pm$ 0.9
Total ($n=78$)	148.2 $\pm$ 3.9	48.9 $\pm$ 12.7	0.40 $\pm$ 1.1	1.53 $\pm$ 0.07	0.09 $\pm$ 0.9	20.7 $\pm$ 4.4	0.47 $\pm$ 1.1

*Reference data: CDC 2000 (National Center for Health Statistics). Available at: http://www.cdc.gov/growthcharts/cdc_charts.htm.

Table 2. Internal consistency coefficients r of the dietary fat screener for child's self-assessment and maternal assessment

Internal consistency measure	Food category	Child's self-assessment ($n=101$)		Maternal assessment ($n=78$)	
		r	P	r	P
Item-total correlation*	Meat	0.55	<0.001	0.48	<0.001
	Eggs	0.41	<0.001	0.09	0.47
	Dairy, milk, high fat	0.54	<0.001	0.63	<0.001
	Dairy, cheese, high fat	0.66	<0.001	0.50	<0.001
	Dairy, dessert, high fat	0.55	<0.001	0.64	<0.001
	Fried foods	0.38	<0.001	0.52	<0.001
	In baked goods	0.64	<0.001	0.61	<0.001
	Convenience foods	0.51	<0.001	0.54	<0.001
	Table fats, high fat	0.35	<0.001	0.50	<0.001
	Snacks, high fat	0.65	<0.001	0.59	<0.001
Cronbach's alpha	All category scores included	0.70	<0.001	0.69	<0.001
	One category deleted†	0.72	<0.001	0.71	<0.001
Split halves*		0.57	<0.001	0.48	<0.001

*Pearson. †Deletion: For children: table fats; For mothers: eggs.

least one missing value and were not included in the child–mother final score comparison because the final scores would reflect less than 10 category scores. Table 1 summarises the anthropometric information of the children for whom self-assessment and maternal fat screeners were available. The positive mean BMI-for-age z -scores confirm the importance of monitoring weight gain in this group.

Internal consistency

In Table 2, the item-total correlations, Cronbach's alpha and the correlations between two random halves of the fat screener are given for the child's self-assessment and for the maternal assessment. Item-total correlations for children ranged from $r=0.35$ (table fats) to $r=0.66$ (cheese). When excluding eggs, the range for mothers was slightly narrower [$r=0.48$ (meat) to $r=0.63$ (milk)].

Comparison of child's self-assessment to maternal assessment

Portion sizes

For all 10 food categories, the medium size was most commonly chosen by the mothers. In the case of the children, the large portion was most frequently selected for dessert, fried foods, convenience foods and table fats. For the rest of the food categories, most of the children also chose the medium option (Table 3). The percentage perfect agreements between mothers and children in terms of the reported usual portion size eaten by the children varied from as low as 18% for fried foods to about 72% for eggs (Table 3), with an average percentage agreement over the 10 food categories of 45%. Chance-corrected agreement based on kappa values was moderate for eggs and fair for cheese. For all other food categories, agreement was poor. Most

Table 3. Child and maternal assessment of portion size and frequency of intake for all food categories ($n = 77$)

Food categories	Portion size*				Frequency of intake			
	Mode portion according to...		Child vs. mother		Median weekly intake according to...		Child vs. mother†	
	...child	...mother	Identical (% pairs)	Kappa statistic‡	...child	...mother	Identical (% pairs)	Kappa statistic‡
	Portion (% children)	Portion (% mothers)		κ	Times per week (range)	Times per week (range)	κ	P
Meat	M (57)	M (64)	46.7	0.07 (Poor)	5 (0–35)	5 (2–11)	0.07 (Poor)	0.36
Eggs	M (51)	M (54)	71.6	0.44 (Moderate)	1.5 (0–7)	2 (1–7)	0.22 (Fair)	0.01
Dairy, milk, high fat	M (44)	M (57)	49.3	0.17 (Poor)	12 (0–49)	5.5 (1–14)	0.21 (Fair)	0.01
Dairy, cheese, high fat	M (63)	M (64)	63.2	0.30 (Fair)	3.5 (0–21)	2.5 (1–7)	0.32 (Fair)	0.00
Dairy, dessert, high fat	L (72)	M (61)	31.7	0.05 (Poor)	1.5 (0–49)	0.5 (1–7)	0.00 (Poor)	0.98
Fried foods	L (79)	M (66)	18.4	–0.08 (Poor)	2.5 (0–28)	2.5 (1–7)	0.21 (Fair)	0.03
In baked goods	M (59)	M (68)	53.7	0.14 (Poor)	2.5 (0.5–35)	1.5 (0.5–7)	0.05 (Poor)	0.56
Convenience foods	L (52)	M (76)	39.7	0.03 (Poor)	2.75 (0–42)	0.75 (0.5–5)	–0.06 (Poor)	0.38
Table fats, high fat	L (44)	M (53)	29.2	–0.02 (Poor)	10.5 (0–63)	2.75 (0.5–20)	0.05 (Poor)	0.34
Snacks, high fat	M (51)	M (63)	46.7	0.14 (Poor)	4.5 (0–28)	1.75 (1–7)	0.18 (Poor)	0.10

*Portion size: S, small; M, medium; L, large. †Scored frequency of weekly consumption: Rarely or never = 0; up to three times = 3; more than three times = 7. ‡Kappa statistic for chance-corrected agreement. §McNemar's test for symmetry.

non-agreeing responses tended to be non-symmetrical (see McNemar's results).

Frequency of intake

From Table 3, it is evident that the percentage child–mother pairs that reported identical frequency of intake ranged from 42% for convenience foods to 74% for milk. This represents an average of 60% across all 10 food categories. For 4 of the 10 food categories (eggs, dessert, cheese and fried foods), the chance-corrected agreement (κ) was fair. For the remaining food categories, this agreement was poor. For seven food categories, non-agreement around the diagonal of perfect matches was not symmetrical (always $P < 0.05$), with eggs, baked foods and snacks being the exception, where $P = 0.91$, $P = 0.10$ and $P = 0.25$, respectively.

Category scores

For all food categories, the mean category score of the children was lower than that of the mothers. A statistically significant (always $P < 0.05$) linear relationship between mothers' and children's category scores was found for meat, milk and cheese. For all other food categories, the correlation coefficient was 0.22 or lower, and in all cases, this was not statistically significant (see Table 4 for exact P -values).

Final scores

The correlation coefficient between the final scores of children and their mothers was 0.23 ($P = 0.04$) (Table 4). For boys, the correlation coefficient was non-significant ($r = 0.13$; $P = 0.46$) while for girls, it was higher than for the group as a whole ($r = 0.33$; $P = 0.04$). The mean difference between children's and mothers' final scores was 32.0 ± 35.1 . The mean difference between children and mothers was very similar for boys (31.6 ± 42.4) and girls (32.4 ± 27.9), but greater variability (see standard deviations) was apparent in the case of boys. Nevertheless, the difference in final scores between children and mothers differed significantly from zero for the group as a whole and for the genders separately ($P < 0.001$ in all three instances).

Table 4. Children's and mothers' category* and final† scores: means and correlations‡

Food category		Mean $\pm$ SD	Correlations§	
			r	P
Meat	Children	13.4 $\pm$ 6.2	0.25	0.03
	Mothers	13.2 $\pm$ 4.9		
Eggs	Children	5.2 $\pm$ 3.9	0.12	0.28
	Mothers	4.4 $\pm$ 3.1		
Dairy, milk, high fat	Children	13.9 $\pm$ 6.2	0.40	0.00
	Mothers	11.2 $\pm$ 6.4		
Dairy, cheese, high fat	Children	10.4 $\pm$ 6.4	0.41	0.00
	Mothers	8.2 $\pm$ 5.5		
Dairy, dessert, high fat	Children	10.8 $\pm$ 5.2	−0.03	0.84
	Mothers	5.3 $\pm$ 3.8		
Fried foods	Children	12.6 $\pm$ 6.9	0.15	0.20
	Mothers	8.4 $\pm$ 4.6		
In baked goods	Children	8.6 $\pm$ 5.2	0.09	0.42
	Mothers	7.0 $\pm$ 5.2		
Convenience foods	Children	10.8 $\pm$ 6.4	0.04	0.70
	Mothers	5.0 $\pm$ 3.3		
Table fats, high fat	Children	16.1 $\pm$ 6.2	0.22	0.05
	Mothers	9.9 $\pm$ 6.2		
Snacks, high fat	Children	12.3 $\pm$ 6.1	0.22	0.06
	Mothers	9.1 $\pm$ 5.0		
Final score	Children	114.4 $\pm$ 30.7	0.23	0.04
	Mothers	82.0 $\pm$ 25.2		

SD, standard deviation. *Frequency score multiplied by portion size score (see Table 3). †Sum of category scores. ‡ $N = 77$ for category scores; $n = 72$ for final score because of incomplete screeners by caregivers. §Spearman.

The data points on the Bland–Altman plot (Fig. 1) emphasise the variability (random error) and the presence of bias (systematic error) by the magnitude of the standard deviation and mean difference, respectively.

Screener classification

Ninety-three per cent of the children's self-assessment resulted in a 'high-fat' intake classification, whereas the maternal assessment of their children led to a figure for high fat consumption of 75%. The percentage identical classifications (values in bold in Table 5) into both, high fat and prudent fat intake, was 76%. When corrected for chance, the overall agreement was, however, poor [$\kappa = 0.16$; 95% confidence interval: (−0.06; 0.39)] and the McNemar's test indicated a departure from symmetry ($P = 0.001$):

Fig. 1. Final score differences (child minus mother) plotted against mean final scores $[(\text{child} + \text{mother})/2]$.

Table 5. Screener classification of fat intake: children vs. mothers ($n=75$)

		Mothers		Total
		High fat intake	Prudent fat intake	n (%)
		n (%)	n (%)	
Children	High fat intake	54 (72)	16 (21)	70 (93)
	Prudent fat intake	2 (3)	3 (4)	5 (7)
Total		56 (75)	19 (25)	75 (100)

Percentage identical classification values are in bold.

more mothers reported prudent fat intake when the children reported a high fat intake, than mothers who reported high fat intake when their child indicated a prudent intake (evident by a non-symmetry around the values not in bold).

Discussion

This study represents a first-time investigation of the 'South Africanised' MEDFICTS dietary fat screener in grade six children and their mothers. Our first

objective was to determine the internal consistency of the screener, i.e. the degree to which the individual food categories measured the same characteristic. As all item-total correlations were well above 0.2 and the Cronbach's alpha was close to 0.7 for children and for mothers (Keller *et al.* 2000), we conclude that all the food categories included in the screener can be retained and that they fully cover the construct that we are interested in, namely fat intake. The alpha obtained in this study was higher than the range of 0.13–0.62 reported in the review by Yaroch *et al.*

(2000) for dietary fat indices, but not as high as the cut-off of 0.80 or higher as recommended by Higgins & Straub (2006) for well-established and widely used instruments.

Apart from measuring internal consistency, item-total correlations have been used as an indication of convergent validity (Heller *et al.* 2000; Johnson *et al.* 2002; Higgins & Straub 2006). Regardless of who completed the dietary fat screener – children or their mothers – and regardless of the statistical technique used for measuring internal consistency (see Table 2), the screener showed characteristics of reliability. This will contribute to validity of measurement when using this instrument, as reliability is a necessary but not a sufficient condition for validity (Higgins & Straub 2006). It has been reasoned that variance among scores in an internally consistent instrument indicates subject differences and not error (McNulty *et al.* 2001). In the original pilot testing of MEDFACTS, Srinath *et al.* (1993) noted that self-administered and interviewer-administered application of the tool among adults resulted in similar findings.

For achieving our second objective, the scores of the mothers were compared with those from the children. We found the following. First, for the two building blocks of the screener (i.e. the reported usual portion size and weekly consumption), there was limited agreement between mothers and children (Table 3). In some cases, the kappa value was even negative, meaning worse than chance-corrected agreement. According to Hoehler (2000), the presence of bias reduces kappa values. The following point shows that systematic error (bias) was present for the reported portion size and weekly consumption, thus explaining some of the poor chance-corrected agreement obtained between children and mothers.

The second finding regarding child vs. parent assessment was that there appeared to be a systematic error: most of the non-agreeing responses were not symmetrical (McNemar's data in Table 3). Typically, mothers reported smaller portion sizes and less-frequent consumption. This would have a carry-over-effect to the category and final scores, as evident from Tables 4 and 5 and Fig. 1. The lower scores of the mothers may be due to under-reporting by the mothers [common among adult females in terms of

their own intake – Scagliusi *et al.* (2009)] or over-reporting by the children [more common among children than among adults (Forrestal 2011)] or both. The difference was more evident for the food categories to which the mothers may have had an 'unhealthy' connotation (e.g. dessert, convenience foods, fried and baked foods, as compared with meat, milk and cheese), whereas among the children, these foods may have been popular and possibly over-reported. This has previously been suggested by Koehler *et al.* (2000). The lack of agreement in our study was more pronounced for portion size compared with frequency of intake. Matheson *et al.* (2002) assessed the validity of 8–12-year-old African-American girls' self-report of food portion estimates and found 'sizable errors in quantitative estimates'. This emphasises the need for cautious interpretation of children's self-reports of portion sizes. The actual consumption of large portions as featured in food advertisements and the availability of super-sized portions of foods (e.g. at the school tuck shop, i.e. kiosk) acquired by children in the absence of their mothers is, however, also a possibility. This creates the potential for food category-specific maternal misreporting about food intake of children of this age group (Livingstone & Robson 2000). Finally, based on the cognitive challenges involved in responding to food frequency questionnaires, the general finding that such questionnaires may overestimate intake (McPherson *et al.* 2000) could also have applied. The above makes it difficult to interpret our findings relative to those of Murakami *et al.* (2012), who associated younger age (ranging from 6 to 15 years), normal weight (International Obesity Task Force classification), low parental education and completion of the assessment alone (as opposed to parental assistance) with over-reporting.

The correlation between the final scores of mothers vs. children was small ($r = 0.23$), yet statistically significant ($P = 0.04$). It is tempting to emphasise the stronger correlation ($r = 0.33$, $P = 0.04$) for girls as subgroup, but our follow-up analyses showed that correlations are not enough: agreement is needed. The mean difference between mother–daughter pairs' final scores also differed significantly from zero, suggesting systematic error and also highlighting consid-

erable variability. Children's self-assessment resulted in higher estimates of fat intake than the assessment of their mothers.

There was a relatively high percentage (76%) of identical classifications into high fat or prudent intake. A percentage perfect match above 70% has sometimes been considered an indication of validity (Snyder *et al.* 2004). However, statistically, the chance-corrected agreement (κ) between mothers' and children's final classifications in our study was poor. According to Hoehler (2000), this may – again – be due to the presence of bias, as noted in this study. It could also be a result of prevalence effects: a high percentage of participants in our study had high fat intakes. The prevalence effect appears to occur when the model is based on an underlying continuous variable: in our case, the final score (Hoehler 2000).

Usually, the aim of a dietary fat screener is to classify fat intakes into high vs. prudent. Using classification agreement and calculating a tool's diagnostic accuracy would thus be appropriate. Because at least one of the two data sources in our study (children or mothers) appeared to be inaccurate and because it is unknown which one of these, it is not meaningful to use either as a 'gold standard' for determining diagnostic accuracy. This study cannot be considered a validation study because the truth regarding fat intake of the children is not known. On the other hand, we believe the study highlights aspects of the methodological and statistical analytical complexity of dietary assessment, for example, how – in spite of good psychometric properties – the credibility of the measurements in a particular target group requires careful attention. Furthermore, a contribution of the study lies in uncovering the nature and extent of some errors that can occur when performing dietary screening based on food frequency-type assessment in children of this age group. At the same time, the potential and practical developmental opportunities of the MEDFICTS dietary fat screener in school children in new contexts have been uncovered, ranging from novel, pictorial presentation [similar to Wallen *et al.* (2011)] to group administration in school, which resulted in good response rates and few missing values, both measures of feasibility (Raat *et al.* 2002).

From a research perspective, it is recommended that refinement of the instrument continues. This includes test–retest reproducibility so as to demonstrate instrument responsiveness, and validation, with particular emphasis on portion size estimation. Should a combination of child and maternal data bring us closer to the truth as suggested by Murakami *et al.* (2012), this would require evidence about the relative quantitative and qualitative contribution of both data sources. Future studies need to consider the prevalence of high fat intakes in the target group during sample planning (Yi *et al.* 2004). More characterisation of the children and their caregivers as potential over- or under-reporters, respectively, in general (Forrestal 2011) or for specific food categories, is needed. The inherent limitations of this instrument as a dietary fat screener (Kris-Etherton *et al.* 2001) need to be kept in mind.

On the practical side, even though accurate screening of usual fat intake in this target group may remain illusive for some time, it is recommended that school-based, comprehensive primary prevention of non-communicable diseases programmes should be developed and implemented. These should target children, their caregivers and schools, as in this age group, many choices are made by the children themselves – for example, in school tuck shops (kiosks) – in the absence of their mothers, while at the same time, children rely on their caregivers to acquire and make food available. Monitoring the impact of such nutrition promotion programmes and achievement of dietary goals, does, however, require reliable and valid tools. It is time for action on both fronts: measurement refinement and intervention.

Acknowledgements

The statistical and programming support of Dr M. van der Linde and Mr J. Grimbeek (STATOMET, University of Pretoria, South Africa) are acknowledged.

Source of funding

None.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Contributions

All three authors contributed substantially to the conception and design of data acquisition, analysis and interpretation of data. They all contributed to the intellectual content and approved the final version of the manuscript.

References

- Altman D.G. (1991) *Practical Statistics for Medical Research*. Chapman & Hall/CRC: London.
- American Dietetic Association (ADA) (2004) Position of the American Dietetic Association: dietary guidance for healthy children ages 2 to 11 years. *Journal of the American Dietetic Association* **104**, 660–677.
- Armstrong M.E.G., Lambert M.I. & Lambert E.V. (2011) Secular trends in the prevalence of stunting, overweight and obesity among South African children (1994–2004). *European Journal of Clinical Nutrition* **65**, 835–840.
- Armstrong M.E.G., Lambert M.I., Sharwood K.A. & Lambert E.V. (2006) Obesity and overweight in South African primary school children – Health of the Nation Study. *South African Medical Journal* **96**, 439–444.
- Burrows T.L., Martin R.J. & Collins C.E. (2010) A systematic review of the validity of dietary assessment methods in children when compared with the methods of doubly labeled water. *Journal of the American Dietetic Association* **110**, 1501–1510.
- Forrestal S.G. (2011) Energy intake misreporting among children and adolescents: a literature review. *Maternal & Child Nutrition* **7**, 112–127.
- Heller L., Fox S., Hell K. & Church J.A. (2000) Development of an instrument to assess nutritional risk factors for children infected with human immunodeficiency virus. *Journal of the American Dietetic Association* **100**, 323–329.
- Higgins P.A. & Straub A.J. (2006) Understanding the error of our ways: mapping the concepts of validity and reliability. *Nursing Outlook* **54**, 23–29.
- Hoehler F.K. (2000) Bias and prevalence effects on kappa viewed in terms of sensitivity and specificity. *Journal of Clinical Epidemiology* **53**, 499–503.
- Holmes A.L., Sanderson B., Maisiak R., Brown A. & Bittner V. (2005) Dietitian services are associated with improved patient outcomes and the MEDFICTS dietary assessment questionnaire is a suitable outcome in cardiac rehabilitation. *Journal of the American Dietetic Association* **105**, 1533–1540.
- Johnson F., Wardle J. & Griffith J. (2002) The Adolescent Food Habits Checklist: reliability and validity of a measure of healthy eating behaviour in adolescents. *European Journal of Clinical Nutrition* **56**, 644–649.
- Keller H.H., Hedley M.R. & Brownlee S.W. (2000) The development of seniors in the community: risk evaluation for eating and nutrition (SCREEN). *Canadian Journal of Dietetic Practice and Research* **61**, 67–72.
- Koehler K.M., Cunningham-Sabo L., Lambert L.C., McCa-lman R., Skipper B.J. & Davis S.M. (2000) Assessing food selection in a health promotion program: validation of a brief instrument for American Indian children in the Southwest United States. *Journal of the American Dietetic Association* **100**, 205–211.
- Kris-Etherton P., Eisenstat B., Jaax S., Srinath U.M.A., Scott L., Rader J. et al. (2001) Validation for MED-FICTS, a dietary assessment instrument for evaluating adherence to total and saturated fat recommendations of the National Cholesterol Education Program Step 1 and Step 2 diets. *Journal of the American Dietetic Association* **101**, 81–86.
- Lee D.L. & Nieman D.C. (2010) *Nutritional Assessment*, 5th edn, McGraw-Hill: Boston.
- Livingstone M.B.E. & Robson P.J. (2000) Measurement of dietary intake in children. *The Proceedings of the Nutrition Society* **59**, 279–293.
- Matheson D.M., Hanson K.A., McDonald T.E. & Robinson T.N. (2002) Validity of children's food portion estimates: a comparison of 2 measurement aids. *Archives of Pediatrics and Adolescent Medicine* **156**, 867–871.
- Mayosi B.M., Flisher A.J., Lalloo U.G., Sitas F., Tollman S.M. & Bradshaw D. (2009) The burden of non-communicable diseases in South Africa. *Lancet* **374**, 934–947.
- McNulty K.Y., Adams C.H., Anderson J.M. & Affenito S.G. (2001) Development and validation of a screening tool to identify eating disorders in female athletes. *Journal of the American Dietetic Association* **101**, 886–892.
- McPherson R.S., Hoelscher D.M., Alexander M., Scanlon K.S. & Serdula M.K. (2000) Dietary assessment methods among school-aged children: validity and reliability. *Preventive Medicine* **31**, S11–S33.
- Mochari H., Gao Q. & Mosca L. (2008) Validation of the MEDFICTS dietary assessment questionnaire in a diverse population. *Journal of the American Dietetic Association* **108**, 817–822.
- Murakami K., Miyake Y., Sasaki S., Tanaka K. & Arakawa M. (2012) Characteristics of under- and over-reporters of energy intake among Japanese children and

- adolescents: the Ryukyus Child Health Study. *Nutrition* **28**, 532–538.
- Norman R., Bradshaw D., Steyn K., Gaziano T. & South African Comparative Risk Assessment Collaborative Group (2007) Estimating the burden of disease attributable to high cholesterol in South Africa in 2000. *South African Medical Journal* **97**, 708–715.
- Raat H., Bonsel G.J., Essink-Bot M.-L., Landgraf J.M. & Gemke R.J.B. (2002) Reliability and validity of comprehensive health status measures in children: the Child Health Questionnaire in relation to the Health Utilities Index. *Journal of Clinical Epidemiology* **55**, 67–76.
- Scagliusi F.B., Ferriolli E., Pfrimer K., Cunha C.S.F., Gualano B., Lourenco B.H. *et al.* (2009) Characteristics of women who frequently under report their energy intake: a doubly labeled water study. *European Journal of Clinical Nutrition* **63**, 1192–1199.
- Schwartz C.G., Torisky D.M., Gloeckner J.W., Marrah G.W. & Nett M. (2000) Adapting MEDFICTS food frequency questionnaire to assess dietary fat intake and educate participants at a cholesterol screening on North Carolina. *Journal of the American Dietetic Association* **100** (Suppl.), A-40.
- Snyder D.C., Sloane R. & Lobach D. (2004) Agreement between a brief mailed screener and an in-depth telephone survey: observations from the Fresh Start study. *Journal of the American Dietetic Association* **104**, 1593–1596.
- Srinath U., Shacklock F., Shannon B.M., Mitchell D.C., Kris-Etherton P.M., Scott L. *et al.* (1993) MEDFICTS – a dietary assessment instrument for evaluating fat, saturated fat and cholesterol intake. *Circulation* **88**, 3414.
- Taylor A.J., Wong H., Wish K., Carrow J., Bell D., Binde-man J. *et al.* (2003) Validation of the MEDFICTS dietary questionnaire: a clinical tool to assess adherence to American Heart Association dietary fat intake guidelines. *Nutrition Journal* **2**, [serial online] 2–4. Available at: <http://www.nutritionj.com/content/2/1/4>.
- Teal C.R., Baham D.L., Gor B.J. & Jones L. (2007) Is the MEDFICTS rapid dietary fat screener valid for premenopausal African-American women? *Journal of the American Dietetic Association* **107**, 773–781.
- Van Horn L., McCain M., Kris-Etherton P.M., Burke F., Carson J.A.S., Champagne C.M. *et al.* (2008) The evidence for dietary prevention and treatment of cardiovascular disease. *Journal of the American Dietetic Association* **108**, 287–331.
- Wallen V., Cunningham-Sabo L., Auld G. & Romaniello C. (2011) Validation of a group-administered pictorial dietary recall with 9- to 11-year-old children. *Journal of Nutrition Education and Behavior* **43**, 50–54.
- Wenhold F.A.M., Kruger S. & Muelhof E. (2008) Nutrition for school-age children and adolescents. In: *Community Nutrition Textbook for South Africa: A Rights-Based Approach* (eds N.P. Steyn & N. Temple), pp 441–478. FAO & MRC: Tygerberg.
- Wolmarans P., Danster N., Dalton A., Rossouw K. & Schönfeldt H. (eds) (2010) *Condensed Food Composition Tables for South Africa*. South African Medical Research Council: Parow Valley, Cape Town.
- Yaroach A.L., Resnicow K. & Khan L.K. (2000) Validity and reliability of qualitative dietary fat index questionnaires: a review. *Journal of the American Dietetic Association* **100**, 240–244.
- Yi Q., Panzarella T. & Corey P. (2004) Incorporating the sampling variation of the disease prevalence when calculating the sample size in a study to determine diagnostic accuracy of a test. *Controlled Clinical Trials* **25**, 417–427.