RESEARCH ARTICLE

Aberrant visual pathway development in albinism: From retina to cortex

Sarim Ather¹ | Frank Anthony Proudlock² | Thomas Welton^{3,4} | Paul S. Morgan^{4,5} |
Viral Sheth² | Irene Gottlob² | Rob A. Dineen^{3,4}

¹Nuffield Department of Surgical Sciences, University of Oxford, Oxford, United Kingdom

²University of Leicester Ulverscroft Eye Unit, Robert Kilpatrick Clinical Sciences Building, Leicester, United Kingdom

³Radiological Sciences, Division of Clinical Neuroscience, University of Nottingham, Queen's Medical Centre, Nottingham, United Kingdom

⁴Sir Peter Mansfield Imaging Centre, University of Nottingham, Queen's Medical Centre, Nottingham, United Kingdom

⁵Medical Physics and Clinical Engineering, Nottingham University Hospitals NHS Trust, Queen's Medical Centre, Nottingham, United Kingdom

Correspondence

Irene Gottlob, University of Leicester Ulverscroft Eye Unit, Robert Kilpatrick Clinical Sciences Building, Leicester Royal Infirmary, Leicester, LE2 7LX, UK.
Email: ig15@le.ac.uk

Abstract

Albinism refers to a group of genetic abnormalities in melanogenesis that are associated neuronal misrouting through the optic chiasm. We perform quantitative assessment of visual pathway structure and function in 23 persons with albinism (PWA) and 20 matched controls using optical coherence tomography (OCT), volumetric magnetic resonance imaging (MRI), diffusion tensor imaging and visual evoked potentials (VEP). PWA had a higher streamline decussation index (percentage of total tractography streamlines decussating at the chiasm) compared with controls ($Z = -2.24$, $p = .025$), and streamline decussation index correlated weakly with inter-hemispheric asymmetry measured using VEP ($r = .484$, $p = .042$). For PWA, a significant correlation was found between foveal development index and total number of streamlines ($r = .662$, $p < .001$). Significant positive correlations were found between peri-papillary retinal nerve fibre layer thickness and optic nerve ($r = .642$, $p < .001$) and tract ($r = .663$, $p < .001$) width. Occipital pole cortical thickness was 6.88% higher ($Z = -4.10$, $p < .001$) in PWA and was related to anterior visual pathway structures including foveal retinal pigment epithelium complex thickness ($r = -.579$, $p = .005$), optic disc ($r = .478$, $p = .021$) and rim areas ($r = .597$, $p = .003$). We were unable to demonstrate a significant relationship between OCT-derived foveal or optic nerve measures and MRI-derived chiasm size or streamline decussation index. Our novel tractographic demonstration of altered chiasmic decussation in PWA corresponds to VEP measured cortical asymmetry and is consistent with chiasmic misrouting in albinism. We also demonstrate a significant relationship between retinal pigment epithelium and visual cortex thickness indicating that retinal pigmentation defects in albinism lead to downstream structural reorganisation of the visual cortex.

KEYWORDS

albinism, diffusion tensor imaging, magnetic resonance imaging, optical coherence tomography, visual pathway

1 | INTRODUCTION

Albinism refers to a group of genetic mutations that lead to abnormalities in the melanin synthesis and transport pathway (Montoliu et al., 2014; Kamaraj & Purohit, 2014). The phenotype of PWA includes

reduced visual acuity, foveal hypoplasia, nystagmus, increased crossing of nerve fibres at the optic chiasm and changes in the visual cortex on MRI (Levin & Stroh, 2011).

In normal foveal development, anti-angiogenic molecules form a molecular barrier to form a 'foveal avascular zone' (FAZ) (Provis, Sandercoe, & Hendrickson, 2000; Provis, 2001; Provis & Hendrickson, 2008). In albinism, this fails to happen, leading to encroachment of the inner retinal layers into the fovea, a deficiency in the formation of the foveal pit and a lack of photoreceptor specialisation (Elschnig, 1913; Naumann, Lerche, & Schroeder, 1976; Akeo et al., 1996; Chong et al.,

Abbreviations: DTI, diffusion tensor imaging; LGN, lateral geniculate nucleus; MRI, Magnetic Resonance Imaging; OCT, optical coherence tomography; ppRNFL, peripapillary nerve fibre layer; PWA, people with albinism; RGC, retinal ganglion cell; ROI, region of interest; RPE, retinal pigment epithelium; VEP, visual evoked potential

2009). Using spectral domain optical coherence tomography (OCT), our group has previously shown that the degree of incursion of the inner retinal layers at the fovea and foveal photoreceptor specialisation are inversely related and together define the degree of foveal development. In addition, we were able to demonstrate a significant relationship between photoreceptor size and best corrected visual acuity (BCVA) (Thomas et al., 2011; Mohammad et al., 2011).

The optic nerve comprises of retinal ganglion cell (RGC) axons which form arcuate bundles that travel in the nerve fibre layer before converging at the optic nerve head (ONH). Albino animal studies have shown a reduction in the number of RGCs (Guillery et al., 1984; Leventhal & Creel, 1985). We have previous shown that reduced RGC numbers shown in animal studies translates to thinner peripapillary nerve fibre layer (ppRNFL) thickness (Mohammad et al., 2015). This in turn leads to smaller optic nerves and chiasm which have been demonstrated in persons with albinism (PWA) using magnetic resonance imaging (MRI) (Schmitz et al., 2003; Mcketton, Kelly, & Schneider, 2014).

Additionally, albinism is associated with an abnormally increased chiasmal decussation of the nerve fibres originating from the temporal hemi-retinae (Guillery, Okoro, & Witkop, 1975). Axon guidance at the chiasm is regulated by a number of molecular mechanisms at the retina (Priour & Rebsam, 2016).

Study of albino mice has shown an increased expression of the transcription factor *Islet2+* which represses the ipsilateral program by reducing the expression of *Zic2* and thus *EphB1* which is a receptor tyrosine kinase important in divergence of axons at the chiasmal midline and plays a key role in stopping axons from crossing at the midline (Garcia-Frigola, Carreres, Vegar, Mason, & Herrera, 2008; Rebsam, Bhansali, & Mason, 2012).

Delayed neurogenesis appears to play a key role in the misrouting seen in albinism (Rachel et al., 2002; Bhansali, Rayport, Rebsam, & Mason, 2014). It has been suggested that the time at which the axon reaches the chiasm determines the fate of decussation. During embryological development, the first RGC axons arrive at the chiasm around the fourth week of gestation. In some mammals such as mice and ferrets, it has been shown that axons that reach the chiasm early during development are more likely to stay ipsilateral (Baker & Reese, 1993). Absence of L-Dopa, a pre-cursor of melanin, in the retinal pigment epithelium delays the point at which cells in the developing albino retina exit the cell cycle (Ilia & Jeffery, 1999; Kralj-Hans, Tibber, Jeffery, & Mobbs, 2006). As uncrossed RGCs are generated earlier than those that project across the midline, a delay in ganglion cell production means that axons from these cells reach the chiasm at a later point and this increases their probability of projecting to the contralateral hemisphere (Erskine & Herrera, 2014).

Diffusion tensor imaging (DTI) is a widely applied quantitative imaging technique for studying white matter anatomy and integrity. By quantifying the magnitude and principle direction of water diffusion within image voxels, DTI data can be used for reconstruction of principle white matter tracts, a technique referred to as tractography (Beaulieu, 2002). Grigorian et al. used the technique to study the optic radiation and found that in albinism, fibres from lateral geniculate nucleus (LGN) to the primary visual cortex (V1) are reduced (Grigorian, Mcketton, & Schneider, 2016).

A number of studies have shown that MRI scanning can detect alterations in visual cortical areas in albinism. Using voxel based morphometry, Von dem Hagen et al. found that people with albinism show a reduction in cortical volume at the occipital pole (Von dem Hagen, Houston, Hoffmann, Jeffery, & Morland, 2005), while Neveu et al. found that the calcarine fissure is shorter. In addition, the latter study also reported a marked asymmetry in the calcarine sulcus between the left and right hemispheres of the majority of PWA. The authors noted that in the presence of a dominant eye, the calcarine sulcus in the contralateral hemisphere is displaced downward (Neveu, von dem Hagen, Morland, & Jeffery, 2008).

Surface based analysis provides an alternative methodology to assess cortical differences in the human brain by generating geometric models of the cortical surface (Dale, Fischl, & Sereno, 1999; Fischl, Sereno, & Dale, 1999; Fischl & Dale, 2000). Using this methodology, Bridge et al. showed reduced gyrification in the occipital cortex of albinism patients that explains the reduced cortical volume reported by Von dem Hagen et al. In addition, they found cortical thickness to be increased at the occipital pole of PWA. This change was more profound in the left hemisphere and cortical thickness was negatively correlated to visual acuity. The authors suggested that these changes are due to a lack of post-natal neuronal pruning as a result of underdevelopment of the fovea seen in albinism and a consequent absence of high-resolution input into V1 (Bridge et al., 2014).

In this study, we conduct a holistic assessment of aberrant visual pathway development in PWA by sampling anatomical variation at multiple points along the visual pathway, from the retina to the visual cortex, using various noninvasive imaging techniques. To study the anterior visual pathway, we perform OCT evaluation of the fovea and optic nerve head structure. The post orbital visual pathway was studied using high-resolution T1-weighted MRI imaging to measure cisternal optic nerves, chiasm, optic tracts and V1 cortical thickness. Our aim was to confirm previous reports regarding altered morphology of these structures in PWA (Schmitz et al., 2003; Bridge et al., 2014; Mcketton et al., 2014).

We employ diffusion tractography to study the chiasmal connectivity in albinism for the first time. Structural connectivity at the chiasm was defined by streamline density measurements based on diffusion tractography. We used this to define a decussation index, describing the proportion of crossing fibres at the chiasm and compared it to chiasmal decussation measured using visual evoked potential (VEP), a functional measure of axonal misrouting though the chiasm in albinism.

This multimodality data, has allowed us to explore whether the anomalous post-orbital optic nerve, chiasm, tract and visual cortex morphology is related to retinal and optic nerve head abnormalities described in albinism. We investigate the hypotheses that:

- Alteration in foveal morphology affects the development of the chiasm.
- Cortical abnormalities in albinism are a result of abnormal visual input due to an underdeveloped fovea.
- Cortical thickness at the occipital pole is related to the degree of melanin present in the foveal RPE.

- Optic nerve head morphology is related to the size and connectivity of the optic chiasm.
- Cortical thickness at the occipital pole is related to optic nerve head morphology in PWA.

2 | MATERIALS AND METHODS

2.1 | Participants and recruitment

The study was performed in accordance with the tenets of the Declaration of Helsinki and was approved by the local UK National Health Service Research Ethics Committee. All participants provided written informed consent prior to participation.

Adult participants with albinism were recruited through the neuro-ophthalmology outpatient clinic at the Leicester Royal Infirmary. Diagnosis of albinism was confirmed by the coexistence of nystagmus, asymmetric VEP responses, foveal hypoplasia and iris transillumination (Gottlob & Proudlock, 2014; Papageorgiou, McLean, & Gottlob, 2014).

Age, gender and ethnicity matched volunteers were recruited for the control group from within the students and faculty at the University of Leicester as well as healthy visitors to the ophthalmology department. For inclusion, potential control group participants had to have no history of eye disease and have had a best corrected visual acuity (BCVA) of better than 0.0 logMAR. Analyses based on this participant cohort have been reported in a previous publication (Welton, Ather, Proudlock, Gottlob, & Dineen, 2017).

All participants underwent MRI scan using the protocol described below. In addition, the albinism group participants underwent a detailed clinical assessment including assessment of best corrected visual acuity, colour vision, stereo-acuity, ocular movements, slit lamp examination and dilated fundus examination as well as OCT and VEP.

2.2 | Optical coherence tomography

Macular and optic nerve OCT scans were acquired using the SOCT Copernicus HR device (OPTOPOL Technology S.A., Zawiercie, Poland). Foveal layers thickness was measured using ImageJ software (National Institutes of Health, MD, USA). Detailed methodology of this analysis has previously been described by our group (Mohammad et al., 2011). As foveal development is a combination of processing layer extrusion and photoreceptor lengthening, both these measures were incorporated in the following formula to calculate a foveal development index.

$$\text{Foveal development index} = 2 - \frac{\text{Processing layer thickness}}{\text{Photoreceptor layer thickness}}$$

This index was used for comparison with chiasmal and cortical measurements.

Work carried out using polarisation sensitive OCT has shown that the reduced melanin in the RPE of patients with albinism alters the reflectivity profile of this layer (Schutze et al., 2014) and therefore the RPE thickness was used as a surrogate for the amount of melanin present.

Optic nerve head analysis has been described in detail previously by our group. In summary, custom written macros were used in ImageJ software (National Institutes of Health, MD, USA) to correct nystagmus related motion artefact. Following this, Copernicus SD-OCT software was used to calculate cup, disc and rim dimensions and peripapillary nerve fibre layer thickness (Mohammad et al., 2015).

2.3 | Visual evoked potentials

VEP testing was carried out in accordance with international society for clinical electrophysiology of vision standards (Odom et al., 2010). The patients were seated and allowed to wear their full spectacle correction throughout the duration of the test. Five electrodes were placed at 10% intervals in a horizontal chain across the posterior part of the scalp left and right of Oz. In addition, a reference electrode was placed in the midline frontally and a ground electrode placed in the midline over the vertex.

The stimulus was a black and white checkerboard pattern, with 100% contrast, a mean luminance of 96 cd/m² and check size of 1° generated on a 17-in. CRT screen positioned at 46 cm distance from the patient, which created a full-field size of 33°. The pattern appeared at a rate of 200 ms onset, 400 ms offset. Patients were asked to fixate on a nonilluminated central spot. The responses for the left and right eyes were recorded separately with the other eye completely occluded using an eye patch. The test was performed twice on each eye and an average of the two sets of results was used for analysis.

VEP asymmetry was calculated by means of an interhemispheric asymmetry index (I_{asym}), based on a methodology described by Apkarian, Reits, Spekreijse, and Van Dorp (1983). The initial step is to calculate response lateralization (A.I.) for each eye by plotting the magnitude of response at each electrode against the electrode position and calculating the area under the graph for each hemisphere (A_L and A_R).

The following formula was used to calculate response lateralization (A.I.) in each eye.

$$\text{for } A_L > A_R, \quad \text{A.I.} = \left(\frac{A_R}{A_L} \right)$$

$$\text{else A.I.} = 2 - \left(\frac{A_L}{A_R} \right)$$

The intra-ocular asymmetry index was calculated by subtracting response lateralization in the right eye from the response lateralization in the left eye.

2.4 | Magnetic resonance imaging

Brain MR imaging was performed using a 3 T Philips Achieva MRI scanner with a 32-channel SENSE head coil (Best, The Netherlands). Sequences performed included axial 3D magnetization-prepared rapid acquisition gradient (3D-MPRAGE, TR = 7.53 ms, TE = 2.22 ms, flip angle = 8°, matrix size 320 × 320, field of view = 256 × 256, 0.8 mm isotropic voxels, SENSE factor = 1.7, 184 slices; acquisition time 6.5 min) and diffusion weighted imaging (axial diffusion-weighted echo-planar imaging, six repeats of the $b = 0$ volume, averaged on the

scanner, and 61 directional diffusion weighted images with $b = 1,000 \text{ s/mm}^2$, TE = 67 ms, TR = 8,270 ms, SENSE factor 3, phase encoding in the anterior–posterior direction, full Fourier, acquisition matrix size 120×120 , 52 contiguous slices, $1.8 \times 1.8 \times 1.8 \text{ mm}$ voxels interpolated to $0.9 \times 0.9 \times 1.8 \text{ mm}$ voxels, acquisition time 9.5 min).

2.5 | Morphometry of optic nerves, chiasm and tracts

The technique for assessing the optic nerve, chiasm and tract dimensions was based on previously described methodology by manually tracing regions of interests (ROIs) around each structure on the MPRAGE images (Mcketton et al., 2014; Schmitz et al., 2003). This was carried out using a custom written macro in ImageJ software (National Institutes of Health, MD, USA) by an assessor who was blinded to patient demographics and group membership. This allowed calculation of the width and area for each structure (Supporting Information Figure S1).

2.6 | Diffusion tensor imaging

DTI data were processed using fMRIB's Diffusion Toolbox in FSL (Behrens, Berg, Jbabdi, Rushworth, & Woolrich, 2007). First, 'eddy_correct' was used to correct artefacts induced by head motion and eddy currents (Andersson & Skare, 2002). We did not need to exclude any volunteer due to excessive artefact.

A binary mask of the brain was created and nonbrain structures were removed with the Brain Extraction Tool. The DTIFit tool in FSL's Diffusion Toolkit, was used to fit tensors to the data and determine a variety of values including the fractional anisotropy, mean diffusivity and the three eigenvector and eigenvalues of each voxel. BEDPOSTX (Bayesian Estimation of Diffusion Parameters Obtained using Sampling Techniques) was used to build sampling distributions on the diffusion parameters at each voxel.

Masks were manually drawn on the FA maps for optic nerve, chiasm and tract. The first axial slices anterior and posterior to the chiasm where two separate nerves and tracts were visible were used to draw the masks. This is demonstrated in Supporting Information Figure S2. Probabilistic fibre tracking was performed using the streamline tractography algorithm, PROTRACKX2 (Behrens et al., 2007) contained within FSL to calculate 5,000 streamlines per seed voxel with a 0.5 mm step length and maximum of 2,000 steps, a 0.2 mm radius of curvature cutoff and an FA threshold of 0.1.

The algorithm propagates streamlines from each voxel in a given seed mask along the path with the largest principal axis of the diffusion tensor until some termination criteria are met (in this case, when the streamline reached the voxels in a termination mask). The number of streamlines generated allows estimation of the strength of connectivity between the seed and target voxels.

To increase the signal to noise ratio, the algorithm was run initially with the optic nerve being the seed and the tract the target and then repeated with the seed and target masks reversed. Results from these two streamline counts were then averaged for subsequent analyses. Wilcoxon signed-rank test was carried out to compare the differences between the number of streamlines generated from the left and right

eyes and there was no significant difference demonstrated in either group ($p > .05$). Intra-class correlation coefficient in the albinism group was 0.898 (95%CI = 0.754–0.958) and in the control group was 0.821 (95%CI = 0.548–0.929).

For comparison of chiasmal decussation estimated using DTI and VEP, a 'streamline decussation index' was deduced by calculating the percentage of streamlines connecting with contralateral regions of interest through the chiasm using the following formula:

$$\text{Streamline decussation index} = \frac{RN_{LT} + LN_{RT}}{(RN_{LT} + RN_{RT} + LN_{LT} + LN_{RT})} \times 100.$$

where, A_B is the number of streamlines between regions of interest A (seed mask) and B (target mask); LN is the left nerve; LT is the left tract; RN is the right nerve; RT is the right tract.

2.7 | Cortical analysis

Cerebral cortical thickness and volume were derived using FreeSurfer version 5.0.0 (<http://surfer.nmr.mgh.harvard.edu>). Detailed methods have been described previously (Dale et al., 1999; Fischl et al., 1999). In summary, the software undertakes a segmentation procedure that identifies white/grey matter interface (white surface) and the grey matter/cerebrospinal fluid interface (pial surface). The distance between these two surfaces is used to calculate cortical thickness and volume. As part of the standard FreeSurfer processing pipeline, an early step in the processing, the 0.9 mm^3 voxels were resampled to 1 mm^3 as part of the co-registration to the MNI305 template. Although the above process is automated, each scan was subject to meticulous manual inspection to check for errors in any of the above steps by an observer masked to the diagnosis. Any inaccuracies were manually corrected and thickness measurements were recalculated. Automatic parcellation of the cortex was performed based on the Destrieux atlas in FreeSurfer (Destrieux, Fischl, Dale, & Hagren, 2010). Measurements for the occipital pole region were used for comparison with the foveal and optic nerve head OCT parameters.

2.8 | Statistical analyses

SPSS software version 22 (SPSS, Inc., Chicago, IL) was used to carry out statistical analyses. Due to non-normality of the data, optic nerve, chiasm and tract parameters between and the albinism and control volunteers were performed using nonparametric tests (Mann–Whitney tests). Spearman's rank correlation co-efficient was used to study the relationship between OCT measurements and the MRI derived measurements of optic pathway structure, cortical thickness and functional data (VEP asymmetry and BCVA). Average values were used for comparison of paired structures.

As the relationship between structures throughout the visual pathway is being assessed, one of the limitations of the study is the number of comparisons that needed to be carried out. To counter this and ensure our results are biologically plausible, a priori hypotheses were defined based on previously published findings. In addition, where testing a hypothesis required multiple statistical comparisons, a Holm–Bonferroni correction is carried out to account for this (Holm, 1979). The corrected p -values have been labelled p' .

3 | RESULTS

3.1 | Group comparison of albinism patients and controls

The albinism group ($n = 23$, 17 males) and control group ($n = 20$, 14 males) were matched for ethnicity and age (mean age = 34.0 $\pm$ 13.6 years, 31.9 $\pm$ 10.6 years, respectively, $t = 0.851$, $p = .400$). The mean BCVA in the albinism group was 0.47 $\pm$ 0.21 logMAR. The control group all had a BCVA of 0.0 logMAR or above and normal stereoscopic vision.

3.2 | OCT parameters

We have previously published detailed OCT analysis of foveal and optic nerve (Mohammad et al., 2015) abnormalities in PWA. In this subset of patients, all PWA displayed some degree of foveal hypoplasia with incursion of inner retinal layers through the foveal zone. The mean value for foveal development index was 0.450 $\pm$ 0.562 while the RPE complex thickness was 28.6 μm $\pm$ 3.28. In our previous work, we have shown that in healthy controls, the mean FDI = 1.96 $\pm$ 0.148 and mean RPE thickness = 29.1 $\pm$ 5.13, respectively (Mohammad et al., 2011).

On OCT analysis of the optic nerve head, eight PWA did not display an optic cup. Mean optic disc, cup and rim areas were 1.82 mm² $\pm$ 0.339, 0.379 mm² $\pm$ 0.349 and 1.44 mm² $\pm$ 0.418, respectively. Mean ppRNFL thickness was 99.3 μm $\pm$ 15.2.

Using a threshold of 0.7 defined by Apkarian et al. (1983) all PWA displayed asymmetric VEP response. The mean VEP asymmetry index was 1.43 $\pm$ 0.32. In controls, this has been previously been shown to be -0.047 ± 0.655 .

4 | MRI ANALYSIS

4.1 | Structural changes to the chiasm region

Two comparisons each were made for the optic nerve and tracts and three for the chiasm. Optic nerve and tract width as well as the chiasm width, area and volume, were significantly smaller in the albinism group compared with controls (Table 1). The optic tract area and the Holm–Bonferroni corrected optic nerve area comparisons were not statistically significant. As the width of the nerve, chiasm and tract

were consistently smaller, this was used in subsequent comparisons with OCT measures.

4.2 | Chiasmal connectivity

Two comparisons were carried out to assess the chiasmal connectivity. Firstly, the total number of streamlines generated between the albinism and control groups were compared, but there was no significant group difference ($p > .05$). However, group comparison of the streamline decussation index showed a significantly higher percentage of decussating streamlines at the chiasm in the albinism group (mean = 42.0% $\pm$ 18.7) compared with the controls (mean = 27.8% $\pm$ 17.5) ($Z = -2.24$, $p = .025$, $p' = .05$) (Figure 1a). The total number of streamlines did not significantly correlate to the size of the ROIs ($p = 0.217$, $r = 0.197$). Figure 2 provides examples of DTI streamline data from albinism and control volunteers demonstrating higher percentage of contralateral streamlines in the PWA. Receiver operator curve (ROC) analysis of the streamline decussation index yielded an area under the curve of 0.727 (95% CI = 0.575–0.880).

4.3 | Cortical changes

Cortical thickness at the occipital pole was 6.88% higher in the albinism group (mean = 2.15 mm $\pm$ 0.16) compared with controls (mean = 2.01 mm $\pm$ 0.12) ($Z = -4.10$, $p < .001$) (Figure 1b).

4.4 | Relationship of orbital OCT measurements to post-orbital MRI-derived measures of visual pathway structure in albinism patients

Table 2 summarises comparisons between foveal and optic nerve measurements obtained via OCT and foveal development index, RPE thickness, ppRNFL thickness and optic disc, cup and rim areas measured using OCT and post orbital optic nerve, chiasm and tract width, diffusion tensor streamlines, decussation and cortical thickness at the occipital pole measured using MRI. Holm–bonferroni correction was applied based on the comparison of the MRI measures with two foveal parameters and four optic nerve parameters.

4.5 | Structural changes to the region

No significant correlation was found between foveal development index and optic nerve ($p = .160$, $r = -.303$), chiasm ($p = .085$, $r = .367$) or optic tract ($p = .241$, $r = .255$) width measured on MRI.

TABLE 1 Comparison of optic nerve, tract and chiasm dimensions

		Optic nerve		Optic tract		Optic chiasm		
		Width (mm)	Area (mm ²)	Width (mm)	Area (mm ²)	Width (mm)	Area (mm ²)	Volume (mm ³)
Albinism	Mean	6.22	16.0	5.54	16.5	11.7	30.7	261
	SD	0.782	2.87	0.993	4.61	1.30	4.97	64.4
Control	Mean	6.86	17.5	6.28	18.3	14.4	36.3	319
	SD	0.781	2.81	1.34	5.21	1.14	5.59	66.33
Mann–Whitney U	p	.001	.026	.010	.103	<.001	.006	.005
	p'	.006	.052	.030	.103	<.001	.024	.025
	Z	−3.26	−2.23	−2.53	−1.631	−4.91	−2.76	−2.79

FIGURE 1 (a) Comparison of the streamline decussation index between albinism and control groups ($Z = -2.24$, $p = .025$, $p' = .05$). (b) Comparison of occipital pole thickness averaged across both hemispheres between the albinism and control groups ($Z = -4.10$, $p < .001$)

The foveal RPE complex thickness was related to optic chiasm width, however, this comparison was not significant once corrected for multiple testing ($r = .413$, $p = .050$, $p' = .100$). The optic nerve and tract width did not relate to the RPE thickness ($p > .05$).

Significant positive correlations were found between ppRNFL thickness and the optic nerve ($r = .642$, $p < .001$, $p' < .001$) and tract ($r = .663$, $p < .001$, $p' < .001$) width. The chiasm width did not correlate to the ppRNFL thickness. The optic disc area was correlated to optic tract width but this relationship did not survive multiple comparison correction ($r = .474$, $p = .023$, $p' = .069$). The disc cup and rim areas did not relate to any of the other structures in the chiasmal region.

4.6 | Chiasmal connectivity

Significant correlations were found between the total number of streamlines and the foveal development index. ($r = .662$, $p < .001$, $p' < .001$, Figure 3). There was no significant relationship between the total number of streamlines or the degree of decussation and foveal RPE or optic nerve head measurements.

4.7 | Cortical thickness

Mean cortical thickness at the occipital pole was inversely correlated with the mean thickness of the foveal RPE ($r = -.579$, $p = .005$, $p' = .010$) (Figure 4). However, there was no relationship of cortical thickness with the foveal development index.

Cortical thickness at the occipital pole was also found to correlate with optic disc and rim areas ($r = .478$, $p = .021$, $p' = .042$, and $r = .597$, $p = .003$, $p' = .009$, respectively, Figure 5a,b). Cortical thickness did not relate to cup area or ppRNFL thickness ($p > .05$).

4.8 | Relationships between structural MRI and measures of visual function in albinism patients

Best-corrected visual acuity was not related to optic nerve, tract or chiasm width or with V1 cortical thickness ($p > .05$). BCVA was not related to total number of streamlines, but BCVA showed a trend toward significant correlation with streamline decussation index ($r = .432$, $p = .051$).

Streamline decussation index correlated weakly with inter-hemispheric asymmetry measured using VEP ($r = .484$, $p = .042$, Figure 6). We did not find any significant relationship between cortical thickness and VEP asymmetry ($p > .05$).

5 | DISCUSSION

This is the first study that comprehensively investigates the relationship between ocular abnormalities, and post orbital chiasmal and cortical abnormalities seen in albinism. To achieve this, the visual pathway was imaged using OCT, structural MRI and DTI. In addition, the

FIGURE 2 Example diffusion tractography streamline data from albinism (left) and control (right) volunteers. Streamlines travelling from the optic nerve to the ipsilateral tract are in orange while streamlines travelling to the contralateral regions of interest are blue. The images were thresholded such that voxels with a streamline density $< 10\%$ of the total streamlines are excluded. The images show a variation in the chiasmal connectivity in both groups [Color figure can be viewed at wileyonlinelibrary.com]

TABLE 2 Comparison of orbital morphology of the fovea and optic nerve head assessed via OCT with post orbital measurements of the optic chiasm and the visual cortex assessed through T1 and diffusion weighted MRI

				Orbital measures (OCT)						
				Foveal region		Peripapillary region				
				FDI	RPE	ppRNFL thickness	Disc area	Cup area	Rim area	
Post-orbital measures (MRI)	Optic chiasm region (structural MRI)	Optic nerve diameter	<i>r</i>	−.303	.277	.642	.181	.133	.032	
			<i>p</i>	.162	.211	<.001	.421	.541	.894	
			<i>p'</i>	.324	.211	<.001	.931	1.00	1.00	
		Optic tract diameter	<i>r</i>	.241	.204	.663	.474	.194	.221	
			<i>p</i>	.255	.358	<.001	.023	.388	.304	
			<i>p'</i>	.510	.358	<.001	.069	.388	.608	
		Optic chiasm diameter	<i>r</i>	.367	.413	−.057	−.002	−.181	.107	
			<i>p</i>	.085	.050	.795	.992	.257	.505	
			<i>p'</i>	.170	.100	1.00	.992	1.00	1.00	
	Optic chiasm region (connectivity: DTI)	Total number of streamlines	<i>r</i>	.536	.069	−.379	−.122	.422	−.444	
			<i>p</i>	.012	.771	.164	.641	.091	.074	
			<i>p'</i>	.024	.771	.328	.641	.273	.296	
		Streamline decussation index	<i>r</i>	.412	.032	.205	−.387	−.367	−.013	
			<i>p</i>	.068	.891	.360	.125	.147	.961	
			<i>p'</i>	.136	.891	.720	.506	.447	.961	
		Cortex (structural MRI)	Cortical thickness	<i>r</i>	−.016	−.579	.368	.478	−.143	.597
				<i>p</i>	.941	.005	.084	.021	.515	.003
				<i>p'</i>	.941	.010	.168	.042	.515	.009

anatomical data were compared with functional measurements such as visual acuity and VEP asymmetry.

5.1 | Chiasmal abnormalities in albinism

Our results agreed with the findings of smaller optic nerve, tract and chiasm in PWA reported by previous studies (Schmitz et al., 2003; Von dem Hagen et al., 2005; Bridge et al., 2014). Our results indicate that DTI tractography may be able to demonstrate group level chiasmal misrouting in albinism. We found that the proportion of

tractography streamlines crossing the chiasm (the streamline decussation index) was significantly higher in PWA compared with healthy controls. These findings are validated by the weakly positive correlation between the chiasmal streamline decussation index and VEP asymmetry. It is possible that the significance of this relationship may be improved using the correlation method of VEP asymmetry assessment developed by Hoffmann, Lorenz, Morland, and Schmidtborn (2005) rather than the Apkarian method we used. This correlation

FIGURE 3 Comparison of total connectivity at the chiasm estimated using diffusion tractography with foveal development index **FIGURE 4** Comparison of foveal retinal pigment epithelium (RPE) thickness measured using OCT with the cortical thickness in patients with albinism

FIGURE 5 Comparison of cortical thickness at the occipital pole with optic disc (a) and rim (b) in patients with albinism

method uses data from the whole time series of VEP traces rather than point measurements and has been demonstrated as a reliable way of estimating the degree of misrouting (Hoffmann et al., 2005). Time series data were unavailable in the current study to allow the correlation method to be performed. Receiver operator curve (ROC) analysis of the streamline decussation index yielded an area under the curve of 0.727 (95% CI = 0.575–0.880). This indicates that while streamline decussation index may demonstrate group level differences between albinism and healthy controls, it cannot be used as a diagnostic tool.

It had been hypothesised by the authors of these earlier studies that the finding of smaller optic nerves, chiasm and tract in albinism could be due to the underdevelopment of the fovea (Von dem Hagen et al., 2005; Mcketton et al., 2014). Although we did not see any relationship between foveal development and the physical size of the chiasm, our DTI data shows significant relationship between foveal development and number of streamlines crossing the chiasm.

While this may seem surprising at first glance, a previous study in albino ferrets reported that a delay in the timing of axonal outgrowth from the retina means that there is a disruption in the distribution of large and small diameter axons within the optic nerve with an abnormal thickening of the myelin sheath (Guibal & Baker, 2009). Consequently, a gross measurement of the optic nerve and chiasm

might not accurately reflect the number of axons within it. Our data suggests that diffusion tractography may reflect the number of axons crossing the chiasm better than morphometry in people with albinism.

We found that the optic nerve size measured using MRI is correlated with the ppRNFL thickness. Using ex-vivo axon tracing studies and through mapping of the visual field to the optic nerve in glaucoma patients, previous studies have indicated that axons from the foveal retinal ganglion cells aggregate in the temporal region of the optic nerve head (Yucel et al., 1998; Zangwill, Bowd, & Weinreb, 2000; Sihota, Sony, Gupta, Dada, & Singh, 2006). Therefore, any variation in the numbers of central ganglion cells would influence the size of the ppRNFL and hence the optic nerve size. Previous animal studies have shown a reduced number of central retinal ganglion cells in albino animals (Stone, Rowe, & Campion, 1978; Guillery et al., 1984; Leventhal & Creel, 1985; Robinson, Horsburgh, Dreher, & McCall, 1987; Donatien, Aigner, & Jeffery, 2002).

However, in our study, the degree of misrouting did not relate to any foveal or optic nerve head abnormalities. Foveal hypoplasia and misrouting of the optic nerve are two cardinal features of albinism. Aberration in the melanin synthesis pathway is believed to be the cause of both these abnormalities (Jeffery, 1997). However, our findings suggest that there is no direct relationship between these two features. These findings agree with previous suggestion by Neveu et al. who compared the retinal findings in PWA and aniridia. Both these conditions are characterised by foveal hypoplasia but patients with aniridia have normal retino-fugal projections. The authors therefore concluded that optic chiasm formation is independent from foveal development (Neveu, Holder, Sloper, & Jeffery, 2005). It is more likely that the misrouting in albinism is a function of delayed cell mitosis in albinism, which is a process regulated by L-dopa, a precursor of melanin (Ilia & Jeffery, 1999). The factor determining whether an axon will decussate is thought to be dependent on the timing at which it reaches the chiasm during the development of the optic nerve. In animal models it has been shown that axons originating in the temporal retina, which develop earlier than those originating in the nasal retina, remain ipsilateral as they grow backward past the chiasm, while the later developing retinally derived axons decussate through the chiasm (Drager, 1985). It is proposed that in albinism, a lack of melanin in the retinal pigment epithelium leads to a delay in the development of

FIGURE 6 Comparison of visual evoked potential and diffusion streamline asymmetry in albinism

the temporal retina and hence a delay in these axons reaching the chiasm leading to increased decussation (Jeffery, 1997, 1998; Ilia & Jeffery, 1999). Albinism patients do however retain some normal projection and the degree of this is related to the amount of pigmentation (Hoffmann & Dumoulin, 2015). Retinal hypopigmentation is not the only cause for chiasmal misrouting since it has also been shown to occur in normally pigmented individuals. It has been hypothesised, therefore, that chiasmal misrouting may indeed be the cause of abnormal development of the fovea (Van Genderen et al., 2006). However, the relationship between chiasmal misrouting and foveal development is complex as foveal hypoplasia can also be present in the absence of chiasmal misrouting (Sloper, 2006; Hingorani, Hanson, & van Heyningen, 2012).

Apart from albinism, abnormally increased chiasmal decussation has recently been reported in foveal hypoplasia, optic nerve decussation defects and anterior segment dysgenesis (FHONDA) syndrome. This is a rare autosomal recessive disorder with 20 reported cases in published literature (Al-Araimi et al., 2013). In 2018, Ahmadi et al. studied two affected individuals using ultra-high field fMRI and found that the degree of misrouting does not vary in FHONDA syndrome unlike albinism where misrouting has been shown to vary between 2° and 15° (Hoffmann et al., 2005), and correlates with the pigment deficit (Von dem Hagen, Houston, Hoffmann, & Morland, 2007). Within the visual cortex of FHONDA patients, the normal representation of the temporal retina seen in albinism is also absent. The authors suggest that the misrouting may be due to complete cessation of uncrossed projections at the optic chiasm which points to a different molecular cascades driving misrouting in FHONDA compared with albinism (Ahmadi et al., 2018).

5.2 | Cortical abnormalities in albinism

Using surface based analysis, we have been able to shed light on a conflict in previous literature regarding the nature of structural changes within the visual cortex of PWA. Using voxel based morphometry, Von dem Hagen et al. found that PWA have a reduction in grey matter volume in the occipital cortex (Von dem Hagen et al., 2005). A more recent study by Bridge et al. used surface based analysis to conclude that visual cortex thickness is increased in PWA (Bridge et al., 2014). Bridge et al. suggested that the difference in results between the two previous studies were due to the two different analysis techniques being employed.

Using a similar technique to Bridge et al., we have found that PWA do indeed have increased cortical thickness at the occipital pole. Bridge et al. found reduced gyrification in PWA, which might explain why the earlier voxel based morphometry study may have reported reduced cortical volume in albinism (Von dem Hagen et al., 2005; Bridge et al., 2014). We previously reported corroboratory evidence using functional MRI that an increased interhemispheric functional connectivity of the visual processing areas is present in albinism, which may be an adaptation to the upstream structural changes in the visual pathway (Welton et al., 2017).

OCT data regarding foveal and optic nerve abnormalities has allowed us to explore possible anterior pathway causes behind cortical

changes seen in albinism. We noted several significant relationships of cortical thickness with the fovea, optic nerve and chiasm.

Comparison of the visual cortex with the fovea showed that cortical thickness was inversely related to the size of the RPE in PWA. The thickness of the RPE measured on OCT is impacted by the amount of melanin present within the RPE cells. This is due to the optical properties of melanin (Wolbarsht, Walsh, & George, 1981), which mean that the light from the OCT device is scattered when it passes through melanin leading to the thick band like appearance of the RPE seen in OCT images (Chauhan & Marshall, 1999). This indicates that the amount of melanin within the RPE of PWA affects the specialisation of the visual cortex. Von dem Hagen et al. have previously found that level of skin pigmentation is related to the degree of functional reorganisation of the visual cortex (Von dem Hagen et al., 2007). Our results suggest that in addition to functional changes, pigmentation defects in albinism also lead to structural changes of the visual cortex.

Bridge et al. noted that the thicker visual cortex in albinism is consistent with findings from early blind (Jiang et al., 2009) and anophthalmic (Bridge, Cowey, Ragge, & Watkins, 2009) individuals and suggested that this is due to a lack of pruning during development. In addition, increased chiasmal decussation means that there is a reduction in binocular competition at V1, which may be another factor in driving axonal pruning. The visual cortex undergoes rapid expansion during foetal and first 4 months of post-natal life and reaches peak levels (~150% of adult) by 7 months gestation (Goswami, 2004). This early post-natal time corresponds with a critical period of foveal (Lee et al., 2015) and visual cortex development (Huttenlocher & de Courten, 1987; Leuba & Garey, 1987).

The rapid growth phase is followed by synaptic revision with loss of the excess 40% of synapses between ages 8 months and 11 years. Subsequently, these synapse numbers remain stable into adulthood (Garey, 1984). The synaptic elimination has shown to be dependent on visual experience (Bourgeois, Jastreboff, & Rakic, 1989). In albinism reduced foveal cone density results in a lack of high-resolution input to V1. However, our results showed no relationship between cortical thickness at the occipital pole and foveal development. We were unable to reproduce the negative correlation between V1 cortical thickness and visual acuity demonstrated by Bridge et al. ($r = .116$, $p = .606$). This may be due to the fact that the visual deficit in albinism is multifaceted with factors such as nystagmus, refractive errors, strabismus, iris transillumination, foveal hypoplasia, optic nerve dysgenesis and chiasmal abnormalities all playing a role.

While comparing the occipital pole to optic nerve head, we found that the disc and rim areas were positively correlated with cortical thickness at the occipital pole. We have previously shown that the rim size is increased in PWA possibly due to arrest in normal embryological development of the optic nerve (Mohammad et al., 2015). The nasal aspect of the rim appears to be composed of glial tissue, which is remnant of the hyaloid vascular system that has failed to fully regress (Jones, 1963; Renz & Vygantas, 1977; Sheth, Sharma, & Chakraborty, 2013). This would support the theory that a lack of pruning is responsible for increased thickness of the visual cortex seen in albinism and that it is a phenomenon that affects more than one location in the visual pathway.

5.3 | Limitations of the study

It is important to point out the inherent limitations in our methodology. Diffusion tractography allows noninvasive in-vivo quantification of white matter structure but many factors including anatomical characteristics of the structure being studied, image acquisition parameters and choice of tract reconstruction algorithm can significantly alter the results. The anterior optic pathways are particularly challenging to study with DTI due to the complex convergence, divergence and crossing of axons as they pass through the chiasm. Within each voxel, there may be multiple fibre orientations of axons making it difficult to distinguish between axons that are kissing, crossing, converging or diverging as they all capable of generating a similar diffusion signal. This means the tractography algorithm may jump between two fibre pathways. There is also potential for partial volume effects of cerebrospinal fluid contamination affecting the tractography algorithm in voxels along the surface of the cisternal segments, and the effect of susceptibility distortions due to the adjacent skull base and paranasal sinuses.

Prior to commencing the study, we undertook optimisation of the DTI protocol by selecting the maximal resolution achievable (1.8 mm isotropic voxel size) while maintaining an acceptable signal to noise ratio and appropriate scan duration. As the structures we were sampling are very small, the mean size of the optic nerve and tract in albinism group for example were 12.3 ± 3.21 and 10.7 ± 3.33 voxels, respectively, we remain cautious regarding the interpretation of absolute streamline counts in our data but feel that expressing the streamline decussation as a percentage of the total streamline count provides a plausible measure of fibre crossing at the chiasm given the positive correlation that we found with VEP asymmetry. The ROC analysis of the streamline decussation index indicated that while the technique was able to show group differences, it should not be used for individual patient diagnosis.

Since we acquired our data between 2010 and 2012, work on the Human Connectome Project and other similar connectomic projects has significantly advanced the image acquisition techniques with the current standard being multi-shell acquisitions and measures to compensate geometric distortions such as acquisition of field maps or dual phase encoding directions. Further work using state-of-the-art image acquisition and analysis techniques such as sparse fascicle model (Rokem et al., 2015) and filtering of streamlines (Pestilli, Yeatman, Rokem, Kay, & Wandell, 2014; Wandell, 2016) is warranted to further investigate chiasmal abnormalities and may account for the unexpected results we encounter such as no group differences in the number of streamlines between albinism and control groups and the underestimation of contralateral streamlines in both groups. The mean decussation with DTI in the control group = $27.8\% \pm 17.5$ rather than the expected 50%.

In conclusion, our study provides novel insights in to the relationship between retinal, chiasmal and cortical abnormalities in albinism through the use of multiple and complimentary noninvasive imaging modalities. We show for the first time that cortical abnormalities are related to pigmentation levels of the RPE and axonal disorganisation of the optic nerve head. Although the study was not designed as a diagnostic accuracy study, we find that diffusion tractography can

demonstrate abnormal chiasmal crossing seen in albinism that relates to VEP asymmetry. Our results suggest that that much like other abnormalities in the anterior visual pathway, the cortical abnormalities in people with albinism represent abnormal embryological and early post-natal development.

ORCID

Sarim Ather <https://orcid.org/0000-0001-9614-5033>

Rob A. Dineen <https://orcid.org/0000-0002-9523-2546>

REFERENCES

- Ahmadi, K., Fracasso, A., van Dijk, J. A., Kruijt, C., van Genderen, M., Dumoulin, S. O., & Hoffmann, M. B. (2018). Altered organization of the visual cortex in FHONDA syndrome. *NeuroImage*, pii, 53.
- Akeo, K., Shirai, S., Okisaka, S., Shimizu, H., Miyata, H., Kikuchi, A., ... Majima, A. (1996). Histology of fetal eyes with oculocutaneous albinism. *Archives of Ophthalmology*, 114, 613–616.
- Al-Araimi, M., Pal, B., Poulter, J. A., van Genderen, M. M., Carr, I., Cudrnak, T., & Toomes, C. (2013). A new recessively inherited disorder composed of foveal hypoplasia, optic nerve decussation defects and anterior segment dysgenesis maps to chromosome 16q23.3-24.1. *Molecular Vision*, 19, 2165–2172.
- Andersson, J. L., & Skare, S. (2002). A model-based method for retrospective correction of geometric distortions in diffusion-weighted EPI. *NeuroImage*, 16, 177–199.
- Apkarian, P., Reits, D., Spekreijse, H., & Van Dorp, D. (1983). A decisive electrophysiological test for human albinism. *Electroencephalography and Clinical Neurophysiology*, 55, 513–531.
- Baker, G. E., & Reese, B. E. (1993). Chiasmatic course of temporal retinal axons in the developing ferret. *The Journal of Comparative Neurology*, 330, 95–104.
- Beaulieu, C. (2002). The basis of anisotropic water diffusion in the nervous system—A technical review. *NMR in Biomedicine*, 15, 435–455.
- Behrens, T. E., Berg, H. J., Jbabdi, S., Rushworth, M. F., & Woolrich, M. W. (2007). Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *NeuroImage*, 34, 144–155.
- Bhansali, P., Rayport, I., Rebsam, A., & Mason, C. (2014). Delayed neurogenesis leads to altered specification of ventrotemporal retinal ganglion cells in albino mice. *Neural Development*, 9, 11.
- Bourgeois, J. P., Jastreboff, P. J., & Rakic, P. (1989). Synaptogenesis in visual cortex of normal and preterm monkeys: Evidence for intrinsic regulation of synaptic overproduction. *Proceedings of the National Academy of Sciences of the United States of America*, 86, 4297–4301.
- Bridge, H., Cowey, A., Ragge, N., & Watkins, K. (2009). Imaging studies in congenital anophthalmia reveal preservation of brain architecture in 'visual' cortex. *Brain*, 132, 3467–3480.
- Bridge, H., von dem Hagen, E. A., Davies, G., Chambers, C., Gouws, A., Hoffmann, M., & Morland, A. B. (2014). Changes in brain morphology in albinism reflect reduced visual acuity. *Cortex*, 56, 64–72.
- Chauhan, D. S., & Marshall, J. (1999). The interpretation of optical coherence tomography images of the retina. *Investigative Ophthalmology & Visual Science*, 40, 2332–2342.
- Chong, G. T., Farsiu, S., Freedman, S. F., Sarin, N., Koreishi, A. F., Izatt, J. A., & Toth, C. A. (2009). Abnormal foveal morphology in ocular albinism imaged with spectral-domain optical coherence tomography. *Archives of Ophthalmology*, 127, 37–44.
- Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I. Segmentation and surface reconstruction. *NeuroImage*, 9, 179–194.
- Destrieux, C., Fischl, B., Dale, A., & Hagren, E. (2010). Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *NeuroImage*, 53, 1–15.
- Donatien, P., Aigner, B., & Jeffery, G. (2002). Variations in cell density in the ganglion cell layer of the retina as a function of ocular pigmentation. *The European Journal of Neuroscience*, 15, 1597–1602.

- Drager, U. C. (1985). Birth dates of retinal ganglion cells giving rise to the crossed and uncrossed optic projections in the mouse. *Proceedings of the Royal Society of London - Series B: Biological Sciences*, 224, 57–77.
- Elschnig, A. (1913). Zur Anatomie des menschlichen Albinoauges. *Graefes Archiv für Ophthalmologie*, 84, 401–419.
- Erskine, L., & Herrera, E. (2014). Connecting the retina to the brain. *ASN Neuro*, 6, 175909141456210.
- Fischl, B., & Dale, A. M. (2000). Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 11050–11055.
- Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *NeuroImage*, 9, 195–207.
- Garcia-Frigola, C., Carreres, M. I., Vegar, C., Mason, C., & Herrera, E. (2008). Zic2 promotes axonal divergence at the optic chiasm midline by EphB1-dependent and -independent mechanisms. *Development*, 135, 1833–1841.
- Garey, L. J. (1984). Structural development of the visual system of man. *Human Neurobiology*, 3, 75–80.
- Goswami, U. (2004). Neuroscience and education. *The British Journal of Educational Psychology*, 74, 1–14.
- Gottlob, I., & Proudlock, F. A. (2014). Aetiology of infantile nystagmus. *Current Opinion in Neurology*, 27, 83–91.
- Grigorian, A., Mcketton, L., & Schneider, K. A. (2016). Measuring connectivity in the primary visual pathway in human albinism using diffusion tensor imaging and Tractography. *Journal of Visualized Experiments*, 23, 1–10.
- Guibal, C., & Baker, G. E. (2009). Abnormal axons in the albino optic tract. *Investigative Ophthalmology & Visual Science*, 50, 5516–5521.
- Guillery, R. W., Okoro, A. N., & Witkop, C. J. (1975). Abnormal visual pathways in the brain of a human albino. *Brain Research*, 96, 373–377.
- Guillery, R. W., Hickey, T. L., Kaas, J. H., Felleman, D. J., Debruyne, E. J., & Sparks, D. L. (1984). Abnormal central visual pathways in the brain of an albino green monkey (*Cercopithecus aethiops*). *The Journal of Comparative Neurology*, 226, 165–183.
- Hingorani, M., Hanson, I., & van Heyningen, V. (2012). Aniridia. *European Journal of Human Genetics*, 20, 1011–1017.
- Hoffmann, M. B., & Dumoulin, S. O. (2015). Congenital visual pathway abnormalities: A window onto cortical stability and plasticity. *Trends in Neurosciences*, 38, 55–65.
- Hoffmann, M. B., Lorenz, B., Morland, A. B., & Schmidtborn, L. C. (2005). Misrouting of the optic nerves in albinism: Estimation of the extent with visual evoked potentials. *Investigative Ophthalmology & Visual Science*, 46, 3892–3898.
- Holm, S. A. (1979). Simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6, 65–70.
- Huttenlocher, P. R., & de Courten, C. (1987). The development of synapses in striate cortex of man. *Human Neurobiology*, 6, 1–9.
- Ilija, M., & Jeffery, G. (1999). Retinal mitosis is regulated by dopa, a melanin precursor that may influence the time at which cells exit the cell cycle: Analysis of patterns of cell production in pigmented and albino retinas. *The Journal of Comparative Neurology*, 405, 394–405.
- Jeffery, G. (1998). The retinal pigment epithelium as a developmental regulator of the neural retina. *Eye (London, England)*, 12(Pt 3b), 499–503.
- Jeffery, G. (1997). The albino retina: An abnormality that provides insight into normal retinal development. *Trends in Neurosciences*, 20, 165–169.
- Jiang, J., Zhu, W., Shi, F., Liu, Y., Li, J., Qin, W., ... Jiang, T. (2009). Thick visual cortex in the early blind. *The Journal of Neuroscience*, 29, 2205–2211.
- Jones, H. E. (1963). Hyaloid remnants in the eyes of premature babies. *The British Journal of Ophthalmology*, 47, 39–44.
- Kamaraj, B., & Purohit, R. (2014). Mutational analysis of oculocutaneous albinism: A compact review. *BioMed Research International*, 2014, 905472.
- Kralj-Hans, I., Tibber, M., Jeffery, G., & Mobbs, P. (2006). Differential effect of dopamine on mitosis in early postnatal albino and pigmented rat retinas. *Journal of Neurobiology*, 66, 47–55.
- Lee, H., Purohit, R., Patel, A., Papageorgiou, E., Sheth, V., Maconachie, G., ... Gottlob, I. (2015). In vivo foveal development using optical coherence tomography. *Investigative Ophthalmology & Visual Science*, 56, 4537–4545.
- Leuba, G., & Garey, L. J. (1987). Evolution of neuronal numerical density in the developing and aging human visual cortex. *Human Neurobiology*, 6, 11–18.
- Leventhal, A. G., & Creel, D. J. (1985). Retinal projections and functional architecture of cortical areas 17 and 18 in the tyrosinase-negative albino cat. *The Journal of Neuroscience*, 5, 795–807.
- Levin, A. V., & Stroh, E. (2011). Albinism for the busy clinician. *Journal of AAPOS*, 15, 59–66.
- Mcketton, L., Kelly, K. R., & Schneider, K. A. (2014). Abnormal lateral geniculate nucleus and optic chiasm in human albinism. *The Journal of Comparative Neurology*, 522, 2680–2687.
- Mohammad, S., Gottlob, I., Sheth, V., Pilat, A., Lee, H., Pollheimer, E., & Proudlock, F. A. (2015). Characterization of abnormal optic nerve head morphology in albinism using optical coherence tomography. *Investigative Ophthalmology & Visual Science*, 56, 4611–4618.
- Mohammad, S., Gottlob, I., Kumar, A., Thomas, M., Degg, C., Sheth, V., & Proudlock, F. A. (2011). The functional significance of foveal abnormalities in albinism measured using spectral-domain optical coherence tomography. *Ophthalmology*, 118, 1645–1652.
- Montoliu, L., Gronskov, K., Wei, A. H., Martinez-Garcia, M., Fernandez, A., Arveiler, B., & Li, W. (2014). Increasing the complexity: New genes and new types of albinism. *Pigment Cell & Melanoma Research*, 27, 11–18.
- Naumann, G. O., Lerche, W., & Schroeder, W. (1976). Foveolar aplasia in tyrosinase-positive oculocutaneous albinism (author's transl). *Albrecht von Graefes Archiv für Klinische und Experimentelle Ophthalmologie*, 200, 39–50.
- Neveu, M. M., von dem Hagen, E., Morland, A. B., & Jeffery, G. (2008). The fovea regulates symmetrical development of the visual cortex. *The Journal of Comparative Neurology*, 506, 791–800.
- Neveu, M. M., Holder, G. E., Sloper, J. J., & Jeffery, G. (2005). Optic chiasm formation in humans is independent of foveal development. *The European Journal of Neuroscience*, 22, 1825–1829.
- Odom, J. V., Bach, M., Brigell, M., Holder, G. E., McCulloch, D. L., & Tormene, A. P. (2010). ISCEV standard for clinical visual evoked potentials (2009 update). *Documenta Ophthalmologica*, 120, 111–119.
- Papageorgiou, E., McLean, R. J., & Gottlob, I. (2014). Nystagmus in childhood. *Pediatrics and Neonatology*, 55, 341–351.
- Pestilli, F., Yeatman, J. D., Rokem, A., Kay, K. N., & Wandell, B. A. (2014). Evaluation and statistical inference for human connectomes. *Nature Methods*, 11, 1058–1063.
- Prieur, D. S., & Rebsam, A. (2016). Retinal axon guidance at the midline: Chiasmatic misrouting and consequences. *Developmental Neurobiology*, 77, 844–860.
- Provis, J. M. (2001). Development of the primate retinal vasculature. *Progress in Retinal and Eye Research*, 20, 799–821.
- Provis, J. M., & Hendrickson, A. E. (2008). The foveal avascular region of developing human retina. *Archives of Ophthalmology*, 126, 507–511.
- Provis, J. M., Sandercoe, T., & Hendrickson, A. E. (2000). Astrocytes and blood vessels define the foveal rim during primate retinal development. *Investigative Ophthalmology & Visual Science*, 41, 2827–2836.
- Rachel, R. A., Dolen, G., Hayes, N. L., Lu, A., Erskine, L., Nowakowski, R. S., & Mason, C. A. (2002). Spatiotemporal features of early neurogenesis differ in wild-type and albino mouse retina. *The Journal of Neuroscience*, 22, 4249–4263.
- Rebsam, A., Bhansali, P., & Mason, C. A. (2012). Eye-specific projections of retinogeniculate axons are altered in albino mice. *The Journal of Neuroscience*, 32, 4821–4486.
- Renz, B. E., & Vygantas, C. M. (1977). Hyaloid vascular remnants in human neonates. *Annals of Ophthalmology*, 9, 179–184.
- Robinson, S. R., Horsburgh, G. M., Dreher, B., & McCall, M. J. (1987). Changes in the numbers of retinal ganglion cells and optic nerve axons in the developing albino rabbit. *Developmental Brain Research*, 35, 161–174.
- Rokem, A., Yeatman, J. D., Pestilli, F., Kay, K. N., Mezer, A., van der Walt, S., & Wandell, B. A. (2015). Evaluating the accuracy of diffusion MRI models in white matter. *PLoS One*, 10, e0123272.
- Schmitz, B., Schaefer, T., Krick, C. M., Reith, W., Backens, M., & Kasmann-Kellner, B. (2003). Configuration of the optic chiasm in

- humans with albinism as revealed by magnetic resonance imaging. *Investigative Ophthalmology & Visual Science*, 44, 16–21.
- Schutze, C., Ritter, M., Blum, R., Zotter, S., Baumann, B., Pircher, M., & Christoph, K. (2014). Retinal pigment epithelium findings in patients with albinism using wide-field polarization-sensitive optical coherence tomography. *Retina*, 34, 2208–2217.
- Sheth, J. U., Sharma, A., & Chakraborty, S. (2013). Persistent hyaloid artery with an aberrant peripheral retinal attachment: A unique presentation. *Oman Journal of Ophthalmology*, 6, 58–60.
- Sihota, R., Sony, P., Gupta, V., Dada, T., & Singh, R. (2006). Diagnostic capability of optical coherence tomography in evaluating the degree of glaucomatous retinal nerve fiber damage. *Investigative Ophthalmology & Visual Science*, 47, 2006–2010.
- Sloper, J. (2006). Chicken and egg. *The British Journal of Ophthalmology*, 90, 1074–1075.
- Stone, J., Rowe, M. H., & Campion, J. E. (1978). Retinal abnormalities in the Siamese cat. *The Journal of Comparative Neurology*, 180, 773–782.
- Thomas, M. G., Kumar, A., Mohammad, S., Proudlock, F. A., Engle, E. C., Andrews, C., ... Gottlob, I. (2011). Structural grading of foveal hypoplasia using spectral-domain optical coherence tomography: A predictor of visual acuity? *Ophthalmology*, 118, 1653–1660.
- Van Genderen, M. M., Riemsdijk, F. C., Schuil, J., Hoebe, F. P., Stijl, J. S., & Meire, F. M. (2006). Chiasmal misrouting and foveal hypoplasia without albinism. *The British Journal of Ophthalmology*, 90, 1098–1102.
- Von dem Hagen, E. A., Houston, G. C., Hoffmann, M. B., & Morland, A. B. (2007). Pigmentation predicts the shift in the line of decussation in humans with albinism. *The European Journal of Neuroscience*, 25, 503–511.
- Von dem Hagen, E. A., Houston, G. C., Hoffmann, M. B., Jeffery, G., & Morland, A. B. (2005). Retinal abnormalities in human albinism translate into a reduction of grey matter in the occipital cortex. *The European Journal of Neuroscience*, 22, 2475–2480.
- Wandell, B. A. (2016). Clarifying human white matter. *Annual Review of Neuroscience*, 39, 103–128.
- Welton, T., Ather, S., Proudlock, F. A., Gottlob, I., & Dineen, R. A. (2017). Altered whole-brain connectivity in albinism. *Human Brain Mapping*, 38, 740–752.
- Wolbarsht, M. L., Walsh, A. W., & George, G. (1981). Melanin, a unique biological absorber. *Applied Optics*, 20, 2184–2186.
- Yucel, Y. H., Gupta, N., Kalichman, M. W., Mizisin, A. P., Hare, W., de Souza Lima, M., ... Weinreb, R. N. (1998). Relationship of optic disc topography to optic nerve fiber number in glaucoma. *Archives of Ophthalmology*, 116, 493–497.
- Zangwill, L. M., Bowd, C., & Weinreb, R. N. (2000). Evaluating the optic disc and retinal nerve fiber layer in glaucoma. II: Optical image analysis. *Seminars in Ophthalmology*, 15, 206–220.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Ather S, Proudlock FA, Welton T, et al. Aberrant visual pathway development in albinism: From retina to cortex. *Hum Brain Mapp*. 2019;40:777–788. <https://doi.org/10.1002/hbm.24411>