RESEARCH ARTICLE

Differences in white matter structure and cortical thickness between patients with traumatic and idiopathic chronic neck pain: Associations with cognition and pain modulation?

I. Coppieters^{1,2,3} | R. De Pauw¹ | K. Caeyenberghs⁴ | D. Lenoir¹ | K. DeBlaere⁵ | E. Genbrugge⁵ | M. Meeus^{1,2,6*} | B. Cagnie^{1*}

¹Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium

²Pain in Motion International Research Group (www.paininmotion.be)

³Department of Physiotherapy, Human Physiology and Anatomy (KIMA), Faculty of Physical Education & Physiotherapy, Vrije Universiteit Brussel, Belgium

⁴School of Psychology, Faculty of Health Sciences, Australian Catholic University, Banyo, Australia

⁵Department of Radiology, Ghent University, Ghent, Belgium

⁶Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium

Correspondence

Iris Coppieters, Rehabilitation Sciences and Physiotherapy Ghent Campus Heymans (UZ) 3 B3, De Pintelaan 185, Ghent, Belgium.
Email: iris.coppieters@vub.ac.be

Funding information

Special Research Fund of Ghent University (BOF-Ghent), Grant Number: BOF13/DOC/276.

Abstract

Brain alterations are hypothesized to be present in patients with chronic whiplash-associated disorders (CWAD). The aim of this case-control study was to examine alterations in cortical thickness and white matter (WM) structure, and the presence of brain microhemorrhages in a patient group encountering chronic neck pain of traumatic origin (i.e., CWAD) when compared with a patient group characterized by nontraumatic chronic neck pain [i.e., chronic idiopathic neck pain (CINP)], and healthy controls. Furthermore, we aimed to investigate associations between brain structure on one hand and cognitive performance and central sensitization (CS) on the other hand. T1-weighted, diffusion-weighted and T2*-weighted magnetic resonance images of the brain were acquired in 105 women (31 controls, 37 CINP, 37 CWAD) to investigate regional cortical thickness, WM structure, and microhemorrhages, respectively. Next, cognitive performance, and CS encompassing distant hyperalgesia and conditioned pain modulation (CPM) efficacy were examined. Cortical thinning in the left precuneus was revealed in CWAD compared with CINP patients. Also, decreased fractional anisotropy, together with increased values of mean diffusivity and radial diffusivity could be observed in the left cingulum hippocampus and tapetum in CWAD compared with CINP, and in the left tapetum in CWAD patients compared with controls. Moreover, the extent of WM structural deficits in the left tapetum coincided with decreased CPM efficacy in the CWAD group. This yields evidence for associations between decreased endogenous pain inhibition, and the degree of regional WM deficits in CWAD. Our results emphasize the role of structural brain alterations in women with CWAD compared with CINP.

KEYWORDS

chronic idiopathic neck pain, chronic whiplash-associated disorders, central sensitization, cognitive performance, cortical thickness, magnetic resonance imaging, white matter

Abbreviations: AD, axial diffusivity; CINP, chronic idiopathic neck pain; CPM, conditioned pain modulation; CS, central sensitization; CWAD, chronic whiplash associated disorders; DTI, diffusion tensor imaging; DWI, diffusion-weighted imaging; FA, fractional anisotropy; GM, grey matter; MD, mean diffusivity; MRI, magnetic resonance imaging; NRS, numeric rating scale; PPT, pressure pain thresholds; RD, radial diffusivity; TBI, traumatic brain injury; TMT, trail making test; WM, white matter.

*Equally contributed last authors

1 | INTRODUCTION

Extensive magnetic resonance imaging (MRI) research has uncovered alterations in grey matter (GM) volume, cortical thickness, and white matter (WM) structure in regions and tracts involved in processing of pain and cognition in various chronic musculoskeletal pain conditions, such as fibromyalgia (Cagnie et al., 2014; Kim et al., 2014; Lieberman et al., 2014; Lutz et al., 2008) and chronic low back pain (Buckalew et al., 2010; Kregel et al., 2015; Mansour et al., 2013). In addition, these brain alterations have been reported in mild traumatic brain injury (TBI)

patients (Govindarajan et al., 2016; Killgore et al., 2016; Kraus et al., 2007; Wang et al., 2015). Also, in mild TBI patients evidence is available for brain microhemorrhages related to the trauma or diffuse axonal injury (Koerte, Hufschmidt, Muehlmann, Lin, & Shenton, 2016). Yet, despite this huge body of work, there have been few attempts to investigate these findings in more detail and to examine alterations in both GM and WM, as well as associations with clinical features in patients with chronic musculoskeletal pain.

Remarkably, structural brain MRI research is limited in two prevalent musculoskeletal pain conditions namely chronic whiplash-associated disorders (CWAD), involving trauma-induced persistent neck pain (Davis, 2013), and chronic idiopathic neck pain (CINP), involving persistent neck pain of nontraumatic origin. However, due to the trauma inducing possible shearing forces through acceleration-deceleration of the brain (Alexander, 1998), it could be hypothesized that subtle structural brain alterations in GM morphology and WM structure are present in CWAD, but not or to a lesser degree in CINP patients. To investigate this hypothesis and to unravel the underlying mechanisms of a wide range of possible structural brain alterations, it seemed valuable to compare patients with CINP and CWAD, who are *a priori* different from each other in the origin of pain. Furthermore, this research could explore pathophysiological differences between both chronic neck pain groups which is necessary to improve their treatment outcome.

Intriguingly, many chronic musculoskeletal pain disorders are characterized by central sensitization (CS) as overlapping pathophysiological mechanism, and often experience cognitive problems (Moriarty, McGuire, & Finn, 2011; Yunus, 2007). In addition, similar to chronic pain patients, mild TBI patients often report cognitive complaints (Bay, Kalpakjian, & Giordani, 2012) and chronic pain (Nampiamparmpil, 2008). These findings indicate that chronic pain is associated with structural brain plasticity, and suggest that cognitive deficits and CS are related to this plasticity (Apkarian, Hashmi, & Baliki, 2011; Davis & Moayed, 2013). Also, it has been shown that GM and WM neuroplasticity is associated with the traumatic factor in mild TBI (Govindarajan et al., 2016; Killgore et al., 2016; Kraus et al., 2007; Wang et al., 2015).

Compelling evidence has shown CS in patients with CWAD (Van Oosterwijck, Nijs, Meeus, & Paul, 2013). In contrast, CS is not a characteristic feature in CINP patients (Malfliet et al., 2015; Scott, Jull, & Sterling, 2005), and cognitive deficits are observed to a lesser extent compared with CWAD (Coppieters et al., 2017b). A recent systematic review concerning brain alterations in WAD and INP, only found three studies investigating structural brain alterations exclusively in WAD (De Pauw, Coppieters, Meeus, Caeyenberghs, & Danneels, 2017), and one study reported altered GM volume in CWAD. However, alterations in cortical thickness and WM structure, and their association with cognitive deficits and CS, have never been investigated in patients with CWAD compared with CINP, which is a prominent research gap.

In order to disentangle subtle structural brain alterations in patients with CWAD compared with nontraumatic CINP, innovative advanced MRI, in particular, diffusion-weighted imaging (DWI) acquisition techniques, and diffusion tensor imaging (DTI) analyses offer opportunities to investigate the brain's WM tissue at a (micro)structural level (Basser, Mattiello, & LeBihan, 1994; Basser & Pierpaoli, 1996; Beaulieu, 2002;

Jones & Leemans, 2011; Le Bihan et al., 2001; Tournier, Mori, & Leemans, 2011). Hence, these techniques will probably be more sensitive for detecting subtle structural brain changes in CWAD (Suleyman Sener & Maas Andrew, 2016). In particular, DWI examines WM structural organization by quantifying the directionality and degree of diffusion of water within tissues (Basser et al., 1994; Basser & Pierpaoli, 1996; Beaulieu, 2002; Jones & Leemans, 2011; Le Bihan et al., 2001; Tournier et al., 2011). In addition, T2*-weighted MRI can assess the presence or absence of brain microhemorrhages related to the trauma in CWAD. Furthermore, MRI techniques investigating cortical thickness could be valuable to extract additional information of the brain's GM morphology. It is suggested that cortical thickness and GM volume reflect different aspects of the neural architecture (Rakic, 1988). Examining cortical thickness could therefore provide additional insights into the macrostructural neural underpinnings of CWAD and CINP.

The first study objective was to investigate alterations in cortical thickness in carefully selected brain regions based on hypothesis-driven predictions regarding the localization of the expected structural brain alterations in specific regions involved in processing of cognition and/or pain in patients with CWAD compared with CINP and healthy controls. The second objective was to examine deficits in WM structure, including alterations in fractional anisotropy (FA), mean diffusivity (MD), axial diffusivity (AD), and radial diffusivity (RD) in hypothesis-driven predefined WM regions/tracts carrying information between regions involved in processing of cognition and/or pain in patients with CWAD compared with CINP, and healthy controls. The third objective was to examine the presence or absence of microhemorrhages related to trauma in CWAD patients compared with the other study groups. The final objective was to explore in each group separately whether alterations in regional cortical thickness and WM structure were associated with cognitive performance and CS.

Based on *a priori* cortical thickness and WM anatomical hypotheses, cortical thinning, and deficits in WM structure of predefined regions/tracts were hypothesized in CWAD due to the trauma, and because of cognitive deficits (Coppieters et al., 2015; Van Oosterwijck et al., 2013) compared with CINP patients and healthy controls. Also, previous MRI studies revealed associations between altered cortical thickness and WM structure, and measures of cognition and pain in various chronic pain conditions (Coppieters et al., 2016; Luerding, Weigand, Bogdahn, & Schmidt-Wilcke, 2008). Therefore, it is hypothesized that alterations in cortical thickness and WM structure in CWAD are associated with worse cognitive performance and signs of CS in CWAD compared with CINP patients.

To guide the selection of cortical regions of interest (ROIs) from the Desikan parcellation scheme (Desikan et al., 2006) and the selection of WM tracts/ROIs from the Mori atlas (Mori et al., 2008) (see Figure 1) evidence was retrieved from our systematic review appraising the evidence for brain alterations in INP and WAD (De Pauw et al., 2017), based on previous studies in patients with chronic pain or mild TBI examining GM morphological alterations (Cagnie et al., 2014; Cauda et al., 2014; Coppieters et al., 2017a; Govindarajan et al., 2016; Kregel et al., 2015; Wang et al., 2015), demonstrating WM structural alterations, and regarding studies exploring associations between cortical thickness or WM

FIGURE 1 Schematic representation of cortical regions of interest (presented in blue) and WM regions/tracts of interest (presented in red) of cognitive and/or pain processing networks and regions, and their proposed function. Target cortical regions of the salience, central executive, default mode and somatosensory network were selected from the Desikan atlas (Desikan et al., 2006). The WM regions/tracts of interest were selected from the ICBM-DTI-81 WM labels atlas, developed by Mori et al. (2008). All regions (except for the corpus callosum) were included from both hemispheres. The careful selection of these regions of interest was based on evidence retrieved from our systematic review appraising the evidence for brain alterations in INP and WAD (De Pauw et al., 2017), on previous studies in patients with chronic pain or mild TBI examining GM morphological alterations (Cagnie et al., 2014; Cauda et al., 2014; Coppieters et al., 2017a; Govindarajan et al., 2016; Kregel et al., 2015; Wang et al., 2015), demonstrating WM structural alterations, and regarding studies exploring associations between cortical thickness or WM structure, and measures of cognition and pain (Baek et al., 2013; Borsook et al., 2016; Chen et al., 2011; Coppieters et al., 2016; De Souza et al., 2014; Erpelding et al., 2012; Kim et al., 2014; Koenig et al., 2015; Kraus et al., 2007; Lieberman et al., 2014; Lin et al., 2015; Ly et al., 2016; Mori et al., 2008; Piche et al., 2013; Sarikcioglu et al., 2007; Schwedt & Chong, 2014; Todd & Lempert, 2007; Upadhyay et al., 2010). In addition, only regions demonstrated to be mainly involved in processing pain or cognition or the combination thereof could be selected [Color figure can be viewed at wileyonlinelibrary.com]

structure, and measures of cognition and pain (Baek et al., 2013; Borsook et al., 2016; Chen, Blankstein, Diamant, & Davis, 2011; Coppieters et al., 2016; DeSouza, Hodaie, & Davis, 2014; Erpelding, Moayed, & Davis, 2012; Kim et al., 2014; Koenig et al., 2015; Kraus et al., 2007; Lieberman et al., 2014; Lin et al., 2015; Ly et al., 2016; Mori et al., 2008; Piche et al., 2013; Sarikcioglu, Ozsoy, & Unver, 2007; Schwedt & Chong, 2014; Todd & Lempert, 2007; Upadhyay et al., 2010). In addition, only regions demonstrated to be mainly involved in processing pain or cognition or the combination thereof could be selected.

2 | METHODS

2.1 | Participants

One hundred and five female participants—37 patients with CINP, 37 patients with CWAD, and 31 healthy pain-free controls—were enrolled.

A subject sample presenting overlap with the current study sample was used in our recent study with regard to GM volume alterations (Coppieters et al., 2017a), and concerning clinical differences between women with CINP and CWAD (Coppieters et al., 2017b). In order to exclude the confounding factor of gender on brain structure and pain modulation, only women were included, as research has demonstrated differences between men and women regarding brain structure, pain sensitivity and processing in healthy persons and pain patients (Bulls et al., 2015; Ruigrok et al., 2014; van Hemmen et al., 2017). All participants were Dutch native speakers between 18 and 65 years old. Participants were recruited by calls on social media and through advertisements placed in health magazines and a patient information brochure, on the Ghent University website, and via local radio. Furthermore, informative flyers and posters were distributed in different medical institutes and associations in Flanders, Belgium (various hospitals, medical physician practices, and physical therapist practices). As general

inclusion criteria, approximately 90% of all participants were right-handed and 10% left-handed. This is a representative sample, because approximately 10% of the general population is ambidextrous or left-handed (Perelle & Ehrman, 1994).

Inclusion criteria for patients with CINP and CWAD were persistent neck pain lasting more than 3 months (Breivik, Collett, Ventafridda, Cohen, & Gallacher, 2006), with a mean pain intensity of more than 3 of 10 on a Numeric Rating Scale (NRS) during the preceding month. All chronic neck pain patients had to report mild/moderate to severe pain-related disability, established by a score of 10 or more of a maximum of 50 on the Neck Disability Index (Vernon, 2008). Additionally, patients with chronic neck pain had to report stability of pain medication intake for at least 4 weeks before study participation.

A specific inclusion criterion for CINP patients was persistent idiopathic neck pain. Patients with CINP were excluded if they ever experienced whiplash trauma, or any other specific cause of neck pain, for example, cervical hernia with clinical symptoms. Patients with CWAD were included only if they had neck pain resulting from a motor vehicle crash or traumatic event and classifiable as WAD II A, B, or C on the modified Quebec Task Force Scale (Sterling, 2004). Patients with CWAD grades I, III (neurological signs), or IV (fracture or dislocation) on the modified Quebec Task Force Scale were excluded. Additionally, CWAD patients who lost consciousness as a result of the traumatic event, and who had suffered distinct posttraumatic amnesia were excluded.

Healthy women could participate only if they were pain-free on each test day (NRS score of $<2/10$), had no history of neck-shoulder-arm pain for longer than eight consecutive days during the preceding year, with a pain intensity of 2 or more of 10 on the NRS, no medical consultation for neck-shoulder-arm pain during the preceding year, and no history of whiplash trauma. Additionally, healthy controls were included only if they had a score of less than 8 of 50 on the Neck Disability Index.

General exclusion criteria for all study groups were the presence of major depression or psychiatric illness; neurologic, or cardiovascular disorders; inflammatory disorders; fibromyalgia; chronic fatigue syndrome; and a history of neck or shoulder girdle surgery. All participants completed the MRI safety checklist and participants who presented contraindications for MRI were excluded. To preclude confounding factors, all participants were asked to discontinue intake of nonopioid analgesics 48 h before study participation. In addition, participants were asked to avoid heavy physical exertion and to refrain from consuming alcohol, caffeine, and nicotine on each testing day.

2.2 | Study design and procedure

This cross-sectional case-control study took place at Ghent University in cooperation with the Ghent Institute for Functional and Metabolic Imaging. The study was performed from February 2014 to September 2015 and was carried out in accordance with the principles of the Declaration of Helsinki. The Ethics Committee of the Ghent University Hospital (EC/2013/1053) approved the research protocol. All participants were informed about the study procedures and signed an informed consent form prior to study enrolment.

First, all participants completed a survey to acquire information on demographics, and completed a series of questionnaires to obtain information on pain-related disability, pain intensity and frequency (as described below). Subsequently, experimental assessments to investigate cognitive performance and CS were performed. On a separate test day (10 ± 7 days apart), T1-weighted MR images, diffusion-weighted MR images, and T2*-weighted gradient echo MR images of the brain were acquired.

2.3 | Self-reported pain and disability measures

On each test day, participants scored current neck pain intensity on a NRS. Scores ranged from 0 to 10, with 0 reflecting 'no pain' and 10 reflecting 'the worst pain imaginable'. Patients reported their neck pain frequency in number of days per week. The Dutch Neck Disability Index was used to investigate self-reported pain-related disability levels (0–50) (Vernon, 2008). Higher scores on the Neck Disability Index indicate higher levels of pain-related disability. The Dutch language version of the Neck Disability Index has been proven to be valid and reliable to assess self-reported disability in patients with chronic neck pain (Jorritsma, de Vries, Geertzen, Dijkstra, & Reneman, 2010; Jorritsma, de Vries, Dijkstra, Geertzen, & Reneman, 2012).

2.4 | Cognitive performance

In order to obtain an objective measure of cognitive performance the Trail Making Test (TMT) was administered. This test consists of two parts, trails A and B. TMT part A requires mainly visuoperceptual (Sanchez-Cubillo et al., 2009) and processing speed (Tombaugh, 2004) abilities, whereas TMT part B reflects primarily working memory and secondarily task-switching ability (Sanchez-Cubillo et al., 2009). In trail A, the participant had to draw lines connecting 25 numbers in ascending order as fast as possible, without lifting the pencil from the paper. In trail B, the participant was instructed to draw lines alternating between numbers and letters in ascending order (going from 1 to A, from A to 2, etc.). The goal of the TMT was to finish part A and part B as accurate and as quickly as possible. The time taken to complete each part of the test was used as outcome measure. Both subtests necessitate graphomotor speed, visual scanning ability, sufficient attention, and numeric sequencing (Misdraji & Gass, 2010; Sanchez-Cubillo et al., 2009; Tombaugh, 2004). The TMT part B further requires letter sequencing, mental double tracking, and alternation (i.e., shifting between number and letter series) (Misdraji & Gass, 2010). The TMT has been proven to be valid for assessing cognitive deficits (Sanchez-Cubillo et al., 2009), and is a quickly administered, widely used neuropsychological test that provides information on a broad range of cognitive skills (Bowie & Harvey, 2006).

2.5 | Measures of central sensitization

2.5.1 | Distant hyperalgesia

Pressure pain thresholds (PPTs) were measured unilaterally with a digital manual algometer (Wagner Instruments, FDX, Greenwich, Connecticut) at a distant asymptomatic region (quadriceps muscle midway

between the anterior superior iliac spine and the basis patellae) to evaluate secondary or distant hyperalgesia (Whiteside, Hansen, & Chaudhuri, 2004). The PPTs were assessed at the most painful side (Kosek, Ekholm, & Hansson, 1999) and in a randomized order (with research randomizer, <https://www.randomizer.org>). If patients experienced the same amount of neck pain at both sides, and in healthy women, PPTs were tested at the dominant handedness side. The participant was seated and pressure was gradually increased at a rate of 1 kgf/s until the participant reported the first sensation of unpleasantness. The PPT was determined as the mean of two consecutive (30 s in between) measurements. Decreased PPTs in the patient groups compared with controls at the quadriceps muscle indicate distant hyperalgesia, and are suggestive for the presence of CS. The described PPT technique has been demonstrated to be reliable (Cathcart & Pritchard, 2006).

2.5.2 | Efficacy of conditioned pain modulation (CPM)

Endogenous pain inhibition was investigated by applying a CPM paradigm. This paradigm relies on the "pain-inhibits-pain" mechanism in which one noxious stimulus is used as a conditioning stimulus to induce a reduction in pain perception from another test stimulus (Yarnitsky, 2010). The conditioning stimulus for eliciting CPM was the cold pressor test. The PPT assessment was used as test stimulus. For the conditioning stimulus, the contralateral hand (of the PPT side) was first immersed in water maintained at room temperature (22 °C) for 1 min to standardize hand temperature (Ng, Pedler, Vicenzino, & Sterling, 2014), before immersing this hand (up to the wrist) in a refrigerated circulating bath (Versacool, Thermo Fisher Scientific, Waltham, MA) with cold water maintained at 12 ± 1 °C (Tousignant-Laflamme, Page, Goffaux, & Marchand, 2008). Participants were asked to keep their hand in the water bath for 2 min (Ng et al., 2014). Meanwhile, the PPT was re-evaluated at the quadriceps muscle, approximately 45 s after immersing the hand (again with an interval of 30 s) (Lewis, Heales, Rice, Rome, & McNair, 2012). If the participant removed the hand from the water before the end of the 2 min, the measurement was registered as missing. For analysis of CPM efficacy, the mean PPT measured before the cold pressor test was subtracted from the mean PPT measured during the cold pressor test. Hence, a lower CPM value reflected less efficient endogenous pain inhibition. The cold pressor test with cold water of 12 °C and immersion of the hand for 2 min is sufficient to induce an endogenous pain inhibitory response (Kennedy, Kemp, Ridout, Yarnitsky, & Rice, 2016).

2.6 | MRI data acquisition

MR images were acquired on a 3T Siemens Magnetom TrioTim MRI scanner (Siemens, Erlangen, Germany) equipped with a 32-channel matrix head coil, at the Ghent University Hospital. First, high-resolution T1-weighted images of the brain were acquired using a three-dimensional magnetization prepared rapid acquisition gradient echo (MP-RAGE) (repetition time [TR] = 2250 ms, echo time [TE] = 4.18 ms, voxel size = $1 \times 1 \times 1$ mm³, field of view (FoV) = 256×256 mm, flip angle = 9°, 176 slices, 1 mm slice

thickness, acquisition time = 5'14"). All T1-weighted anatomical scans were visually checked for overall quality and motion artefacts.

Second, DW images of the brain were acquired using single-shot echo planar imaging with a twice-refocused spin echo sequence with FoV = 240×240 mm, TR = 10,800 ms, TE = 83 ms, and 60 contiguous sagittal slices (slice thickness = 2.5 mm; voxel size = $2.5 \times 2.5 \times 2.5$ mm³), and an acquisition time of 12' 36". Diffusion sensitizing gradients were applied at a *b* value of 1200 s/mm², along 64 noncollinear directions. Additionally, one set of images with no diffusion weighting (*b*-value = 0 s/mm²) was acquired as a reference image.

Finally, axial T2*-weighted brain images were acquired using a T2*-weighted acquisition gradient echo with TR = 839 ms, TE = 18.60 ms, voxel size = $1 \times 0.7 \times 3$ mm³, FoV = 230×230 mm, flip angle = 20°, 3 mm slice thickness, and an acquisition time of 3' 48". All T2*-weighted images were visually inspected by two expert neuroradiologists (KD, EG) to evaluate possible microhemorrhages or hemorrhagic shearing lesions related to trauma or diffuse axonal injury.

2.7 | T1-weighted MRI data processing

T1-weighted anatomical scans were analyzed utilizing the FreeSurfer v5.3.0 software package (<http://surfer.nmr.mgh.harvard.edu>). The analyses were performed using additional computing resources from the high performance computing TIER1 cluster at the University of Ghent (<http://www.ugent.be/hpc/>). The FreeSurfer analysis suite was used to extract cortical thickness using an automated approach described in detail in prior publications (see Fischl, 2012). Previous research has shown that this automated procedure yields accurate and reliable results (Fischl et al., 2002). Briefly, image processing included (1) removal of nonbrain tissue using a hybrid watershed/surface deformation procedure (skull stripping) (Segonne et al., 2004), (2) automated Talairach transformations, (3) segmentation of the subcortical WM and deep GM volumetric structures (Fischl et al., 2002), (4) intensity normalization (Sled, Zijdenbos, & Evans, 1998), (5) tessellation of the boundary between GM and WM, automated topology correction (Fischl, Liu, & Dale, 2001; Segonne, Pacheco, & Fischl, 2007) and (6) surface deformation along intensity gradients for optimal placement of the borders between GM, WM and cerebrospinal fluid (Dale, Fischl, & Sereno, 1999; Fischl & Dale, 2000). Automated parcellation of the cerebral cortex into units with respect to gyral and sulcal structures, and calculation of cortical thickness from all vertices within the cortical parcellations was performed per hemisphere using the Desikan atlas (Desikan et al., 2006). Also, estimates of total intracranial volume, and mean total thickness of the left and right hemisphere were obtained for each subject.

The T1.mgz (i.e., the FreeSurfer T1 image) and aparc + aseg.mgz (i.e., image containing ROIs constructed by the FreeSurfer pipeline) files were converted to the Neuroimaging Informatics Technology Initiative (NIFTI) format (T1.nii and aparc + aseg.nii) to be used in the further DWI analyses.

Two independent researchers (IC, RDP) visually checked the data quality of the FreeSurfer processing output including the accuracy of skull stripping, registration, segmentation, and cortical surface reconstruction. Poor data quality, such as inclusion of dura in the pial surface

FIGURE 2 Lateral (left) and medial (right) view of the cortical parcellation of the Desikan atlas (Desikan et al., 2006) displayed on an inflated template (<https://surfer.nmr.mgh.harvard.edu>). Numbered regions indicate the cortical regions of interest: (1) lateral orbitofrontal cortex; (2) pars orbitalis; (3) postcentral cortex; (4) superior parietal cortex; (5) supramarginal cortex; (6) precuneus; (7) posterior cingulate cortex; (8) caudal anterior cingulate cortex; (9) parahippocampal cortex [Color figure can be viewed at wileyonlinelibrary.com]

after skull stripping, and surface deformations, was revealed in 12 participants (controls = 3, CINP = 3, CWAD = 6). These cortical thickness datasets were excluded from further analyses.

2.8 | Cortical regions of interest

Cortical thickness was extracted from 9 ROIs from the Desikan atlas (Desikan et al., 2006). The selection of these ROIs was based on our systematic review appraising the evidence for brain alterations in INP and WAD (De Pauw et al., 2017), on previous studies in patients with chronic pain or mild TBI examining GM morphology alterations (Cagnie et al., 2014; Cauda et al., 2014; Coppieters et al., 2017a; Govindarajan et al., 2016; Kregel et al., 2015; Wang et al., 2015), and regarding studies exploring associations between cortical thickness, and measures of cognition and pain (Coppieters et al., 2016; Erpelding et al., 2012; Piche et al., 2013; Schwedt & Chong, 2014). In addition, only regions demonstrated to be mainly involved in processing pain and/or cognition were selected. Specifically, the following nine ROIs were selected from the Desikan atlas (Desikan et al., 2006): caudal anterior cingulate cortex, posterior cingulate cortex, lateral orbitofrontal cortex, superior parietal cortex, postcentral cortex, precuneus, pars orbitalis of the inferior frontal gyrus, parahippocampal cortex and supramarginal cortex (see Figure 2 for cortical ROIs). For each ROI, cortical thickness was calculated for the right and left hemisphere separately.

2.9 | DWI processing

The DWI data were analyzed and processed in ExploreDTI v4.8.6 with MATLAB (Leemans, Sijbers, & Jones, 2009) using the following procedure: DWI volumes were looped at a high frame rate to check for obvious artefacts in the data, such as large signal dropouts and geometric distortions. Next, we toggled between the views of the first and last acquired DW image to observe subtle system drifts. Furthermore, we inspected the images in different orthogonal views (coronal, sagittal, axial) to check for interslice and intravolume instabilities, and visualized various image maps to check for artefacts. Then, we checked the residual map for outliers, reflecting the difference between the modeled and the measured signal (Tournier et al., 2011). Next, the DW data were

corrected for distortions induced by the DW gradients, artefacts due to head motion, eddy current-induced geometric distortions, and EPI distortions (Leemans & Jones, 2009). EPI distortions were corrected by co-registering the DW images to the T1-weighted anatomical images, which were normalized for intensity using FreeSurfer. Moreover, we performed appropriate reorientation of the encoding vectors. Next, a tensor estimation procedure was performed on the preprocessed DW data.

Each scan was visually checked for accuracy after both the motion correction and co-registration steps by two independent researchers. To check the accuracy of the co-registration, the preprocessed DWI was overlaid on the normalized T1-weighted anatomical image. Poor data quality was observed in one healthy subject because of too much head translation due to severe head motion (exceeding the size of 1 voxel), and two patients with CINP, one due to a too high percentage of outliers in the preprocessed DW images and 1 because of a general low data quality profile (ghosting, spikes). These WM structural data were excluded from further analyses.

Translational motions (average of axial, coronal and sagittal) did not exceed the size of one voxel, that is, $2.5 \times 2.5 \times 2.5 \text{ mm}^3$ (mean $\pm$ SE: controls: $0.85 \pm 0.03 \text{ mm}$; CINP patients: $0.79 \pm 0.03 \text{ mm}$; $0.87 \pm 0.03 \text{ mm}$ for CWAD patients). We reinspected the data in three orthogonal planes to ensure that the motion/distortion correction was performed correctly and that no additional artefacts were introduced into the data. Finally, DWI derived metrics FA, MD, AD, and RD were extracted from the preprocessed DWI data using an automated approach based on the ICBM DTI-81 WM atlas, which will be further explained in the following paragraph (Mori et al., 2008).

2.10 | White matter regions of interest

The ICBM-DTI-81 WM labels atlas, developed by Mori et al. (2008), was used for automated parcellation of the ROIs. This is a stereotaxic probabilistic WM atlas that fuses DTI-based WM information with the standard MNI anatomical template (ICBM-152). The WM parcellations were applied to the preprocessed images and the DWI derived metrics were calculated in each ROI, by warping the atlas template to each individual data set.

FIGURE 3 White matter regions of interest masks. Regions are depicted on mean diffusivity maps of the white matter atlas of Mori et al. (2008). (a) Axial view of bilateral anterior limb of internal capsule (red), bilateral posterior limb of internal capsule (blue), bilateral anterior corona radiate (green), bilateral posterior corona radiate (violet); (projection fibers). (b) Axial view of bilateral superior cerebellar peduncles (blue); (tract in the brain stem). (c) Axial view of the cingulum cingulate gyrus (yellow), and the cingulum hippocampus (cyan); (association fibers). (d) Sagittal view of the fornix (green) (association fibers) and superior cerebellar peduncle (blue). (e) Sagittal view of the fornix (green) (association fibers) and superior cerebellar peduncle (blue). (f) Splenium of the corpus callosum (green) (commissural fibers) [Color figure can be viewed at wileyonlinelibrary.com]

The selection of WM ROIs was based on previous MRI research demonstrating WM structural alterations in chronic pain or mild TBI patients, or revealing associations between WM structure, and measures of cognition and pain. Based on our anatomical hypotheses, only WM atlas labels that are mainly involved in pain or cognition or the combination thereof were selected. Hereby, the following 10 WM regions/tracts were defined: projection fibers namely (1) the superior cerebellar peduncle (Chong, Plasencia, Frakes, & Schwedt, 2017; Drijckoning et al., 2015) (2) the anterior corona radiata, (3) the posterior corona radiata (Baek et al., 2013; DeSouza et al., 2014; Koenig et al., 2015; Kraus et al., 2007; Lin et al., 2015; Ly et al., 2016), (4) the anterior limb of internal capsule, (5) the posterior limb of internal capsule (Lieberman et al., 2014). Association fibers, namely (6) the cingulum cingulate gyrus (i.e., subgenual and retrosplenial part of the cingulum) and (7) the cingulum hippocampus (i.e., parahippocampal part of the cingulum) (Lieberman et al., 2014; Mori et al., 2008) and (8) fornix and stria terminalis (Borsook et al., 2016; Chen et al., 2011; Upadhyay et al., 2010). Finally, commissural fibers, that is (9) the tapetum of the corpus callosum (Kim et al., 2014; McAllister, 1992) and (10) the splenium of the corpus callosum (Lieberman et al., 2014). Acceleration-deceleration of the brain is believed to affect the superior cerebellar peduncles, and periventricular WM (McAllister, 1992), for example, tapetum of the corpus callosum (Kim et al., 2014; Sarikcioglu et al.,

2007; Todd & Lempert, 2007). For visualization purposes, masks of these ROIs are displayed on MD maps of the WM Mori atlas (Mori et al., 2008) in Figure 3. To examine WM structural organization, average FA, MD, AD, and RD values were computed in all WM ROI for each subject for the right and left hemisphere separately.

2.11 | Statistical analyses

All statistical analyses were performed with IBM SPSS Statistics 22.0. First, normality of variables was assessed with the Shapiro-Wilk test and by visual evaluation of histograms and quantile-quantile plots. Additionally, the equality of variance was examined with the Levene's test. If the assumptions of normality and equal between-group variances were met, data were analyzed with parametric tests. Otherwise, nonparametric tests were applied.

The comparability of study groups for demographics was explored with a one-way ANOVA with *post hoc* pairwise comparisons using Bonferroni correction [family wise error rate (FWER) < .05], or with the Kruskal-Wallis test with *post hoc* pairwise comparisons using the Mann-Whitney *U* test. Differences measured with the Mann-Whitney *U* test were only assumed significant below the significance threshold of 0.017 (Bonferroni correction: 0.05/3) to correct for the number of groups. Categorical data were analyzed with the Fisher's exact test.

Group differences for cognitive performance and CS were explored with one-way ANOVA (*post hoc* pairwise comparisons using Bonferroni correction, FWER < .05) or the Kruskal–Wallis test (*post hoc* pairwise comparisons using the Mann–Whitney *U* test, $p < .017$).

An ANCOVA model was fitted, controlling for age, to examine differences in total intracranial volume and total mean thickness of the left and right hemisphere (*post hoc* pairwise comparisons using Bonferroni correction, FWER < .05).

A MANCOVA model controlling for age was performed to determine group differences in cortical thickness of the 9 ROIs if the assumptions for performing MANCOVA were verified (multivariate normality in the data, homogeneity of variance between groups, absence of multicollinearity). Four MANCOVA models including age as covariate were performed investigating WM structure with respectively FA, MD, AD, and RD of 10 WM ROIs in separate models each, as dependent variables, in patients with CWAD, CINP, and controls, as independent variables. The significance threshold was Bonferroni corrected for the number of DWI derived metrics, resulting in an adjusted p value of $< .0125$ ($.05/4$) for the multivariate test and for the individual WM tracts. Next, *post hoc* pairwise comparison using Bonferroni correction was applied for the group comparisons. Because there were significant effects of age on various diffusion metrics in WM regions, age was included as covariate in the MANCOVA model. For each MANCOVA model, the partial eta square η^2 was calculated. This measure shows how much variance is explained by the independent variable (study group) and is used as the effect size for the MANCOVA model.

Finally, group-specific partial correlations (controlling for age) were conducted between cognitive performance and CS measures on one hand, and cortical thickness or WM structure on the other hand in cortical regions or WM tracts displaying significant group differences. To correct for multiple comparisons (for the number of clinical variables), partial correlations were deemed significant only below the $.0125$ ($0.05/4$) level.

3 | RESULTS

3.1 | Demographic characteristics and self-reported pain and disability measures

The demographic characteristics and self-reported pain and disability measures of 102 participants (30 healthy controls, 35 CINP, 37 CWAD) are presented in Table 1. No significant differences in demographic characteristics were found between all study groups ($p > .05$), except for age (healthy controls were younger compared with CINP; $p = .010$). Furthermore, both patient groups were comparable in use of medications, neck pain duration, and frequency of neck pain complaints ($p > .05$). In contrast, higher neck pain intensity at the MRI test day ($p < .001$), and higher pain-related disability were reported by patients with CWAD compared with CINP ($p = .001$).

3.2 | Control analyses

The MANCOVA with age and handedness as covariates, revealed no significant main effect of handedness on regional cortical thickness

($p = .284$) or WM structure ($p = .349$ for FA; $p = .215$ for MD; $p = .170$ for AD; $p = .094$ for RD). Therefore, the cortical thickness and WM structure results of the left- and right-handed women were analyzed together. The ANCOVA controlling for age showed no significant group differences for total intracranial volume ($p = .137$), and total mean thickness of the left ($p = .563$) and right ($p = .404$) hemisphere. The assumptions for all applied models were verified.

3.3 | Group differences in cognitive performance and central sensitization

As shown in Table 1, the completion time of TMT part A ($p < .001$) and part B ($p = .002$) was significantly longer in the CWAD group compared with healthy controls, denoting worse performance in CWAD patients. In addition, the time needed to perform TMT part A ($p < .001$) was significantly longer in patients with CWAD compared with CINP.

Decreased PPTs were demonstrated at the quadriceps muscle, reflecting distant hyperalgesia in patients with CWAD compared with healthy controls ($p = .002$). The CPM value, measured at the quadriceps muscle, was significantly lower in CWAD patients compared with controls ($p = .005$), indicating diminished endogenous pain inhibition in patients with CWAD (Table 1).

3.4 | Group differences in cortical thickness

The MANCOVA model examining cortical thickness in 9 ROIs between individuals with CWAD, CINP, and healthy controls, including age as covariate, demonstrated significant differences for cortical thickness ($p = .015$, $\eta^2 = 0.312$), based on the multivariate test (Supporting Information Table 1). Only the left precuneus showed significant group differences in cortical thickness ($p = .037$). *Post hoc* pairwise comparisons between study groups using Bonferroni correction revealed significant decreased cortical thickness in the left precuneus ($p = .032$) (Figure 4; Supporting Information Table 1) in patients with CWAD compared with CINP. Hence, no significant differences in regional cortical thickness could be found between either the CWAD or CINP group and healthy controls.

3.5 | Group differences in white matter structure

As can be seen in Supporting Information Table 2, four MANCOVA models with age as covariate investigating WM structure with respectively FA, MD, AD, and RD of 10 WM tracts in separate models in patients with CWAD, CINP, and controls, showed significant differences for FA ($p = .010$, $\eta^2 = 0.293$), MD ($p = .007$, $\eta^2 = 0.274$), and RD ($p = 0.007$, $\eta^2 = 0.297$) based on the multivariate tests. The MANCOVA of AD could not detect significant differences between study groups ($p = .499$), hence regional WM differences in AD were not further analyzed.

The following WM tracts did retain significance after correcting for multiple comparisons [$p < .0125$ ($.05/4$): left cingulum hippocampus ($p = .002$ for FA, $p = .004$ for MD, $p = .002$ for RD) and left tapetum ($p = .003$ for FA, $p = .005$ for MD, $p = .004$ for RD) (Supporting Information Table 2). Subsequent, *post hoc* pairwise comparisons between study groups, using Bonferroni correction, revealed significantly decreased FA

TABLE 1 Demographic characteristics, self-reported pain and disability measures, cognitive performance, distant hyperalgesia and conditioned pain modulation efficacy in patients with CWAD, CINP and healthy controls

		Median	IQR	Range (min–max)	Test statistic (p value)	p value post hoc
Demographic characteristics	Age (years) ^b	25	22–44	18–62	8.020 (.018)	.010 ^e .017 ^f .769 ^g
Body mass Index (kg/m ²) ^{b,a}	HCON	20.94	20.13–22.90	17.99–26.75	3.462 (.177)	.069 ^e .438 ^f .242 ^g
	CINP	23.05	20.12–25.62	18.34–29.07		
	CWAD	22.31	19.60–23.88	16.65–31.46		
Frequencies						
Education level, n (%) ^d	HCON	0 (0); 1 (3.3); 7 (23.3); 21 (70)			2.172 (1.000)	NA
	CINP	0 (0); 1 (2.9); 8 (22.9); 23 (65.7)				
Smoker, n (%) ^d Smoker; former smoker; nonsmoker	CWAD	1 (2.7); 1 (2.7); 9 (24.3); 25 (67.6)				
	HCON	1 (3.3); 3 (10); 25 (83.3)			7.908 (.066)	NA
Handedness, n (%) ^d (LH; RH)	CINP	0 (0); 9 (25.7); 18 (51.4)				
	CWAD	4 (10.8); 10 (27); 23 (62.2)				
	HCON	5 (17.2); 24 (82.8)			2.241 (.344)	NA
Demographic characteristics: Medication use	CINP	2 (5.7); 33 (94.3)				
	CWAD	3 (8.8); 31 (91.2)				
	HCON	0 (0)			10.962 (.003)	.069 ^g
Analgesics–antipyretics, n (%) ^d	CINP	3 (8.6)				
	CWAD	10 (27)				
	HCON	0 (0)			2.261 (.330)	NA
Narcotic analgesics, n (%) ^d	CINP	0 (0)				

(Continues)

TABLE 1 (Continued)

		Median	IQR	Range (min-max)	Test statistic (p value)	p value post hoc
Benzodiazepines, n (%) ^d	CWAD	2 (5.4)				
	HCON	1 (3.3)			5.768 (.053)	NA
	CINP	1 (2.9)				
	CWAD	7 (18.9)				
Antidepressants, n (%) ^d	HCON	0 (0)			2.479 (.316)	NA
	CINP	3 (8.6)				
	CWAD	2 (5.4)				
		Median	IQR	Range (min-max)	Test statistic (p value)	p value post hoc
Self-reported pain measures	Neck pain duration (months) ^b	NA	NA	NA	531.000 (.882)	NA
	CINP	60	22-120	4-288		
	CWAD	60	30-120	3-444		
	HCON	NA	NA	NA	228.000 (.035)	NA
Days/week neck pain? ^b	CINP	5	3.25-7	3-7		
	CWAD	7	5-7	2-7		
	HCON	0	0-0	0-0	35.582 (<.001)	<.001 ^e <.001 ^f .026 ^g
Current neck pain intensity (VNRS/10) _{C^{b,a}}	CINP	4	2-7	0-8		
	CWAD	7	5-7	0-10		
	HCON	0	0-0	0-0	71.625 (<.001)	<.001 ^e <.001 ^f <.001 ^g
Self-reported disability	Neck Disability Index (/50) ^{b,a}	2	1-4	0-6	65.109 (<.001)	<.001 ^e <.001 ^f .001 ^g
	CINP	3	1-5	0-6		
	CWAD	6	4-8	1-10		
	HCON	16	12-20	10-27		
	CWAD	22	17-27	10-37		

(Continues)

TABLE 1 (Continued)

		Median	IQR	Range (min–max)	Test statistic (p value)	p value post hoc	
Objective cognitive performance	TMT part A (s) ^{b,a}	HCON	18.615	18.615	10.030–29.750	19.247 (<.001)	.339 ^e <.001 ^f <.001 ^g
		CINP	19.370	19.370	11.560–28.400		
		CWAD	27.090	27.090	15.060–81.000		
	TMT part B (s) ^{b,a}	HCON	34.370	27.255–45.895	21.440–128.000	11.727 (.003)	.121 ^e .002 ^f .017 ^g
		CINP	38.360	31.455–44.365	26.600–148.000		
	CWAD	46.000	36.710–77.500	25.860–251.000			
Distant HA		Mean	SD	Range (min–max)	Test statistic (p value)	p value post hoc	
	PPT quadriceps (kgf) ^c	HCON	5.087	1.633	2.940–8.790	6.587 (.002)	.382 ^e .002 ^f .092 ^g
		CINP	4.383	1.945	1.700–9.720		
		CWAD	3.432	1.840	0.300–7.720		
	CPM quadriceps (PPT quadriceps during CPT minus PPT quadriceps before CPT) ^c	HCON	1.146	0.703	–0.140–3.000	5.698 (.005)	1.000 ^e .005 ^f .061 ^g
CPM efficacy		CINP	0.937	0.929	–0.590–3.290		
		CWAD	0.427	0.853	–1.050–1.870		

Abbreviations (y = years; HCON = healthy pain-free controls; CWAD = chronic whiplash-associated disorders; CINP = chronic idiopathic neck pain; VNRS = verbal numeric rating scale; No degr = no degree; Lower second = lower secondary; Higher edu = higher education; HA = hyperalgesia; CPM = conditioned pain modulation; CPT = cold pressor test; TMT = trail making test; kgf = kilogram force; PPT = pressure pain thresholds; VNRS = verbal numeric rating scale; IQR = interquartile range.

Data of 102 subjects were analyzed (30 HCON, 35 CINP, 37 CWAD) for the demographic variables, the self-reported pain and disability measures. Data of 97 subjects were analyzed (28 HCON, 35 CINP, 34 CWAD) for the current neck pain intensity at the clinical test day, TMT, PPT, and CPM efficacy.

^aVariances were not equally distributed across groups, Levene's test $p < .05$.

^bData which were not normally distributed and subsequently group differences were analyzed using the Kruskal-Wallis test and for post hoc pairwise comparisons the Mann-Whitney U test. To correct for multiple comparisons, differences measured with the Mann-Whitney U test were only deemed significant below the .017 level (Bonferroni's correction: 0.05/3).

^cData which were assumed to be normally distributed and where variances were equally distributed across groups were analyzed with one-way ANOVA (F test) and post hoc pairwise comparisons were applied using Bonferroni's correction ($p < .05$).

^dCategorical data was analyzed by performing the Fisher's exact test. Significant differences were presented in bold.

^e p value for significant differences between CON-CINP.

^f p value for significant differences between CON-CWAD.

^g p value for significant differences between CINP-CWAD.

FIGURE 4 Significant differences in cortical thickness of grey matter ROIs between patients with CWAD ($n = 31$), CINP ($n = 32$), and healthy controls ($n = 26$). Abbreviations: HCON = healthy pain-free controls, CINP = chronic idiopathic neck pain, CWAD = chronic whiplash-associated disorders, SE = standard error, ROIs = regions of interest. Data were analyzed using MANCOVA with age as covariate and *post hoc* pairwise comparisons were applied using Bonferroni correction; * $p = .032$ [Color figure can be viewed at wileyonlinelibrary.com]

($p = .007$ and $p = .013$), increased MD ($p = .010$ and $p = .025$), and increased RD ($p = .009$ and $p = .020$) in the left tapetum in CWAD patients compared with controls and CINP patients, respectively (Figure 5, Supporting Information Table 2). Also, significantly decreased FA ($p = .002$), increased MD ($p = .004$), and increased RD ($p = .001$) in the left cingulum hippocampus were found in CWAD patients compared with CINP patients (Figure 5, Supporting Information Table 2).

3.6 | Results of T2*-weighted brain imaging analyses

Based on detailed visual inspection of the axial T2*-weighted brain images, no hemorrhagic shearing lesions or microhemorrhages related to trauma or diffuse axonal injury were detected. In one CWAD patient a small a-specific T2* hypointensity left parietal without clinical relevance was observed. Furthermore, in one CINP patient the neuroradiologists detected a small round T2* hypointensity in the left thalamus and left globus pallidus suggesting microhemorrhages related to hypertension.

3.7 | Associations between cortical thickness alterations, and cognitive performance and central sensitization

Associations (partial correlations corrected for age) between cognitive performance and CS, and cortical thickness were investigated only in ROIs showing significant group differences (Table 2). The time to complete TMT part B was negatively correlated with cortical thickness of the

left precuneus ($r = -.520$, $p = .005$) within the CINP group ($n = 27$). Specifically, worse performance on the TMT part B coincided with decreased cortical thickness in the left precuneus in patients with CINP, however worse TMT performance could not be revealed in CINP patients compared with healthy controls. In the CWAD ($n = 26$), and control group ($n = 26$), no significant correlations could be found ($p > .017$).

3.8 | Associations between white matter structural alterations, and cognitive performance and central sensitization

As presented in Table 3, associations (partial correlations corrected for age) between WM structure (only FA, MD, RD), and cognitive performance and CS measures were investigated only in WM tracts demonstrating significant group differences. In the CWAD group ($n = 34$), efficacy of CPM was negatively correlated with MD ($r = -.478$, $p = .010$) and RD ($r = -.477$, $p = .010$) in the left tapetum. In other words, decreased efficacy of CPM was moderately correlated with increased MD and RD in the left tapetum in CWAD patients. In the CINP ($n = 33$) and control group ($n = 28$), no significant correlations could be revealed [p value $> .0125$ (0.05/4)].

4 | DISCUSSION

The present study has demonstrated cortical thinning in the left precuneus, a region crucially involved in cognitive functioning, in patients

FIGURE 5 Significant differences in DTI-derived metrics of white matter ROIs between CWAD patients ($n = 37$), CINP patients ($n = 35$), and healthy women ($n = 30$). Abbreviations: HCON = healthy pain-free controls, CINP = chronic idiopathic neck pain, CWAD = chronic whiplash-associated disorders, SE = standard error, DTI = diffusion tensor imaging, ROIs = regions of interest. Data were analyzed using MANCOVA with age as covariate [$p < .0125$ (.05/4)] and *post hoc* pairwise comparisons were applied using Bonferroni correction; * $p < .05$, ** $p < .01$ [Color figure can be viewed at wileyonlinelibrary.com]

with CWAD compared with CINP. Furthermore, abnormalities in WM structure encompassing decreased FA coinciding with increased MD and RD in the left cingulum hippocampus were revealed in patients with CWAD compared with CINP, and in the left tapetum in patients with CWAD compared with CINP and healthy controls. Interestingly,

brain microhemorrhages related to trauma were not observed. This is the first study revealing alterations in GM macrostructure and WM structure in women with CWAD compared with CINP or healthy women, indicating a potential association between these brain alterations and the signs and symptoms induced by the traumatic event in

TABLE 2 Partial correlations (controlling for age) between grey matter ROIs with significant group differences, and experimental measures of pain and cognition in patients with CINP and CWAD

GM ROI cortical thickness (mm)	Visuoperceptual and processing speed abilities	Working memory and task-switching ability	Distant hyperalgesia	CPM efficacy
	TMT A	TMT B	PPT quadriceps	CPM quadriceps
CINP (n = 27)				
Precuneus left	−0.164	−0.520	−0.241	0.030
	0.414	0.005	0.208	0.886
CWAD (n = 26)				
Precuneus left	−0.022	−0.107	0.154	0.411
	0.914	0.596	0.433	0.051

Abbreviations (CINP = chronic idiopathic neck pain; CWAD = chronic whiplash associated disorders; TMT = trail making test; PPT = pressure pain thresholds; CPM = conditioned pain modulation; ROIs = regions of interest).

To correct for multiple comparisons, correlations significant at a statistical threshold level of $p < .0125$ level (0.05/4) (two-tailed) were deemed significant, and are presented in bold. p values are presented below the correlation coefficient. In healthy pain-free controls, no significant correlations could be revealed (data not presented).

patients with CWAD. Moreover, decreased CPM efficacy (denoting CS) was associated with the extent of WM deficits in the left tapetum in CWAD patients. The latter yields novel evidence for underlying structural neural correlates of disturbed endogenous pain inhibition.

4.1 | Group differences in cognitive performance and central sensitization

CWAD patients displayed worse processing speed abilities compared with CINP patients and controls based on the results of the TMT part A. Decreased working memory span was revealed in the CWAD group compared with controls, utilizing the TMT part B. Consistent with these findings, two recent studies revealed worse cognitive performance in CWAD patients compared with healthy volunteers, specifically on domains of sustained attention and working memory evaluated with computer-based cognitive tasks (Coppieters et al., 2015; Meeus et al., 2015). Hence, our study highlights the importance of evaluating cognitive performance in patients with CWAD in clinical practice.

Furthermore, CWAD patients showed decreased PPTs at the quadriceps muscle compared with healthy controls reflecting distant hyperalgesia. This indicates significant indirect signs of CS and thus enhanced central pain sensitivity in CWAD. In accordance with our expectations, distant hyperalgesia was not demonstrated in CINP patients. Likewise, Scott and colleagues reported decreased PPTs at distant regions in CWAD but not in CINP patients (Scott et al., 2005). Furthermore, decreased CPM efficacy was only observed in the CWAD group, implying disturbed endogenous pain inhibitory mechanisms in CWAD but not in CINP patients. Consistent with these findings, reduced endogenous pain inhibition has previously been reported in CWAD (Daenen et al., 2013; Van Oosterwijck, Nijs, Meeus, Van Loo, & Paul, 2012) but could not be found in patients with CINP (Chua, Timmerman, Vissers, & Wilder-Smith, 2012). The lack of signs for CS in CINP patients is furthermore in line with the results of a recent systematic review concluding that CS is not a characteristic feature of CINP

(Malfliet et al., 2015). Our results add evidence to support that CS is present, at a group level, in CWAD and usually not in CINP patients, which further elucidates difference in underlying mechanisms.

4.2 | Group differences in cortical thickness

The cortical thinning in the left precuneus in CWAD supports our hypothesis of decreased regional cortical thickness in patients with traumatic compared with nontraumatic chronic neck pain. The observed cortical thinning in the precuneus in CWAD compared with CINP patients has a valuable meaning because it demonstrates macrostructural differences between both patient groups. Moreover, this region plays a crucial role in a wide range of cognitive and mental processes, and is involved in neurodegenerative processes (Cavanna & Trimble, 2006; Zhang & Li, 2012). The precuneus is part of the structural core of the brain (Hagmann et al., 2008), and is a core hub (i.e., highly interconnected nodes) of the default mode network (Fransson & Marrelec, 2008). The latter network is a constellation of brain regions involved in self-referential mental activity, emotional processing, and recollection of prior experiences, and is deactivated during externally focused tasks (Raichle, 2015). It is reported that damage to network hub regions connecting different subnetworks, such as the precuneus, causes the largest disturbances in network organization (Aerts, Fias, Caeyenberghs, & Marinazzo, 2016). Also, functional MRI studies have demonstrated altered resting-state functional connectivity in the precuneus in patients with chronic musculoskeletal pain (Ichesco et al., 2016), and in mild TBI patients (Xiong et al., 2016). Furthermore, Baliki, Mansour, Baria, and Apkarian (2014) have demonstrated reorganization of the default mode network dynamics in different chronic pain conditions including alterations in precuneus connectivity.

Nevertheless, opposite to our hypothesis, we could not detect significant correlations between cortical thinning in the precuneus and cognitive performance or experimental CS measures in CWAD patients. Furthermore, no cortical thickness differences were detected

TABLE 3 Partial correlations (controlling for age) between DTI-derived metrics in WM ROIs with significant group differences, and measures of cognition and central sensitization in patients with CINP and CWAD

WM ROI	DTI-derived metric	Visuo-perceptual and processing speed abilities	Working memory and task switching ability	Distant hyperalgesia	CPM efficacy
		TMT A	TMT B	PPT quadriceps	CPM quadriceps
CINP (n = 33)					
CgH left	FA	−0.113	−0.212	0.068	−0.038
		0.552	0.260	0.712	0.848
	MD	0.330	0.356	−0.093	−0.082
		0.075	0.053	0.612	0.679
	RD	0.304	0.364	−0.103	−0.042
		0.102	0.048	0.576	0.831
TAP left	FA	−0.376	−0.330	−0.093	0.068
		0.040	0.075	0.614	0.730
	MD	0.001	0.197	0.059	−0.076
		0.998	0.297	0.748	0.700
	RD	0.078	0.245	0.079	−0.081
		0.684	0.193	0.668	0.682
CWAD (n = 34)					
CgH left	FA	−0.198	−0.025	0.213	−0.021
		0.278	0.891	0.235	0.915
	MD	−0.006	−0.077	−0.142	0.418
		0.974	0.677	0.432	0.027
	RD	0.022	−0.065	−0.150	0.379
		0.903	0.725	0.405	0.047
TAP left	FA	−0.295	−0.240	0.197	0.397
		0.102	0.187	0.272	0.037
	MD	0.332	0.287	−0.152	−0.478
		0.063	0.112	0.397	0.010
	RD	0.342	0.290	−0.171	−0.477
		0.055	0.107	0.340	0.010

Abbreviations: CINP = chronic idiopathic neck pain; CWAD = chronic whiplash associated disorders; DTI = diffusion tensor imaging; TMT = trail making test; PPT = pressure pain thresholds; CPM = conditioned pain modulation; WM = white matter; ROI = region of interest; CgH = cingulum hippocampus; TAP = tapetum; FA = fractional anisotropy; MD = mean diffusivity; RD = radial diffusivity.

Correlations significant at a statistical threshold level of $p < .0125$ level (.05/4) (two-tailed) were deemed significant, and are presented in bold. p values are presented below the correlation coefficient. In healthy pain-free controls, no significant correlations could be revealed (data is not presented).

in all investigated brain regions between women with CINP and healthy women. Also, in contrast to our hypothesis and to the results of previous studies in patients with chronic musculoskeletal pain (Jensen et al., 2013; Kregel et al., 2015) and mild TBI (Govindarajan et al., 2016), differences in regional cortical thickness between CWAD patients and healthy controls could not be demonstrated. Yet, the cortical thickness MANCOVA model comprising all ROIs showed a partial eta squared of 0.31, which means that 31% of the variance in cortical thickness in the 9 ROIs can be explained by study group, which corresponds with a large effect size.

Possible explanations for the lack of significant cortical thickness differences between CINP or CWAD patients and controls may be that the MRI scan and analyzing procedures used in the present study were not sensitive enough to detect subtle differences or perhaps there are cortical thickness differences in other noninvestigated brain regions. On the other hand, it can be that these patient conditions are just not characterized by cortical thickness changes compared with healthy controls and thus display different underlying pathophysiological mechanisms compared with for example fibromyalgia patients who present regional cortical thickness changes. Nevertheless, further research is

warranted before drawing solid conclusions. Noteworthy, we recently demonstrated decreased regional GM volume in CWAD patients compared with healthy controls (Coppieters et al., 2017a). These contrasting findings underscore the idea that cortical thickness and GM volume reflect different aspects of the neural architecture (Rakic, 1988).

4.3 | Group differences in white matter structure

Results showed a consistent pattern of decreased FA coinciding with increased MD and RD in the left cingulum hippocampus, and in the left tapetum in patients with CWAD compared with healthy controls (tapetum) or compared with CINP patients (cingulum and tapetum). In contrast, AD values were not different between all study groups.

The observed WM structural differences between CWAD patients compared with healthy controls were in line with our hypothesis. In addition, differences in WM structure were revealed between CINP and CWAD patients in the left cingulum hippocampus and the left tapetum. The tapetum of the corpus callosum is periventricular WM, which is believed to be affected by traumatic acceleration–deceleration of the brain (Sarikioglu et al., 2007; Todd & Lempert, 2007). To date, limited studies are available on WM abnormalities in the tapetum as separate region. The tapetum is the temporal component of the corpus callosum (Mori et al., 2008) formed by decussating fibers in the splenium that arch over the atrium of the lateral ventricle and course inferiorly in the posterior and temporal horns of this ventricle (Sarikioglu et al., 2007). In contrast, more evidence exists for WM deficits in the corpus callosum (genu, rostrum, body, splenium, tapetum) in chronic musculoskeletal pain patients (Lieberman et al., 2014) and mild TBI patients (Xiong et al., 2014).

The revealed pattern of WM structural deficits encompassing decreased FA, increased MD and increased RD in CWAD is consistent with results in other chronic musculoskeletal pain conditions. For example, Lieberman et al. (2014) observed decreased FA and increased RD in the left cingulum hippocampus in chronic musculoskeletal pain patients compared with healthy volunteers. In patients with complex regional pain syndrome decreased FA in the left cingulum bundle has also been demonstrated (Geha et al., 2008). Furthermore, the observed WM abnormalities are in accordance with findings of increased MD in the cingulum in mild TBI patients compared with healthy persons (Veeramuthu et al., 2015). In literature, studies tend to show decreased FA associated with numerous neurological and neurodegenerative diseases (Curran & Leemans, 2016; Horsfield & Jones, 2002).

Additionally, WM structural abnormalities were found in the cingulum hippocampus in CWAD compared with CINP patients but not compared with healthy persons. The cingulum bundle is a prominent WM tract extending longitudinally above the corpus callosum (Jones, Christiansen, Chapman, & Aggleton, 2013), and carrying information from the cingulate gyrus to the hippocampus. Specifically, the cingulum contains many afferent and efferent fibers associated with the cingulate cortices (Morris, Pandya, & Petrides, 1999; Mufson & Pandya, 1984; Vogt, Pandya, & Rosene, 1987). These fibers include connections with the thalamus, dorsolateral prefrontal cortex, and insula (Mufson & Pandya, 1984). Other cingulum fibers are connected to the

parahippocampal cortices, the inferior component of the hippocampal formation, and amygdala (Morris et al., 1999; Mufson & Pandya, 1984).

Although this study has found innovative WM structural differences in the left tapetum in CWAD patients compared with healthy controls, no WM structural differences could be revealed in all other investigated WM ROIs between the CWAD or CINP group and healthy controls. Hence, the present study only found minimal and subtle differences in WM structure between women with CWAD and healthy women. The latter raises the question if investigating other WM regions/tracts or using other analyzing techniques would have revealed more differences and subsequently requires further research. It can be hypothesized that both chronic neck pain groups differ in terms of pain duration, report less maladaptive pain cognitions (e.g., pain catastrophizing), depressive and anxiety symptoms, cognitive problems, and medication use compared with other chronic pain conditions such as fibromyalgia in which more extensive WM structural alterations have been revealed compared with healthy controls (Coppieters et al., 2015, 2016; Lutz et al., 2008). Indeed, the role of longer pain duration, maladaptive pain cognitions, depressive and anxiety symptoms, cognitive deficits and medication use on structural brain alterations in chronic pain conditions has been reported in literature (Coppieters et al., 2016; Diaz-Piedra, Guzman, Buela-Casal, & Catena, 2016; Hsu et al., 2009; Luerding et al., 2008; Malfliet et al., 2017).

In CINP patients there is less evidence for maladaptive pain cognitions, cognitive problems and frequent medication use (e.g., narcotic analgesics) (Coppieters et al., 2017a,b). It could be that the extent of debilitating associated symptoms beside the chronic pain is associated with the more extensive structural brain alterations shown in other chronic pain conditions (Coppieters et al., 2016; Malfliet et al., 2017). In addition, no evidence for CS or objective cognitive deficits, and the nontraumatic origin of pain in CINP patients could play a role in the lack of GM and WM structural differences between women with CINP and controls.

We can very carefully suggest that our findings are indicative of WM structural abnormalities in the left tapetum and the left cingulum hippocampus in the CWAD group, maybe to some extent reflecting WM demyelination evidenced by the unchanged AD together with increased RD (Janve et al., 2013; Song et al., 2002; Sun, Liang, Schmidt, Cross, & Song, 2007). Interestingly, in mild TBI patients research has demonstrated that long associative WM tracts such as the cingulum bundle are frequently affected (Rutgers et al., 2008; Wu et al., 2010). As such, the underlying mechanisms of the revealed WM deficits in CWAD patients could be associated with the traumatic whiplash injury, which perhaps induced a cascade of events eventually leading to WM (micro)structural deficits. Nevertheless, based on visual inspection of the T2*-weighted brain images, microhemorrhages related to trauma or diffuse axonal injury were not observed in patients with CWAD. In patients with mild TBI, previous studies could detect subtle microhemorrhagic lesions suggesting differences in underlying pathophysiological mechanisms between patients with CWAD and mild TBI (Liu, Ghimire, Pang, Wu, & Shi, 2015). However, in future research we should include susceptibility weighted imaging or quantitative

susceptibility mapping to detect possible microhemorrhages related to trauma in CWAD patients.

4.4 | Decreased cortical thickness in the precuneus: Associations with cognitive performance and central sensitization

In CINP patients, decreased precuneus thickness coincided with worse performance on the TMT part B, however decreased cortical thickness and worse cognitive performance could not be revealed in CINP patients compared with controls. The observed association between working memory capacity and cortical thickness in CINP patients is in line with studies reporting associations between regional GM morphology and working memory capacity or other features of cognition in patients with fibromyalgia (Luerding et al., 2008), patients with complex regional pain syndrome (Lee et al., 2015), and in mild TBI patients (Urban et al., 2017).

4.5 | Structural white matter abnormalities in the cingulum hippocampus and tapetum: Associations with cognitive performance and central sensitization

Consistent with our hypothesis, deficits in WM structure in the left tapetum, encompassing increased MD and RD, were associated with decreased CPM efficacy (denoting CS) in the CWAD group. However, associations between cognitive performance or CS, and the observed WM alterations in the cingulum hippocampus could not be detected. Also, no associations were demonstrated between cognitive performance or distant hyperalgesia, and WM structural abnormalities in CWAD patients, which was in contrast with our expectations. Interestingly, we provide novel evidence for associations between structural WM deficits and decreased CPM efficacy in CWAD. Recently, a longitudinal MRI study uncovered structural WM vulnerabilities (i.e., FA differences) to develop chronic back pain (Mansour et al., 2013). Also, moderate evidence exists to support that higher pain intensity is associated with FA alterations in WM tracts involved in pain and cognition (Coppieters et al., 2016).

4.6 | Clinical implications

Our findings support a role of WM structural abnormalities in the left tapetum in the observed dysfunctional CPM response in CWAD. This yields innovative evidence for underlying WM structural correlates of disturbed endogenous pain inhibition in CWAD but not in CINP patients. Accordingly, these results emphasize the role of WM abnormalities in patients with CWAD compared with CINP and maybe reflect one piece of the puzzle underlying the observed clinical differences between both neck pain conditions with possibly a role of the traumatic injury mediating the structural brain differences.

Our results furthermore indicate that chronic pain in CWAD patients should be interpreted, at least in part, as a result of structural plasticity of the central nervous system, associated with experimental signs of CS, alterations in cortical thickness and WM structure in

regions involved in various aspects of pain and cognitive processing. Accordingly, it can be recommended that therapy approaches for CWAD should address the brain and take into account neuroplasticity of the central nervous system. As such, clinicians should take into account the observed CS, and WM structural differences associated with this CS when treating patients with CWAD.

4.7 | Strengths, limitations, and recommendations for further research

Several strengths can be outlined. This is the first study examining alterations in cortical thickness and WM structure, and their relationships with cognitive performance and CS in patients with a traumatic origin of pain (CWAD) compared with patients with nontraumatic CINP and healthy controls. This research is important because it unraveled differences between both pain conditions, and provided evidence for CS and structural brain alterations as underlying pathophysiological mechanisms in CWAD. Another strength is the evaluation of alterations in both GM macrostructure (cortical thickness) and WM structure (FA, MD, AD, RD).

The following limitations have to be taken into account when interpreting our results. Metrics derived from DWI data using the tensor model are indirect measures that relate to but do not directly quantify tissue features and are influenced by various methodological and biological factors (Curran & Leemans, 2016). These metrics cannot disentangle the individual microscopic contributions at the voxel level and therefore should be interpreted with caution. It remains unclear whether decreased FA is due to changes in membrane permeability, organelles, axon thickness, fiber density, or degree of myelination (Curran & Leemans, 2016). It has been suggested that alterations in RD combined with unchanged AD reflect demyelination (Song et al., 2002), but the interpretation of these metrics has been a topic of controversy (Wheeler-Kingshott & Cercignani, 2009).

A second limitation of DTI analyses is the inability of the tensor model to adequately characterize diffusion in regions of complex fiber architecture. Single-shell data reconstructed with the diffusion tensor model assumes a single straight fiber orientation within each voxel and is inadequate to model more than one fiber orientation per voxel. In the brain however voxels often contain fiber populations with more than one dominant orientation, such as crossing fibers (Curran & Leemans, 2016). Therefore, advanced models based on the high angular resolution DWI acquisition strategy to provide more robust estimates of the fiber orientation are recommended for further research in CINP and CWAD.

A third limitation is related to the ROI-based approach which implies a well-considered anatomical hypothesis-driven selection of investigated brain regions but on the other hand also implies that other nonselected brain regions/tracts could have revealed alterations in GM or WM structure in patients with CINP and CWAD compared with controls. Analyzing possible cortical alterations in for example the insula or the rostral anterior cingulate cortex could also have been interesting, hence the latter regions are recommended to include in future research regarding structural brain alterations in these patient

conditions to increase our insight in the involved brain regions in CINP and CWAD. Furthermore, whole brain and structural network analysis could be valuable for further research in these patient conditions.

Noteworthy, the selection of CWAD and CINP patients implies that we compared two patient groups that are *a priori* different from each other in the origin of pain. In order to interpret the cortical thickness and WM structural results correctly it is important to emphasize that CWAD patients relate to a traumatic event in contrast to CINP patients. Interestingly, traumatic chronic neck pain patients reported higher neck pain intensity and more severe pain-related disability compared with nontraumatic CINP patients and demonstrated cognitive problems and CS compared with controls. Subsequently, the cortical thickness and WM structural differences between the study groups could also in part be interpreted in light of these more severe symptoms revealed in the traumatic CWAD group.

A final limitation pertains to the cross-sectional nature of this study implying that no conclusions can be drawn on the causality of the observed associations. To test causal inference, future longitudinal studies are necessary.

Further research investigating associations between abnormalities in WM structure in CWAD, and other experimental features of CS such as temporal summation of second pain could add valuable insights into the brain structural correlates of CS. Finally, it could be recommended to disentangle possible functional brain alterations using (resting-state) functional MRI techniques and perform functional network analyses in patients with CWAD and CINP compared with healthy controls.

5 | CONCLUSION

In conclusion, cortical thinning in the left precuneus, a core hub of the default mode network and part of the structural brain core, was found in women with CWAD compared with CINP. Additionally, abnormalities in WM structure were revealed in 2 WM tracts carrying information between regions involved in affective–cognitive dimensions of pain processing and cognition in women with CWAD compared with CINP or healthy women. This study provided novel evidence for associations between dysfunctional CPM and the degree of WM deficits in the left tapetum in patients with CWAD. This yields innovative evidence for underlying WM structural correlates of CS and in particular of disturbed endogenous pain inhibition in CWAD. Accordingly, these results emphasize the role of structural brain alterations in patients with CWAD compared with CINP.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the study participants, without whom this study could have never been completed successfully. For its financial support, the authors' gratitude goes out to the Special Research Fund of Ghent University (BOF-Ghent), which made this experimental research possible. We acknowledge the University of Ghent for the possibility to use the additional computing resources from the high performance computing (HPC) TIER1 cluster (<http://www.ugent.be/hpc/>) for performing the FreeSurfer analyses. Finally, we acknowledge Stephanie Bogaert for her radiological assistance with the T2*-weighted MRI analyses.

www.ugent.be/hpc/) for performing the FreeSurfer analyses. Finally, we acknowledge Stephanie Bogaert for her radiological assistance with the T2*-weighted MRI analyses.

CONFLICT OF INTEREST

The authors declare that no conflicts of interest exist.

ORCID

I. Coppieters <http://orcid.org/0000-0003-1598-632X>

REFERENCES

- Aerts, H., Fias, W., Caeyenberghs, K., & Marinazzo, D. (2016). Brain networks under attack: Robustness properties and the impact of lesions. *Brain: A Journal of Neurology*, 139, 3063–3083.
- Alexander, M. P. (1998). In the pursuit of proof of brain damage after whiplash injury. *Neurology*, 51, 336–340.
- Apkarian, A. V., Hashmi, J. A., & Baliki, M. N. (2011). Pain and the brain: Specificity and plasticity of the brain in clinical chronic pain. *Pain*, 152, S49–S64.
- Baek, S. O., Kim, O. L., Kim, S. H., Kim, M. S., Son, S. M., Cho, Y. W., ... Jang, S. H. (2013). Relation between cingulum injury and cognition in chronic patients with traumatic brain injury; diffusion tensor tractography study. *Neurorehabilitation*, 33, 465–471.
- Baliki, M. N., Mansour, A. R., Baria, A. T., & Apkarian, A. V. (2014). Functional reorganization of the default mode network across chronic pain conditions. *PLoS One*, 9, e106133.
- Basser, P. J., & Pierpaoli, C. (1996). Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. *Journal of Magnetic Resonance: Part B*, 111, 209–219.
- Basser, P. J., Mattiello, J., & LeBihan, D. (1994). MR diffusion tensor spectroscopy and imaging. *Biophysical Journal*, 66, 259–267.
- Bay, E., Kalpakjian, C., & Giordani, B. (2012). Determinants of subjective memory complaints in community-dwelling adults with mild-to-moderate traumatic brain injury. *Brain Injury*, 26, 941–949.
- Beaulieu, C. (2002). The basis of anisotropic water diffusion in the nervous system: A technical review. *NMR & Biomedicine*, 15, 435–455.
- Borsook, D., Linnman, C., Faria, V., Strassman, A. M., Becerra, L., & Elman, I. (2016). Reward deficiency and anti-reward in pain chronification. *Neuroscience & Biobehavioral Reviews*, 68, 282–297.
- Bowie, C. R., & Harvey, P. D. (2006). Administration and interpretation of the Trail Making Test. *Nature Protocols*, 1, 2277–2281.
- Breivik, H., Collett, B., Ventafridda, V., Cohen, R., & Gallacher, D. (2006). Survey of chronic pain in Europe: Prevalence, impact on daily life, and treatment. *European Journal of Pain*, 10, 287–333.
- Buckalew, N., Haut, M. W., Aizenstein, H., Morrow, L., Perera, S., Kuwabara, H., & Weiner, D. K. (2010). Differences in brain structure and function in older adults with self-reported disabling and nondisabling chronic low back pain. *Pain Medicine*, 11, 1183–1197.
- Bulls, H. W., Freeman, E. L., Anderson, A. J., Robbins, M. T., Ness, T. J., & Goodin, B. R. (2015). Sex differences in experimental measures of pain sensitivity and endogenous pain inhibition. *Journal of Pain Research*, 8, 311–320.
- Cagnie, B., Coppieters, I., Denecker, S., Six, J., Danneels, L., & Meeus, M. (2014). Central sensitization in fibromyalgia? A systematic review on structural and functional brain MRI. *Seminars in Arthritis & Rheumatism*, 44, 68–75.

- Cathcart, S., & Pritchard, D. (2006). Reliability of pain threshold measurement in young adults. *Journal of Headache & Pain*, 7, 21–26.
- Cauda, F., Palermo, S., Costa, T., Torta, R., Duca, S., Vercelli, U., ... Torta, D. M. (2014). Gray matter alterations in chronic pain: A network-oriented meta-analytic approach. *Neuroimage Clinics*, 4, 676–686.
- Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: A review of its functional anatomy and behavioural correlates. *Brain: A Journal of Neurology*, 129, 564–583.
- Chen, J. Y., Blankstein, U., Diamant, N. E., & Davis, K. D. (2011). White matter abnormalities in irritable bowel syndrome and relation to individual factors. *Brain Research*, 1392, 121–131.
- Chong, C. D., Plasencia, J. D., Frakes, D. H., & Schwedt, T. J. (2017). Structural alterations of the brainstem in migraine. *Neuroimage: Clinical*, 13, 223–227.
- Chua, N. H. L., Timmerman, H., Vissers, K. C., & Wilder-Smith, O. H. (2012). Multi-modal quantitative sensory testing in patients with unilateral chronic neck pain: An exploratory study. *Journal of Musculoskeletal Pain*, 20, 292–299.
- Coppieters, I., De Pauw, R., Caeyenberghs, K., Danneels, L., Kregel, J., Pattyn, A., Meeus, M., & Cagnie, B. (2017a). Decreased regional grey matter volume in women with chronic whiplash-associated disorders: Relationships with cognitive deficits and disturbed pain processing. *Pain Physician*, 20, E1025–E1051.
- Coppieters, I., De Pauw, R., Kregel, J., Malfliet, A., Goubert, D., Lenoir, D., ... Meeus, M. (2017b). Differences between women with traumatic and idiopathic chronic neck pain and women without neck pain: Interrelationships among disability, cognitive deficits, and central sensitization. *Physical Therapy*, 97, 338–353.
- Coppieters, I., Ickmans, K., Cagnie, B., Nijs, J., Pauw, R., Noten, S., & Meeus, M. (2015). Cognitive performance is related to central sensitization and health-related quality of life in patients with chronic whiplash-associated disorders and fibromyalgia. *Pain Physician*, 18, E389–E401.
- Coppieters, I., Meeus, M., Kregel, J., Caeyenberghs, K., Pauw, R., Goubert, D., & Cagnie, B. (2016). Relations between brain alterations and clinical pain measures in chronic musculoskeletal pain: A systematic review. *Journal of Pain*, 17, 949–962.
- Curran, K. M., & Leemans, L. E. A. (2016). Quantitative DTI measures. In L. E. Wim van Hecke & S. Stefan (Eds.), *Diffusion tensor imaging: A practical handbook* (pp. 65–85). New York: Springer.
- Daenen, L., Nijs, J., Roussel, N., Wouters, K., Van Loo, M., & Cras, P. (2013). Dysfunctional pain inhibition in patients with chronic whiplash-associated disorders: An experimental study. *Clinical Rheumatology*, 32, 23–31.
- Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage*, 9, 179–194.
- Davis, C. G. (2013). Mechanisms of chronic pain from whiplash injury. *Journal of Forensic & Legal Medicine*, 20, 74–85.
- Davis, K. D., & Moayed, M. (2013). Central mechanisms of pain revealed through functional and structural MRI. *Journal of Neuroimmune Pharmacology*, 8, 518–534.
- De Pauw, R., Coppieters, I., Meeus, M., Caeyenberghs, K., & Danneels, L. (2017). B C. Is traumatic and non-traumatic neck pain associated with brain alterations? A systematic review. *Pain Physician*, 20, 245–260.
- Desikan, R. S., Segonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., ... Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*, 31, 968–980.
- DeSouza, D. D., Hodaie, M., & Davis, K. D. (2014). Abnormal trigeminal nerve microstructure and brain white matter in idiopathic trigeminal neuralgia. *Pain*, 155, 37–44.
- Diaz-Piedra, C., Guzman, M. A., Buela-Casal, G., & Catena, A. (2016). The impact of fibromyalgia symptoms on brain morphometry. *Brain Imaging & Behaviour*, 10, 1184–1197.
- Drijkoningen, D., Caeyenberghs, K., Leunissen, I., Vander Linden, C., Leemans, A., Sunaert, S., ... Swinnen, S. P. (2015). Training-induced improvements in postural control are accompanied by alterations in cerebellar white matter in brain injured patients. *Neuroimage Clinics*, 7, 240–251.
- Erpelding, N., Moayed, M., & Davis, K. D. (2012). Cortical thickness correlates of pain and temperature sensitivity. *Pain*, 153, 1602–1609.
- Fischl, B. (2012). FreeSurfer. *Neuroimage*, 62, 774–781.
- Fischl, B., & Dale, A. M. (2000). Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of National Academic Sciences USA*, 97, 11050–11055.
- Fischl, B., Liu, A., & Dale, A. M. (2001). Automated manifold surgery: Constructing geometrically accurate and topologically correct models of the human cerebral cortex. *IEEE Transactions on Medical Imaging*, 20, 70–80.
- Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., ... Dale, A. M. (2002). Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron*, 33, 341–355.
- Fransson, P., & Marrelec, G. (2008). The precuneus/posterior cingulate cortex plays a pivotal role in the default mode network: Evidence from a partial correlation network analysis. *Neuroimage*, 42, 1178–1184.
- Geha, P. Y., Baliki, M. N., Harden, R. N., Bauer, W. R., Parrish, T. B., & Apkarian, A. V. (2008). The brain in chronic CRPS pain: Abnormal gray-white matter interactions in emotional and autonomic regions. *Neuron*, 60, 570–581.
- Govindarajan, K. A., Narayana, P. A., Hasan, K. M., Wilde, E. A., Levin, H. S., Hunter, J. V., Miller, E. R., Patel, V. K., Robertson, C. S., & McCarthy, J. J. (2016). Cortical thickness in mild traumatic brain injury. *Journal of Neurotrauma*, 33, 1809–1817.
- Hagmann, P., Cammoun, L., Gigandet, X., Meuli, R., Honey, C. J., Wedeen, V. J., & Sporns, O. (2008). Mapping the structural core of human cerebral cortex. *PLoS Biology*, 6, e159.
- Horsfield, M. A., & Jones, D. K. (2002). Applications of diffusion-weighted and diffusion tensor MRI to white matter diseases: A review. *NMR in Biomedicine*, 15, 570–577.
- Hsu, M. C., Harris, R. E., Sundgren, P. C., Welsh, R. C., Fernandes, C. R., Clauw, D. J., & Williams, D. A. (2009). No consistent difference in gray matter volume between individuals with fibromyalgia and age-matched healthy subjects when controlling for affective disorder. *Pain*, 143, 262–267.
- Ichesco, E., Puui, T., Hampson, J. P., Kairys, A. E., Clauw, D. J., Harte, S. E., ... Schmidt-Wilcke, T. (2016). Altered fMRI resting-state connectivity in individuals with fibromyalgia on acute pain stimulation. *European Journal of Pain*, 20, 1079–1089.
- Janve, V. A., Zu, Z., Yao, S. Y., Li, K., Zhang, F. L., Wilson, K. J., ... Gochberg, D. F. (2013). The radial diffusivity and magnetization transfer pool size ratio are sensitive markers for demyelination in a rat model of type III multiple sclerosis (MS) lesions. *Neuroimage*, 74, 298–305.
- Jensen, K. B., Srinivasan, P., Spaeth, R., Tan, Y., Kosek, E., Petzke, F., ... Kong, J. (2013). Overlapping structural and functional brain changes in patients with long-term exposure to fibromyalgia pain. *Arthritis & Rheumatology*, 65, 3293–3303.
- Jones, D. K., & Leemans, A. (2011). Diffusion tensor imaging. *Methods in Molecular Biology (Clifton, New Jersey)*, 711, 127–144.
- Jones, D. K., Christiansen, K. F., Chapman, R. J., & Aggleton, J. P. (2013). Distinct subdivisions of the cingulum bundle revealed by diffusion

- MRI fibre tracking: Implications for neuropsychological investigations. *Neuropsychologia*, 51, 67–78.
- Jorritsma, W., de Vries, G. E., Dijkstra, P. U., Geertzen, J. H., & Reneman, M. F. (2012). Neck Pain and Disability Scale and Neck Disability Index: Validity of Dutch language versions. *European Spine Journal*, 21, 93–100.
- Jorritsma, W., de Vries, G. E., Geertzen, J. H., Dijkstra, P. U., & Reneman, M. F. (2010). Neck Pain and Disability Scale and the Neck Disability Index: Reproducibility of the Dutch Language Versions. *European Spine Journal*, 19, 1695–1701.
- Kennedy, D. L., Kemp, H. I., Ridout, D., Yarnitsky, D., & Rice, A. S. (2016). Reliability of conditioned pain modulation: A systematic review. *Pain*, 157, 2410–2419.
- Killgore, W. D., Singh, P., Kipman, M., Pisner, D., Fridman, A., & Weber, M. (2016). Gray matter volume and executive functioning correlate with time since injury following mild traumatic brain injury. *Neuroscience Letters*, 612, 238–244.
- Kim, D. J., Lim, M., Kim, J. S., Son, K. M., Kim, H. A., & Chung, C. K. (2014). Altered white matter integrity in the corpus callosum in fibromyalgia patients identified by tract-based spatial statistical analysis. *Arthritis & Rheumatology*, 66, 3190–3199.
- Koenig, K. A., Sakaie, K. E., Lowe, M. J., Lin, J., Stone, L., Bermel, R. A., ... Phillips, M. D. (2015). The relationship between cognitive function and high-resolution diffusion tensor MRI of the cingulum bundle in multiple sclerosis. *Multiple Sclerosis Journal*, 21, 1794–1801.
- Koerte, I. K., Hufschmidt, J., Muehlmann, M., Lin, A. P., & Shenton, M. E. (2016). Frontiers in neuroscience advanced neuroimaging of mild traumatic brain injury. In D. Laskowitz & G. Grant (Eds.), *Translational research in traumatic brain injury*. Boca Raton (FL): CRC Press/Taylor and Francis Group (c) 2016 by Taylor & Francis Group, LLC.
- Kosek, E., Ekholm, J., & Hansson, P. (1999). Pressure pain thresholds in different tissues in one body region. The influence of skin sensitivity in pressure algometry. *Scandinavian Journal of Rehabilitation Medicine*, 31, 89–93.
- Kraus, M. F., Susmaras, T., Caughlin, B. P., Walker, C. J., Sweeney, J. A., & Little, D. M. (2007). White matter integrity and cognition in chronic traumatic brain injury: A diffusion tensor imaging study. *Brain: A Journal of Neurology*, 130, 2508–2519.
- Kregel, J., Meeus, M., Malfliet, A., Dolphens, M., Danneels, L., Nijs, J., & Cagnie, B. (2015). Structural and functional brain abnormalities in chronic low back pain: A systematic review. *Seminars in Arthritis & Rheumatology* 45, 229–237.
- Le Bihan, D., Mangin, J. F., Poupon, C., Clark, C. A., Pappata, S., Molko, N., & Chabriat, H. (2001). Diffusion tensor imaging: Concepts and applications. *Journal of Magnetic Resonance Imaging*, 13, 534–546.
- Lee, D. H., Lee, K. J., Cho, K. I., Noh, E. C., Jang, J. H., Kim, Y. C., & Kang, D. H. (2015). Brain alterations and neurocognitive dysfunction in patients with complex regional pain syndrome. *Journal of Pain*, 16, 580–586.
- Leemans, A. J. B., Sijbers, J., & Jones, D. K. (2009). ExploreDTI: A graphical toolbox for processing, analyzing, and visualizing diffusion MR data. *Paper presented at the 17th Annual Meeting of the International Society for Magnetic Resonance in Medicine (ISMRM)* (pp. 35–37). Hawaii, HI, April.
- Leemans, A., & Jones, D. K. (2009). The B-matrix must be rotated when correcting for subject motion in DTI data. *Magnetic Resonance in Medicine*, 61, 1336–1349.
- Lewis, G. N., Heales, L., Rice, D. A., Rome, K., & McNair, P. J. (2012). Reliability of the conditioned pain modulation paradigm to assess endogenous inhibitory pain pathways. *Pain Research & Management*, 17, 98–102.
- Lieberman, G., Shpaner, M., Watts, R., Andrews, T., Filippi, C. G., Davis, M., & Naylor, M. R. (2014). White matter involvement in chronic musculoskeletal pain. *Journal of Pain: Official Journal of the American Pain Society*, 15, 1110–1119.
- Lin, L., Xue, Y., Duan, Q., Sun, B., Lin, H., Chen, X., ... Zhang, Z. (2015). Microstructural white matter abnormalities and cognitive dysfunction in subcortical ischemic vascular disease: An atlas-based diffusion tensor analysis study. *Journal of Molecular Neuroscience*, 56, 363–370.
- Liu, G., Ghimire, P., Pang, H., Wu, G., & Shi, H. (2015). Improved sensitivity of 3.0 Tesla susceptibility-weighted imaging in detecting traumatic bleeds and its use in predicting outcomes in patients with mild traumatic brain injury. *Acta Radiologica*, 56, 1256–1263.
- Luerding, R., Weigand, T., Bogdahn, U., & Schmidt-Wilcke, T. (2008). Working memory performance is correlated with local brain morphology in the medial frontal and anterior cingulate cortex in fibromyalgia patients: Structural correlates of pain-cognition interaction. *Brain: A Journal of Neurology*, 131, 3222–3231.
- Lutz, J., Jager, L., de Quervain, D., Krauseneck, T., Padberg, F., Wichnalek, M., ... Schelling, G. (2008). White and gray matter abnormalities in the brain of patients with fibromyalgia: A diffusion-tensor and volumetric imaging study. *Arthritis & Rheumatology*, 58, 3960–3969.
- Ly, M., Adluru, N., Destiche, D. J., Lu, S. Y., Oh, J. M., Hoscheidt, S. M., ... Bendlin, B. B. (2016). Fornix microstructure and memory performance is associated with altered neural connectivity during episodic recognition. *Journal of the International Neuropsychological Society*, 22, 191–204.
- Malfliet, A., Coppieters, I., Van Wilgen, P., Kregel, J., Pauw, R., Dolphens, M., & Ickmans, K. (2017). Brain changes associated with cognitive and emotional factors in chronic pain: A systematic review. *European Journal of Pain*, 21, 769–786.
- Malfliet, A., Kregel, J., Cagnie, B., Kuipers, M., Dolphens, M., Roussel, N., ... Nijs, J. (2015). Lack of evidence for central sensitization in idiopathic, non-traumatic neck pain: A systematic review. *Pain Physician*, 18, 223–236.
- Mansour, A. R., Baliki, M. N., Huang, L., Torbey, S., Herrmann, K. M., Schnitzer, T. J., & Apkarian, A. V. (2013). Brain white matter structural properties predict transition to chronic pain. *Pain*, 154, 2160–2168.
- McAllister, T. W. (1992). Neuropsychiatric sequelae of head injuries. *Psychiatric Clinics of North America*, 15, 395–413.
- Meeus, M., Van Oosterwijck, J., Ickmans, K., Baert, I., Coppieters, I., Roussel, N., ... Nijs, J. (2015). Interrelationships between pain processing, cortisol and cognitive performance in chronic whiplash-associated disorders. *Clinical Rheumatology*, 34, 545–553.
- Misdraji, E. L., & Gass, C. S. (2010). The Trail Making Test and its neuro-behavioral components. *Journal of Clinical & Experimental Neuropsychology*, 32, 159–163.
- Mori, S., Oishi, K., Jiang, H., Jiang, L., Li, X., Akhter, K., ... Mazziotta, J. (2008). Stereotaxic white matter atlas based on diffusion tensor imaging in an ICBM template. *Neuroimage*, 40, 570–582.
- Moriarty, O., McGuire, B. E., & Finn, D. P. (2011). The effect of pain on cognitive function: A review of clinical and preclinical research. *Progress in Neurobiology*, 93, 385–404.
- Morris, R., Pandya, D. N., & Petrides, M. (1999). Fiber system linking the mid-dorsolateral frontal cortex with the retrosplenial/presubicular region in the rhesus monkey. *Journal of Comparative Neurology*, 407, 183–192.
- Mufson, E. J., & Pandya, D. N. (1984). Some observations on the course and composition of the cingulum bundle in the rhesus monkey. *Journal of Comparative Neurology*, 225, 31–43.

- Nampiarampil, D. E. (2008). Prevalence of chronic pain after traumatic brain injury: A systematic review. *JAMA*, 300, 711–719.
- Ng, T. S., Pedler, A., Vicenzino, B., & Sterling, M. (2014). Less efficacious conditioned pain modulation and sensory hypersensitivity in chronic whiplash-associated disorders in Singapore. *Clinical Journal of Pain*, 30, 436–442.
- Perelle, I. B., & Ehrman, L. (1994). An international study of human handedness: The data. *Behavioral Genetics*, 24, 217–227.
- Piche, M., Chen, J. I., Roy, M., Poitras, P., Bouin, M., & Rainville, P. (2013). Thicker posterior insula is associated with disease duration in women with irritable bowel syndrome (IBS) whereas thicker orbito-frontal cortex predicts reduced pain inhibition in both IBS patients and controls. *Journal of Pain*, 14, 1217–1226.
- Raichle, M. E. (2015). The brain's default mode network. *Annual Review of Neuroscience*, 38, 433–447.
- Rakic, P. (1988). Defects of neuronal migration and the pathogenesis of cortical malformations. *Progress in Brain Research*, 73, 15–37.
- Ruigrok, A. N., Salimi-Khorshidi, G., Lai, M. C., Baron-Cohen, S., Lombardo, M. V., Tait, R. J., & Suckling, J. (2014). A meta-analysis of sex differences in human brain structure. *Neuroscience Biobehaviour & Review*, 39, 34–50.
- Rutgers, D. R., Toulgoat, F., Cazejust, J., Fillard, P., Lasjaunias, P., & Ducreux, D. (2008). White matter abnormalities in mild traumatic brain injury: A diffusion tensor imaging study. *AJNR American Journal of Neuroradiology*, 29, 514–519.
- Sanchez-Cubillo, I., Perianez, J. A., Adrover-Roig, D., Rodriguez-Sanchez, J. M., Rios-Lago, M., Tirapu, J., & Barcelo, F. (2009). Construct validity of the Trail Making Test: Role of task-switching, working memory, inhibition/interference control, and visuomotor abilities. *Journal of the International Neuropsychological Society*, 15, 438–450.
- Sarikcioglu, L., Ozsoy, U., & Unver, G. (2007). Tapetum corporis callosi: Carpet of the brain. *Journal of the History of the Neurosciences*, 16, 432–434.
- Schwedt, T. J., & Chong, C. D. (2014). Correlations between brain cortical thickness and cutaneous pain thresholds are atypical in adults with migraine. *PLoS One*, 9, e99791.
- Scott, D., Jull, G., & Sterling, M. (2005). Widespread sensory hypersensitivity is a feature of chronic whiplash-associated disorder but not chronic idiopathic neck pain. *Clinical Journal of Pain*, 21, 175–181.
- Segonne, F., Dale, A. M., Busa, E., Glessner, M., Salat, D., Hahn, H. K., & Fischl, B. (2004). A hybrid approach to the skull stripping problem in MRI. *Neuroimage*, 22, 1060–1075.
- Segonne, F., Pacheco, J., & Fischl, B. (2007). Geometrically accurate topology-correction of cortical surfaces using nonseparating loops. *IEEE Transactions on Medical Imaging*, 26, 518–529.
- Sled, J. G., Zijdenbos, A. P., & Evans, A. C. (1998). A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE Transactions on Medical Imaging*, 17, 87–97.
- Song, S. K., Sun, S. W., Ramsbottom, M. J., Chang, C., Russell, J., & Cross, A. H. (2002). Dysmyelination revealed through MRI as increased radial (but unchanged axial) diffusion of water. *Neuroimage*, 17, 1429–1436.
- Sterling, M. (2004). A proposed new classification system for whiplash associated disorders—Implications for assessment and management. *Manual Therapy*, 9, 60–70.
- Suleyman Sener, P. P. M., & Maas Andrew, I. R. (2016). Diffusion tensor imaging in traumatic brain injury. In E. L. Van Hecke Wim & S. Sunaert (Eds.), *Diffusion tensor imaging: A practical handbook* (pp. 373–380). New York: Springer.
- Sun, S. W., Liang, H. F., Schmidt, R. E., Cross, A. H., & Song, S. K. (2007). Selective vulnerability of cerebral white matter in a murine model of multiple sclerosis detected using diffusion tensor imaging. *Neurobiological Diseases*, 28, 30–38.
- Todd, E., & Lempert, E. H. L. G. (2007). Pediatric neuroimaging. In A. Clyde & W. E. B. Helms (Eds.), *Fundamentals of diagnostic radiology* (pp. 217). Philadelphia, USA: Lippincott Williams & Wilkins and Wolters Kluwer Business.
- Tombaugh, T. N. (2004). Trail Making Test A and B: Normative data stratified by age and education. *Archives on Clinical Neuropsychology*, 19, 203–214.
- Tournier, J. D., Mori, S., & Leemans, A. (2011). Diffusion tensor imaging and beyond. *Magnetic Resonance in Medicine*, 65, 1532–1556.
- Tousignant-Laflamme, Y., Page, S., Goffaux, P., & Marchand, S. (2008). An experimental model to measure excitatory and inhibitory pain mechanisms in humans. *Brain Research*, 1230, 73–79.
- Upadhyay, J., Maleki, N., Potter, J., Elman, I., Rudrauf, D., Knudsen, J., ... Borsook, D. (2010). Alterations in brain structure and functional connectivity in prescription opioid-dependent patients. *Brain: A Journal of Neurology*, 133, 2098–2114.
- Urban, K. J., Riggs, L., Wells, G. D., Keightley, M., Chen, J. K., Ptito, A., Fait, P., Taha, T., & Sinopoli, K. J. (2017). Cortical thickness changes and their relationship to dual-task performance following mild traumatic brain injury in youth. *Journal of Neurotrauma*, 34, 816–823.
- van Hemmen, J., Saris, I. M., Cohen-Kettenis, P. T., Veltman, D. J., Pouwels, P. J., & Bakker, J. (2017). Sex differences in white matter microstructure in the human brain predominantly reflect differences in sex hormone exposure. *Cerebral Cortex* 27, 2994–3001.
- Van Oosterwijck, J., Nijs, J., Meeus, M., & Paul, L. (2013). Evidence for central sensitization in chronic whiplash: A systematic literature review. *European Journal of Pain (London, England)*, 17, 299–312.
- Van Oosterwijck, J., Nijs, J., Meeus, M., Van Loo, M., & Paul, L. (2012). Lack of endogenous pain inhibition during exercise in people with chronic whiplash associated disorders: An experimental study. *Journal of Pain*, 13, 242–254.
- Veeramuthu, V., Narayanan, V., Kuo, T. L., Delano-Wood, L., Chinna, K., Bondi, M. W., ... Ramli, N. (2015). Diffusion tensor imaging parameters in mild traumatic brain injury and its correlation with early neuropsychological impairment: A longitudinal study. *Journal of Neurotrauma*, 32, 1497–1509.
- Vernon, H. (2008). The Neck Disability Index: State-of-the-art, 1991–2008. *Journal of Manipulative & Physiological Therapeutics*, 31, 491–502.
- Vogt, B. A., Pandya, D. N., & Rosene, D. L. (1987). Cingulate cortex of the rhesus monkey: I. Cytoarchitecture and thalamic afferents. *Journal of Comparative Neurology*, 262, 256–270.
- Wang, X., Xie, H., Cotton, A. S., Tamburrino, M. B., Brickman, K. R., Lewis, T. J., ... Liberzon, I. (2015). Early cortical thickness change after mild traumatic brain injury following motor vehicle collision. *Journal of Neurotrauma*, 32, 455–463.
- Wheeler-Kingshott, C. A., & Cercignani, M. (2009). About “axial” and “radial” diffusivities. *Magnetic Resonance in Medicine*, 61, 1255–1260.
- Whiteside, A., Hansen, S., & Chaudhuri, A. (2004). Exercise lowers pain threshold in chronic fatigue syndrome. *Pain*, 109, 497–499.
- Wu, T. C., Wilde, E. A., Bigler, E. D., Yallampalli, R., McCauley, S. R., Troyanskaya, M., ... Levin, H. S. (2010). Evaluating the relationship between memory functioning and cingulum bundles in acute mild traumatic brain injury using diffusion tensor imaging. *Journal of Neurotrauma*, 27, 303–307.
- Xiong, K. L., Zhang, J. N., Zhang, Y. L., Zhang, Y., Chen, H., & Qiu, M. G. (2016). Brain functional connectivity and cognition in mild traumatic brain injury. *Neuroradiology*, 58, 733–739.

- Xiong, K., Zhu, Y., Zhang, Y., Yin, Z., Zhang, J., Qiu, M., & Zhang, W. (2014). White matter integrity and cognition in mild traumatic brain injury following motor vehicle accident. *Brain Research*, 1591, 86–92.
- Yarnitsky, D. (2010). Conditioned pain modulation (the diffuse noxious inhibitory control-like effect): Its relevance for acute and chronic pain states. *Current Opinion in Anaesthesiology*, 23, 611–615.
- Yunus, M. B. (2007). Fibromyalgia and overlapping disorders: The unifying concept of central sensitivity syndromes. *Seminars in Arthritis & Rheumatology*, 36, 339–356.
- Zhang, S., & Li, C. S. (2012). Functional connectivity mapping of the human precuneus by resting state fMRI. *Neuroimage*, 59, 3548–3562.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Coppieters I, De Pauw R, Caeyenberghs K, et al. Differences in white matter structure and cortical thickness between patients with traumatic and idiopathic chronic neck pain: Associations with cognition and pain modulation? *Hum Brain Mapp*. 2018;39:1721–1742. <https://doi.org/10.1002/hbm.23947>