Desynchronization of Autonomic Response and Central Autonomic Network Connectivity in Posttraumatic Stress Disorder

Janine Thome,¹ Maria Densmore,² Paul A. Frewen,^{2,3,4}
Margaret C. McKinnon,⁵ Jean Théberge,^{2,6} Andrew A. Nicholson,⁴
Julian Koenig,^{7,8} Julian F. Thayer,⁷ and Ruth A. Lanius^{2,4*}

¹Medical Faculty Mannheim, Institute of Psychiatric and Psychosomatic Psychotherapy, Central Institute of Mental Health, University of Heidelberg, Germany

²Department of Psychiatry, University of Western Ontario, London, Ontario, Canada

³Department of Psychology, University of Western Ontario, London, Ontario, Canada

⁴Department of Neuroscience, University of Western Ontario, London, Ontario, Canada

⁵Department of Psychiatry and Behavioral Neurosciences, Mc Master University, Hamilton, Ontario, Canada

⁶Department of Medical Imaging, Lawson Health Research Institute, London, Ontario, Canada

⁷Department of Psychology, Ohio State University, Columbus, Ohio

⁸Section for Translational Psychobiology in Child and Adolescent Psychiatry, Department of Child and Adolescent Psychiatry, Centre for Psychosocial Medicine, University of Heidelberg, Heidelberg, Germany

Abstract: *Objectives:* Although dysfunctional emotion regulatory capacities are increasingly recognized as contributing to posttraumatic stress disorder (PTSD), little work has sought to identify biological markers of this vulnerability. Heart rate variability (HRV) is a promising biomarker that, together with neuroimaging, may assist in gaining a deeper understanding of emotion dysregulation in PTSD. The objective of the present study was, therefore, to characterize autonomic response patterns, and their related neuronal patterns in individuals with PTSD at rest. *Methods:* PTSD patients ($N = 57$) and healthy controls ($N = 41$) underwent resting-state fMRI. Connectivity patterns of key regions within the central autonomic network (CAN)—including the ventromedial prefrontal cortex (vmPFC), amygdala, and periaqueductal gray (PAG)—were examined using a seed-based approach. Observed connectivity patterns were then correlated to resting HRV. *Results:* In contrast to controls, individuals with PTSD exhibited lower HRV. In addition, whereas controls engaged a localized connectivity pattern of CAN-related brain regions, in PTSD, key CAN regions were associated with widespread connectivity patterns in regions related to emotional reactivity (vmPFC and amygdala to insular cortex and lentiform

Additional Supporting Information may be found in the online version of this article.

Contract grant sponsors: Canadian Institute of Health Research, Physician-Scientist Fellowship provided by the Medical School of Heidelberg University (to J.K.), and Collaborative Research Center SFB 636 (DFG; to J.T.)

Julian F. Thayer and Ruth A. Lanius shared last co-author.

*Correspondence to: Ruth A. Lanius, M.D., Ph.D., Department of Psychiatry, University of Western Ontario, University Hospital

(Room C3-103), 339 Windermere Road, London, ON, Canada N6A 5A5. E-mail: ruth.lanius@lhsc.on.ca

Received for publication 4 February 2016; Revised 26 May 2016; Accepted 25 July 2016.

DOI: 10.1002/hbm.23340

Published online 20 September 2016 in Wiley Online Library (wileyonlinelibrary.com).

nucleus; PAG to insula) and motor readiness (vmPFC and amygdala to precentral gyrus; PAG to precentral gyrus and cerebellum). Critically, whereas CAN connectivity in controls was strongly related to higher HRV (insula, mPFC, superior frontal cortex, thalamus), HRV covariation was absent in PTSD subjects. **Conclusions:** This study provides the first evidence for a specific psychophysiological-neuronal profile in PTSD individuals characterized by lower resting HRV and a lack of HRV covariation with CAN-related brain connectivity. *Hum Brain Mapp* 38:27–40, 2017. © 2016 Wiley Periodicals, Inc.

Key words: posttraumatic stress disorder; fMRI; heart rate variability; central autonomic network; periaqueductal gray; top-down modulation

INTRODUCTION

As evidenced by the newly added symptoms of “negative alterations in cognition and mood” and the “inability to experience positive emotions” in the DSM-5 (2013), posttraumatic stress disorder (PTSD) is increasingly acknowledged as involving persistent negative emotional states. The integration of heightened negative states and the decreased ability to experience positive emotions into these diagnostic guidelines highlights an emerging focus on emotion regulatory capacities in PTSD [Friedman et al., 2011; Lanius et al., 2011, 2015; Nawijn et al., 2015; Powers et al., 2015a,b; Resick et al., 2008; Sadeh et al., 2015; Taylor, 2015; van Wingen et al., 2011]. Accordingly, numerous studies have begun to characterize emotional dysregulation associated with contrasting states of emotional under- and over-modulation. These opposing states have been linked to the experience of heightened or depressed emotionality, including re-experiencing symptoms and a detachment from emotional experience that occurs during emotional numbing and states of depersonalization and derealization, respectively [Etkin and Wager, 2007; Lanius et al., 2010; Nicholson et al., 2015; Reinders et al., 2014; Wolf et al., 2014].

Heart rate variability (HRV) is one promising psychophysiological indicator that may assist in gaining a deeper understanding of emotion regulatory capacity in PTSD. HRV reflects vagal activity that indexes the state of the autonomic nervous system (ANS), enabling the organism to calibrate bodily, emotional, and cognitive reactions to contextual demands. The ANS is separated into two branches, normally behaving in a balanced fashion: the parasympathetic nervous system (PNS), associated with restorative and vegetative functioning, and the sympathetic nervous system, linked to energy mobilization. Imbalance of these two branches is associated with blunted flexibility to react to environmental changes, increased vulnerability to somatic diseases, and has been associated with mental disorders [Carney and Freedland, 2009; Stein et al., 2000; Thayer and Brosschot, 2005]. Thus, HRV is thought to guide the individuals ability to organize emotional, cognitive, and behavioral responses, and higher levels of HRV to be a physiological index of the

ability to respond in a context appropriate manner [Gillie and Thayer, 2014; Melzig et al., 2009; Thayer and Lane, 2000].

To date, HRV investigations in PTSD have revealed that indices of high-frequency heart rate variability (HF-HRV) and root mean squared successive differences (RMSSD) of the inter-beat interval time series at rest, both indexing activity of the PNS, are reduced as compared to both trauma and non-trauma exposed healthy control subjects [for meta-analyses see Chalmers et al., 2014; Sammito et al., 2015; but see Agorastos et al., 2013; Keary et al., 2009 for null findings]. Fewer studies have addressed cardiac response to stressful tasks, revealing heterogeneous findings [Cohen et al., 2000, 1998; Hauschildt et al., 2011; Keary et al., 2009; Norte et al., 2013]. Whereas three studies failed to identify differential effects of affective cues or stressful tasks on HRV response in PTSD as compared to control subjects [Cohen et al., 2000, 1998; Hauschildt et al., 2011], two studies reported a greater decrease of HRV in response to stressful tasks or trauma script exposure [Keary et al., 2009; Norte et al., 2013].

A growing number of studies have investigated the relation between HRV and neural activation, where the central autonomic network (CAN) is thought to serve as a critical link between the brain and the ANS [for meta-analyses see Beissner et al., 2013; Thayer et al., 2012]. The CAN is comprised of the ventromedial prefrontal cortex (vmPFC), anterior cingulate (ACC), insular cortex, amygdala, hypothalamus, periaqueductal gray (PAG), parabrachial complex, nucleus of the tractus solitarius, and the ventrolateral medulla [Benarroch, 1993; Cersosimo and Benarroch, 2013; Palkovits, 1999; Thayer and Brosschot, 2005]. These components are reciprocally interconnected, with higher-order cortical functions associated with cognitive functioning regulating the response of subcortical structures, which in turn regulate autonomic input to the heart and allows for the complex variability that characterizes the healthy HR time series [Thayer and Brosschot, 2005]. Despite this knowledge, to date, no studies to date have examined connectivity in CAN regions in association with HRV in PTSD.

The aim of the present study was, therefore, to investigate the functional neural correlates of autonomic activity

in individuals with PTSD as compared to controls. As studies in PTSD have suggested not only lower HRV in PTSD but also alterations in top-down modulatory processes on a neuronal level [Etkin and Wager, 2007; Lanius et al., 2010; Nicholson et al., 2015; Reinders et al., 2014; Wolf et al., 2014], combining resting HRV and neural activation will facilitate a greater understanding of altered emotion regulatory processes in this disorder. Specifically, we investigated resting state neural connectivity patterns using fMRI and their HRV-related associations with three seed regions that comprise key nodes of the CAN: (1) vmPFC; (2) amygdala; and (3) the PAG [Beissner et al., 2013; Thayer et al., 2012; Thayer and Lane, 2000]. Consistent with prior findings indicating impaired emotion regulation capacity, we hypothesized that individuals with PTSD would exhibit diminished HRV [Chalmers et al., 2014; Sammito et al., 2015] and show reduced correlations between HRV scores and connectivity between key structures of the CAN as compared to healthy controls [Beissner et al., 2013; Thayer, 2006].

METHODS AND MATERIALS

Sample Description

Ninety-eight participants were included in the current study: 57 individuals with a primary diagnosis of PTSD and 41 healthy control subjects. Subjects were recruited between 2009 and 2015 from the Department of Psychiatry London Health Services Center (LHSC), through family physician, mental health professionals, psychology/psychiatric clinics, community programs for traumatic-stress survivors, and posters/advertisements, all within the London, ON community. PTSD diagnosis was confirmed via the Clinician-Administered PTSD scale, with a cut-off score of 50 for moderate PTSD [Blake et al., 1995]. Co-morbid Axis I disorders were diagnosed with the Structured Clinical Interview for DSM-IV Axis I Disorders [First, 1997]. Childhood trauma history was assessed with the Childhood Trauma Questionnaire (CTQ) [Bernstein et al., 2003], as well as depressive symptoms with the Beck Depression Inventory (BDI) [Beck et al., 1997]. Subjects also completed the Multiscale Dissociation Inventory (MDI) [Briere et al., 2005], capturing dissociative experiences. Immediately after the resting state scanning procedure, subjects were instructed to rate their state anxiety (3 items of the State-Trait Anxiety Inventory: STAI-S) [Kvaal et al., 2005] and their level of state dissociation, re-experiencing, avoidance, and hyperarousal symptoms by means of the Response to Script Driven Imagery (RSDI) Scale [Hopper et al., 2007a]. Clinical and socio-demographic variables are presented in Table I.

Exclusion criteria for all participants comprised implants or metals that do not comply with 3T fMRI safety standards for research, a history of head injury with any loss of consciousness, significant untreated medical illness, a history of neurological disorders, history of any pervasive

developmental disorders, pregnancy, and current use of any psychotropic or cardiovascular medications within one month prior to study (except 1 subject). Further exclusion criteria for individuals with PTSD were a history of bipolar disorder or schizophrenia, alcohol, or substance abuse/dependence not in sustained full remission within 6 month prior to participation of the study, while healthy control subjects had to be free of any lifetime Axis-I mental disorder. Scanning took place either at the Robarts Research Institute's Center for Functional and Metabolic Mapping or at the Lawson Health Research Institute for Imaging in London, ON, Canada. The study was approved by the research ethics board at Western University of Canada. All subjects provided written informed consent.

fMRI Data Acquisition

All images were collected using a 3.0, whole-body MRI scanner (Magnetom Tim Trio, Siemens Medical Solutions, Erlangen, Germany) with a manufacturer's 32-channel phased array head coil. BOLD fMRI images were obtained with the standard gradient-echo EPI pulse sequence. EPI volumes were acquired with 2 mm isotropic resolution with the following parameters: FOV = 192 mm × 192 mm × 128 mm (94 × 94 matrix, 64 slices), TR/TE = 3,000 ms/20 ms, flip angle = 90°, 120 volumes. Participants were instructed to close their eyes, relax, and let their minds wander during the 6-min resting scan.

Statistical Analyses

HRV analysis

Pulse data were recorded using a finger-tip pulse oximeter (Powerlab 8/35, LabChart 7 Pro) during the 6-min resting functional MRI scanning procedure and sampled at 200 Hz (Robarts Research Institute) or 40 Hz (Lawson Health Science Institute). Pulse data acquired at 40 Hz were re-sampled at 200 Hz by linear interpolation. In-house software provided by Stefanie Lis was applied for peak detection and visual inspection of the pulse signal. The inter-beat time series were saved as text files and imported to the KUBIOS Heart Rate Variability Software Package (Kuopio, Finland; version 2.2) [Tarvainen et al., 2014]. Data were then re-sampled at 4 Hz and common time domain measures were computed (RMSSD). Frequency domain estimates of HF-HRV and LF-HRV were calculated following the Task Force Guidelines (1996). Spectral analysis using a Fast Fourier transform algorithm, with a window width of 256 s and a window overlap of 50% was applied to generate heart period power spectrum. Frequency bands were set to 0.04–0.15 for the LF component and 0.15–0.4 for the HF component, respectively (Kuopio, Finland; version 2.2) [Tarvainen et al., 2014]. Raw measurements, including RMSSD, HF-HRV, and LF-HRV were natural log transformed before the analysis to normalize their distributions. Further analyses were

TABLE I. Sociodemographic, clinical, and physiological sample characteristics

	PTSD N = 57		HC N = 41		F/ χ value	P
Demographics						
Age mean (SD)	33.98	(11.57)	37.11	(12.80)	0.96	0.33
Gender (N)						
Female	18		26		0.27	0.61
Male	39		15			
Clinical Characteristics						
CAPS Re-experiencing mean (SD)	21.96	(6.73)	—		—	—
CAPS Avoidance/ Numbing mean (SD)	26.22	(7.00)	—		—	—
CAPS Hyperarousal mean (SD)	22.41	(5.99)	—		—	—
CAPS Total mean (SD)	69.16	(15.47)	—		—	—
MDI Depersonalization mean (SD)	8.11	(3.59)	5.24	(0.69)	22.45	<0.001
MDI Derealization mean (SD)	9.89	(4.12)	5.27	(0.66)	47.18	<0.001
MDI Total mean (SD)	61.96	(20.07)	34.15	(4.04)	64.31	<0.001
CTQ mean (SD)	62.47	(23.29)	31.28	(8.37)	65.61	<0.001
BDI mean (SD)	26.73	(9.79)	1.54	(2.60)	234.37	<0.001
State Characteristics						
STAI-S (SD)	1.97	(0.76)	1.22	(0.42)	32.66	<0.001
RSDI Dissociation (SD)	1.51	(0.61)	1.14	(0.21)	17.60	<0.001
RSDI Re-Experiencing (SD)	1.46	(0.67)	1.06	(0.19)	16.77	<0.001
RSDI Avoidance (SD)	2.04	(1.06)	1.35	(0.76)	14.33	<0.001
RSDI Hyperarousal (SD)	1.67	(0.56)	1.18	(0.26)	34.95	<0.001
Comorbidities (%)						
Alcohol dependence	0		0			
MDD	18		0			
Panic Disorder w/wo Agoraphobia	11		0			
Social Phobia	7		0			
Specific Phobia	3		0			
Generalized Anxiety Disorder	0		0			
Obsessive Compulsive Disorder	2		0			
Somatization Disorder	5		0			
Somatoform Disorder	8		0			
Eating Disorder	1		0			
HRV characteristics						
LN RMSDD mean (SD)	3.76	(0.09)	4.06	(0.11)	4.79	0.031
LN LF power value mean (SD)	6.05	(0.20)	6.91	(0.23)	8.34	0.005
LN HF power value mean (SD)	6.35	(0.19)	6.93	(0.25)	4.14	0.045

Significance threshold $P < 0.05$.

Abbreviations: PTSD = posttraumatic stress disorder group; HC = healthy control group; CAPS= Clinician-Administered PTSD scale; MDI = Multiscale Dissociation Inventory; CTQ = Childhood Trauma Questionnaire; BDI = Beck Depression Inventory; STAI-S = State-Trait Anxiety Inventar ; RSDI = Response to Script Driven Imagery Scale; MDD = major depressive disorder; RMSDD = root-mean square differences of successive R-R intervals; LF-HRV = low frequency heart rate variability, HF-HRV = high frequency heart rate variability, ln = natural logarithm.

performed using SPSS (version 22; SPSS). Group differences in HRV measurements were assessed with multivariate analyses of variance.

fMRI analysis

Image preprocessing and statistical analyses of the BOLD signal were conducted using Statistical Parametric Mapping (SPM 8, Wellcome Trust Center of Neuroimaging, London, UK: <http://www.fil.ion.ucl.ac.uk/spm>) implemented in Matlab 8.3 and 11b (MathWorks).

Preprocessing

For each subject, all functional images were realigned to the first image in their series, re-sliced and the mean functional image created. ART software (Gabrieli Lab; McGovern Institute for Brain Research, Cambridge, MA) was used to compute motion outlier regressors, which were applied within the 1st level analysis as a covariate of no interest. Given that an increasing body of literature points to an influence of head movement on functional connectivity measurements [Power et al., 2012, 2014; Pujol et al., 2014; Satterthwaite et al., 2013, 2012; Van Dijk et al., 2012],

we wanted to further ensure that groups did not differ with regard to their frequency of defined outliers. We, therefore, coded whether the ART software detected any outliers within each individual scan and subsequently ran a Fisher's exact test to examine potential group differences. Here, group differences were not observed ($P = 0.520$). The images were further spatially normalized and the mean image was co-registered to the SPM EPI template. The resulting deformation matrix was applied to the functional images. All images were then smoothed using a 6 mm full-width-half-maximum isotropic Gaussian filter and band-pass filtered within a 0.012–0.1 Hz range with in house software by Jean Théberge.

Connectivity analysis

For each subject, the mean signal intensity time course was extracted from seeds defined in WFU PickAtlas (Functional MRI Laboratory, Wake Forest University School of Medicine) using in-house software by co-author Jean Théberge for each seed region, including the vmPFC [2 22 –8] [Thayer et al., 2012], right and left amygdala (WFU PickAtlas), and PAG [–2 –27 –3] [Napadow et al., 2008]. All seed regions were based on previous publications and defined as 6 mm spheres for each reported coordinate. Extracted time courses were then used as a regressor within a first level multiple regression model for each seed region. Connectivity was thus indicative of a correlation between each seed and other brain areas, where both positive correlations and anti-correlations were examined.

To explore connectivity patterns within groups, one-sample t -tests were conducted for each group separately. To control for the effects of a current major depression, we re-ran the within-group PTSD analyses by applying a multivariate analysis of covariance and including the comorbid diagnosis as a covariate of no interest. Two-sample t -tests were used to compare PTSD with healthy control subjects for each seed region. Statistical significance was set at a P -value of < 0.001 corrected for whole brain FDR and an extent threshold of $k = 10$ voxels.

Correlations between HRV and BOLD functional connectivity

To examine associations between HRV and CAN-related connectivity patterns, separate multiple regression analyses for RMSSD, LF-HRV, and HF-HRV scores were examined; each of these scores revealed significant differences between groups. PAG seed correlations were masked with a CAN mask. The CAN mask was created in WFU PickAtlas based on 10 mm spheres on prominent coordinates published in previous literature [Beissner et al., 2013; Thayer et al., 2012], that is, bilateral amygdala [–20 –6 –18] and [20 –6 –18] [Beissner et al., 2013], right medial prefrontal cortex [10 54 18; Thayer et al., 2012], right dorsal cingulate [2 10 40] [Beissner et al., 2013]; left thalamus [–4 –16 8] [Beissner et al., 2013]. Two additional masks were created, comprising the bilateral insula from the WFU PickAtlas applied to the

Figure 1.

Between group differences in HRV. Figure represents significant lower Mean Natural Log RMSSD, LF-HRV, and HF-HRV in PTSD as compared to the control group (all $F_{(1,96)}$'s > 4.14 , all P 's < 0.048). Statistical threshold $P < 0.05$. Abbreviations: PTSD = posttraumatic stress disorder group; HC = healthy control group; RMSSD = root-mean square differences of successive R-R intervals; LF-HRV = low frequency heart rate variability, HF-HRV = high frequency heart rate variability, $\ln$ = natural logarithm.

vmPFC seed correlation, as well as a PAG mask [Napadow et al., 2008] applied to the right and left amygdala seed correlations, each with a 10-mm radius. A second level multiple regression model was established for each seed in combination with each HRV score, respectively. Simple multiple regression models examining both positive and negative associations between scores were conducted and the results masked with the CAN, PAG, and insula mask. Two-samples t -tests were used to compare PTSD with healthy controls for each seed region and each HRV score respectively, with a covariate (HRV score), and a term that computes the interaction between the covariate and factor group. The interaction effect indicates regions for which the functional connectivity with the seed region is greater in people that have greater HRV scores in controls than individuals with PTSD (and vice versa). Significance was set to $P < 0.05$, ROI FDR corrected and an extent threshold of $k = 10$ voxels.

RESULTS

Cardiovascular Responding (HRV)

As compared to controls, individuals with PTSD showed lower RMSSD values ($F_{(1,96)} = 4.79$, $P = 0.031$, $\eta^2 = 0.048$), lower LF power ($F_{(1,96)} = 8.34$, $P = 0.005$, $\eta^2 = 0.080$) and lower HF power ($F_{(1,96)} = 4.14$, $P = 0.045$, $\eta^2 = 0.041$) (see also Fig. 1, Table I).

Seed-Based Functional Connectivity

Ventromedial prefrontal cortex

Resting-state activity in the vmPFC predicted a localized connectivity pattern within control subjects. In contrast, vmPFC activity predicted widespread connectivity to multiple cortical and subcortical regions in PTSD. Adding

Figure 2.

Between group differences in key CAN seed resting state functional connectivity, including vmPFC, amygdala, PAG. Illustration represents greater connectivity between key CAN seed regions, including the vmPFC, amygdala, and PAG with multiple cortical and subcortical regions in PTSD as compared to healthy controls. By contrast, HC individuals did not exhibit any significantly greater connectivity to any of the CAN seed regions as

compared to PTSD individuals. Statistical threshold $P < 0.001$ corrected for whole brain FDR, $k = 10$ for all two-sample t -tests. Abbreviations: PTSD = posttraumatic stress disorder group; HC = healthy control group; vmPFC = ventromedial prefrontal cortex; PAG = periaqueductal gray; FDR = false discovery rate. [Color figure can be viewed at wileyonlinelibrary.com.]

current major depression as a covariate did not change the results (for details on each seed regions see Supporting Information Table I, Supporting Information Figure I).

Between-group comparisons indicated that HC individuals did not exhibit any significantly greater connectivity to the vmPFC as compared to PTSD individuals. By contrast, PTSD subjects exhibited greater functional connectivity between the vmPFC and cingulate cortex, frontal cortex, bilateral precentral gyrus, left insula, left parahippocampal gyrus, bilateral lentiform nuclei, and left thalamus (see also Fig. 2A, Supporting Information Table IV).

Amygdala

Resting state connectivity with the left and right amygdala predicted a restricted connectivity pattern in controls.

In contrast, among individuals with PTSD, left and right amygdala activity predicted widespread connectivity patterns in multiple cortical and subcortical regions. Adding current major depression as a covariate did not change the results (for details on each seed regions see Supporting Information Table II, Supporting Information Figure I).

Between-group comparisons indicated that HCs did not exhibit significantly greater connectivity to either the left or the right amygdala as compared to individuals with PTSD. By contrast, PTSD individuals exhibited greater functional connectivity between the left amygdala and the cingulate cortex, frontal cortex, bilateral precentral gyrus, bilateral insula, bilateral parahippocampal gyrus, bilateral thalamus, and bilateral lentiform nuclei (see also Fig. 2B, Supporting Information Table VA). Similarly, individuals

Figure 3.

HRV correlations related connectivity to key CAN seed regions within healthy control subjects, including vmPFC, amygdala, PAG. Illustration represents increased functional connectivity patterns to all CAN regions in controls, which is predicted by HRV. In PTSD, analyses did not reveal significant clusters; only healthy control subjects are, therefore, presented. Significance threshold

$P < 0.05$, ROI FDR corrected, $k = 10$ voxels for all one-sample t -tests. Abbreviations: PTSD = posttraumatic stress disorder group; HC = healthy control group; vmPFC = ventromedial prefrontal cortex; PAG = periaqueductal gray; ROI = region of interest analysis. [Color figure can be viewed at wileyonlinelibrary.com.]

with PTSD showed greater functional connectivity between the right amygdala and the cingulate cortex, frontal cortex, bilateral precentral gyrus, right parahippocampal gyrus, bilateral insula, and bilateral lentiform nuclei (see also Fig. 2B, Supporting Information Table VB).

Periaqueductal gray

Resting-state activity in the PAG predicted a localized connectivity pattern in control individuals. In contrast, PAG activity predicted widespread connectivity in multiple cortical and subcortical regions in PTSD subjects. Adding current major depression as a covariate did not change the results (for details on each seed regions see Supporting Information Table III, Supporting Information Figure I).

HC subjects did not show any greater connectivity to the PAG as compared to PTSD subjects. In contrast, individuals with PTSD showed greater functional connectivity between the PAG seed and the bilateral mid cingulate cortex, left dorsal cingulate cortex, bilateral medial frontal gyrus, left superior frontal gyrus, left precentral gyrus, left

insula, bilateral putamen, left thalamus, hippocampus, and right cerebellar vermis (see also Fig. 2C, Supporting Information Table VI).

HRV Correlations to CAN Related Seed-Based Functional Connectivity

Ventromedial prefrontal cortex

In control subjects, higher RMSSD, LF-HRV, and HF-HRV values predicted increased functional connectivity between the vmPFC and the right insula (see also Fig. 3A, Table II). In contrast, none of the HRV scores were related to increased or decreased connectivity between the vmPFC and CAN-related brain regions among individuals with PTSD (Table II).

Direct group comparisons revealed that none of the HRV scores predicted increased functional connectivity between the vmPFC and CAN related brain regions in neither controls as compared to PTSD, nor within the reversed contrast (see also Table III for additional information regarding FWE corrected results).

TABLE II. HRV correlations related connectivity to key CAN seed regions within healthy control subjects, as well as within PTSD individuals, including the vmPFC, amygdala, PAG

HRV score	Brain regions	BA	Hemisphere	<i>x</i>	<i>y</i>	<i>z</i>	<i>k</i>	<i>T</i> -value	pFDR	pFWE	
vmPFC											
HC											
RMSSD	Insula	13	R	34	−2	14	55	4.24	0.057	0.031	
HF-HRV	Insula	13	R	34	−2	14	116	4.12	0.022	0.023	
LF-HRV	Insula	13	R	38	−12	14	49	4.05	0.058	0.049	
PTSD											
n.s.											
Left Amygdala											
HC											
RMSSD	PAG, Pulvinar		L	−4	−26	4	15	4.02	0.010	0.013	
HF-HRV	PAG, Pulvinar		L	−4	−28	4	14	4.05	0.010	0.012	
LF-HRV	PAG, Pulvinar		L	−6	−28	4	8	3.15	0.055	0.092	
PTSD											
n.s.											
Right Amygdala											
HC											
n.s.											
PTSD											
n.s.											
PAG											
HC											
RMSSD	Dorsal cingulate cortex	32	R	6	8	44	87	4.50	0.028	0.021	
	Medial frontal cortex	9	R	4	52	24	55	4.33	0.028	0.033	
	Medial frontal cortex	10	R	4	56	12	Of55	3.25	0.038	0.346	
	Medial frontal cortex	9	R	4	46	18	Of55	2.85	0.044	0.616	
	Thalamus	md nucl.	L	−4	−8	10	18	3.97	0.028	0.080	
	Superior frontal cortex	10	R	18	58	22	14	3.23	0.038	0.358	
	HF-HRV	Medial frontal cortex	9	R	4	52	24	63	4.44	0.036	0.025
		Thalamus	md nucl.	L	−4	−8	10	26	4.25	0.036	0.040
		Dorsal cingulate cortex	32	R	6	8	44	32	3.59	0.038	0.181
	Superior frontal cortex	10	R	18	58	22	12	3.26	0.049	0.341	
LF-HRV	n.s.										
PTSD											
n.s.											

Significance threshold $P < 0.05$, ROI FDR corrected, $k = 10$ voxels for all one-sample t -tests.

Abbreviations: PTSD = posttraumatic stress disorder group; HC = healthy control group; vmPFC = ventromedial prefrontal cortex; PAG = periaqueductal gray; md nucl = mediodorsal nucleus; FDR = false discovery rate; FWE = family wise error, RMSSD = root-mean square differences of successive R-R intervals; LF-HRV = low frequency heart rate variability, HF – HRV = high frequency heart rate variability; BA = Brodmann Areal.

Amygdala

In control subjects, higher RMSSD, as well as higher LF-HRV and HF-HRV values predicted increased connectivity between the left amygdala and the left PAG, while none of the HRV scores were related to the functional connectivity of the right amygdala in controls (see also Fig. 3B, Table II). In individuals with PTSD, none of the HRV scores were related to increased or decreased connectivity between the left or right amygdala and CAN-related brain regions (Table II).

Direct group comparisons revealed that none of the HRV scores predicted increased connectivity between the left amygdala and CAN related brain regions in controls as compared to PTSD, nor within the reversed contrast

(see also Table III for additional information regarding FWE corrected results). Direct group comparisons revealed that higher RMSSD and HF-HRV values predicted increased connectivity between the right amygdala and the right insula in controls as compared to PTSD. In contrast, none of the HRV scores predicted increased connectivity patterns to the right amygdala in PTSD patients as compared to controls (see also Table III for additional information regarding FWE corrected results).

Periaqueductal gray

In control subjects, higher RMSSD and HF-HRV values also predicted increased functional connectivity between the PAG and the right dorsal cingulate cortex, right medial

TABLE III. HRV correlations related connectivity to key CAN seed regions for healthy control subjects greater than PTSD, as well as for the reversed contrast including the vmPFC, amygdala, PAG

HRV Score	Brain regions	BA	Hemisphere	<i>x</i>	<i>y</i>	<i>z</i>	<i>k</i>	<i>T</i> -value	pFDR	pFWE
vmPFC										
HC>PTSD										
RMSSD	Insula	13	R	54	2	-6	20	3.16	0.147	0.050
HF-HRV	Insula	13	R	54	2	-6	24	3.19	0.094	0.049
LF-HRV		n.s.								
PTSD>HC										
n.s.										
Left Amygdala										
HC>PTSD										
RMSSD	Insula	13	R	54	2	-6	5*	3.08	0.208	0.055
HF-HRV		n.s.								
LF-HRV		n.s.								
PTSD>HC										
n.s.										
Right Amygdala										
HC>PTSD										
RMSSD	Insula	13	R	54	2	-8	28	3.95	0.015	0.007
HF-HRV	Insula	13	R	54	2	-8	23	3.60	0.034	0.020
LF-HRV		n.s.								
PTSD>HC										
n.s.										
PAG										
HC>PTSD										
RMSSD	Insula	13	R	48	16	-2	20	2.85	0.212	0.049
HF-HRV	Insula	13	R	48	16	-2	33	3.07	0.036	0.065
LF-HRV		n.s.								
PTSD>HC										
n.s.										

Significance threshold $P < 0.05$, ROI FDR corrected, $k = 10$ voxels for all two-sample *t*-tests. * did not meet *k* threshold.

Abbreviations: PTSD = posttraumatic stress disorder group; HC = healthy control group; vmPFC = ventromedial prefrontal cortex; PAG = periaqueductal gray; FDR = false discovery rate; FWE = family wise error, RMSSD = root-mean square differences of successive R-R intervals; LF-HRV = low frequency heart rate variability, HF-HRV = high frequency heart rate variability; BA = Brodmann Areal.

prefrontal cortex, right superior frontal gyrus and left thalamus. In contrast, LF-HRV did not predict functional connectivity between the PAG and CAN-related brain regions in HC subjects, nor did any of the HRV scores predicted increased functional connectivity patterns to the PAG in PTSD (see also see also Fig. 3C, Table II).

Direct group comparisons revealed that higher HF-HRV values predicted increased functional connectivity between the PAG and the right insula in controls as compared to PTSD. In contrast, none of the HRV scores predicted increased connectivity patterns to the PAG in PTSD patients as compared to controls (see also Table III for additional information regarding FWE corrected results).

DISCUSSION

To the best of our knowledge, this is the first study to investigate autonomic response and its relation to resting state central autonomic activity in individuals with PTSD. As hypothesized, as compared to controls, individuals

with PTSD showed reduced resting HRV. At the neuronal level, resting-state activity in CAN-associated brain regions, including the vmPFC, amygdala, and PAG predicted a localized and restricted connectivity pattern in control subjects. In stark contrast, individuals with PTSD exhibited widespread functional connectivity to brain regions involved in emotional reactivity [Kirby and Robinson, 2015; White et al., 2015], motor readiness [Herzfeld et al., 2014; Mendoza and Merchant, 2014; Stoodley and Schmahmann, 2010; Stoodley et al., 2012], self-referential processing [Bluhm et al., 2011, 2012; Buckner et al., 2008; Spreng et al., 2009], and stimuli salience detection [Craig, 2009; Sripada et al., 2012; Thome et al., 2014] (emotional reactivity: all seed regions to the insula, vmPFC, and amygdala to the lentiform nucleus; motor readiness: all seed regions to the precentral gyrus, PAG to the cerebellum, self-referential processing: vmPFC and PAG to PCC, dmPFC, amygdala to rostral ACC, salience detection: all seed regions to anterior insula and dorsal ACC). Critically, whereas higher HRV predicted increased functional connectivity within key regions of the CAN, no such

correlations were uncovered when examining patients with PTSD, pointing to a relative uncoupling of the ANS from the CAN in this group.

As compared to controls, individuals with PTSD exhibited lower HRV at rest as reflected by all HRV parameters. Indeed, the present findings are consistent with previous cross-sectional studies that reported overall lower resting HRV in PTSD even after controlling for important confounders such as incidence of traumatic brain injury and severity of depression [Minassian et al., 2014; Moon et al., 2013; Norte et al., 2013; Shah et al., 2013; Shaikh al arab et al., 2012; Wahbeh and Oken, 2013], a pattern confirmed by a recent meta-analysis [Chalmers et al., 2014]. Blunted HRV in PTSD is also in line with previous theoretical models, such as the neurovisceral integration model [Thayer et al., 2009], which proposes that prefrontal regions moderate parasympathetic activity and vagal nerve inhibition, and that inadequate vagus nerve regulation is associated with a variety of somatic and mental diseases [Danesh et al., 2000; Duncan et al., 2003; Gao and Hong, 2008]. In this model, whereas healthy functioning is characterized by a high level of adaptive HRV [Thayer et al., 2012, 2009; Thayer and Lane, 2000], lower HRV is thought to reflect decreased vagal output leading to behavioral inflexibility.

Consistent with the neurovisceral integration model, the polyvagal theory [Porges, 2011] further posits that a safe environment promotes an increase of vagal outflow, thereby promoting regeneration and homeostatic functions. It is, therefore, probable that difficulties in detecting safe environments result in an increase in sympathetic tone and concomitant decreased HRV. Indeed, several studies reported a failure of contextual learning in PTSD resulting in an inability to differentiate between threat and safety contexts [Acheson et al., 2015; Levy-Gigi et al., 2016; Steiger et al., 2015; van Wingen et al., 2011]. For example, a recent study demonstrated that individuals with PTSD did not differentiate between threat and safety contexts on a behavioral level (contingency ratings) during the acquisition phase of a differential context and cue conditioning paradigm; this was reflected in increased hippocampal activation in response to both contexts [Steiger et al., 2015]. This decreased capacity for safety perception may, therefore, be associated with sympathetic over reactivity and/or parasympathetic insufficiency, and decreased HRV in PTSD [Tulloch et al., 2014]. However, it is important to note that emerging evidence of blunted HRV and reduced flexibility in PTSD is based primarily on cross-sectional studies. Longitudinal studies are, therefore, urgently needed to determine whether an inflexible ANS represents a vulnerability factor for PTSD or instead emerges as the result of heightened stress states. Interestingly, Minassian et al. [2015] recently demonstrated that, in active-duty marines, higher LF/HF ratios at a pre-deployment visit (1 to 2 months before combat exposure) were positively correlated with the risk of developing PTSD as measured at a

post-deployment visit (4 to 6 months after return) [Minassian et al., 2015]. This study points to the fact that ANS functioning may contribute to an individual's proneness, as well as resiliency, in reacting to stress and may, therefore, represent a risk factor for the development of trauma-related disorders [see also Shaikh al arab et al., 2012].

Although numerous studies have investigated HRV in PTSD, here we describe the first study directly linking HRV to brain regions that modulate the variability of HR time series (CAN). While higher HRV scores in healthy controls predicted increased functional connectivity within regions of the CAN (vmPFC to bilateral insula; left amygdala to the left PAG; PAG to right medial and superior PFC dorsal cingulate cortex, and left thalamus; for meta-analyses see also [Beissner et al., 2013; Thayer et al., 2012], measures of HRV within the PTSD group were unrelated with functional connectivity within CAN-related brain regions. The absence of a correlation between HRV and functional connectivity of the CAN within PTSD subjects may reflect an uncoupling of the ANS from the CAN in PTSD where top-down modulation of cardiac function by higher-order brain regions fails to occur. Such a pattern has indeed been reported across a number of different paradigms including emotion-provoking pictures and film scenes, facial processing, gambling tasks, and traumatic script-driven imagery [Cohen et al., 2013; Etkin and Wager, 2007; Felmingham et al., 2008; Hopper et al., 2007b; Lanius et al., 2012, 2015; Mickleborough et al., 2011; Reiser et al., 2014; Shin et al., 2006; van Wingen et al., 2011]. Interestingly, a recent positron emission tomography study in PTSD found an increased correlation between HR and activation in orbitofrontal, precentral and occipital regions in response to traumatic scripts solely among individuals with PTSD [Barkay et al., 2012], suggesting that sensorimotor regions might regulate the stress response induced by traumatic scripts in PTSD. In comparison, the absence of HRV-CAN covariation at rest within the present sample demonstrates a lack of top-down CAN regulation of autonomic responses.

It is important to note that the present sample of individuals with PTSD displayed widespread resting state functional connectivity between CAN-related regions and other brain regions associated with emotional reactivity, motor readiness, self-referential processing, and stimuli salience detection. However, the examined regions appear neither associated with HRV nor associated with anatomical connections to the brain stem, which have been directly related to the vagus nerve. Future studies examining brain stem structures directly are, therefore, warranted.

There is a growing literature suggesting that low resting HRV and related psychopathological states are associated with undifferentiated threat responses to a wide range of conditions and situations [Melzig et al., 2009; Ruiz-Padial and Thayer, 2014; Wendt et al., 2015]. Thus, the present findings of widespread connectivity in PTSD may reflect

the neural concomitants of a response pattern, reflecting difficulties in differentiating between threat and safety contexts, which has been observed in other studies [e.g., see Steiger et al., 2015]. However, in contrast, context appropriate activation of localized brain networks has been identified with context appropriate psychophysiological and behavioral responses in persons with high resting HRV, which is associated with a flexibly recruited network of loosely coupled bio-oscillators that has been linked to healthy dynamical systems [Thayer, 2006; Thayer and Lane, 2000]. The widespread connectivity exhibited by PTSD patients in the present study might point to a response pattern that is suboptimal in generating context appropriate responses.

The observed pattern of resting state brain activation in PTSD is also consistent with Panksepp and Biven's [2012] proposal of a basic affective system. Here, the PAG is thought to gate various affective responses or emotional systems through its connections with higher brain structures whose activity contributes to the functioning of all basic emotional systems. Given the increased widespread PAG resting functional connectivity observed in PTSD subjects, it is further probable that this pattern is reflective of a hypersensitive affective system observed during rest. However, future research is needed to more fully investigate this notion.

Several limitations of the current study are worth noting. Due to the cross-sectional design of the present study, these findings provide no indication of whether decreased HRV is caused by traumatic stress or represents a pre-existing vulnerability factor. We did not include trauma-exposed individuals without PTSD in our analyses, as screened individuals matched for trauma severity met exclusion criteria for current or past psychiatric disorders. In addition, given that only one related study links HR to cerebral blood flow during traumatic script exposure [Barkay et al., 2012], future studies using symptom provocation are expected to be useful in identifying potential differential functional connectivity patterns in relation to HRV in PTSD. Here, comparing peripheral physiological and neuronal responses during resting state to those induced during symptom provocation will be necessary to gain insights into emotional regulatory capacity in response to stressors.

In conclusion, the present study represents an important first step in linking peripheral (HRV) and central (CAN) measures of autonomic response in PTSD. HRV was reduced in PTSD, while the CAN evidenced increased functional connectivity across myriad of brain regions, including those associated with emotional processing and motor readiness. Moreover, whereas CAN-related functional connectivity was strongly correlated with HRV in controls, a striking lack of covariation between HRV and CAN-functional connectivity was observed in PTSD. The present investigation, therefore, represents a first step in demonstrating the relative absence of the regulatory

capacity of the CAN on the ANS functioning at rest in PTSD, which may partially mediate the negative alterations in cognition and mood observed in this disorder.

ACKNOWLEDGMENTS

Dr. Lanius was the principal investigator of the study and supervised the collection and analysis of data. fMRI data acquisition was supervised by Dr. Thèberge. Mrs. Densmore and Ms. Thome processed and analyzed the data; Ms. Thome prepared the paper, with review and contributions by Dr. Lanius, and the rest of the authors. We thank Stefanie Lis for her support in analyzing the pulse data and providing the software. Conflict of interest/financial disclosure: The following statement covers the last three years: All the authors have no interests to declare generally and in relation to the present study.

REFERENCES

- Acheson DT, Geyer MA, Baker DG, Nievergelt CM, Yurgil K, Risbrough VB (2015): Conditioned fear and extinction learning performance and its association with psychiatric symptoms in active duty Marines. *Psychoneuroendocrinology* 51:495–505.
- Agorastos A, Boel JA, Heppner PS, Hager T, Moeller-Bertram T, Haji U, Motazedi A, Yanagi MA, Baker DG, Stiedl O (2013): Diminished vagal activity and blunted diurnal variation of heart rate dynamics in posttraumatic stress disorder. *Stress* 16: 300–310.
- Barkay G, Freedman N, Lester H, Louzoun Y, Sapoznikov D, Luckenbaugh D, Shalev AY, Chisin RG, Bonne O (2012): Brain activation and heart rate during script-driven traumatic imagery in PTSD: Preliminary findings. *Psychiatry Res* 204:155–160.
- Beck AT, Guth D, Steer RA, Ball R (1997): Screening for major depression disorders in medical inpatients with the Beck Depression Inventory for Primary Care. *Behav Res Ther* 35: 785–791.
- Beissner F, Meissner K, Bar KJ, Napadow V (2013): The autonomic brain: An activation likelihood estimation meta-analysis for central processing of autonomic function. *J Neurosci* 33: 10503–10511.
- Benarroch EE (1993): The central autonomic network: Functional organization, dysfunction, and perspective. *Mayo Clin Proc* 68: 988–1001.
- Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, Stokes J, Handelsman L, Medrano M, Desmond D, Zule W (2003): Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl* 27:169–190.
- Blake DD, Weathers FW, Nagy LM, Kaloupek DG, Gusman FD, Charney DS, Keane TM (1995): The development of a clinician-administered PTSD scale. *J Trauma Stress* 8:75–90.
- Bluhm RL, Clark CR, McFarlane AC, Moores KA, Shaw ME, Lanius RA (2011): Default network connectivity during a working memory task. *Hum Brain Mapp* 32:1029–1035.
- Bluhm RL, Frewen PA, Coupland NC, Densmore M, Schore AN, Lanius RA (2012): Neural correlates of self-reflection in post-traumatic stress disorder. *Acta Psychiatr Scand* 125:238–246.

- Briere J, Weathers FW, Runtz M (2005): Is dissociation a multidimensional construct? data from the multiscale dissociation inventory. *J Trauma Stress* 18:221–231.
- Buckner RL, Andrews-Hanna JR, Schacter DL (2008): The brain's default network: Anatomy, function, and relevance to disease. *Ann N Y Acad Sci* 1124:1–38.
- Carney RM, Freedland KE (2009): Depression and heart rate variability in patients with coronary heart disease. *Cleve Clin J Med* 76(Suppl 2):S13–S17.
- Cersosimo MG, Benarroch EE (2013): Central control of autonomic function and involvement in neurodegenerative disorders. *Handb Clin Neurol* 117:45–57.
- Chalmers JA, Quintana DS, Abbott MJ, Kemp AH (2014): Anxiety disorders are associated with reduced heart rate variability: A meta-analysis. *Front Psychiatry* 5:80.
- Cohen H, Kotler M, Matar MA, Kaplan Z, Loewenthal U, Miodownik H, Cassuto Y (1998): Analysis of heart rate variability in posttraumatic stress disorder patients in response to a trauma-related reminder. *Biol Psychiatry* 44:1054–1059.
- Cohen H, Benjamin J, Geva AB, Matar MA, Kaplan Z, Kotler M (2000): Autonomic dysregulation in panic disorder and in post-traumatic stress disorder: Application of power spectrum analysis of heart rate variability at rest and in response to recollection of trauma or panic attacks. *Psychiatry Res* 96:1–13.
- Cohen JE, Shalev H, Admon R, Hefetz S, Gasho CJ, Shachar LJ, Shelef I, Hendler T, Friedman A (2013): Emotional brain rhythms and their impairment in post-traumatic patients. *Hum Brain Mapp* 34:1344–1356.
- Craig AD (2009): How do you feel—now? The anterior insula and human awareness. *Nat Rev Neurosci* 10:59–70.
- Danesh J, Whincup P, Walker M, Lennon L, Thomson A, Appleby P, Gallimore JR, Pepys MB (2000): Low grade inflammation and coronary heart disease: Prospective study and updated meta-analyses. *BMJ* 321:199–204.
- Duncan BB, Schmidt MI, Pankow JS, Ballantyne CM, Couper D, Vigo A, Hoogeveen R, Folsom AR, Heiss G (2003): Low-grade systemic inflammation and the development of type 2 diabetes: The atherosclerosis risk in communities study. *Diabetes* 52:1799–1805.
- Etkin A, Wager TD (2007): Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 164:1476–1488.
- Felmingham K, Kemp AH, Williams L, Falconer E, Olivieri G, Peduto A, Bryant R (2008): Dissociative responses to conscious and non-conscious fear impact underlying brain function in post-traumatic stress disorder. *Psychol Med* 38:1771–1780.
- First MB (1997): User's guide for the structured clinical interview for DSM-IV axis I disorders SCID-I: Clinician version. Washington, DC: American Psychiatric Press.
- Friedman MJ, Resick PA, Bryant RA, Brewin CR (2011): Considering PTSD for DSM-5. *Depress Anxiety* 28:750–769.
- Gao HM, Hong JS (2008): Why neurodegenerative diseases are progressive: Uncontrolled inflammation drives disease progression. *Trends Immunol* 29:357–365.
- Gillie BL, Thayer JF (2014): Individual differences in resting heart rate variability and cognitive control in posttraumatic stress disorder. *Front Psychol* 5:758.
- Hauschildt M, Peters MJ, Moritz S, Jelinek L (2011): Heart rate variability in response to affective scenes in posttraumatic stress disorder. *Biol Psychol* 88:215–222.
- Herzfeld DJ, Pastor D, Haith AM, Rossetti Y, Shadmehr R, O'Shea J (2014): Contributions of the cerebellum and the motor cortex to acquisition and retention of motor memories. *NeuroImage* 98:147–158.
- Hopper JW, Frewen PA, Sack M, Lanius RA, Van der Kolk BA (2007a): The Responses to Script-Driven Imagery Scale (RSDI): Assessment of state posttraumatic symptoms for psychobiological and treatment research. *J Psychopathol Behav Assess* 29:249–268.
- Hopper JW, Frewen PA, van der Kolk BA, Lanius RA (2007b): Neural correlates of reexperiencing, avoidance, and dissociation in PTSD: Symptom dimensions and emotion dysregulation in responses to script-driven trauma imagery. *J Trauma Stress* 20:713–725.
- Keary TA, Hughes JW, Palmieri PA (2009): Women with posttraumatic stress disorder have larger decreases in heart rate variability during stress tasks. *Int J Psychophysiol* 73:257–264.
- Kirby LA, Robinson JL (2015): Affective mapping: An activation likelihood estimation (ALE) meta-analysis. *Brain Cogn* (in press).
- Kvaal K, Ulstein I, Nordhus IH, Engedal K (2005): The Spielberger State-Trait Anxiety Inventory (STAI): The state scale in detecting mental disorders in geriatric patients. *Int J Geriatr Psychiatry* 20:629–634.
- Lanius RA, Vermetten E, Loewenstein RJ, Brand B, Schmahl C, Bremner JD, Spiegel D (2010): Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *Am J Psychiatry* 167:640–647.
- Lanius RA, Bluhm RL, Frewen PA (2011): How understanding the neurobiology of complex post-traumatic stress disorder can inform clinical practice: A social cognitive and affective neuroscience approach. *Acta Psychiatr Scand* 124:331–348.
- Lanius RA, Brand B, Vermetten E, Frewen PA, Spiegel D (2012): The dissociative subtype of posttraumatic stress disorder: Rationale, clinical and neurobiological evidence, and implications. *Depress Anxiety* 29:701.
- Lanius RA, Frewen PA, Tursich M, Jetly R, McKinnon MC (2015): Restoring large-scale brain networks in PTSD and related disorders: A proposal for neuroscientifically-informed treatment interventions. *Eur J Psychotraumatol* 6:27313.
- Levy-Gigi E, Richter-Levin G, Okon-Singer H, Keri S, Bonanno GA (2016): The hidden price and possible benefit of repeated traumatic exposure. *Stress* 19:1–7.
- Melzig CA, Weike AI, Hamm AO, Thayer JF (2009): Individual differences in fear-potentiated startle as a function of resting heart rate variability: Implications for panic disorder. *Int J Psychophysiol* 71:109–117.
- Mendoza G, Merchant H (2014): Motor system evolution and the emergence of high cognitive functions. *Prog Neurobiol* 122:73–93.
- Mickleborough MJ, Daniels JK, Coupland NJ, Kao R, Williamson PC, Lanius UF, Hegadoren K, Schore A, Densmore M, Stevens T, Lanius RA (2011): Effects of trauma-related cues on pain processing in posttraumatic stress disorder: An fMRI investigation. *J Psychiatry Neurosci* 36:6–14.
- Minassian A, Geyer MA, Baker DG, Nievergelt CM, O'Connor DT, Risbrough VB (2014): Heart rate variability characteristics in a large group of active-duty marines and relationship to posttraumatic stress. *Psychosom Med* 76:292–301.
- Minassian A, Maihofer AX, Baker DG, Nievergelt CM, Geyer MA, Risbrough VB (2015): Association of predeployment heart rate variability with risk of postdeployment posttraumatic stress disorder in active-duty marines. *JAMA Psychiatry* 72:979–986.
- Moon E, Lee SH, Kim DH, Hwang B (2013): Comparative study of heart rate variability in patients with schizophrenia, bipolar

- disorder, post-traumatic stress disorder, or major depressive disorder. *Clin Psychopharmacol Neurosci* 11:137–143.
- Napadow V, Dhond R, Conti G, Makris N, Brown EN, Barbieri R (2008): Brain correlates of autonomic modulation: combining heart rate variability with fMRI. *NeuroImage* 42:169–177.
- Nawijn L, van Zuiden M, Frijling JL, Koch SB, Veltman DJ, Olff M (2015): Reward functioning in PTSD: A systematic review exploring the mechanisms underlying anhedonia. *Neurosci Biobehav Rev* 51:189–204.
- Nicholson AA, Densmore M, Frewen PA, Theberge J, Neufeld RW, McKinnon MC, Lanius RA (2015): The dissociative subtype of posttraumatic stress disorder: Unique resting-state functional connectivity of basolateral and centromedial amygdala complexes. *Neuropsychopharmacology* 40:2317–2326.
- Norte CE, Souza GG, Vilete L, Marques-Portella C, Coutinho ES, Figueira I, Volchan E (2013): They know their trauma by heart: An assessment of psychophysiological failure to recover in PTSD. *J Affect Disord* 150:136–141.
- Palkovits M (1999): Interconnections between the neuroendocrine hypothalamus and the central autonomic system. Geoffrey Harris Memorial Lecture, Kitakyushu, Japan, October. *Front Neuroendocrinol* 20:270–295.
- Panksepp J, Biven L (2012): *The Archaeology of Mind: Neuroevolutionary Origins of Human Emotions*. New York, NY: W.W. Norton & Company.
- Porges SW (2011): *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-Regulation*. New York, NY: W.W. Norton & Company.
- Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE (2012): Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage* 59:2142–2154.
- Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE (2014): Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage* 84:320–341.
- Powers A, Cross D, Fani N, Bradley B (2015a): PTSD, emotion dysregulation, and dissociative symptoms in a highly traumatized sample. *J Psychiatr Res* 61:174–179.
- Powers A, Etkin A, Gyurak A, Bradley B, Jovanovic T (2015b): Associations between childhood abuse, posttraumatic stress disorder, and implicit emotion regulation deficits: Evidence from a low-income, inner-city population. *Psychiatry* 78:251–264.
- Pujol J, Macia D, Blanco-Hinojo L, Martinez-Vilavella G, Sunyer J, de la Torre R, Caixas A, Martin-Santos R, Deus J, Harrison BJ (2014): Does motion-related brain functional connectivity reflect both artifacts and genuine neural activity? *NeuroImage* 101:87–95.
- Reinders AA, Willemsen AT, den Boer JA, Vos HP, Veltman DJ, Loewenstein RJ (2014): Opposite brain emotion-regulation patterns in identity states of dissociative identity disorder: A PET study and neurobiological model. *Psychiatry Res* 223:236–243.
- Reiser EM, Weiss EM, Schuster G, Holmes EA, Fink A, Papousek I (2014): Prefrontal-posterior coupling while observing the suffering of other people, and the development of intrusive memories. *Psychophysiology* 51:546–555.
- Resick PA, Galovski TE, O'Brien Uhlmansiek M, Scher CD, Clum GA, Young-Xu Y (2008): A randomized clinical trial to dismantle components of cognitive processing therapy for posttraumatic stress disorder in female victims of interpersonal violence. *J Consult Clin Psychol* 76:243–258.
- Ruiz-Padial E, Thayer JF (2014): Resting heart rate variability and the startle reflex to briefly presented affective pictures. *Int J Psychophysiol* 94:329–335.
- Sadeh N, Spielberg JM, Miller MW, Milberg WP, Salat DH, Amick MM, Fortier CB, McGlinchey RE (2015): Neurobiological indicators of disinhibition in posttraumatic stress disorder. *Hum Brain Mapp* 36:3076–3086.
- Sammito S, Thielmann B, Zimmermann P, Bockelmann I (2015): [Influence of post-traumatic stress disorder on heart rate variability as marker of the autonomic nervous System. A systematic review]. *Fortschr Neurol Psychiatr* 83:30–37.
- Satterthwaite TD, Wolf DH, Loughhead J, Ruparel K, Elliott MA, Hakonarson H, Gur RC, Gur RE (2012): Impact of in-scanner head motion on multiple measures of functional connectivity: Relevance for studies of neurodevelopment in youth. *NeuroImage* 60:623–632.
- Satterthwaite TD, Elliott MA, Gerraty RT, Ruparel K, Loughhead J, Calkins ME, Eickhoff SB, Hakonarson H, Gur RC, Gur RE, Wolf DH (2013): An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *NeuroImage* 64:240–256.
- Shah AJ, Lampert R, Goldberg J, Veledar E, Bremner JD, Vaccarino V (2013): Posttraumatic stress disorder and impaired autonomic modulation in male twins. *Biol Psychiatry* 73:1103–1110.
- Shaikh al arab A, Guedon-Moreau L, Ducrocq F, Molenda S, Duhem S, Salleron J, Chaudieu I, Bert D, Libersa C, Vaiva G (2012): Temporal analysis of heart rate variability as a predictor of post traumatic stress disorder in road traffic accidents survivors. *J Psychiatr Res* 46:790.
- Shin LM, Rauch SL, Pitman RK (2006): Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. *Ann N Y Acad Sci* 1071:67–79.
- Spreng RN, Mar RA, Kim AS (2009): The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: A quantitative meta-analysis. *J Cogn Neurosci* 21:489–510.
- Sripada RK, King AP, Welsh RC, Garfinkel SN, Wang X, Sripada CS, Liberzon I (2012): Neural dysregulation in posttraumatic stress disorder: Evidence for disrupted equilibrium between salience and default mode brain networks. *Psychosom Med* 74:904–911.
- Steiger F, Nees F, Wicking M, Lang S, Flor H (2015) Behavioral and central correlates of contextual fear learning and contextual modulation of cued fear in posttraumatic stress disorder. *Int J Psychophysiol* 98:584–593.
- Stein PK, Carney RM, Freedland KE, Skala JA, Jaffe AS, Kleiger RE, Rottman JN (2000): Severe depression is associated with markedly reduced heart rate variability in patients with stable coronary heart disease. *J Psychosom Res* 48:493–500.
- Stoodley CJ, Schmahmann JD (2010): Evidence for topographic organization in the cerebellum of motor control versus cognitive and affective processing. *Cortex* 46:831–844.
- Stoodley CJ, Valera EM, Schmahmann JD (2012): Functional topography of the cerebellum for motor and cognitive tasks: An fMRI study. *NeuroImage* 59:1560–1570.
- Tarvainen MP, Niskanen JP, Lipponen JA, Ranta-Aho PO, Karjalainen PA (2014): Kubios HRV—heart rate variability analysis software. *Comput Methods Programs Biomed* 113:210–220.
- Taylor TF (2015): The influence of shame on posttrauma disorders: Have we failed to see the obvious? *Eur J Psychotraumatol* 6:28847.

-
- Thayer JF (2006): On the importance of inhibition: Central and peripheral manifestations of nonlinear inhibitory processes in neural systems. *Dose Response* 4:2–21.
- Thayer JF, Brosschot JF (2005): Psychosomatics and psychopathology: Looking up and down from the brain. *Psychoneuroendocrinology* 30:1050–1058.
- Thayer JF, Lane RD (2000): A model of neurovisceral integration in emotion regulation and dysregulation. *J Affect Disord* 61: 201–216.
- Thayer JF, Hansen AL, Saus-Rose E, Johnsen BH (2009): Heart rate variability, prefrontal neural function, and cognitive performance: The neurovisceral integration perspective on self-regulation, adaptation, and health. *Ann Behav Med* 37:141–153.
- Thayer JF, Ahs F, Fredrikson M, Sollers JJ III, Wager TD (2012): A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev* 36:747–756.
- Thome J, Frewen P, Daniels JK, Densmore M, Lanius RA (2014): Altered connectivity within the salience network during direct eye gaze in PTSD. *Borderline Personality Disord Emot Dysregul* 1:17.
- Tulloch H, Greenman PS, Tasse V (2014): Post-traumatic stress disorder among cardiac patients: Prevalence, risk factors, and considerations for assessment and treatment. *Behav Sci* 5: 27–40.
- Van Dijk KR, Sabuncu MR, Buckner RL (2012): The influence of head motion on intrinsic functional connectivity MRI. *NeuroImage* 59:431–438.
- van Wingen GA, Geuze E, Vermetten E, Fernandez G (2011): Perceived threat predicts the neural sequelae of combat stress. *Mol Psychiatry* 16:664–671.
- Wahbeh H, Oken BS (2013): Peak high-frequency HRV and peak alpha frequency higher in PTSD. *Appl Psychophysiol Biofeedback* 38:57–69.
- Wendt J, Neubert J, Koenig J, Thayer JF, Hamm AO (2015): Resting heart rate variability is associated with inhibition of conditioned fear. *Psychophysiology* 52:1161–1166.
- White SF, Costanzo ME, Blair JR, Roy MJ (2015): PTSD symptom severity is associated with increased recruitment of top-down attentional control in a trauma-exposed sample. *NeuroImage* 7: 19–27.
- Wolf EJ, Rasmusson AM, Mitchell KS, Logue MW, Baldwin CT, Miller MW (2014): A genome-wide association study of clinical symptoms of dissociation in a trauma-exposed sample. *Depress Anxiety* 31:352–360.
-