Evidence That Smooth Pursuit Velocity, Not Eye Position, Modulates Alpha and Beta Oscillations in Human Middle Temporal Cortex

Benjamin T. Dunkley,^{1,2*} Tom C.A. Freeman,²
Suresh D. Muthukumaraswamy,³ and Krish D. Singh²

¹Department of Diagnostic Imaging, the Hospital for Sick Children, Toronto, Canada

²School of Psychology, CUBRIC (Cardiff University Brain Research Imaging Centre),
Cardiff University, Park Place, Cardiff, United Kingdom

³Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand

Abstract: Suppression of 5–25 Hz oscillations have been observed in MT+ during pursuit eye movements, suggesting oscillations that play a role in oculomotor control and/or the integration of extraretinal signals during pursuit. The amplitude of these rhythms appears to covary with head-centered eye position, but an alternative is that they depend on a velocity signal that lags the movement of the eyes. To investigate, we explored how alpha and beta amplitude changes related to ongoing eye movement depended on pursuit at different eccentricities. The results revealed largely identical patterns of modulation in the alpha and beta amplitude, irrespective of the eccentricity at which the pursuit eye movement was performed. The signals we measured therefore do not depend on head-centered position. A second experiment was designed to investigate whether the alpha and beta oscillations depended on the direction of pursuit, as opposed to just speed. We found no evidence that alpha or beta oscillations depended on direction, but there was a significant effect of eye speed on the magnitude of the beta suppression. This suggests distinct functional roles for alpha and beta suppression in pursuit behavior. *Hum Brain Mapp* 36:5220–5232, 2015. © 2015 Wiley Periodicals, Inc.

Key words: smooth pursuit; magnetoencephalography; extra-retinal signals; MT+/MST; neuronal oscillations

INTRODUCTION

Cortical oscillations are a pervasive phenomenon in neuroscience and are thought to subserve a number of cognitive, sensory, and motor processes. Oscillatory changes in human middle temporal cortex (MT+) have been shown to

accompany smooth eye pursuit, but their functional role is still not completely understood. In medial superior temporal cortex (MST), a majority of neurons that display a pursuit-related response (i.e., those that do not show a response to retinal motion) have been shown to code for eye velocity, whilst only a small minority display have any sensitivity to head-centered eye position [Ilg and Thier, 2003]. We previously reported [Dunkley et al., 2013] that suppression of the 5–25 Hz response in MT+, a correlate of increased cortical activity, covaried with eye position, but other studies have shown that eye speed was the dominant signal in animal [Churchland and Lisberger, 2005] and human studies [Nagel et al., 2006, 2008]. However, given the paradigm we used, it was not possible to explain this discrepancy because eye position and speed covaried.

*Correspondence to: Ben Dunkley, Department of Diagnostic Imaging, The Hospital for Sick Children, Toronto, Canada. E-mail: ben.dunkley@sickkids.ca

Received for publication 24 June 2015; Revised 11 September 2015; Accepted 16 September 2015.

DOI: 10.1002/hbm.23006

Published online 29 September 2015 in Wiley Online Library (wileyonlinelibrary.com).

The first experiment presented here was therefore designed to tease these two stimulus parameters apart by presenting pursuit targets with identical velocity profiles at different mean eccentricities (center, left, or right). If oscillatory activity coded primarily for eye position during ongoing pursuit, we might expect to see distinct amplitude envelope profiles for alpha and beta activity dependent on the eccentricity of the pursuit cycle. For example, when engaging in a pursuit cycle in the contralateral hemifield with respect to the recording site, relatively *greater* suppression (decrease in amplitude) of the activity in MT+ might be expected; and equally, *less* suppression might be expected when pursuing a target in the ipsilateral visual hemifield. Conversely, if the changes in the amplitude envelopes do not depend on eye position, then these macroscopic signals might reflect some other aspect of the pursuit movement, such as speed, that is common to all eccentricities tested.

To our knowledge, there have been only a handful of neuroimaging articles on how pursuit velocity modulates activity in MT+. For example, Nagel et al. [2008] reported an increasing cortical activation [measured via the blood oxygen level-dependent (BOLD) effect] observed for increasing stimulus velocity [Nagel et al., 2006, 2008]. However, they in fact demonstrated *speed*-related activity because BOLD signals were collapsed across the direction of pursuit.

Extraretinal signals in MST characterized by activity in neurons with a preference for the ipsilateral pursuit are thought to make up the majority of neurons in MST [Ilg and Thier, 2003; Ono and Mustari, 2006]. We therefore performed a second experiment to test whether oscillatory changes in human MT+ depend on relative direction or just speed, by examining potential biases in oscillatory amplitude for ipsi- or contra-verse pursuit. Furthermore, we examined whether there is any evidence of a directional bias in alpha and beta for different pursuit directions, which might reveal a specific mechanistic role for these rhythms.

Given the link between spike-rate, the BOLD effect and low-frequency suppression [Mukamel et al., 2005], and animal data that show the existence of pursuit-sensitive velocity-dependent neurons in MT+ [Ono and Mustari, 2006], we hypothesized that alpha and beta suppression in

unilateral MT+ would increase with increasing ipsiversive pursuit velocity.

METHODS

Experiment 1

Participants

Eighteen healthy participants (13 females, mean age = 23 years) completed Experiment 1. All had given prior informed consent, had normal or correct-to-normal visual acuity and no history of neurological disorders. All experimental procedures were approved by the Cardiff University School of Psychology Ethics Committee.

Design and procedure

Figure 1 shows the experimental protocol for Experiment 1, which consisted of three conditions. In the first condition, "central pursuit," participants were required to make sinusoidal horizontal pursuit eye movements by tracking a faint dot in the central portion of the screen (0.5 Hz frequency, amplitude $\pm 5^\circ$). In the second "right pursuit" condition, participants made horizontal pursuit eye movements that took place completely in the right hemifield; that is, in a head-centered reference frame, starting at 7.5° eccentricity, with an amplitude of $\pm 5^\circ$ and a frequency of 0.5 Hz (otherwise identical to condition 1, "central pursuit"). In the third condition, "left pursuit" participants engaged in sinusoidal pursuit with amplitude of $\pm 5^\circ$ and a frequency of 0.5 Hz, except this time the pursuit cycle starts at -7.5° eccentricity.

The experiment comprised of a boxcar design, with 15×60 s epochs consisting of a 10 s central fixation baseline period immediately followed by the 10 s "central pursuit" condition (active period), then a 10 s right fixation period, 10 s "right pursuit" condition, 10 s left field fixation, and finally 10 s of "left pursuit." Each run lasted for about 15 min.

Apparatus and data acquisition

Visual stimuli were generated on a GeForce graphics card (Nvidia Corp.) and back-projected (Sanyo XP41 LCD) onto a screen at 60 Hz (size $34 \text{ cm} \times 24.7 \text{ cm}$, total visual angle $25.6^\circ \times 19.2^\circ$, resolution of $1,024 \times 768$) at a distance of 71 cm. The visual stimuli and experimental protocol were programmed in Pascal (Delphi 7, Borland Software Corporation) using the OpenGL software library for graphics hardware.

Magnetoencephalography (MEG) data was recorded using a 275 channel whole-head system (CTF Systems Inc., a subsidiary of VSM MedTech Ltd.) in a magnetically shielded room at a sample rate of 1.2 kHz using a radial gradiometer configuration, with the primary sensors analyzed as synthetic third order gradiometers. During data

Abbreviations

BOLD	blood oxygen level dependent;
EEG	electroencephalography;
EOG	electrooculography;
MEG	magnetoencephalography;
MRI	magnetic resonance imaging;
MT/MT+	middle temporal cortex;
MST	medial superior temporal cortex;
ROI	region-of-interest;
SAM	synthetic aperture magnetometry;
SPM	statistical parametric map

Figure 1.

Experimental protocol for the excursive pursuit paradigm (Experiment I). Sinusoidal pursuit eye movements were performed at three horizontal eccentricities (central, right, and left hemifield) in the sequence shown.

acquisition, head position and motion were monitored using three fiducial markers placed on the nasion and 1 cm anteriorly from both the left and right tragi. Each participant's MEG data was then co-registered with their anatomical data, based on the position of these easily identifiable anatomical landmarks from the MR scan. These points were verified using high-resolution digital photographs taken during fiducial placement.

Electrooculographic (EOG) recordings were taken at the same time as MEG data collection, which allowed us to characterize eye movements while observers wore polarizing goggles designed to eliminate any retinal illumination other than from the faintly visible pursuit target. Pairs of electrodes were placed above and below the eye to record vertical displacement, and to the lateral corner of each eye to record horizontal displacement. The impedance of each electrode was then measured to ensure satisfactory conductance of the EOG signal. Skin preparation was performed again and the electrodes were re-applied if the electrical resistance was found to be above a pre-defined threshold of 5 k Ω .

Data analysis

For localization of the oscillatory response, a multiple, local-spheres forward-model was derived by fitting overlapping spheres [Huang et al., 1999] to the brain surface extracted by the Brain Extraction Tool (BET) [Smith, 2002].

Source analysis was performed using Synthetic Aperture Magnetometry (SAM), a non-linear beamformer spatial filter that generates statistical parametric maps of oscillatory power changes [Pfurtscheller and Silva, 1999] between baseline/passive (fixation) and active (stimulus tracking/retinal motion) periods within specific frequency bands.

SAM images were constructed on a 5 mm $\times$ 5 mm $\times$ 5 mm grid throughout the brain for each condition and each participant. Oscillatory changes between the passive and active periods for all 30 epochs were mapped for alpha (5–15 Hz), beta (15–25 Hz), and gamma (30–80 Hz) frequency ranges. Peak coordinates for activation in the specified frequency band in the SAM images were visualized using mri3dX [Singh, 2009] and chosen on the basis of their locality within extra-striate cortex, for peaks with a predefined threshold of pseudo *t*-values greater than 0.5 and with an exclusion radius of 5 mm. These coordinates were used to compute suitable weights for virtual electrode generation on an individual basis and used in all subsequent analyses (performed using custom-scripts in Matlab). Virtual sensor weights were estimated using a covariance matrix filtered between 15 and 25 Hz (hence, modulations seen in this spectrogram can only be guaranteed to be coming from the desired location for this frequency range). Computation of these time-frequency spectrograms were performed at peak voxel locations based on the Hilbert transform, in 0.5 Hz steps. The percentage change in oscillatory amplitude during the active period was baselined against the 0–10 s passive period.

Figure 2.

Experimental protocol for the pursuit velocity experiment (Experiment 2). Participants fixate a central stimulus for 3 s (baseline condition), followed by either a fixation to the left or right at -12.5° or 12.5° respectively. Active tracking then followed where participants pursue the target stimulus at one of three speeds (5, 10, or $15^\circ/\text{s}$) in either direction, given six total velocities (three levels per factor).

Experiment 2

Participants

Twenty participants with normal or correct-to-normal visual acuity completed Experiment 2 (12 females, mean age = 22.3 years). All participants had given prior informed consent and no history of neurological disorders. All experimental procedures were approved by the Cardiff University School of Psychology Ethics Committee.

Design and procedure

Figure 2 depicts the protocol for Experiment 2. Participants begin by fixating a central point (baseline period, 3 s). This was followed by the appearance of a target in the right hemifield at 12.5° eccentricity, which observers were instructed to make an immediate eye movement toward and then fixate. After 3 s, this target stimulus then moved leftwards at either -5 , -10 , or $-15^\circ/\text{s}$ for 25° of visual angle. This period is then immediately followed by another central fixation (3 s), which in turn is followed by a left field fixation (amplitude -12.5° , gaze held for 3 s). A period of rightward smooth pursuit at 5, 10, or $15^\circ/\text{s}$ for 25° then follows. Thus, there were six conditions in total, each comprising of pursuit along the horizontal meridian, with 15 trials of each velocity. Condition presentation was randomized and counter-balanced within the whole run.

Data analysis

SAM images were constructed for each velocity versus the central fixation baseline period. The percentage change in the amplitude for pursuit was referenced to the baseline fixation period. Alpha and beta amplitude changes relative to baseline were sampled over a 1 s time window during the central portion of the pursuit phase for use in later analyses. By sampling amplitude changes partway through the pursuit phase, smooth pursuit gain was likely to be closer to 1, thus reducing the likelihood that any measured changes in alpha and beta amplitude would be related to significant retinal slip or catch-up saccades. To further ensure this was the case, any trials that were found to be severely contaminated with saccadic eye movements, or blinks, were removed from any further imaging analysis, resulting in an average trial exclusion rate of 13.3% per participant per session.

RESULTS

Experiment 1

Eye movement data

Figure 3 shows the group mean pursuit gain. Visual inspection of the EOG traces revealed that observers were able to track the moving stimulus reasonably accurately. This was confirmed by the group mean pursuit gains

Figure 3.

Group-averaged pursuit gains values for the three eccentricities examined. Error bars are ± 1 SE throughout.

shown in Figure 3. Crucially, a repeated measures ANOVA revealed no significant difference between the mean gains found in the three pursuit conditions ($F(2,17) = 1.722$, $P > 0.05$).

SAM source analysis

Figure 4a shows the group mean SAM analysis for 15–25 Hz activity. This revealed bilateral extra-striate suppression of the beta response for the “central pursuit” versus baseline central fixation condition in all participants. As this frequency band displayed the most consistent peaks localized to putative MT+ [similar to Dunkley et al., 2013], these coordinates were used to compute the weights required for the virtual sensors (over a 0.1–80 Hz range), and the subsequent time–frequency analysis using the Hilbert transfer (in 0.5 Hz steps). Additionally, areas of the posterior parietal cortex (PPC), frontal eye fields (FEF), and cuneus showed evidence of beta rhythm amplitude decreases. The group average beta response with a peak in left association cortex is indicative of the individual source localization results. Furthermore, alpha changes for central pursuit eye movements were, for the most part, spatially coincident with the beta response shown above, that is, located in association cortex consistent with MT+, PPC, FEF, and cuneus. However, the beamformer reconstruction of the beta rhythm proved to be more reliable in its estimation of MT+. Finally, there were no gamma (30–70 Hz) responses evident in any individual or Group SAM images.

Time–frequency analysis

Figure 4b shows the group-averaged spectrogram from the virtual electrode in left MT+ during eye movement for all three pursuit eccentricities. Trials began with the cen-

tral fixation baseline window, which was then followed by the “central pursuit” condition, right fixation, “right pursuit” condition, left fixation, and finally “left pursuit.” The time–frequency analysis from left MT+ revealed sustained, low frequency (5–25 Hz) amplitude decreases for the duration of tracking for all three conditions. A similar response profile alpha–beta changes was also observed in right MT+, with sustained suppression for the duration of all three conditions. There is no evidence in the spectrograms of any discernable difference in the alpha or beta response as a function of pursuit cycle eccentricity.

Additionally, there appears to be demarcation in the oscillatory amplitude changes of the 5–25 Hz frequency band (denoted by the white bounded box in the spectrograms of Fig. 4b), showing sustained amplitude decreases in two distinct frequency bands during pursuit eye movements in MT+. One of these is in the range of the alpha band, and the other in the higher beta band. The pattern of these responses is similar to those found in our previous article [Dunkley et al., 2013]. This effect appeared more prominent in the left MT+ voxel (right MT+ time course not shown for brevity).

To assess the spectral characteristics of these pursuit-related responses, amplitude versus frequency plots were computed from the averaged, active time-windows, and all participants (Fig. 4c). These revealed two peaks; the largest peak being around the 5–15 Hz band, and a second, smaller, peak at around 18–22 Hz (approximately beta frequency). On this basis, the broadband 5–25 Hz signal was decomposed into two smaller frequencies for further analysis. These 5–15 and 15–25 Hz amplitude envelopes from MT+ were examined for differences in the modulation of this activity at the three pursuit eccentricities. This was then compared across conditions during the sinusoidal pursuit cycle, discussed in the next section.

Oscillatory amplitude envelope analysis

Figure 5 shows the group-averaged amplitude envelopes in the alpha (*top panels*) and beta bands (*bottom panels*) relative to the baseline central fixation condition, in both left and right MT+ during a pursuit cycle. For reference, the dashed line in each figure shows the phase of the relative eye position during each cycle (the eye position units are arbitrary). Different colors denote the three different eccentricities (center, left and right). For both alpha and beta responses, amplitude appears to be independent of eccentricity throughout the pursuit cycle. The modulation of low-frequency oscillations in MT+ shows no evidence of ipsi- or contralateral eye position-dependence (relative to the MT+ recording site).

There does appear to be a consistent modulation of the alpha and beta amplitude envelope within the pursuit cycle. In the left MT+ recording site, alpha and beta seems to covary with eye position in the contraversive phase of the pursuit cycle. Hence, there is a salient trough in the first half of the pursuit cycle, when the eyes are moving

Figure 4.

(a) Group-averaged SAM images for beta overlaid on a template brain. Cold-color map indicates task-induced amplitude decreases, with the greatest changes localized to bilateral extra-striate regions. (b) Group averaged spectrogram for left MT+. Low-frequency suppression (5–25 Hz decreases, delineated by the white-bounded box) was observed for all three pursuit conditions as denoted by the cold-color map (indicating percentage

change from baseline). (c) Group-mean spectral power from left and right MT+ during pursuit phase. Normalized absolute amplitude changes in frequency show a peak at around 5–15 Hz, and a second, smaller peak at around 18–22 Hz, consistent with the alpha and beta bands, respectively. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

away from the left recording site, but this then flattens in the second half of the cycle. For right MT+ however, the shape resembles more of a symmetrical “W” shape, with two minima in the alpha and beta amplitude corresponding to the maximum amplitude of the eyes during the pursuit cycle. Therefore, it is possible these amplitude envelope profiles could reflect a process that reflects either eye speed or direction, which was the motivation of Experiment 2.

Experiment 2

Eye movement data

Figure 6a shows the group-averaged horizontal eye position trace when tracking the stimulus in both the left- and rightward directions and the mean pursuit gain for each condition (note the eye movements were analyzed over the middle 1-s window during tracking of the

Figure 5.

Group-averaged amplitude envelopes for alpha (top row) and beta (bottom row). A representative sinusoidal pursuit cycle showing the relative phase of the ongoing eye movement is denoted by the dashed gray line (arbitrary units). The “central pursuit” condition is depicted by the red trace, “right pursuit”

condition in green, and the “left pursuit” condition in blue. Shaded regions define ± 1 SE. The percentage amplitude changes shown are all relative to the baseline central fixation period. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

stimulus). Participants with mean gain values not falling within the predefined range of 0.8–1.2 across all conditions were excluded from further behavioral and imaging analysis, in order to minimize the amount of retinal slip affecting subsequent analyses. This resulted in the exclusion of

one participant, whose average gain value was 0.78, leaving a cohort of 19 participants.

One-way ANOVA on eye velocity gains for leftward and rightward pursuit direction revealed no significant difference in velocity for leftward ($F(2,36) = 0.30$, $P > 0.05$)

Figure 6.

(a) Group-averaged horizontal eye position traces for pursuit in both the leftward (left panel) and rightward (right panel) direction. (b) Group mean pursuit gain for leftward (left panel) and rightward (right panel) eye movements conditions. Gain was calculated over the central 1 s window midway through the pursuit eye movement. (c)

Group mean pursuit gain, collapsed across all velocity conditions, for both the left and rightward direction. Error bars are ± 1 SE throughout. (* $P < 0.05$). [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

or rightward pursuit ($F(2,36) = 0.20$, $P > 0.05$). However, when mean pursuit gain was collapsed across conditions to give a mean measure of pursuit gain based on direc-

tionality, there was a significant if small difference in the mean eye velocity gain for left- versus rightward pursuit (paired t -test, $t(56) = -2.62$, $P < 0.05$).

Figure 7.

Single-subject SAM analysis for beta during leftward pursuit at $-10^\circ/\text{s}$ and group-averaged spectrograms for right MT+. (a) SAM analysis revealed beta decreases over right extra-striate cortex, consistent with MT+, along with a second peak identified in left MT+. Additionally, there were peaks identified in FEF, SEF, the cerebellum, and the PPC. (b) Grand-averaged spectrograms from right MT+ during leftward pursuit. The baseline period (central fixation: CF) was followed by right field fixation

(RF) at 12.5° from -3 to 0 s. Subsequently, there was leftward pursuit (LP) at either -5 , -10 , -15 , or $-20^\circ/\text{s}$, terminating at -12.5° eccentricity. White box denotes the time-period over which the average amplitude change was quantified and subsequently used when examining speed modulation of alpha and beta in MT+. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

SAM source reconstruction

Figure 7a shows a representative single-subject SAM analysis for beta during stimulus tracking at $-10^\circ/\text{s}$, versus the central fixation baseline period. This revealed bilateral 15–25 Hz suppression over extra-striate cortices, consistent with MT+, in addition to peaks over the PPC, FEF and supplementary eye fields (SEF) regions. Similar patterns of activity to this were observed for this subject in the alpha band for all other pursuit velocities in both the left- and rightward directions. No gamma peaks were identified in any of the participants in the MT+ region. The beta response observed during pursuit, as in previous experiments, was shown to be the most consistent in the SAM source estimation for localizing MT+ for the majority of subjects. Therefore, the location of this beta band peak response was used to define the coordinates for the virtual sensor reconstruction at 0.1–80 Hz, and subsequent time-frequency analysis using the Hilbert transform.

Time-frequency analysis

Grand-averaged time-frequency plots of the oscillatory changes in right MT+ for leftwards pursuit can be seen in Figure 7b. Both the left- and rightward direction induced sustained low-frequency (5–25 Hz) suppression for the duration of object tracking starting 0 s onward. Interestingly, there was also consistent suppression of the 5–25 Hz response in MT+ when the eyes are fixated and stationary in the right visual hemifield following saccadic eye movements to the eccentric fixation point.

In the central fixation baseline period (the time window -6 to -3 s denoted by CF), there is 1–2 s of low-frequency suppression preceding termination of pursuit and the subsequent saccade to the central fixation point. This appears prominent in the alpha band, with a greater decrease in amplitude and often sustained for longer than the suppression in beta. Following this, alpha and beta amplitude returns to baseline levels after 1–2 s.

Following the time–frequency analysis, the mean alpha and beta activity in the central 1 s time window of the pursuit phase was extracted from each individual’s spectral data and used to examine pursuit velocity modulation activity from this region (denoted by the white box in the spectrograms).

Effect of eye velocity on low-frequency suppression

Figure 8a depicts the mean alpha response for each stimulus velocity in left (top row of panels) and right (bottom row) MT+. A One-way Repeated Measures ANOVA was run for each hemisphere and pursuit direction separately. Results indicated there was no significant effect of stimulus velocity on alpha (all ANOVA tests $P > 0.05$).

Figure 8b shows the mean beta amplitude for each of the stimulus velocity conditions, in both the leftward and rightward direction, in left and right MT+. For beta amplitude change from the left hemisphere during leftward pursuit (top left panel), there was no significant difference in the mean induced beta response ($P > 0.05$). There was a significant effect of stimulus velocity on the beta amplitude measured from the left hemisphere during rightward pursuit (top right panel, $F(2,32) = 4.39$, $P < 0.05$), as well as a significant effect of stimulus velocity on beta for leftward pursuit in the right hemisphere (bottom left panel, $F(2,32) = 6.74$, $P < 0.01$). Finally, for beta from right MT+ during rightward pursuit (bottom right panel), there was a significant effect of stimulus velocity on the magnitude of the beta amplitude change in MT+ ($F(2,31) = 3.862$, $P < 0.05$).

In summary, although we did not find *direction*-specific changes in the response as originally predicted, there does however appear to be a trend for increased beta suppression with increasing eye *speed*, irrespective of the direction of the pursuit eye movement. This would suggest that beta amplitude decreases might be linked to activity in these regions that is related to the processing of oculomotor and/or extraretinal eye movement signals coding for eye speed. A further set of analyses was therefore performed to compare alpha and beta activity with changes in eye speed. This was achieved by simply reversing the sign of the leftward stimulus velocity and collapsing the left- and rightward data together, resulting in four tests (for both alpha and beta responses, and each hemisphere). The results of this analysis are discussed in the next section.

Effect of eye speed on low-frequency suppression

Figure 9 shows the mean alpha (top panels) and the beta (bottom panels) activity for each eye speed condition, from both the left and right MT+. Repeated-measured ANOVA for the alpha found there was no significant effect of eye speed on amplitude in either the left or right hemispheres (top panels, $P > 0.05$).

There was a significant effect of eye speed on beta in the left hemisphere (bottom left panel, $F(2,65) = 4.73$, $P = 0.01$). Post hoc comparisons found beta suppression in the $15^\circ/\text{s}$ ($M = -10.15$, $SD = 7.95$) condition was significantly greater than the $5^\circ/\text{s}$ ($M = -14.60$, $SD = 7.61$) condition ($P = 0.015$).

For the right hemisphere MT+ beta, there was a significant effect of eye speed on the amplitude ($F(2,65) = 10.65$, $P < 0.001$), with post hoc comparisons indicating a significant difference between the $5^\circ/\text{s}$ ($M = -11.84$, $SD = 8.69$) and $15^\circ/\text{s}$ ($M = -19.29$, $SD = 12.87$) conditions ($P < 0.001$). Additionally, there was also a significant difference between the $10^\circ/\text{s}$ ($M = -14.27$, $SD = 9.57$) and $15^\circ/\text{s}$ ($M = -19.29$, $SD = 12.87$) conditions ($P = 0.012$), but not for the other conditions ($P > 0.05$). Therefore, this might suggest some functional distinction in the role of alpha and beta suppression in MT+ during pursuit.

DISCUSSION

Two experiments were carried out to further explore the link between oscillatory suppression in MT+ and smooth pursuit eye movement [Dunkley et al., 2013]. In Experiment 1, we investigated whether the modulation reflects head-centered eye position or velocity. In Experiment 2, we tested whether modulation was related to the speed or direction of the eye movement.

The results of Experiment 1 showed that the alpha and beta amplitude envelopes measured during pursuit showed no differential response related to eye position. Activity was virtually identical across the three eccentricities tested, suggesting that responses from MT+ during eye movements reflect the activity of neurons that code for some other aspect of pursuit, such as eye speed. In Experiment 2, we found little connection between alpha and beta activity, and the direction of pursuit (although there was a trend for slightly greater suppression for contraversive eye movements). However, we did find a significant effect of *eye speed* on beta suppression, whereas there was no similar effect on alpha. This suggests that beta suppression represents a distinct mechanism related to pursuit behavior, reflecting some fundamental aspect of the eye movement, while alpha suppression does not. We speculate that the latter is more likely to reflect the hypothesized “inhibitory gating” mechanism [Handel et al., 2011; Jensen and Mazaheri, 2010], or perhaps superordinate functions such as visuo-spatial attention that have been implicated in intentional pursuit eye movements [Lovejoy et al., 2009; Ono and Mustari, 2006].

The beta suppression we found could be related to neuronal activity involved in oculomotor control [Ilg, 1997; Krauzlis, 2004] or it could be related to the integration of extraretinal information in the perception of motion during pursuit [Freeman et al., 2010; Furman and Gur, 2012; Spering and Montagnini, 2011]. It is not possible to draw a definitive conclusion from this experiment on either of

Figure 8.

(a) Mean alpha responses from both left and right MT+ for each of the speed conditions in the leftward and rightward pursuit directions. Negative amplitude values indicate a decrease with respect to baseline. (b) Mean beta responses. Error bars are $\pm$ SE throughout. (* $P < 0.05$, ** $P < 0.01$).

Figure 9.

Mean magnitude of alpha and beta suppression in both the left and right hemispheres at various pursuit speeds. Error bars are $\pm$ SE throughout. (* $P < 0.05$, ** $P < 0.01$).

these issues, as this task was performed in the presence of visual feedback in the form of the retinal slip that accompanies closed-loop pursuit. Given this, it is not possible to isolate purely “extraretinal” components of a reference signal without looking at pursuit activity in the absence of any visual input (thereby fully isolating nonretinal mechanisms). Additionally, without an objective measure of perceived motion during pursuit, it is also not possible to conclude whether alpha and beta oscillations are used in this process either. However, this experiment does suggest a fundamental role of beta suppression in MT+ during ongoing pursuit, whether it be in the generation of oculomotor signals or the integration of an internally generated reference signal that codes eye speed estimates.

It has been shown that alpha and beta decreases in striate and association cortices might reflect the engagement of visuo-spatial attention and anticipatory processing of any predictable sensory change and incoming information [Worden et al., 2000; Yamagishia et al., 2008]. In other words, alpha and beta amplitude decreases occur over the area of cortex known to be active when covert visuo-

spatial attention mechanisms are engaged, and this effect is enhanced in this region of the brain by anticipatory signals awaiting an impending sensory change). It is possible the sustained alpha and beta suppression witnessed during gaze fixation reflects an “inhibitory filter mechanism” [Jensen and Mazaheri, 2010] that helps focus attention (or perhaps attention itself modulates this change; causality cannot be implied in this case) and prevents distracting stimuli from interfering with task-related processing, given that the onset of the motion stimulus can be anticipated. Therefore, it seems likely that the oscillatory response observed in MT+ during excursive gaze fixation could represent either the allocation of selective visuo-spatial attention or anticipatory priming of the brain for stimulus motion onset, the macroscopic signature of neuronal activity reflecting eye position sensitivity, or a combination of both these processes.

In Experiment 2, this was examined by utilizing a paradigm similar to that used in other studies [Ilg and Thier, 2003; Nagel et al., 2008], to test whether the neuronal response characteristics, defined as oscillatory amplitude

changes, reflect a velocity-based signal. Additionally, the data was examined further to see whether a response for a preferred direction could be extracted from the macroscopic signature, given that animal research suggests the majority of neurons in MST that display a pursuit-related response code for ipsiversive pursuit [Ilg and Thier, 2003].

Consistent beta suppression was found over extra-striate cortices for all stimulus conditions and a significant effect of pursuit eye speed on the magnitude of the beta decreases (increased suppression with increasing eye speed). Furthermore, no effect of eye speed on alpha in MT+ was found. This suggests there might be some functionally distinct role that these frequency bands play in the maintenance of pursuit and/or the integration of extra-retinal eye movement signals in this area.

The findings presented here do not conform to our original predictions regarding a speed-sensitive bias in alpha and beta suppression for ipsiversive pursuit directionality (relative to the recording site). Given these results, we speculate that the relationship between beta suppression and eye speed demonstrates functionally relevant role with regards to smooth pursuit. Specifically, it appears that the brain may engage in active beta band suppression in MT+ during pursuit, which could either reflect the generation of oculomotor signals for pursuit maintenance in the presence of ongoing visual feedback and/or the use of extraretinal signals for perceived motion during pursuit. The lack of any clear relationship between alpha activity and the eye movement also suggests that suppression mediates, or is perhaps in turn mediated by, some other sensory or cognitive process associated with pursuit.

REFERENCES

- Churchland AK, Lisberger SG (2005): Relationship between extra-retinal component of firing rate and eye speed in area MST of macaque monkeys. *J Neurophysiol* 94:2416–2426.
- Dunkley BT, Freeman TCA, Muthukumaraswamy SD, Singh KD (2013): Cortical oscillatory changes in human middle temporal cortex underlying smooth pursuit eye movements. *Hum Brain Mapp* 34:837–851.
- Freeman TCA, Champion RA, Warren PA (2010): A Bayesian model of perceived head-centered velocity during smooth pursuit eye movement. *Curr Biol* 20:757–762.
- Furman M, Gur M (2012): And yet it moves: Perceptual illusions and neural mechanisms of pursuit compensation during smooth pursuit eye movements. *Neuroscience and Biobehavioral Reviews* 36:143–151.
- Handel BF, Haarmeier T, Jensen O (2011): Alpha oscillations correlate with the successful inhibition of unattended stimuli. *J Cogn Neurosci* 23:2494–2502.
- Huang MX, Mosher JC, Leahy RM (1999): A sensor-weighted overlapping-sphere head model and exhaustive head model comparison for MEG. *Phys Med Biol* 44:423–440.
- Ilg UJ (1997): Slow eye movements. *Progress in Neurobiology*, 53, 293–329.
- Ilg UJ, Thier P (2003): Visual tracking neurons in primate area MST are activated by smooth-pursuit eye movements of an “imaginary” target. *J Neurophysiol* 90:1489–1102.
- Jensen O, Mazaheri A (2010): Shaping functional architecture by oscillatory alpha activity: Gating by inhibition. *Front Hum Neurosci* 4:1–8.
- Krauzlis RJ (2004): Recasting the smooth pursuit eye movement system. *J Neurophysiol* 91:591–603.
- Lovejoy LP, Fowler GA, Krauzlis RJ (2009): Spatial allocation of attention during smooth pursuit eye movements. *Vision Res* 49:1275–1285.
- Mukamel R, Gelbard H, Arieli A, Hasson U, Fried I, Malach R (2005): Coupling between neuronal firing, field potentials, and fMRI in human auditory cortex. *Science* 309:951–954.
- Nagel M, Sprenger A, Zapf S, Erdmann C, Kömpf D, Heide W, Binkofski F, Lencer R (2006): Parametric modulation of cortical activation during smooth pursuit with and without target blanking. An fMRI study. *NeuroImage* 29:1319–1325.
- Nagel M, Sprenger A, Hohagen F, Binkofski F, Lencer R (2008): Cortical mechanisms of retinal and extraretinal smooth pursuit eye movements to different target velocities. *NeuroImage* 41: 483–492.
- Ono S, Mustari MJ (2006): Extraretinal signals in MSTd neurons related to volitional smooth pursuit. *J Neurophysiol* 96:2819–2825.
- Pfurtscheller G, Silva FHLd (1999): Event-related EEG/MEG synchronisation and desynchronisation: Basic principles. *Clin Neurophysiol* 110:1842–1857.
- Singh KD (2009): mri3dX. Available at: <http://cubic.psych.cf.ac.uk/Documentation/mri3dX/features>. [Accessed September 2011].
- Smith SM (2002): Fast robust automated brain extraction. *Hum Brain Mapp* 17:143–155.
- Spering M, Montagnini A (2011): Do we track what we see? Common versus independent processing for motion perception and smooth pursuit eye movements: A review. *Vision Res* 51:836–852.
- Worden MS, Foxe JJ, Wang N, Simpson GV (2000): Anticipatory biasing of visuospatial attention indexed by retinotopically specific α -band electroencephalography increases over occipital cortex. *J Neurosci* 20:RC63.
- Yamagishia N, Callana DE, Anderson SJ, Kawatob M (2008): Attentional changes in pre-stimulus oscillatory activity within early visual cortex are predictive of human visual performance. *Brain Res* 1197:115–122.