Adult Attachment Style is Associated With Cerebral μ -Opioid Receptor Availability in Humans

Lauri Nummenmaa,^{1,2,3*} Sandra Manninen,¹ Lauri Tuominen,^{1,2}
Jussi Hirvonen,¹ Kari K. Kalliokoski,¹ Pirjo Nuutila,^{1,4} Iiro P. Jääskeläinen,²
Riitta Hari,² Robin I.M. Dunbar,^{2,5} and Mikko Sams²

¹Turku PET Centre, University of Turku, Turku, 20520 Finland

²Department of Neuroscience and Biomedical Engineering, Aalto University School of Science, 00076 AALTO, Espoo, Finland

³Department of Psychology, University of Turku, 20014 Turku, Finland

⁴Department of Endocrinology, Turku University Hospital, 20521 Turku, Finland

⁵Department of Experimental Psychology, University of Oxford, Oxford, OX1 3UD, UK

Abstract: Human attachment behavior mediates establishment and maintenance of social relationships. Adult attachment characteristically varies on anxiety and avoidance dimensions, reflecting the tendencies to worry about the partner breaking the social bond (anxiety) and feeling uncomfortable about depending on others (avoidance). In primates and other mammals, the endogenous μ -opioid system is linked to long-term social bonding, but evidence of its role in human adult attachment remains more limited. We used in vivo positron emission tomography to reveal how variability in μ -opioid receptor (MOR) availability is associated with adult attachment in humans. We scanned 49 healthy subjects using a MOR-specific ligand [¹¹C]carfentanil and measured their attachment avoidance and anxiety with the Experiences in Close Relationships-Revised scale. The avoidance dimension of attachment correlated negatively with MOR availability in the thalamus and anterior cingulate cortex, as well as the frontal cortex, amygdala, and insula. No associations were observed between MOR availability and the anxiety dimension of attachment. Our results suggest that the endogenous opioid system may underlie interindividual differences in avoidant attachment style in human adults, and that differences in MOR availability are associated with the individuals' social relationships and psychosocial well-being. *Hum Brain Mapp* 36:3621–3628, 2015. © 2015 Wiley Periodicals, Inc.

Key words: attachment; emotion; opioids; neurotransmitters; social interaction; positron emission tomography

Contract grant sponsor: aivoAALTO Project of the Aalto University; Contract grant sponsor: Academy of Finland; Contract grant numbers: 265917 (to L.N.), 138145 (to I.P.J.), and 131483 (to R.H.); Contract grant sponsor: European Research Council (ERC); Contract grant numbers: 313000 (to L.N.), 232946 (to R.H.), and 295663 (to R.D.)

*Correspondence to: Lauri Nummenmaa; Department of Neuroscience and Biomedical Engineering, School of Science, Aalto Uni-

versity, FI 00076 AALTO, Espoo, Finland.

E-mail: lauri.nummenmaa@aalto.fi

Received for publication 22 April 2015; Revised 21 May 2015; Accepted 21 May 2015.

DOI: 10.1002/hbm.22866

Published online 5 June 2015 in Wiley Online Library (wileyonlinelibrary.com).

INTRODUCTION

Human adults have enduring affective relationships with each other in all areas of life, yet they differ markedly in the way they establish and maintain such social bonds. Attachment theory [Bowlby, 1982] posits an innate bonding system—presumably analogous with the attachment circuitry in other mammals—whose function is to maintain infants' proximity to significant others in the presence of threats and to regulate seeking for support when needed [Nelson and Panksepp, 1998]. In human adulthood, this attachment mechanism continues to modulate social interactions in close relationships [Hazan and Shaver, 1987].

Adult attachment behavior characteristically varies on two dimensions—anxiety and avoidance. These dimensions reflect the tendencies to worry about the partner breaking the social bond (anxiety) and feeling uncomfortable about depending on others (avoidance), with low scores on both dimensions reflecting attachment security [Brennan et al., 1998; Fraley et al., 2000]. Avoidantly attached individuals experience less pleasure from social interactions and avoid intimate interpersonal exchanges with potential of social rejection, thus leading to preference for being alone [Bartholomew, 1990]. Because large social network size [Holt-Lunstad et al., 2010] and the availability of social support [Broadhead et al., 1983] are associated with beneficial effects for somatic health and psychological well-being, it is not surprising that individual differences in adult attachment influence psychosocial and somatic well-being. Specifically, insecure attachment has been associated with psychological adjustment problems, substance abuse, and psychopathology [Brennan and Shaver, 1995; Griffin and Bartholomew, 1994; Mikulincer and Shaver, 2007].

Individual differences in adult attachment modulate activation of the corticolimbic circuits that encode social and affective information [see review in Vrticka and Vuilleumier, 2012]. At subjective level, avoidant attachment is associated with lowered experience of pleasantness when viewing positive social scenes, whereas anxious attachment is associated with heightened experience of arousal when viewing negative social scenes [Vrticka et al., 2012]. In line with these findings, haemodynamic responses in the striatal reward circuit to signals conveying positive emotional signals are negatively associated with attachment avoidance [Vrticka et al., 2008], whereas amygdala responses to negative emotional signals have been positively associated with both attachment avoidance [Norman et al., 2015] and anxiety [Vrticka et al., 2008]. Finally, the putative human social distress circuit (dorsal anterior cingulate cortex [dACC] and insula) is more responsive to social rejection in anxiously attached and less responsive in avoidantly attached individuals [DeWall et al., 2012]. In sum, attachment anxiety seems to increase responsiveness to negative social signals, whereas attachment avoidance lowers responsivity of brain circuits involved in socioemotional processing.

The neurochemical basis of human attachment is less well understood. Oxytocin and vasopressin systems contribute significantly to attachment behavior [Johnson and Young, 2015; Young et al., 2001], yet animal studies suggest that the endogenous opioid system could be another candidate neurochemical pathway for human attachment circuit. The opioidergic system modulates reward functions, drives motivated behaviors [Van Ree et al., 2000] and endogenous opiates widely modulate social behavior in rodents [Panksepp et al., 1980]. Exogenous opioid agonists alleviate separation distress in puppies [Panksepp et al., 1978], whereas μ -opioid receptor (MOR) gene knock-out mice pups display deficit maternal attachment behavior [Moles et al., 2004]. The A118G polymorphism of the MOR gene OPRM1 is associated with enhanced dispositional and neural sensitivity to social rejection in humans [Way et al., 2009], whereas rhesus infants carrying a gain-of-function OPRM1 77G allele experience increased reward from maternal contact [Barr et al., 2008].

Pharmacological manipulations in primates further show that opioid receptor antagonists increase the frequency of both social grooming [Fabre-Nys et al., 1982; Graves et al., 2002] and grooming solicitations [Keverne et al., 1989], suggesting that endogenous opioid tone modulates attachment behavior. Furthermore, activation of the endogenous MORs is essential for bonding in monogamous adult voles [Burkett et al., 2011; Resendez et al., 2013], suggesting the role of MOR in long-term adult social bonds. In line with this, genetic studies in human adults suggest that the minor allele (G) of the OPRM1 A118G polymorphism is associated self-reported avoidant attachment [Troisi et al., 2011]. Finally, human behavioral studies provide indirect evidence for the role of MOR system in bonding, by showing that pain threshold—a proxy of endogenous opioid release—is increased by social behavior promoting intragroup affiliation [Dunbar et al., 2012].

These converging lines of evidence suggest that the MOR-mediated neurotransmission could underlie individual differences in human adult attachment. However, this hypothesis currently lacks empirical support. Here, we use positron emission tomography (PET) coupled with behavioral attachment-style measures to show that avoidant but not anxious attachment style is negatively associated with cerebral MOR availability.

MATERIALS AND METHODS

Participants

The study protocol was approved by the ethics board of the South-western Finland hospital district, and the study was conducted in accordance with the Declaration of Helsinki. A priori power analysis based on effect sizes in previous PET studies on associations between personality trait variables and MOR availability suggested that sample

sizes exceeding $n = 45$ would have power higher than 0.95 for detecting statistically significant effects. Consequently, the final sample size was set at 50. Altogether, 50 healthy adults (20 females, ages 19–58, mean age 32 years, SD 6.4 years) volunteered for the study. Exclusion criteria (in addition to standard PET and MRI exclusion criteria) were poor compliance, smoking, excessive alcohol consumption (over 8 weekly doses), substance abuse determined by interview and blood tests, a history of or current neurological or psychiatric disease, and current medication affecting the central nervous system. All subjects were compensated for their time and travel costs, and they signed ethics-committee-approved, informed consent forms. One participant was removed from the sample because their MRI revealed previously nondiagnosed neurological disease.

Questionnaires

The participants completed the Experiences in Close Relationships-Revised (ECR-R) questionnaire addressing adult attachment style [Fraley et al., 2000]. The questionnaire consists of 36 Likert-scale questions and has been psychometrically validated to capture anxiety and avoidance dimensions of adult attachment style. As endogenous opioid systems is associated with both mood and anxiety related processes [Colasanti et al., 2011; Lutz and Kieffer, 2013], participants complete the Beck Depression Inventory II [Beck et al., 1988] and State-Trait-Anxiety Inventory [Spielberger et al., 1983] to rule out anxious and depressive symptoms and their potential association with MOR availability.

PET Imaging and Analysis

Data were acquired with the Philips Ingenuity PET-MR and GE Healthcare Discovery TM 690 PET/CT scanners in Turku PET Centre. After intravenous radioligand injection (targeted 250 MBq; mean 251 MBq, SD = 11 MBq), radioactivity in the brain was measured with the PET camera for 51 min with in-plane resolution of 3.75 mm. The subjects were lying in a supine position throughout the studies. Data were corrected for dead-time, decay, and measured photon attenuation and dynamic PET-scans were reconstructed with vendor provided standard MRAC and MRP methods [Alenius and Ruotsalainen, 1997]. High-resolution anatomical MR reference images (1 mm³ resolution) were acquired using a T1-weighted sequence (TR 25 ms, TE 4.6 ms, flip angle 30°, scan time 376 s).

To correct for head motion, dynamic PET images were first realigned frame-to-frame. The individual T1-weighted MR images were coregistered to the summation images calculated from the realigned frames. Occipital cortex was delineated manually on MRI images with PMOD 3.4 software (PMOD Technologies, Zurich, Switzerland) and used as reference region. Receptor binding was expressed in

terms of BP_{ND}, which is the ratio of specific to nondisplaceable binding in brain using the occipital cortex as the reference region. BP_{ND} was calculated for each voxel using the simplified reference tissue model with reference tissue time activity curves as input data [Gunn et al., 1997]. The subject-wise parametric BP_{ND} images were normalized to the MNI space using the T1-weighted MR images, and smoothed with a Gaussian kernel of 8-mm FWHM. The effects of anxious and avoidant attachment styles on MOR availability were then assessed in SPM8 (<http://www.fil.ion.ucl.ac.uk/spm/>) using linear regression model. Statistical threshold was set at $P < 0.05$, FDR corrected at cluster level.

In addition to its putative role in social bonding, the endogenous opioid system is intimately involved in processing both rewards and nociceptive information [Van Ree et al., 2000; Zubieta et al., 2001], and both mechanisms have been proposed to contribute to the maintenance of social ties [Eisenberger, 2012]. To specify the role of these circuits in adult attachment, anatomical regions of interest (ROIs) were drawn around components of the reward (ventral striatum, dorsal striatum, amygdala, and orbitofrontal cortex [OFC]) and pain circuits (ACC, insula, and thalamus). This functional division is obviously not exclusive, as most of these regions subserve a multitude of functions related to social, emotional, and nociceptive processing. The ROIs were defined using the AAL [Tzourio-Mazoyer et al., 2002] and Anatomy [Eickhoff et al., 2005]. Subsequently, associations between regional BP_{ND} and attachment avoidance and anxiety were analyzed in each ROI. In a complementary methodological approach, linear regression analysis with attachment avoidance (or anxiety) as the dependent variable and backward elimination of ROI-wise BP_{ND} predictors was used to pinpoint regions whose BP_{ND}'s had the strongest associations with attachment anxiety and avoidance.

RESULTS

Figure 1 shows the mean [11C]carfentanil binding distribution in the brain. Avoidance scores correlated negatively ($P < 0.05$ FDR corrected) with MOR availability in the thalamus, anterior (ACC), middle (MCC) and posterior (PCC) cingulate cortices, and medial (mPFC) and lateral prefrontal cortex (Figs. 2 and 3). Additional associations were observed in regions spanning ventral striatum, amygdala, and insula. There was no association between MOR availability and anxiety scores. The effect of attachment avoidance (or anxiety) on MOR availability did not differ across males and females; accordingly avoidance and anxiety scores did not differ between males and females, $t_{s(47)} < 1.67$, $P_s > 0.11$ and were not correlated with each other ($r = 0.08$, n.s.) Finally, neither depression nor anxiety scores were associated with MOR availability.

ROI analysis paralleled the overall pattern of the results from whole-brain GLM in a subset of the tested

Figure 1.
Mean [11C]carfentanil BP_{ND} distribution in the brain.

ROIs: Attachment avoidance (but not anxiety) was negatively associated with BP_{ND} in OFC, amygdala, dorsal striatum, and thalamus ($P_s < 0.05$ in one-tailed test). However, associations were not significant in insula, ACC, ventral, and dorsal striatum ($P_s > 0.05$). Finally, lin-

ear regression analysis revealed that [11C]carfentanil BP_{ND} in the OFC emerged as the sole predictor for attachment avoidance ($\beta = -0.89$, $t = -2.02$, $P < 0.05$), with no significant predictors observed for attachment anxiety.

Figure 2.
Brain regions showing a statistically significant association between avoidant attachment and MOR availability. The data are thresholded at $P < 0.05$, FDR corrected. ACC = anterior cingulate cortex, MCC = Middle cingulate cortex, mPFC = medial prefrontal cortex, OFC = orbitofrontal cortex, PCC = Posterior cingulate cortex, vSTR = ventral striatum.

Figure 3.

Association between avoidant attachment style and MOR availability in orbitofrontal cortex (**A**) and thalamus (**B**). Note: The scatterplots are for visualization only and the statistical reasoning is based on the full-volume SPM analysis. Cook's distance for all observations is <1 suggesting that no single data point or their removal significantly biases the correlation.

DISCUSSION

We show for the first time that individual differences in the attachment style of human adults is associated with baseline MOR availability, with high attachment avoidance being associated with low MOR availability both in regions belonging to the putative social distress circuit [ACC and thalamus; Panksepp, 2003] and in the medial frontocortical regions that support functions such as emotions and theory-of-mind that are critical for social interaction [mPFC, OFC; Amodio and Frith, 2006]. Subset of the regions showing attachment-dependent MOR availability also overlap with the fronto-insular saliency network [Seeley et al., 2007], possibly highlighting the role of the this system in subserving alert functions during intimate interpersonal situations. Our finding accords with primate work showing that the opioid system modulates pair bonding and attachment in primates and other mammals [Machin and Dunbar, 2011; Nelson and Panksepp, 1998], in addition to the well-known oxytocin and vasopressin systems [Young et al., 2001].

These findings suggest that variation in focal MOR availability may provide an important neurochemical mechanism explaining individual differences in avoidant adult attachment behavior. Our results emphasize that this is a quantitative relationship with receptor density, and animal studies have indeed linked endogenous MOR availability with attachment behavior. Preclinical

experiments and postmortem work in humans suggest OPRM1-modulated MOR expression [Mague et al., 2009; Zhang et al., 2005], also confirmed by human PET studies [Pecina et al., 2015; Weerts et al., 2013]. Whereas low endogenous MOR availability seen in gene knock-out mice leads to a deficit in maternal attachment in mice [Moles et al., 2004], monkey infants with gain-of-function OPRM1 77G allele express enhanced maternal attachment [Barr et al., 2008]. In human adults, the minor allele (G) of the OPRM1 A118G polymorphism also predicts self-reported avoidant attachment [Troisi et al., 2011]. In line with these studies, prior PET work in humans shows that MOR activity increases in the amygdala and anterior insula during social acceptance [Hsu et al., 2013], whereas MOR-mediated neurotransmission is decreased when individuals recall past events involving termination of social bonds [Zubieta et al., 2003]. Together with these findings, our data suggest that the MOR system is intimately involved in maintenance of social connections with significant others.

Pharmacological studies further corroborate the role of MOR in attachment behavior by showing that exogenous opioid antagonists increase social grooming in monkeys [Fabre-Nys et al., 1982; Keverne et al., 1989], whereas opioid agonists alleviate separation distress in puppies [Panksepp et al., 1978]. These effects may reflect the well-known role the opioid system plays in reward and pain processing

[Van Ree et al., 2000; Zubieta et al., 2001]. Low availability of endogenous MORs may dilute the calming effects of endogenously released opioid peptides acting on the MOR during social interaction, thus inhibiting establishment and reinforcement of safe attachment bonds [see e.g. Nelson and Panksepp, 1998]. This also accords with findings from functional imaging studies showing that attachment avoidance is associated with lowered responses to particularly positive social signals [see review in Vrticka and Vuilleumier, 2012]. However, even though the anxiety-alleviating effects of exogenous opioid agonists are well-known [Colasanti et al., 2011], individual differences in attachment anxiety or trait anxiety within the subclinical range were not mediated by endogenous MOR availability in our dataset. These findings suggest that distinct neuromolecular pathways support avoidant and anxious attachment behavior, and also imply that individual differences in subclinical trait anxiety are not manifested in MOR availability.

Prior functional magnetic resonance imaging (fMRI) work demonstrates that avoidant attachment is associated with lowered responsiveness in the brain's reward circuit during social encounters [Vrticka et al., 2008]. Our ROI analyses support these findings by showing that MOR availability was significantly associated with avoidant attachment in ROIs belonging to the reward circuit (dorsal striatum, amygdala, and OFC). Regression analysis further highlighted that MOR availability in the OFC was the best predictor for avoidant attachment. Even though significant in full-volume analysis, associations between BPND and attachment avoidance in regions involved in pain processing (insula, anterior cingulate cortex) failed to reach significance in the ROI analyses. However, whether individual differences in avoidant attachment are more strongly mediated by MOR-dependent brain circuits encoding the rewarding rather than punishing value of social encounters needs to be qualified in studies using active challenge paradigms [see DeWall et al., 2012 for related fMRI work; Vrticka et al., 2008].

The observed linkage between MOR availability and avoidant attachment also accords with clinical data on patients with substance abuse. Disruption of the endogenous opioid system by opiate addiction is associated with antisocial behavior [Ross et al., 2005]. Chronic opiate addiction leading to opioid tolerance [Koch and Holtt, 2008] may thus render MOR-dependent attachment circuitries unresponsive to social interaction. Importantly, insecure maternal—offspring attachment is associated with higher likelihood of opioid use [Cerdá et al., 2014; Schindler et al., 2005], while, critically, abuse of heroin (but not drugs that do not influence the opioidergic system, such as ecstasy or cannabis) is associated with avoidant adult attachment [Schindler et al., 2009]. Together with animal work showing how opioid agonists alleviate separation distress [Panksepp et al., 1978], these data suggest that exogenous opiates may be used as a chemical substitutes of secure attachment in avoidantly attached human individuals, further highlighting the

specific role of the opioid system in governing human avoidant attachment behavior.

Secure adult attachment style is an important protective psychosocial factor [Griffin and Bartholomew, 1994]. While secure attachment leads to relationship satisfaction, insecure and avoidant attachment is associated with psychological adjustment problems including loneliness and substance abuse [Brennan and Shaver, 1995; Kassel et al., 2007]. Moreover, avoidant attachment is associated with avoidance of intimate interpersonal exchanges that may make them vulnerable to social rejection, ultimately leading to preference for being alone [Bartholomew, 1990]. The orbitofrontal and medial prefrontal regions that show negative association between MOR availability and attachment avoidance are also crucially involved in the maintenance of human social networks [Lewis et al., 2011; Powell et al., 2012]. Insecure attachment is also associated with smaller social networks [Fiori et al., 2011], and both large social network size [Holt-Lunstad et al., 2010] and the availability of social support [Broadhead et al., 1983] are associated with beneficial effects for somatic health. Accordingly, the reciprocal links between MOR availability, attachment avoidance, and social networks could in part explain the beneficial psychosocial effects of secure attachment.

Our cross-sectional study cannot, however, establish a causal link between MOR system and attachment behavior. Although it is biologically plausible to assume that genetically determined individual differences in opioidergic neurotransmission influence social behavior [Barr et al., 2008; Moles et al., 2004; Way et al., 2009] and attachment [Troisi et al., 2010], it is also possible that individual differences in engaging with different types of social relationships could trigger neuroplastic changes in the opioid system during development. Perpetual overstimulation of the MOR system may lead to its downregulation [Karlsson et al., 2015], and similarly recurrent presence of threats and consequent support seeking could downregulate the attachment-related components of the MOR system. Even though the well-established stability of attachment security from infancy to early adulthood [Fraley, 2002] suggests that MOR availability may causally influence attachment behavior, longitudinal studies will be needed to resolve this relationship and to assess the specific contributions of genetically determined MOR neurotransmission versus experience-dependent plasticity of infant and adult attachment systems.

Ultimately, however, these findings are important because they demonstrate a direct quantitative relationship between psychological attachment style and the opioidergic system. Such a relationship might be expected to generalize across primates as a function of species differences in the intensity of the bonding system [Dunbar and Shultz, 2010] and perhaps other mammalian taxa [Shultz and Dunbar, 2010]. Differences in the degree of “bondedness” between different species or orders may be correlated with the density of the μ -receptor system and the underlying genetics.

Finally, we stress that our study was strictly focused on the individual differences in attachment and baseline receptor availability. In future studies, it would be important to use challenge paradigms where acute changes in MOR availability are measured when the attachment system is activated. Such an approach would help elucidating how individual differences in attachment styles translate to task-dependent activation of the MOR system.

CONCLUSIONS

We show that low endogenous MOR levels are associated with avoidant attachment style in human adults, suggesting that the endogenous opioid system is an important candidate for the neuromolecular mechanism underlying individual differences in attachment. Endogenous opioids may significantly contribute to human attachment behavior, in addition to the well-known effects of oxytocin and vasopressin systems [Young et al., 2001]. Yet future studies are needed to delineate the causal link between MOR expression and attachment behavior. Altogether, our findings suggest that individual differences in MOR availability could have a profound impact on the social relationships individuals establish, thus potentially having a strong impact on psychological well-being.

REFERENCES

- Alenius S, Ruotsalainen U (1997): Bayesian image reconstruction for emission tomography based on median root prior. *Eur J Nucl Med* 24:258–265.
- Amodio DM, Frith CD (2006): Meeting of minds: The medial frontal cortex and social cognition. *Nat Rev Neurosci* 7:268–277.
- Barr CS, Schwandt ML, Lindell SG, Higley JD, Maestripieri D, Goldman D, Suomi SJ, Heilig M (2008): Variation at the mu-opioid receptor gene (OPRM1) influences attachment behavior in infant primates. *Proc Natl Acad Sci USA* 105:5277–5281.
- Bartholomew K (1990): Avoidance of intimacy—An attachment perspective. *J Soc Pers Relat* 7:147–178.
- Beck AT, Steer RA, Garbin MG (1988): Psychometric properties of the beck depression inventory—25 years of evaluation. *Clin Psychol Rev* 8:77–100.
- Bowlby J (1982): *Attachment and Loss*, 2nd Ed. New York, NY: Basic Books.
- Brennan KA, Shaver PR (1995): Dimensions of adult attachment, affect regulation, and romantic relationship functioning. *Pers Soc Psychol Bull* 21:267–283.
- Brennan KA, Clark CL, Shaver PR (1998): Self-report measurement of adult romantic attachment: An integrative overview. In: Simpson JA, Rholes WS, (Eds.), *Attachment Theory and Close Relationships*. New York: Guilford Press. pp 46–76.
- Broadhead WE, Kaplan BH, James SA, Wagner EH, Schoenbach VJ, Grimson R, Heyden S, Tibblin G, Gehlbach SH (1983): The epidemiologic evidence for a relationship between social support and health. *Am J Epidemiol* 117:521–537.
- Burkett JP, Spiegel LL, Inoue K, Murphy AZ, Young LJ (2011): Activation of mu-opioid receptors in the dorsal striatum is necessary for adult social attachment in monogamous prairie voles. *Neuropsychopharmacology* 36:2200–2210.
- Cerdá M, Bordelais P, Keyes KM, Roberts AL, Martins SS, Reiser SL, Austin SB, Corliss HL, Koenen KC (2014): Family ties: Maternal-offspring attachment and young adult nonmedical prescription opioid use. *Drug Alcohol Depend* 142:231–238.
- Colasanti A, Rabiner E, Lingford-Hughes A, Nutt D (2011): Opioids and anxiety. *J Psychopharmacol* 25:1415–1433.
- DeWall CN, Masten CL, Powell C, Combs D, Schurtz DR, Eisenberger NI (2012): Do neural responses to rejection depend on attachment style? An fMRI study. *Soc Cogn Affect Neurosci* 7:184–192.
- Dunbar RIM, Shultz S (2010): Bondedness and sociality. *Behaviour* 147:775–803.
- Dunbar RIM, Baron R, Frangou A, Pearce E, van Leeuwen EJC, Stow J, Partridge G, MacDonald I, Barra V, van Vugt M (2012): Social laughter is correlated with an elevated pain threshold. *Proc R Soc B-Biol Sci* 279:1161–1167.
- Eickhoff S, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, Zilles K, (2005): A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *NeuroImage* 25:1325–1335.
- Eisenberger NI (2012): The pain of social disconnection: Examining the shared neural underpinnings of physical and social pain. *Nat Rev Neurosci* 13:421–434.
- Fabre-Nys C, Meller RE, Keverne EB (1982): Opiate antagonists stimulate affiliative behaviour in monkeys. *Pharmacol Biochem Behav* 16:653–659.
- Fiori KL, Consedine NS, Merz EM (2011): Attachment, social network size, and patterns of social exchange in later life. *Res Aging* 33:465–493.
- Fraley RC (2002): Attachment stability from infancy to adulthood: Meta-analysis and dynamic modeling of developmental mechanisms. *Pers Soc Psychol Rev* 6:123–151.
- Fraley RC, Waller NG, Brennan KA (2000): An item response theory analysis of self-report measures of adult attachment. *J Pers Soc Psychol* 78:350–365.
- Graves FC, Wallen K, Maestripieri D (2002): Opioids and attachment in rhesus macaque (*Macaca mulatta*) abusive mothers. *Behav Neurosci* 116:489–493.
- Griffin D, Bartholomew K (1994): Models of the self and other—Fundamental dimensions underlying measures of adult attachment. *J Pers Soc Psychol* 67:430–445.
- Gunn RN, Lammertsma AA, Hume SP, Cunningham VJ (1997): Parametric imaging of ligand-receptor binding in PET using a simplified reference region model. *Neuroimage* 6:279–287.
- Hazan C, Shaver P (1987): Romantic love conceptualized as an attachment process. *J Pers Soc Psychol* 52:511–524.
- Holt-Lunstad J, Smith TB, Layton JB (2010): Social relationships and mortality risk: A meta-analytic review. *PLoS Med* 7:e1000316.
- Hsu DT, Sanford BJ, Meyers KK, Love TM, Hazlett KE, Wang H, Ni L, Walker SJ, Mickey BJ, Korycinski ST, Koeppe RA, Crocker JK, Langenecker SA, Zubieta JK (2013): Response of the mu-opioid system to social rejection and acceptance. *Mol Psychiatry* 18:1211–1217.
- Johnson ZV, Young LJ (2015): Neurobiological mechanisms of social attachment and pair bonding. *Curr Opin Behav Sci* 3:38–44.
- Karlsson HK, Tuominen L, Tuulari JJ, Hirvonen J, Parkkola R, Helin S, Salminen P, Nuutila P, Nummenmaa L (2015): Obesity is associated with decreased mu-opioid but unaltered dopamine D-2 receptor availability in the brain. *J Neurosci* 35:3959–3965.

- Kassel JD, Wardle M, Roberts JE (2007): Adult attachment security and college student substance use. *Addict Behav* 32:1164–1176.
- Keverne EB, Martensz ND, Tuite B (1989): Beta-endorphin concentrations in cerebrospinal-fluid of monkeys are influenced by grooming relationships. *Psychoneuroendocrinology* 14:155–161.
- Koch T, Holtt V (2008): Role of receptor internalization in opioid tolerance and dependence. *Pharmacol Ther* 117:199–206.
- Lewis PA, Rezaie R, Brown R, Roberts N, Dunbar RIM (2011): Ventromedial prefrontal volume predicts understanding of others and social network size. *Neuroimage* 57:1624–1629.
- Lutz PE, Kieffer BL (2013): Opioid receptors: Distinct roles in mood disorders. *Trends Neurosci* 36:195–206.
- Machin AJ, Dunbar RIM (2011): The brain opioid theory of social attachment: A review of the evidence. *Behaviour* 148:985–1025.
- Mague SD, Isiegas C, Huang P, Liu-Chen LY, Lerman C, Blendy JA (2009): Mouse model of OPRM1 (A118G) polymorphism has sex-specific effects on drug-mediated behavior. *Proc Natl Acad Sci USA* 106:10847–10852.
- Mikulincer M, Shaver PR (2007): *Attachment in Adulthood*. New York: Guilford.
- Moles A, Kieffer BL, D'Amato FR (2004): Deficit in attachment behavior in mice lacking the mu-opioid receptor gene. *Science* 304:1983–1986.
- Nelson EE, Panksepp J (1998): Brain substrates of infant-mother attachment: Contributions of opioids, oxytocin, and norepinephrine. *Neurosci Biobehav Rev* 22:437–452.
- Norman L, Lawrence N, Iles A, Benattayallah A, Karl A (2015): Attachment-security priming attenuates amygdala activation to social and linguistic threat. *Soci Cogn Affect Neurosci* 10:832–839.
- Panksepp J (2003): Feeling the pain of social loss. *Science* 302:237–239.
- Panksepp J, Herman B, Conner R, Bishop P, Scott JP (1978): Biology of social attachments—Opiates alleviate separation distress. *Biol Psychiatry* 13:607–618.
- Panksepp J, Herman BH, Vilberg T, Bishop P, Deeskinazi FG (1980): Endogenous opioids and social behavior. *Neurosci Biobehav Rev* 4:473–487.
- Pecina M, Love T, Stohler CS, Goldman D, Zubieta JK (2015): Effects of the mu opioid receptor polymorphism (OPRM1 A118G) on pain regulation, placebo effects and associated personality trait measures. *Neuropsychopharmacology* 40:957–965.
- Powell J, Lewis PA, Roberts N, Garcia-Finana M, Dunbar RIM (2012): Orbital prefrontal cortex volume predicts social network size: An imaging study of individual differences in humans. *Proc R Soc B-Biol Sci* 279:2157–2162.
- Resendez SL, Dome M, Gormley G, Franco D, Nevarez N, Hamid AA, Aragona BJ (2013): Mu-opioid receptors within subregions of the striatum mediate pair bond formation through parallel yet distinct reward mechanisms. *J Neurosci* 33:9140–9149.
- Ross J, Teesson M, Darke S, Lynskey M, Ali R, Ritter A, Cooke R (2005): The characteristics of heroin users entering treatment: Findings from the Australian Treatment Outcome Study (ATOS). *Drug Alcohol Rev* 24:411–418.
- Schindler A, Thomasius R, Sack PM, Gemeinhardt B, KÜstner U, Eckert J (2005): Attachment and substance use disorders: A review of the literature and a study in drug dependent adolescents. *Attach Hum Dev* 7:207–228.
- Schindler A, Thomasius R, Petersen K, Sack PM (2009): Heroin as an attachment substitute? Differences in attachment representations between opioid, ecstasy and cannabis abusers. *Attach Hum Dev* 11:307–330.
- Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, Reiss AL, Greicius MD (2007): Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* 27:2349–2356.
- Shultz S, Dunbar RIM (2010): Encephalisation is not a universal macroevolutionary phenomenon in mammals but is associated with sociality. *Proc Natl Acad Sci USA* 107:21582–21586.
- Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA (1983): *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting Psychologists Press.
- Troisi A, Alcini S, Coviello M, Nanni RC, Siracusano A (2010): Adult attachment style and social anhedonia in healthy volunteers. *Pers Individ Differ* 48:640–643.
- Troisi A, Frazzetto G, Carola V, Di Lorenzo G, Coviello M, D'Amato FR, Moles A, Siracusano A, Gross C (2011): Social hedonic capacity is associated with the A118G polymorphism of the mu-opioid receptor gene (OPRM1) in adult healthy volunteers and psychiatric patients. *Soc Neurosci* 6:88–97.
- Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B, Joliot M (2002): Automatic anatomical labelling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage* 15:273–289.
- Van Ree JM, Niesink RJM, Van Wolfswinkel L, Ramsey NF, Kornet M, Van Furth WR, Vanderschuren L, Gerrits M, Van den Berg CL (2000): Endogenous opioids and reward. *Eur J Pharmacol* 405:89–101.
- Vrticka P, Vuilleumier P (2012): Neuroscience of human social interactions and adult attachment style. *Front Hum Neurosci* 6:212.
- Vrticka P, Andersson F, Grandjean D, Sander D, Vuilleumier P (2008): Individual attachment style modulates human amygdala and striatum activation during social appraisal. *PLoS One* 3:11.
- Vrticka P, Sander D, Vuilleumier P (2012): Influence of adult attachment style on the perception of social and non-social emotional scenes. *J Soc Pers Relat* 29:530–544.
- Way BM, Taylor SE, Eisenberger NI (2009): Variation in the mu-opioid receptor gene (OPRM1) is associated with dispositional and neural sensitivity to social rejection. *Proc Natl Acad Sci USA* 106:15079–15084.
- Weerts EM, McCaul ME, Kuwabara H, Yang XJ, Xu XQ, Dannals RF, Frost JJ, Wong DF, Wand GS (2013): Influence of OPRM1 Asn(40)Asp variant (A118G) on C-11 carfentanil binding potential: Preliminary findings in human subjects. *Int J Neuropsychopharmacol* 16:47–53.
- Young LJ, Lim MM, Gingrich B, Insel TR (2001): Cellular mechanisms of social attachment. *Horm Behav* 40:133–138.
- Zhang Y, Wang DX, Johnson AD, Papp AC, Sadee W (2005): Allelic expression imbalance of human mu opioid receptor (OPRM1) caused by variant A118G. *J Biol Chem* 280:32618–32624.
- Zubieta JK, Smith YR, Bueller JA, Xu YJ, Kilbourn MR, Jewett DM, Meyer CR, Koeppe RA, Stohler CS (2001): Regional mu opioid receptor regulation of sensory and affective dimensions of pain. *Science* 293:311–315.
- Zubieta JK, Ketter TA, Bueller JA, Xu YJ, Kilbourn MR, Young EA, Koeppe RA (2003): Regulation of human affective responses by anterior cingulate and limbic mu-opioid neurotransmission. *Arch Gen Psychiatry* 60:1145–1153.