Longitudinal Resting State fMRI Analysis in Healthy Controls and Premanifest Huntington's Disease Gene Carriers: A Three-Year Follow-Up Study

Omar F.F. Odish,^{1*} Annette A. van den Berg-Huysmans,²
Simon J.A. van den Bogaard,¹ Eve M. Dumas,¹ Ellen P. Hart,¹
Serge A.R.B. Rombouts,^{2,3,4} Jeroen van der Grond,²
and Raymund A.C. Roos¹ on behalf of the TRACK-HD Investigator Group

¹Department of Neurology, Leiden University Medical Center, Leiden, The Netherlands

²Department of Radiology, Leiden University Medical Center, Leiden, The Netherlands

³Leiden University – Institute of Psychology, Leiden, The Netherlands

⁴Leiden Institute for Brain and Cognition (LIBC), Leiden, The Netherlands

Abstract: Background: We previously demonstrated that in the premanifest stage of Huntington's disease (preHD), a reduced functional connectivity exists compared to healthy controls. In the current study, we look at possible changes in functional connectivity occurring longitudinally over a period of 3 years, with the aim of assessing the potential usefulness of this technique as a biomarker for disease progression in preHD. Methods: Twenty-two preHD and 17 healthy control subjects completed resting state functional magnetic resonance imaging (fMRI) scans in two visits with 3 years in between. Differences in resting state connectivity were examined for eight networks of interest using FSL with three different analysis types: a dual regression method, region of interest approach, and an independent component analysis. To evaluate a possible combined effect of gray matter volume change and the change in blood oxygenation level dependent signal, the analysis was performed with and without voxel-wise correction for gray matter volume. To evaluate possible correlations between functional connectivity change and the predicted time to disease onset, the preHD group was classed as preHD-A if ≥ 10.9 years and preHD-B if < 10.9 years from predicted disease onset. Possible correlations between burden of pathology score and functional connectivity change in preHD were also assessed. Finally, longitudinal change in whole brain and striatal volumetric measures was assessed in the studied cohort. Results: Longitudinal analysis of the resting state-fMRI (RS-fMRI) data revealed no differences in the degree of connectivity change between the groups over a period of 3 years, though a significantly higher rate of striatal atrophy was found in the preHD group compared to controls in the same period. Discussion: Based on the results found in this study, the provisional conclusion is that

Contract grant sponsor: CHDI Foundation, Inc.; Contract grant sponsor: The Netherlands Organisation for Scientific Research (NWO; to S.R.)

*Correspondence to: Omar F.F. Odish, Department of Neurology (J3-R-162), Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, The Netherlands. E-mail: o.odish@lumc.nl

Received for publication 5 March 2013; Revised 3 July 2014; Accepted 12 August 2014.

DOI: 10.1002/hbm.22616

Published online 19 August 2014 in Wiley Online Library (wileyonlinelibrary.com).

RS-fMRI lacks sensitivity in detecting changes in functional connectivity in HD gene carriers prior to disease manifestation over a 3-year follow-up period. *Hum Brain Mapp* 36:110–119, 2015. © 2014 Wiley Periodicals, Inc.

Key words: Huntington's disease; premanifest; resting state functional magnetic resonance imaging; longitudinal biomarker; functional connectivity; TRACK-HD; movement disorders

INTRODUCTION

Huntington's disease (HD) is an autosomal dominantly inherited neurodegenerative disorder characterized by motor, cognitive, and psychiatric symptoms with a mean age at onset between 30 and 50 years [Roos, 2010]. It is caused by an expanded cytosine-adenine-guanine (CAG) trinucleotide repeat in the huntingtin (HTT) gene on the short arm of chromosome 4 [The Huntington's Disease Collaborative Research Group, 1993]. Magnetic resonance imaging (MRI) studies in HD have revealed extensive brain atrophy, most notably in the striatum [Aylward et al., 2012; Dumas et al., 2012; Hadzi et al., 2012; Stoffers et al., 2010; Tabrizi et al., 2009; Van den Bogaard et al., 2011; Vonsattel et al., 2011].

A current challenge in HD research is establishing reliable biomarkers for measuring disease progression in HD, both before and after disease manifestation. This is crucial for assessing the efficacy of future proposed therapies. Several large longitudinal studies are currently being conducted for the purpose of establishing such biomarkers [Aylward et al., 2011; Paulsen et al., 2010; Tabrizi et al., 2012, 2013]. Using MRI, these studies have shown that atrophy of different structures in the brains of premanifest gene carriers (preHD), and of the caudate nucleus in particular, is correlated with the estimated years to disease onset (YTO) as calculated by the formula of Langbehn et al. [2004] [Aylward et al., 2011; Langbehn et al., 2004; Paulsen et al., 2010; Tabrizi et al., 2012, 2013]. This is of particular interest, as these subjects have yet to present clinically with the hallmark motor symptoms of HD.

As the correlations found up to this point only partially predict the rate of clinical deterioration, combining imaging modalities might increase the predictive validity of a potential biomarker. With resting state functional MRI (RS-fMRI), interregional correlations of blood oxygenation level dependent (BOLD) signal fluctuations between brain regions that are spatially distinct are measured in the wakeful brain, without challenging it with a particular task. The patterns acquired with this technique are usually referred to as "functional connectivity." RS-fMRI has the theoretical potential of revealing changes occurring in the brain before changes on the structural imaging level are evident, which could be important in targeting the disease in its earliest stages. It may in addition help to unravel compensatory mechanisms responsible for apparently normal brain function despite ongoing neurodegeneration.

The technique has already been shown to be a valuable marker for tracking disease progression in Alzheimer's disease, and in mild cognitive impairment [Bai et al., 2011; Damoiseaux et al., 2012].

In a previous report, our group has reported functional connectivity differences among controls, preHD, and manifest HD subjects, cross sectionally. The results showed preHD subjects already exhibiting altered functional connectivity with different structures in the brain compared to the matched control group. Importantly, this was still valid after correction for atrophy [Dumas et al., 2013]. The first report detailing reduced corticostriatal functional connectivity findings in preHD when compared to controls was by Unschuld et al. [2012]. A recent report by Poudel et al. [2014] further confirms findings of functional connectivity reductions in both preHD and manifest HD subjects.

In the current longitudinal study, we aim to assess the potential usefulness of this technique as a biomarker for disease progression in the premanifest stage of the disease. We investigate possible changes in functional connectivity occurring longitudinally over a follow-up period of 3 years. With the aim of having a comprehensive interpretation of the acquired data, three separate data analysis methods were applied.

METHODS

Subjects

Of the 28 premanifest HD carriers (preHD) and 28 healthy age-matched control subjects who completed RS-fMRI scans during their first visit at the Leiden University Medical Center (LUMC) study site of the TRACK-HD study [Tabrizi et al., 2009], 23 preHD and 19 control subjects completed the resting state scans at the second visit, with a 3-year interval between visits. Excluded from analysis were one preHD subject due to missing scan volumes and two control subjects due to excessive motion artifacts (maximum motion during scan < 4 mm [Jenkinson et al., 2012]). This resulted in 22 preHD and 17 healthy control subject that were included in this study (Table I).

Inclusion criteria for study participation for preHD subjects comprise a positive genetic test with ≥ 40 CAG repeats, the absence of motor disturbances on the total motor score (TMS) of the Unified Huntington's Disease Rating Scale (UHDRS) of more than five points and a burden of pathology score greater than 250 ((CAG repeat

TABLE I. Group characteristics and clinical scores

	Healthy controls	preHD (A and B)
N	18	22 ^a
Gender M/F	7/11	10/12
Age in years (V1), mean (SD)	46.7 (6.9)	43.3 (8.5)
Handedness R/L	18/0	18/4
Level of education (ISCED), median (range)	4 (3)	4 (3)
DART-IQ, mean (SD)	105.3 (9.3)	100.3 (11.6)
BMI in kg/m ² (V1), mean (SD)	26.9 (6.6)	24.9 (4.1)
CAG repeat length, mean (SD)	n/a	42.6 (2.6)
Estimated years to onset (YTO), mean (SD)	n/a	11.6 (4.4)
Total functional capacity, mean (SD)		
V1	13.0 (0.0)	12.7 (0.8)
V2	13.0 (0.0)	12.6 (0.9) ^b
UHDRS-TMS, mean (SD)		
V1	2.4 (2.5)	2.4 (1.5)
V2	2.2 (3.0)	5.4 (5.7) ^b
SDMT, mean (SD)		
V1	53.7 (8.9)	48.7 (9.7)
V2	58.4 (8.0)	49.4 (10.5) ^b
BDI-II, mean (SD)		
V1	4.4 (6.3)	5.1 (5.7)
V2	4.8 (5.1)	5.3 (6.0)
Between-scan interval in months, mean (SD)	35.6 (1.20)	35.3 (0.94)

N, number of participants; SD, standard deviation; n/a, not applicable; ISCED, International Standard Classification of Education; DART-IQ, Dutch Adult Reading Test Intelligence Quotient; CAG, cytosine-adenine-guanine; UHDRS-TMS, Unified Huntington's Disease Rating Scale total motor score; SDMT, symbol digit modalities test; BDI-II, Beck Depression Inventory-II; BMI, body mass index; V1, visit 1; and V2, visit 2.

^aIncluding four subjects progressing to the manifest stage during the 3-year follow-up period.

^bA significant difference at $P < 0.05$.

length $- 35.5) \times \text{age}$) [Penney et al., 1997; Tabrizi et al., 2009]. Age- and gender-matched gene-negative relatives of HD gene carriers and spouses were included as healthy controls. Exclusion criteria for all participants included significant previous head trauma, any neurological or major psychiatric disorder or unwillingness to undergo MRI scanning [Tabrizi et al., 2009]. Medical history taking, an interview-based assessment and questionnaires, was used to ascertain that no major psychiatric disorder could be classified at the time of inclusion and scanning. Consequently, the use of neuroleptic medications or antidepressants was sparse and considered to be of no influence.

For preHD subjects, the estimated number of years until disease onset was calculated based on their current age and the CAG repeat length, by means of the formula developed by Langbehn et al. [2004].

As previously applied by Tabrizi et al. [2009], for a second analysis, the preHD group was divided at baseline

according to the median (10.9 years) for the predicted years to onset into preHD-A (≥ 10.9 years from predicted onset) and preHD-B (< 10.9 years). This resulted in two groups each consisting of 11 subjects (Table II). In a further analysis performed within the preHD group, possible associations between functional connectivity change and burden of pathology score were assessed.

The study was approved by the ethics committee of the LUMC and written informed consent was obtained from all participants following a complete description of the study and procedures. For full details of study parameters, see Tabrizi et al. [2009].

Clinical Measures

To monitor disease state, the following clinical measures were collected longitudinally for all groups: Unified Huntington's Disease Rating Scale total motor score (UHDRS-TMS), total functional capacity (TFC), symbol digit modalities test (SDMT), and Beck Depression Inventory-II (BDI-II) scores.

The UHDRS-TMS is the traditional measure which defines disease state in HD. The SDMT in particular has been shown to be a sensitive longitudinal cognitive measure in HD, independent of disease related motor effects [Tabrizi et al., 2011].

MRI Acquisition

MRI acquisition was performed on a 3-T whole body scanner (Philips Achieva, Healthcare, Best, The Netherlands) with an eight channel receive array head coil. An anatomical T1-weighted scan was acquired using an ultra-fast gradient echo 3D acquisition sequence with the following imaging parameters: repetition time (TR) = 7.7 ms, echo time (TE) = 3.5 ms, field-of-view = $24 \times 24 \times 16.4 \text{ cm}^3$, matrix size 224×224 , with a duration of 9 min. For postprocessing registration purposes, a high-resolution T2*-weighted scan, with the following parameters was collected: repetition time (TR) = 2,200 ms, echo time (TE) = 30 ms, field-of-view = $220 \times 220 \times 168 \text{ mm}^3$, flip angle = 80° , matrix size = $112 \times 109 \text{ mm}^2$, with a duration of 46 s. A RS-fMRI scan with the following parameters was obtained: 200 EPI volumes, repetition time (TR) = 2,200 ms, echo time (TE) = 30 ms, field-of-view = $220 \times 220 \times 10.4$, resolution = $2.75 \times 2.75 \times 2.75$, no slice gap, flip angle = 80° , matrix size 80×79 , with a duration of 7.5 min. No background music was played during the RS-fMRI scan and to ensure a wakeful disposition participants were asked to keep their eyes open with normal background light.

Preprocessing of Resting State Data

RS-fMRI images were analyzed using FSL 5.0 (fMRIB Software Library; available at www.fmrib.ox.ac.uk/fsl). Preprocessing consisted of motion correction [Jenkinson et al., 2002], removal of nonbrain tissue [Smith, 2002], spatial smoothing using a Gaussian kernel of 6 mm full width at

TABLE II. preHD-A vs. preHD-B, visit 1

	preHD-A	preHD-B
N	11	11
Gender M/F	3/8	7/4
Age in years, mean (SD)	43.8 (5.8)	43.0 (10.9)
Handedness R/L	9/2	9/2
Level of education (ISCED), median (range)	4 (3)	4 (3)
DART-IQ, mean (SD)	102.3 (9.9)	98.3 (13.2)
BMI in kg/m ² , mean (SD)	25.6 (3.0)	23.1 (2.3)
CAG repeat length, mean (SD)	41.5 (1.4)	43.8 (3.1) ^a
Estimated years to onset (YTO), mean (SD)	14.4 (4.5)	8.8 (1.6) ^a
Total functional capacity, mean (SD)	12.7 (0.7)	12.6 (0.9)
UHDRS-TMS, mean (SD)	1.9 (1.5)	2.9 (1.3)
SDMT, mean (SD)	51.6 (9.9)	45.9 (9.1)
BDI-II, mean (SD)	4.5 (6.0)	5.6 (5.7)
Between-scan interval in months, mean (SD)	35.6 (1.0)	34.9 (0.7)

N, number of participants; SD, standard deviation; ISCED, International Standard Classification of Education; DART-IQ, Dutch Adult Reading Test Intelligence Quotient; CAG, cytosine-adenine-guanine; UHDRS-TMS, Unified Huntington's Disease Rating Scale total motor score; SDMT, symbol digit modalities test; BDI-II, Beck Depression Inventory-II; and BMI, body mass index.

^aA significant difference at $P < 0.05$.

half maximum (FWHM) and high-pass temporal filtering equivalent to 100 s (0.01 Hz). After preprocessing, the functional images were registered to the high-resolution T2*-weighted images. These high-resolution images were subsequently registered to the anatomical T1-weighted images. Finally, the anatomical scan was registered to the 2 mm isotropic MNI152 standard space image [Jenkinson et al., 2002]. These three registration matrices were combined to obtain a matrix for transforming fMRI data from native space to standard space and its inverse (from MNI space to native space). Visual quality control was performed by two qualified raters to ensure correct registration.

Statistical Analysis

Statistical analysis of group demographics and clinical measures was performed using IBM SPSS Statistics (version 20.0, IBM Corp., USA). Where appropriate either an independent samples *t*-test or chi-squared tests were applied. Potential longitudinal change in clinical measures between the groups was also investigated. Difference values were computed and independent samples *t*-tests on these delta-scores evaluated whether preHD subjects experienced a greater change from visit 1 to visit 2 than control subjects.

Striatal and whole brain volumes were obtained from the TRACK-HD study database [Tabrizi et al., 2009, 2013]. These measures were calculated using the Iowa BRAINS method as previously described [Ghayoor et al., 2013; Tab-

rizi et al., 2009, 2013; Young and Johnson, 2013]. Assessment of possible longitudinal volumetric change was performed using a general linear model with age, gender, and total brain volume (the latter only for assessing striatal volumes) as covariates in the model.

The functional connectivity analysis was performed in three ways using the dual regression method of FSL, a technique that allows a voxel-wise comparison of resting state functional connectivity [previously described by Filippini et al., 2009]. To assess possible associations between the burden of pathology score and functional connectivity change, a regression analysis was performed within the preHD group only.

Network of Interest Analysis

First, resting state functional connectivity was determined in terms of similarity of the BOLD fluctuations in the brain in relation to characteristic fluctuations in predefined resting state networks or networks of interest (NOIs). Our choice of resting state networks was based on high reproducibility of these networks from independent component analysis (ICA) of different datasets [Beckmann et al., 2005; Damoiseaux et al., 2006]. These standardized resting state networks parcellate the brain into eight templates that represent more than 80% of the total brain volume [Khalili-Mahani et al., 2012]: (1) medial visual network, (2) lateral visual network, (3) auditory network, (4) sensorimotor system, (5) default mode network, (6) executive control network, and (7 and 8) dorsal visual stream networks (Fig. 1) [Beckmann et al., 2005]. To account for noise, a white matter (WM) and a cerebrospinal fluid (CSF) template were included in the analysis [Birn, 2012; Cole et al., 2010; Fox et al., 2005].

Dual regression analysis (part of FSL 5.0) was performed to identify subject-specific time course and spatial maps. To create the average time course within each network for every subject, the eight resting state networks [Beckmann et al., 2005] and the two additional WM and CSF maps [Birn, 2012; Cole et al., 2010; Fox et al., 2005] were used in a linear model fit against each individual subject's fMRI dataset (spatial regression). Hence, WM and CSF activities were included in the regression model as proxy measures for non-neuronal noise. The personalized time courses were subsequently regressed back onto that subject's fMRI dataset to create personal spatial maps (temporal regression). This gives ten 3D images per individual per visit, with voxel-wise the *z*-scores of functional connectivity to each of the templates. The higher the absolute value of the *z*-score, the stronger the connectivity to a network.

Independent Component Analysis

In a second approach, large-scale patterns of functional connectivity were identified by ICA using probabilistic ICA as implemented in the MELODIC tool of FSL [Beckmann et al., 2005; Beckmann and Smith, 2004]. The original concatenated 4D RS-fMRI dataset was decomposed into sets of

Figure 1.

Sagittal, coronal and axial views of the dominant BOLD fluctuations within the eight predefined networks of interest [Beckmann et al., 2005]. All images have been coregistered into the MNI152 standard space template. Numbers at the top of the images denote the MNI coordinates (xyz) and images are shown in radiological orientation. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

time courses and associated spatial maps, to identify different activation components without any model being specified [Beckmann and Smith, 2004; McKeown et al., 1998]. The number of components was fixed to 25 to limit independent component splitting into subcomponents [Damoiseaux et al., 2012; Filippini et al., 2009].

Subsequently, the dual regression analysis as described above was repeated for the group ICA results. This time the 25 independent components were used as spatial regressors, ultimately resulting in 25 z-score maps per individual per visit, reflecting the connectivity strength of each voxel in the brain to each of the 25 independent components.

Region of Interest Analysis

Given the overwhelming volume of evidence indicating the striatum as the prime and earliest region affected within the brain in HD, we chose the striatum as a region

of interest (ROI) in our analysis. A mask was created to analyze the change in connectivity with the 8 NOIs and the 25 independent components of the voxels within this ROI. The mask was based on the probabilistic atlas incorporated in FSL provided by the Harvard Center for Morphometric Analysis and contained the striatum from both hemispheres (Fig. 2) [Desikan et al., 2006; Frazier et al., 2005; Goldstein et al., 2007; Makris et al., 2006].

Longitudinal change in connectivity per subject and per predefined network/independent component was the main parameter of interest. To assess this change, the individual functional connectivity maps (z-score) from the second visit were subtracted from the corresponding functional connectivity maps from the first visit.

For the between-group analysis, the z-score maps created by dual regression and the maps containing the differences in z-score were collected across subjects into single 4D maps (one per NOI or original independent component, with the

Figure 2.

Axial view of the region of interest (ROI) mask of the striatum shown superimposed on a MNI152 standard image. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

fourth dimension being subject identification) and submitted to voxel-based statistical testing. To obtain group averages of maps containing the differences in z-score, a one-sample nonparametric *t*-test was used and a two-sample *t*-test was applied to obtain group differences for each of the eight NOIs and each independent component, using a general linear modeling (GLM) approach as implemented in FSL. Age and gender were included as covariates in the model. To statistically account for potential effects of local structural differences within and between the two groups, gray matter volume of each voxel was included as subject-wise and voxel-wise covariates in the GLM design [Oakes et al., 2007]. To evaluate a possible combined effect of gray matter volume change and the change in BOLD signal, the analysis was also performed without voxel-wise correction for gray matter volume.

Voxel-wise nonparametric permutation testing was performed using FSL-randomize (5,000 permutations) [Nichols and Holmes, 2002]. All statistical maps were family-wise error (FWE) corrected using $P < 0.05$, based on the TFCE statistic image [Smith and Nichols, 2009].

As multiple comparison correction method only corrects the results at the predefined network/independent component level but does not adjust for the risk of Type 1 error (false positives) induced by increasing the number of components tested simultaneously at high model orders, additional correction for multiple comparisons was done using Bonferroni correction. The multiple comparisons consisted of two comparisons (either connectivity increase or decrease as compared to healthy controls) for 8 NOIs and 25 independent components.

RESULTS

Group characteristics are shown in Table I. Age, gender, handedness, and level of education did not differ signifi-

cantly between controls and preHD subjects. At baseline, no differences were found in UHDRS-TMS, TFC, SDMT, BDI-II, and Dutch Adult Reading Test Intelligence Quotient (DART-IQ) scores. There also was no difference in body mass index (BMI) at baseline. Repeated assessment at 3-year follow-up revealed significantly higher UHDRS-TMS and lower TFC and SDMT scores in the preHD group (Table I). Four of the twenty-two preHD subjects began to exhibit typical HD motor symptoms during the 3-year follow-up period, therefore reaching the definition of early manifest disease stage. The cross-sectional difference in UHDRS-TMS and TFC score between the groups at the second visit was negated after exclusion of these four converter subjects, yet the difference in SDMT score remained significant ($P = 0.07$, $P = 0.36$, and $P = 0.01$, respectively). The difference in SDMT comprises higher mean scores within the control group when compared to their first visit, while the scores of the preHD group remained stagnant.

The longitudinal change in the UHDRS-TMS was significant when all participants were included ($P = 0.03$), yet this result was only reached as a result of outlier scores: when the four converters were excluded from analysis, this difference vanished ($P = 0.25$).

The longitudinal change in the SDMT score was significant when all participants were included ($P = 0.04$). While the mean SDMT difference in the preHD group remained essentially the same when the four converters were excluded (+0.64 vs. +0.67 difference points, respectively), statistical significance could no longer be reached ($P = 0.06$). See Table III for a view of the mean longitudinal change of the different measures.

No differences in any of the scores outlined above were found while comparing the preHD-A and preHD-B groups, neither at the first or second visit nor longitudinally. The CAG trinucleotide repeat count was significantly higher in the preHD-B relative to the preHD-A group ($P = 0.03$) (Table II; longitudinal change data not shown).

All scans were analyzed with and without inclusion of the four converters. All scan analyses were also repeated with exclusion of the four left-handed subjects to avoid any possible lateralization effects. The reported results are with and without voxel-wise correction for gray matter volume, as described in the Methods section. No difference was found in the amount of motion between the groups.

RS-fMRI Network Analyses

In the eight designated NOIs, longitudinal analysis of the RS-fMRI data revealed no statistically significant differences in the degree of connectivity change between controls and the preHD group. There also were no statistically significant differences between controls and preHD-A and controls and preHD-B subjects. No association could be demonstrated between the degree of connectivity change

TABLE III. Longitudinal change in clinical scores,^a mean difference

	Healthy controls	preHD (A and B)
N	18	22 ^b
Total functional capacity, MD (SD)	0.0 (0.0)	-0.1 (0.6)
UHDRS-TMS, MD (SD)	-0.2 (2.9)	3.0 (5.4) ^c
SDMT, MD (SD)	4.7 (5.7)	0.6 (6.1) ^c
BDI-II, MD (SD)	0.4 (3.6)	0.2 (5.1)
BMI in kg/m ² , MD (SD)	0.5 (2.3)	-0.4 (1.6)

N, number of participants; MD, mean difference; SD, standard deviation; UHDRS-TMS, Unified Huntington's Disease Rating Scale total motor score; SDMT, symbol digit modalities test; BDI-II, Beck Depression Inventory-II; and BMI, body mass index.

^aLongitudinal change denotes scores from visit 2 subtracted from scores from visit 1.

^bIncluding four subjects progressing to the manifest stage during the 3-year follow-up period.

^cA significant difference at $P < 0.05$.

in the different networks and the groups designated as *far* and *near* from expected onset of motor symptoms, nor with the burden of pathology score.

RS-fMRI ICA

Using the ICA method, 25 components were extracted from the data per person per visit and the differences between the two visits compared across the above outlined groups. There were no statistically significant differences in the degree of connectivity change between any of the groups. Dividing the preHD group according to the expected time of motor symptom onset again revealed no significant differences in the degree of connectivity change. Regression analysis using the burden of pathology score revealed no associations with the degree of functional connectivity change within the preHD group.

RS-fMRI ROI Analysis

Using the described mask to assess the change of connectivity strength in the voxels within the striatum, no statistically significant differences could be demonstrated between any of the groups described above.

When comparing results from the outlined analysis methods, the ROI analysis provided the proximity to achieving a significant longitudinal reduction in functional connectivity in preHD when compared to controls. This was the case with the lateral visual network (NOI 2; $P = 0.08$) and default mode network (NOI 5; $P = 0.11$) (Fig. 3). Power analysis using these results show that a minimum of 23 subjects per group would be needed to detect a significant longitudinal reduction in functional connectivity in 3 years within the striatum with the lateral visual

network for preHD compared to controls (at 5% FWE rate with a power of 80%).

Table IV provides an overview of significance levels for longitudinal reduction of functional connectivity within the striatum over 3 years in preHD subjects compared to controls with the eight NOIs.

Longitudinal Volumetric Analysis

In the 3-year follow-up period, no statistically significant difference in whole brain volume decline was found between controls (0.33%) and preHD (0.58%) ($P = 0.35$).

The striatal volume showed a significantly higher rate of decline over the 3-year period in preHD as compared to controls: 1.45% in the control group versus 7.29% in the preHD group ($P < 0.001$). Striatal volume decline over the 3 years was significantly higher in both preHD-A (6.62%) and preHD-B (8.15%) when compared to controls ($P < 0.001$). The difference in striatal volume decline rate between preHD-A and preHD-B was not statistically significant over this time period ($P = 0.31$).

DISCUSSION

This study showed no longitudinal difference in functional connectivity change between preHD and healthy control subjects over a period of 3 years. This was also the case when preHD subjects were divided in a preHD-A and preHD-B group based on the expected time to disease onset and when using burden of pathology score as a regressor for functional connectivity change. These conclusions are based on results obtained from three different analysis methods. Results remained the same with and

Figure 3.

P-value maps of the nonsignificant longitudinal reductions in functional connectivity in preHD compared to controls in the striatum with the lateral visual and default mode networks in the 3-year study period.

TABLE IV. Statistical parameters for longitudinal reduction of functional connectivity within the striatum over 3 years in preHD subjects compared to controls with the eight networks of interest

Network of interest (NOI)		Minimal P -value	x	y	z	t -Stat
1	Medial visual	0.356	39	68	42	3.527
2	Lateral visual	0.078	53	69	35	4.175
3	Auditory	0.686	36	56	49	3.280
4	Sensorimotor	0.804	54	62	45	2.491
5	Default mode	0.112	53	70	36	3.845
6	Executive control	0.734	31	69	32	2.327
7	Dorsal visual stream	0.502	59	66	39	3.355
8	Dorsal visual stream	0.262	36	68	29	3.754

x , y , and z denote MNI152 standard space coordinates.

without voxel-wise correction for gray matter volume and while running the analysis with the inclusion and/or exclusion of converters and left-handed subjects.

This result, taken together with clinical parameters like the UHDRS-TMS and SDMT showing longitudinal change between the included subjects, and significantly higher longitudinal striatal atrophy rate in preHD compared to controls, alludes to a lack of sensitivity of RS-fMRI in detecting concomitant changes in functional connectivity occurring longitudinally in preHD. This statement should be considered as tentative, as future studies with greater numbers of participants, improved signal-to-noise ratio, different analysis methods and/or a longer follow-up period might be able to demonstrate longitudinal differences in functional connectivity change. That being said, results from this study suggest that even if there is functional connectivity change occurring in the 3-year follow-up period, this is too small to detect with this technique using the highlighted methods with this cohort size, which is a relevant finding in light of longitudinal biomarker research in preHD.

Our study confirms the results found by Seibert et al. [2012]. Their study reported no change in functional connectivity over a 1 year period. The differences between the study of Seibert et al. [2012] and our own were the methodology used, where seeds instead of a priori spatial NOIs were used and subject-native space registration instead of the MNI152 standard space template was applied. The number of subjects examined in that report was higher than in our study: 22 controls and 34 preHD subjects.

Our earlier cross-sectional results suggested that functional connectivity, at the group level, was a fairly sensitive measure to differentiate preHD subjects from controls [Dumas et al., 2013]. As such, we were quite hopeful to demonstrate a divergent longitudinal functional connectivity evolution between the groups, which in turn could serve as a measure for disease progression. We were however unable to reproduce these results within our baseline cohort, most likely due to the smaller number of subjects that were included, as only those with scans at both time points could be assessed longitudinally. This study can therefore not account for the functional connectivity of the

dropouts, as no data are available. Furthermore, the discrepancy in baseline findings might involve deteriorating health prompting more severely affected subjects to dropout prematurely of the study, thus leaving a relatively fitter group for this study. As such, selection bias disproportionately affecting subjects with the fastest rate of clinical deterioration is a possible reason for not finding different functional connectivities between the groups. This spurred using a more comprehensive approach and to base the hypothesis-driven part of the analysis solely by singling out the striatum as the primary region where possible changes in resting state activity are expected, given the fact that it is the region first affected in HD, as was again demonstrated by the volumetric study of the striatum within this cohort. Despite using three different analysis methods, no longitudinal change could be demonstrated in our cohort in a time frame of 3 years with two measurement points. The combination of a highly significant difference in striatal atrophy rate between preHD and controls with a total lack of significant difference in the rate of functional connectivity change between these groups strongly points to a lower sensitivity of RS-fMRI in demonstrating longitudinal change in the preHD population.

A similar sequence of results was found by the study of Wolf et al. [2011], where task-based fMRI showed significantly lower activity cross sectionally in the left prefrontal cortex in preHD, yet failed to demonstrate a significant decline of that activity over a 2-year follow-up period. In that study, the baseline and longitudinally examined cohort consisted of the same subjects. Despite the obvious differences in methodology and spatial parameters used in measuring the BOLD signals, the longitudinal study by Wolf et al. [2011] may further consolidate the notion of a lack of sensitivity in detecting BOLD signal changes occurring during a time frame that can be considered feasible for assessing the efficacy of an intervention in preHD.

The strength of our study lies in the application of three different analysis methods which provide for a more comprehensive interpretation of the data. This strength is complemented by the acquisition methodology used: the duration of the RS-scans (>6 min) and acquisition while

the patients have their eyes open provide the most robust estimates of functional connectivity as demonstrated by different studies [Van Dijk et al., 2010; Yan et al., 2009].

A limitation of this study is the loss of power due to the expansive testing of various networks and independent components. This expansive testing is however justified given the goal of finding robust and specific functional connectivity changes in preHD for usage as biomarker candidate in a clinical trial setting. Other possible limitations include transforming the data to an atlas volume instead of subject-native space, the relatively small number of tested subjects and possible confounding effects of dropouts, the conceivably short follow-up period in the preHD stage setting and not accounting for possibly confounding covariables such as depression scores in the analysis model.

Based on the results found in this study, the provisional conclusion is that RS-fMRI seems to lack sensitivity in detecting changes in functional connectivity in HD gene carriers prior to disease manifestation over a 3-year follow-up period. This conclusion applies to this selective group of participants and the particular analysis methods used in this study. Results from future longitudinal studies, such as the ongoing Track-On HD which has larger groups and more time points measured, should be awaited before articulating a definite recommendation on the possible utility of RS-fMRI as a biomarker tracking disease progression in preHD.

ACKNOWLEDGMENTS

TRACK-HD is supported by the CHDI Foundation, Inc., a not for profit organization dedicated to finding treatments for Huntington's disease. The authors thank Sarah Tabrizi, University College London, who is the global PI for TRACK-HD and clinical site PI for London, Hans Johnson, University of Iowa, responsible for producing morphometric regional measures of the brain and analysis protocols for medical imaging, and Doug Langbehn, University of Iowa, who is a biostatistician and played a key role in statistical design and data analysis of the TRACK-HD data. The authors also extend their gratitude to the TRACK-HD investigators responsible for collecting the data and to the study participants and their families. The authors declare that there are no conflicts of interest.

REFERENCES

- Aylward EH, Nopoulos PC, Ross CA, Langbehn DR, Pierson RK, Mills JA, Johnson HJ, Magnotta VA, Juhl AR, Paulsen JS (2011): Longitudinal change in regional brain volumes in prodromal Huntington disease. *J Neurol Neurosurg Psychiatry* 82: 405–410.
- Aylward EH, Liu D, Nopoulos PC, Ross CA, Pierson RK, Mills JA, Long JD, Paulsen JS (2012): Striatal volume contributes to the prediction of onset of Huntington disease in incident cases. *Biol Psychiatry* 71:822–828.
- Bai F, Watson DR, Shi Y, Wang Y, Yue C, YuhuanTeng, Wu D, Yuan Y, Zhang Z (2011): Specifically progressive deficits of brain functional marker in amnesic type mild cognitive impairment. *PLoS One* 6:e24271.
- Beckmann CF, Smith SM (2004): Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans Med Imaging* 23:137–152.
- Beckmann CF, DeLuca M, Devlin JT, Smith SM (2005): Investigations into resting-state connectivity using independent component analysis. *Philos Trans R Soc Lond B: Biol Sci* 360:1001–1013.
- Birn RM (2012): The role of physiological noise in resting-state functional connectivity. *Neuroimage* 62:864–870.
- Cole DM, Smith SM, Beckmann CF (2010): Advances and pitfalls in the analysis and interpretation of resting-state FMRI data. *Front Syst Neurosci* 4:8.
- Damoiseaux JS, Rombouts SA, Barkhof F, Scheltens P, Stam CJ, Smith SM, Beckmann CF (2006): Consistent resting-state networks across healthy subjects. *Proc Natl Acad Sci USA* 103: 13848–13853.
- Damoiseaux JS, Prater KE, Miller BL, Greicius MD (2012): Functional connectivity tracks clinical deterioration in Alzheimer's disease. *Neurobiol Aging* 33:828–830.
- Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ (2006): An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 31:968–980.
- Dumas EM, van den Bogaard SJ, Ruber ME, Reilman RR, Stout JC, Craufurd D, Hicks SL, Kennard C, Tabrizi SJ, van Buchem MA, van der Grond J, Roos RA (2012): Early changes in white matter pathways of the sensorimotor cortex in premanifest Huntington's disease. *Hum Brain Mapp* 33:203–212.
- Dumas EM, van den Bogaard SJ, Hart EP, Soeter RP, van Buchem MA, van der Grond J, Rombouts SA, Roos RA (2013): Reduced functional brain connectivity prior to and after disease onset in Huntington's disease. *NeuroImage: Clin* 2:377–384.
- Filippini N, MacIntosh BJ, Hough MG, Goodwin GM, Frisoni GB, Smith SM, Matthews PM, Beckmann CF, Mackay CE (2009): Distinct patterns of brain activity in young carriers of the APOE-epsilon4 allele. *Proc Natl Acad Sci USA* 106:7209–7214.
- Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005): The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci USA* 102:9673–9678.
- Frazier JA, Chiu S, Breeze JL, Makris N, Lange N, Kennedy DN, Herbert MR, Bent EK, Koneru VK, Dieterich ME, Hodge SM, Rauch SL, Grant PE, Cohen BM, Seidman LJ, Caviness VS, Biederman J (2005): Structural brain magnetic resonance imaging of limbic and thalamic volumes in pediatric bipolar disorder. *Am J Psychiatry* 162:1256–1265.
- Ghayoor A, Vaidya JG, Johnson HJ (2013): Development of a novel constellation based landmark detection algorithm. *Proc SPIE* 8669: id. 86693; doi:10.1117/12.2006471.
- Goldstein JM, Seidman LJ, Makris N, Ahern T, O'Brien LM, Caviness VS Jr, Kennedy DN, Faraone SV, Tsuang MT (2007): Hypothalamic abnormalities in schizophrenia: Sex effects and genetic vulnerability. *Biol Psychiatry* 61:935–945.
- Hadzi TC, Hendricks AE, Latourelle JC, Lunetta KL, Cupples LA, Gillis T, Mysore JS, Gusella JF, MacDonald ME, Myers RH, Vonsattel JP (2012): Assessment of cortical and striatal

- involvement in 523 Huntington disease brains. *Neurology* 79: 1708–1715.
- Jenkinson M, Bannister P, Brady M, Smith S (2002): Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* 17:825–841.
- Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM (2012): FSL. *Neuroimage* 62:782–790.
- Khalili-Mahani N, Zoethout RM, Beckmann CF, Baerends E, de Kam ML, Soeter RP, Dahan A, van Buchem MA, van Gerven JM, Rombouts SA (2012): Effects of morphine and alcohol on functional brain connectivity during “resting state”: A placebo-controlled crossover study in healthy young men. *Hum Brain Mapp* 33:1003–1018.
- Langbehn DR, Brinkman RR, Falush D, Paulsen JS, Hayden MR (2004): A new model for prediction of the age of onset and penetrance for Huntington’s disease based on CAG length. *Clin Genet* 65:267–277.
- Makris N, Goldstein JM, Kennedy D, Hodge SM, Caviness VS, Faraone SV, Tsuang MT, Seidman LJ (2006): Decreased volume of left and total anterior insular lobule in schizophrenia. *Schizophr Res* 83:155–171.
- McKeown MJ, Makeig S, Brown GG, Jung TP, Kindermann SS, Bell AJ, Sejnowski TJ (1998): Analysis of fMRI data by blind separation into independent spatial components. *Hum Brain Mapp* 6:160–188.
- Nichols TE, Holmes AP (2002): Nonparametric permutation tests for functional neuroimaging: A primer with examples. *Hum Brain Mapp* 15:1–25.
- Oakes TR, Fox AS, Johnstone T, Chung MK, Kalin N, Davidson RJ (2007): Integrating VBM into the General Linear Model with voxelwise anatomical covariates. *Neuroimage* 34:500–508.
- Paulsen JS, Nopoulos PC, Aylward E, Ross CA, Johnson H, Magnotta VA, Juhl A, Pierson RK, Mills J, Langbehn D, Nance M (2010): Striatal and white matter predictors of estimated diagnosis for Huntington disease. *Brain Res Bull* 82:201–207.
- Penney JB Jr, Vonsattel JP, MacDonald ME, Gusella JF, Myers RH (1997): CAG repeat number governs the development rate of pathology in Huntington’s disease. *Ann Neurol* 41:689–692.
- Poudel GR, Egan GF, Churchyard A, Chua P, Stout JC, Georgiou-Karistianis N (2014): Abnormal synchrony of resting state networks in premanifest and symptomatic Huntington disease: The IMAGE-HD study. *J Psychiatry Neurosci* 39:87–96.
- Roos RA (2010): Huntington’s disease: A clinical review. *Orphanet J Rare Dis* 5:40.
- Seibert TM, Majid DS, Aron AR, Corey-Bloom J, Brewer JB (2012): Stability of resting fMRI interregional correlations analyzed in subject-native space: A one-year longitudinal study in healthy adults and premanifest Huntington’s disease. *Neuroimage* 59: 2452–2463.
- Smith SM (2002): Fast robust automated brain extraction. *Hum Brain Mapp* 17:143–155.
- Smith SM, Nichols TE (2009): Threshold-free cluster enhancement: Addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* 44:83–98.
- Stoffers D, Sheldon S, Kuperman JM, Goldstein J, Corey-Bloom J, Aron AR (2010): Contrasting gray and white matter changes in preclinical Huntington disease: An MRI study. *Neurology* 74: 1208–1216.
- Tabrizi SJ, Langbehn DR, Leavitt BR, Roos RA, Durr A, Craufurd D, Kennard C, Hicks SL, Fox NC, Scahill RI, Borowsky B, Tobin AJ, Rosas HD, Johnson H, Reilmann R, Landwehrmeyer B, Stout JC (2009): Biological and clinical manifestations of Huntington’s disease in the longitudinal TRACK-HD study: Cross-sectional analysis of baseline data. *Lancet Neurol* 8:791–801.
- Tabrizi SJ, Scahill RI, Durr A, Roos RA, Leavitt BR, Jones R, Landwehrmeyer GB, Fox NC, Johnson H, Hicks SL, Kennard C, Craufurd D, Frost C, Langbehn DR, Reilmann R, Stout JC (2011): Biological and clinical changes in premanifest and early stage Huntington’s disease in the TRACK-HD study: The 12-month longitudinal analysis. *Lancet Neurol* 10:31–42.
- Tabrizi SJ, Reilmann R, Roos RA, Durr A, Leavitt B, Owen G, Jones R, Johnson H, Craufurd D, Hicks SL, Kennard C, Landwehrmeyer B, Stout JC, Borowsky B, Scahill RI, Frost C, Langbehn DR (2012): Potential endpoints for clinical trials in premanifest and early Huntington’s disease in the TRACK-HD study: Analysis of 24 month observational data. *Lancet Neurol* 11:42–53.
- Tabrizi SJ, Scahill RI, Owen G, Durr A, Leavitt BR, Roos RA, Borowsky B, Landwehrmeyer B, Frost C, Johnson H, Craufurd D, Reilmann R, Stout JC, Langbehn DR (2013): Predictors of phenotypic progression and disease onset in premanifest and early-stage Huntington’s disease in the TRACK-HD study: Analysis of 36-month observational data. *Lancet Neurol* 12: 637–649.
- The Huntington’s Disease Collaborative Research Group (1993): A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington’s disease chromosomes. *Cell* 72: 971–983.
- Unschuld PG, Joel SE, Liu X, Shanahan M, Margolis RL, Biglan KM, Bassett SS, Schretlen DJ, Redgrave GW, van Zijl PC, Pekar JJ, Ross CA (2012): Impaired cortico-striatal functional connectivity in prodromal Huntington’s Disease. *Neurosci Lett* 514: 204–209.
- Van den Bogaard SJ, Dumas EM, Ferrarini L, Milles J, van Buchem MA, van der Grond J, Roos RA (2011): Shape analysis of subcortical nuclei in Huntington’s disease, global versus local atrophy—Results from the TRACK-HD study. *J Neurol Sci* 307:60–68.
- Van Dijk KR, Hedden T, Venkataraman A, Evans KC, Lazar SW, Buckner RL (2010): Intrinsic functional connectivity as a tool for human connectomics: Theory, properties, and optimization. *J Neurophysiol* 103:297–321.
- Vonsattel JP, Keller C, Cortes Ramirez EP (2011): Huntington’s disease—Neuropathology. *Handb Clin Neurol* 100:83–100.
- Wolf RC, Sambataro F, Vasic N, Wolf ND, Thomann PA, Landwehrmeyer GB, Orth M (2011): Longitudinal functional magnetic resonance imaging of cognition in preclinical Huntington’s disease. *Exp Neurol* 231:214–222.
- Yan C, Liu D, He Y, Zou Q, Zhu C, Zuo X, Long X, Zang Y (2009): Spontaneous brain activity in the default mode network is sensitive to different resting-state conditions with limited cognitive load. *PLoS One* 4:e5743.
- Young KE, Johnson HJ (2013): Robust multi-site MR data processing: Iterative optimization of bias correction, tissue classification, and registration. *Front Neuroinform* 7:29.