Common Structural Correlates of Trait Impulsiveness and Perceptual Reasoning in Adolescence

Christina Schilling,^{1,2*} Simone Kühn,^{1,3} Alexander Romanowski,¹
Tobias Banaschewski,⁴ Alexis Barbot,⁵ Gareth J. Barker,⁶ Rüdiger Brühl,⁷
Christian Büchel,⁸ Katrin Charlet,¹ Patricia J. Conrod,⁶ Katharina Czech,¹
Jeff W. Dalley,^{9,10} Herta Flor,¹¹ Ines Häke,¹ Bernd Ittermann,⁷ Nikolay Ivanov,¹
Karl Mann,¹² Katharina Lüdemann,¹ Jean-Luc Martinot,^{13,14} Carla Palafox,¹
Tomas Paus,^{15,16,17} Jean-Baptiste Poline,⁵ Jan Reuter,¹ Marcella Rietschel,¹⁸
Trevor W. Robbins,¹⁰ Michael N. Smolka,^{19,20} Andreas Ströhle,¹
Bernadeta Walaszek,⁷ Norbert Kathmann,² Gunter Schumann,⁶ Andreas Heinz,¹
Hugh Garavan,²¹ and Jürgen Gallinat¹; and the IMAGEN consortium

¹Department of Psychiatry and Psychotherapy, Charité University Medicine Campus Mitte, Berlin, Germany

²Department of Psychology, Humboldt-Universität zu Berlin, Berlin, Germany

³Department of Experimental Psychology and Ghent, Institute for Functional and Metabolic Imaging, Ghent, Belgium

⁴Department of Child and Adolescent Psychiatry, Central Institute of Mental Health, Mannheim, Germany

⁵Neurospin, Commissariat à l'Energie Atomique, Paris, France

⁶Centre for Neuroimaging Sciences, King's College London, Institute of Psychiatry, London, United Kingdom

⁷Physikalisch-Technische Bundesanstalt, Berlin, Germany

⁸Institut für Systemische Neurowissenschaften, Universitätsklinikum Hamburg Eppendorf, Hamburg, Germany

⁹Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom

¹⁰Department of Experimental Psychology, Behavioural and Clinical Neurosciences Institute, University of Cambridge, Cambridge, United Kingdom

¹¹Department of Cognitive and Clinical Neuroscience, Central Institute of Mental Health, Mannheim, Germany

¹²Department of Addictive Behaviour and Addiction Medicine, Central Institute of Mental Health, Mannheim, Germany

¹³Neuroimaging & psychiatry, Institut National de la Santé et de la Recherche Médicale, INSERM Unit 1000 "Imaging and Psychiatry," University Paris Sud, Orsay, France

¹⁴AP-HP Department of Adolescent Psychopathology and Medicine, Maison de Solenn, University Paris Descartes, Paris, France

Contract grant sponsors: Studienstiftung des Deutschen Volkes, European Community's Sixth Framework Programme; Contract grant number: LSHM-CT-2007-037286; Contract grant sponsors: UK Department of Health NIHR-Biomedical Research Centre "Mental Health", The MRC programme grant "Developmental pathways into adolescents' substance abuse."

The IMAGEN study receives research funding from the European Community's Sixth Framework Programme (LSHM-CT-2007-037286) and is supported by the UK Department of Health NIHR-Biomedical Research Centre 'Mental Health' and the MRC programme grant "Developmental pathways into adolescents' substance abuse."

*Correspondence to: Christina Schilling, Charité Medicine Berlin, Clinic for Psychiatry and Psychotherapy, St. Hedwig Krankenhaus, Große Hamburger Str. 5-11, 10115 Berlin, Germany.

E-mail: christina.schilling@charite.de

Received for publication 20 February 2011; Revised 8 June 2011; Accepted 27 July 2011

DOI: 10.1002/hbm.21446

Published online 11 November 2011 in Wiley Online Library (wileyonlinelibrary.com).

¹⁵Population Neuroscience, Rotman Research Institute, University of Toronto, Toronto, Canada

¹⁶Cognitive Neuroscience Unit/Neuropsychology Department, Montreal Neurological Institute, McGill University, Montreal, Canada

¹⁷School of Psychology, University of Nottingham, Nottingham, United Kingdom

¹⁸Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Mannheim, Germany

¹⁹Department of Psychiatry and Psychotherapy, Technische Universität Dresden, Dresden, Germany

²⁰Neuroimaging Center, Systems Neuroscience, Department of Psychology, Technische Universität Dresden, Dresden, Germany

²¹Department of Psychology, Institute of Neuroscience, Trinity College Dublin, Dublin, Ireland

Abstract: *Background:* Trait impulsiveness is a potential factor that predicts both substance use and certain psychiatric disorders. This study investigates whether there are common structural cerebral correlates of trait impulsiveness and cognitive functioning in a large sample of healthy adolescents from the IMAGEN project. *Methods:* Clusters of gray matter (GM) volume associated with trait impulsiveness, Cloningers' revised temperament, and character inventory impulsiveness (TCI-R-I) were identified in a whole brain analysis using optimized voxel-based morphometry in 115 healthy 14-year-olds. The clusters were tested for correlations with performance on the nonverbal tests (Block Design, BD; Matrix Reasoning, MT) of the Wechsler Scale of Intelligence for Children IV reflecting perceptual reasoning. *Results:* Cloningers' impulsiveness (TCI-R-I) score was significantly inversely associated with GM volume in left orbitofrontal cortex (OFC). Frontal clusters found were positively correlated with performance in perceptual reasoning tasks (Bonferroni corrected). No significant correlations between TCI-R-I and perceptual reasoning were observed. *Conclusions:* The neural correlate of trait impulsiveness in the OFC matches an area where brain function has previously been related to inhibitory control. Additionally, orbitofrontal GM volume was associated with scores for perceptual reasoning. The data show for the first time structural correlates of both cognitive functioning and impulsiveness in healthy adolescent subjects. *Hum Brain Mapp* 34:374–383, 2013. © 2011 Wiley Periodicals, Inc.

Key words: trait; impulsiveness; structural correlates; frontal cortex; perceptual reasoning; adolescence

INTRODUCTION

Impulsiveness is a fundamental behavioral trait with substantial heritability [Hur and Bouchard, 1997; Pedersen et al., 1988; Seroczynski et al., 1999] and has a considerable impact on individual as well as on social life. For instance, impulsiveness is a predictor of behaviors such as aggression, delinquency, substance use, and the number of sexual partners [Flory et al., 2006; Miller et al., 2004]. In addition, impulsiveness has been associated with borderline personality disorder [Bagge et al., 2004; Conrod et al., 2000; Fossati et al., 2004; Links et al., 1999], substance abuse [Sher et al., 2000], attention-deficit/hyperactivity disorder [ADHD; Avila et al., 2004; Lijffijt et al., 2005], and suicidal behavior [Dougherty et al., 2004; Esposito et al., 2003; Swann et al., 2005; Yen et al., 2004]. Some evidence indicates that impulsiveness is a stronger predictor for antisocial behavior than other major personality dimensions such as extraversion/sociability and neuroticism/emotionality [Cale and Lilienfeld, 2006].

An extensive literature has focused on the association of impulsiveness with cognitive processing and intelligence

[Baron, 1982; Brebner, 1995], because Barratt [1967] described perceptual-motor performance as being related to impulsiveness. The link between high impulsiveness and impaired cognitive functioning has been related to deficits in attention, efficiency of information processing, and reasoning. Empirical evidence indicates that high impulsive individuals compared to low impulsive ones show worse performance in reaction time tasks in terms of reaction time and errors [Dickman, 1993, 2000; Marsh et al., 2002], working memory [Whitney et al., 2004], figural and numeric/alphabetical reasoning tasks [Schweizer, 2002], planning strategy [de Wit, 2007], and fluid or crystalline intelligence [Corr, 1998; de Wit, 2007; Harmon-Jones et al., 1997; Russo et al., 2008], although negative findings have also been published [Ashton et al., 2000; Austin et al., 2002]. For instance, despite high intelligence quotients (IQ) children with an impulsiveness-related disorder (ADHD) showed a poorer performance on the Block Design (BD), a subtest of the Wechsler Scale of Intelligence for Children (WISC) III [Antshel et al., 2007]. The effect of impulsiveness is also observed in animal models with several studies indicating that animals with pharmacologically

increased impulsiveness perform worse in cognitive tasks, which require them to wait or to process a large amount of information [Bizot and Thiebot, 1996]. Thus, impulsiveness appears to be an important modulatory factor in cognitive function.

The multifaceted construct of impulsiveness has been investigated as a personality measure using self-report questionnaire data as well as in the behavioral domain investigating disinhibition. Several subfactors have been described at the level of personality measures, but most evidence indicates three dimensions termed motor, attention, and nonplanning impulsiveness [Patton et al., 1995], which may form an underlying common trait in healthy individuals [Barratt, 1994; Cloninger, 1986]. This article focuses on impulsiveness as a trait, because personality-related dimensions of impulsiveness predict individual behavior [Perales et al., 2009; Romer et al., 2009], psychopathological syndromes, and mental disorders [Berlin et al., 2005; Solanto et al., 2009] and further provide the best empirical evidence for a high heritability [Hur and Bouchard, 1997; Pedersen et al., 1988; Seroczynski et al., 1999] as well as associations with genetic variants of particular neurotransmitter systems [Kazantseva et al., 2009; Munafo et al., 2008].

Functional neuroimaging studies on impulsiveness have tended to focus on behavioral inhibition, while many other dimensions of impulsiveness have remained unexplored. For instance, one popular but rather narrow conceptualization of impulsiveness, motor response inhibition as tested by Go/NoGo and stop signal tasks, has frequently shown activity in the right inferior frontal cortex and the subthalamic nucleus [STN; Congdon and Canli, 2008]. The inferior frontal cortex has been hypothesized as being essential for controlling behavioral inhibition and the STN for playing a central role in braking ongoing motor commands [Aron and Poldrack, 2005; Gerfen, 2000; Mink, 1996; Nambu et al., 2002]. Hence, when impulsiveness as a trait is targeted in fMRI, the results are restricted to correlational analyses with brain activation patterns related to specific disinhibition tasks. In this context, brain morphometry may be advantageous, because it is not restricted neither to certain paradigms tailored to specific regions of interest nor to specific operationalizations of impulsiveness.

Brain morphometry studies in psychiatric patients have reported associations between impulsiveness and gray matter (GM) volume of the orbitofrontal, cingulate, bilateral parietal, and temporal cortex as well as of the ventral striatum [Antonucci et al., 2006; Carmona et al., 2005; Lotfipour et al., 2009; Monkul et al., 2007; Schwartz et al., 2010]. Studies in healthy subjects are rare. Gardini and colleagues [Gardini et al., 2009] have shown a positive association with GM volume in frontal and posterior cingulate regions. Conversely, negative associations between nonplanning as well as motor impulsiveness and right ventromedial prefrontal cortex volume have been reported [Boes et al., 2009]. Similarly, identical aspects of impulsiveness

were negatively correlated with bilateral orbitofrontal cortex (OFC) GM volume [Matsuo et al., 2009]. In particular, middle frontal cortex volume has been shown to be related not only to inhibition [Boehler et al., 2010; Kuhn et al., 2009; Sharp et al., 2010], but also to attention [Andersson et al., 2009]. Thus, cerebral regions related to the inhibitory aspects of the impulsiveness trait are also critical for attention and executive components of cognitive processing. This structural link may provide an explanatory framework for the association between trait impulsiveness and cognitive performance.

This study focuses on the common structural underpinnings of both trait impulsiveness and related cognitive functions using optimized voxel-based morphometry (VBM) in a large sample of adolescents. This may be clinically relevant, because personality traits and intelligence in children explain to some degree the drinking and smoking behavior of the adults they become [Kubicka et al., 2001]. We investigated healthy 14-year-old subjects ($N = 115$), for whom mechanisms of impulsiveness may not be confounded with disease-specific variables or life style factors such as alcohol, nicotine, and illegal drug use, which, in turn, affect brain morphology [Gallinat et al., 2006; Lotfipour et al., 2009]. Impulsiveness was operationalized using a specific personality facet instead of broad personality concepts like novelty-seeking [Gardini et al., 2009] or a questionnaire targeting behavioral aspects such as the Pediatric Behavior Scale [Boes et al., 2009]. Cloninger's TCI-R-I was used, because it is based on a psychobiological model derived from empirical findings on personality, neurobiology and genetics [Cloninger et al., 1993]. The TCI-R-I appeared to be particularly suitable for our study, because it is a precisely defined personality facet based on a comprehensive model of personality. To assess cognitive functioning, while controlling for nuisance effects caused by verbal or numerical testing material [Goel, 2007], the nonverbal core tasks of the abbreviated WISC IV were chosen. More importantly, these tests are conceptualized to assess in particular abstract reasoning and executive dysfunctions [Szeremi et al., 2005], which were hypothesized to be affected by impulsiveness. Both nonverbal tasks (BD; Matrix Reasoning, MT) require attention, spatial reasoning, and response selection [Petermann and Ulrike, 2007; Wechsler, 2003].

Although there is some consensus for a prominent role of frontal and parietal structures in impulsiveness and cognitive function [Greening et al., 2011; Schneider et al., 2010; Westlye et al., 2011], we aimed to identify relevant structures by means of a whole brain analysis for trait impulsiveness and a subsequent analysis of the identified clusters with perceptual reasoning.

METHODS AND MATERIALS

Participants

One hundred and fifteen 14-year-olds (SD 0.32 years; 70 females) volunteered for this study within the scope of the

IMAGEN project, an European multicenter genetic-neuroimaging study in adolescence [Schumann et al., 2010]. Written informed consent was obtained from all participants as well as from their legal guardians. The adolescents were recruited from secondary schools in Berlin. The assessment was approved by both the ethics committee and by each head teacher of the school. All interested students were screened for MR contraindications. Furthermore, any participant with a medical condition such as a tumor, neurological disorders like epilepsy or mental-health problems like affective disorders was excluded. All participating students were assessed by means of both a self rating and two external ratings (parents; psychiatrists specialized in pediatrics) as part of a scale tailored to adolescents and based on ICD-10 as well as DSM-IV (The Development and Well-Being Assessment Interview, DAWBA; [Goodman et al., 2000]. Additionally, data on life time substance use were collected (*number of occasions in your whole lifetime you have had any alcoholic drinks* M 1.41, SD 1.48; [...] *used marijuana* M 0.04, SD 0.25; [...] *smoked cigarettes* M 0.78, SD 1.43, Scale: European School Survey Project on Alcohol and Drugs, ESPAD, www.espad.org).

Materials

Personality measure

Participants filled in the TCI-R [Cloninger, 1999] a computerized, self-rating personality questionnaire with the scale impulsiveness versus reflection as one of the novelty seeking facets (administered in German, [Richter et al., 1999]). Answers were given by means of a five-point Likert scale. By adding up the scores of the nine items of interest, a total was computed with higher values signifying higher impulsiveness.

Performance measures

Perceptual reasoning was assessed by means of two subscale scores according to the nonverbal core tests included in the abbreviated WISC IV [Wechsler, 2003] in a German version [Petermann and Ulrike, 2007]: BD and MT. Both tasks require attention, spatial reasoning, and response selection. BD involves putting together red-and-white blocks in a pattern so as to match to a displayed model. Speed is stressed, and some of the more difficult puzzles award bonuses for speed. MT asks participants to pick out of five images one fitting a shown array of pictures with one missing square. Sum scores, both in BD and MT, were converted to the more precise age equivalent scores, higher ones indicate higher performance.

MR Imaging

Structural MRI was performed on a General Electric SIGNA EXCITE 3T scanner (Milwaukee, WI) with a standard eight-channel head coil. High-resolution anatomical magnetic resonance images were obtained using a three-

dimensional T1-weighted magnetization prepared gradient-echo sequence (MPRAGE) based on the ADNI protocol (sequence version = adni14m4; www.adni-info.org; repetition time = 7.16 ms; echo time = 3.02 ms; flip angle = 8°; 256 × 256 × 166 matrix, 1.1 × 1.1 × 1.1 mm resolution).

Image Analysis

First, anatomical images were visually controlled for motion artifacts and gross structural abnormalities. The image processing followed the optimized VBM protocol [Good et al., 2001] implemented in Statistical Parametric Mapping (SPM) 8 (Wellcome Department of Neuroimaging, London, United Kingdom; <http://www.fil.ion.ucl.ac.uk/spm>) running on Matlab R2007a (MathWorks, Natick, MA). Images were preprocessed using the SPM8 modules segmentation and smoothing involving bias correction, spatial normalization, and smoothing with an 8-mm full-width-at-half-maximum Gaussian kernel by means of SPM8. In line with Good and colleagues [2001], a recursive process of segmentation, extraction, and normalization integrated in SPM8 was applied in our study. So, initially, structural MR images were segmented into gray and white matter images. Following, unconnected nonbrain voxels were removed from the segmented images, before images were normalized to gray/white matter templates. Resulting images were then subject to segmentation into gray and white matter, CSF, and non-CSF-partition, a second extraction step. Individual brains were normalized to Montreal Neurological Institute (MNI) space. Moreover, images were modulated in order to focus on volumetric differences instead of tissue concentration [Ashburner and Friston, 2000].

Statistical Analysis

At first, whole brain analysis on the association of GM volumes with impulsiveness (TCI-R-I) was modeled by means of voxelwise multiple regression. To control for nuisance effects arising from interindividual differences in local GM content due to modulation [Mechelli et al., 2005] total GM volume was included as a covariate of no interest besides age in days and gender [Good et al., 2001]. The total GM volume was estimated from the modulated and normalized GM images using Matlab (MathWorks) code (http://www.cs.ucl.ac.uk/staff/G.Ridgway/vbm/get_totals.m). Cluster criteria were a voxel threshold of $P < 0.001$ without correction for multiple comparisons but a cluster volume of >25 voxels. Standardized regression coefficients and t values are reported for the maximum voxel within the resulting brain region. Although such a liberal approach has been shown to be sensitive for detecting subtle structural differences, we additionally applied the more conservative cluster level threshold of $P < 0.05$ provided by SPM.

To explore their relationship to perceptual reasoning, for all clusters identified in the above analysis data were extracted by means of MarsBaR (<http://marsbar.sourceforge.net>). Respectively, for all clusters, the GM values of all

Figure 1.

The regions inversely (blue) and positively (red) correlated with trait impulsiveness; $z = -18$; $z = 57$; co-varied for total gray matter, age, and gender; ($p < 0.001$ uncorrected; $N = 115$). [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

voxels were averaged within each area. Subsequently, the resulting values were correlated separately with BD and MT scores (WISC IV) with a statistical threshold of $P < 0.05$, applying a Bonferroni correction for all seven clusters that were included in these analyses at $P < 0.003$ ($0.05/14$), in PASW Statistics 18 (SPSS, Chicago, IL).

RESULTS

Descriptive Data on Personality and Performance Measures

The participants' trait impulsiveness scores (TCI-R-I; $M = 26.14$, $SD = 4.35$) were normally distributed (Kolmogorov–Smirnov test $P = 0.38$) and comparable to the average in the general population [Elovainio et al., 2004]. Total scores achieved in the nonverbal tasks of the WISC IV indicated an average level of performance (BD $M = 10.73$, $SD = 2.75$; MT $M = 10.97$, $SD = 2.29$). Neither BD ($r = 0.04$, $P = 0.69$) nor MT ($r = 0.06$, $P = 0.55$) were correlated with TCI-R-I.

Whole Brain Analysis—Correlation Between GM Volume and Impulsiveness

The whole brain regression analysis showed a significant inverse relationship between GM volume in frontal cortex

areas and impulsiveness (TCI-R-I). In particular, the impulsiveness score was negatively correlated with both left and right OFC, right IFG as well as left middle frontal cortex at a threshold of $P < 0.001$ uncorrected (Fig. 1 and Table I). The more conservative, cluster level threshold of $P < 0.05$ confirmed the negative correlation for a cluster in the left orbitofrontal area ($-32, 36, -24$, MNI coordinates; Fig. 2).

Additionally, the whole brain analysis revealed that impulsiveness (TCI-R-I) was positively correlated with GM volume in the parietal, precentral, and temporal cortex (see Table II for details; Fig. 1). The positive associations were significant at $P < 0.001$ uncorrected, but did not survive the correction for multiple comparisons.

Association Between Neuronal Correlates of Impulsiveness and WISC

Although none of the pre-, postcentral, and temporal clusters resulting from the whole brain analysis of impulsiveness (TCI-R-I) were significantly correlated with the WISC subtests, all orbital frontal and the middle frontal regions showed a significant positive correlation. Although MT was statistically significantly ($P < 0.05$) associated with two out of three regions ($-32, 60, 2$ $r = 0.19$; $-32, 36, -24$ $r = 0.21$, MNI coordinates), BD was linked to all three

TABLE I. The Regions Inversely Correlated With Impulsiveness ($N = 115$)

	MNI coordinate ^a	Brodmann area	Cluster size	<i>t</i>	<i>P</i> uncorrected
Left middle frontal cortex	($-32, 60, 2$)	10	34	4.21	<0.001
Left OFC	($-32, 36, -24$)	11	82	4.06	<0.001
Right OFC/IFG orbitalis	($46, 42, -2$)	47/45	60	3.63	<0.001

Note: Analyses are covaried for total gray matter, age, and gender. OFC, orbitofrontal cortex; IFG, inferior frontal gyrus.

^aMNI coordinates of the voxel of maximal statistical significance.

Figure 2.

Correlation between average gray matter volume of the cluster in the left orbitofrontal area ($-32, 36, -24$, MNI coordinates) and trait impulsiveness. Co-varied for total gray matter, age, and gender; ($p < 0.05$ cluster level threshold; $N = 115$).

clusters ($-32, 60, 2$ $r = 0.31$; $-32, 36, -24$ $r = 0.32$; $46, 42, -2$ $r = 0.27$, MNI coordinates). Two, out of the three, correlations of BD with GM survived a Bonferroni correction at $P < 0.003$, indicating two left frontal clusters including the left orbitofrontal area ($-32, 60, 2$; $-32, 36, -24$, MNI coordinates; Fig. 3). The explained variance ranged from 4 to 10%.

DISCUSSION

The present study is the first investigation on morphometric correlates of trait impulsiveness in a large sample of healthy 14-year-olds without psychiatric disorders, nicotine, alcohol (max. one drink during their whole life), or any illegal drug use (IMAGEN, [Schumann et al., 2010]). The results provide evidence for a link between trait impulsiveness and structures in the left OFC. Exploratory analyses further suggest associations with inferior as well as middle frontal, pre-, and postcentral cortical areas, which have previously been attributed to inhibitory control of action or for motor and somatosensory processing respectively. In particular for OFC, an association with perceptual reasoning was found

indicating a structural correlate of impaired cognitive function in high impulsive subjects.

The inverse correlation between trait impulsiveness and GM volume of the left OFC is compatible with reports in nonclinical adult samples [Kumari et al., 2009; Matsuo et al., 2009]. GM volumes of the bilateral OFC have been found to be inversely correlated with the BIS total score as well as the subfactors nonplanning and motor impulsiveness [Matsuo et al., 2009]. However, because impulsiveness and substance use are associated in adult samples, those investigations cannot rule out a confounding effect on GM volume as described previously [Gallinat et al., 2006; Kühn et al., 2010]. Consistently, MRI studies in clinical, adult populations stress a link between reduced OFC GM volume and disorders comprising impulsive symptoms [Hesslinger et al., 2002; Ogura et al., 2011]. Thus, conversely, our findings may indicate that reduced GM volume in adults might not necessarily be a consequence of their illness/substance use [Schwartz et al., 2010]. In the fMRI literature, the OFC has been reported to be active during response inhibition [Horn et al., 2003]. Beyond its involvement in motor impulsiveness, OFC has been found to be associated with higher cognitive functions such as decision making [Schoenbaum et al., 2006; Wallis, 2007].

Supplementary considering the results of the exploratory analyses may help to understand the role of frontal structures related to the phenomenon impulsiveness. In addition to the OFC, the IFG has also been associated with the inhibitory control of action. Aron and Poldrack [2005] have shown that particularly the activity of the right IFG was correlated with response inhibition, a causal role for which had been suggested by previous lesion studies [Aron et al., 2004]. Similarly, the left middle frontal cortex, which we also found to be correlated negatively with impulsiveness in an exploratory analysis, is known to be important for both inhibitory control [Sharp et al., 2010] and cognitive functions such as the regulation of executive attention [Andersson et al., 2009]. However, in the current analysis, only correlations with left frontal clusters survived the cluster level threshold. The observed correlations and possible consequences for brain function should be interpreted with caution. Moreover, studies of neural plasticity in humans (e.g., hippocampal volumes on taxi drivers [Maguire et al., 2000]) suggest that such plastic changes may sometimes be a consequence of experience rather than a causal factor in biologically predetermined behavior.

TABLE II. The Regions Positively Correlated With Impulsiveness ($N = 115$)

	MNI coordinate ^a	Brodmann area	Cluster size	<i>t</i>	<i>P</i> uncorrected
Right postcentral gyrus	(48, -28, 58)	1	45	3.97	<0.001
Right precentral gyrus	(32, -22, 52)	4	32	3.90	<0.001
Left precentral gyrus	(-36, -18, 58)	4	60	3.89	<0.001
Left superior temporal cortex	(-60, -10, 10)	42	34	3.78	<0.001

Note: Analyses are covaried for total grey matter, age, and gender.

^aMNI coordinates of the voxel of maximal statistical significance.

Figure 3.

Correlation between average grey matter volume of the cluster in the left orbitofrontal area ($-32, 36, -24$, MNI coordinates) and Block Design. Covaried for total gray matter, age, and gender; ($p < 0.003$ Bonferroni corrected; $N = 115$).

In line with the general role that prefrontal structures play in directing attention [Mesulam, 1981], Frangou and colleagues [2004] have reported a positive association between IQ and GM density of the OFC in healthy adolescents and young adults. Consistently, Russo and colleagues [2008] showed that highly impulsive individuals had difficulties to inhibit irrelevant information in intelligence tests. However, despite some evidence for the involvement of the OFC and of the middle frontal cortex in both impulsiveness and cognitive functions as discussed earlier, relatively little is known about overlapping brain structures of personality dimensions of impulsiveness and mental abilities. The positive correlations between the subtests of the WISC IV and the frontal clusters of GM reduction identified in this study support the hypothesis of brain areas being involved in both impulsiveness and cognitive functioning. Both subscale scores, BD and MT, have been conceptualized as performance measures of perceptual reasoning. In particular, MT is thought to assess attention and executive functioning, which have been considered to engage aspects of fluid intelligence by some researchers. Even though Tranel and colleagues [2008] did not find a connection between fluid intelligence and prefrontal lobe damage, our results are in line with studies relating perceptual reasoning to OFC function [Sabbagh, 2004] and the medial frontal cortex [Prado and Noveck, 2007]. Similarly to MT, BD has been conceptualized as assessing abstract reasoning, namely spatial cognitive ability. Interestingly, impulsiveness was found to be especially associated with spatial cognitive function. For instance, similar to the negative correlation between BD and right

IFG orbitalis shown in our study, spatial working memory performance has been found to be associated with right IFG damage in adults with an impulsiveness-related disorder (ADHD; [Clark et al., 2007]. With regard to spatial attention, the activity of the OFC, as part of a circuitry critical for selective allocation of attentional resources, has especially been found to be associated with the speed of attentional shifts [Mohanty et al., 2008]. However, in additionally performed whole brain analyses, no correlative associations between the perceptual reasoning subtests and those prefrontal areas were found in our study. In sum, the correlations between the frontal clusters and the WISC subscales discussed might suggest frontal structures associated both with trait impulsiveness and aspects of the participants' behavior related to impulsiveness [Szeremi et al., 2005].

Furthermore, in uncorrected whole brain analyses, we found higher impulsiveness to be associated with larger GM volumes of primarily the bilateral precentral gyrus as well as the parietal and the temporal cortex. In particular, the correlations with primary motor cortex (BA 4) GM volume suggest an association with motor functions [Ellermann et al., 1998; Gelnar et al., 1999]. Cloninger [1986] has defined impulsiveness as one of the facets of the personality trait novelty seeking, described as a so-called temperament actively searching for particularly stimulating, novel environments, as shown for adolescents' impulsiveness predicting risky behavior [Zimmerman, 2010]. Furthermore, it has been shown that motor activity, such as learning to juggle, can result in GM volume increase in the motor cortex [Driemeyer et al., 2008]. The greater exploratory activity of the high impulsives is further presumably associated with more stimulating experiences from other different modalities besides kinaesthetic and proprioceptive feedback. Structural neuroimaging studies have provided evidence for the fact that exposure to acoustic stimuli can alter brain structure [Golestani et al., 2002]. Hence, it appears plausible, even though speculative, that the clusters of the somatosensory cortex and secondary auditory cortex found in our study might suggest a similar association driven by heightened impulsiveness and hence increased exploratory behavior.

Some methodological limitations of this study shall be noted. Although, there is a more recent junior TCI scale for children and adolescents [Luby et al., 1999], the TCI-R was used. This revised scale has been specifically modified for the purposes of neuroimaging and repeatedly applied previously [Cloninger, 2008]. Furthermore, participants have not been assessed with the standard instrument, the Structural Criteria Interview for Diagnosis nonpatient version. Yet, based on both a self-rating and external ratings (parents; psychiatrists specialized in pediatrics) as part of a scale developed for children and adolescents (DAWBA, [Goodman et al., 2000], volunteers with psychiatric disorders according to ICD-10 and DSM-IV were excluded. Moreover, instead of more objective measures such as urine drug screenings the well established self-rating scale

ESPAD (www.espad.org) was chosen, because it provides information on, for example, life time alcohol use. To minimize social desirability biases, participants got an individual, but anonymous online login to fill in the computerized questionnaire on their own in an undisturbed environment.

In summary, our findings support an association between higher trait impulsiveness and smaller GM volume of the left OFC. For the first time, morphological correlates of the impulsiveness personality facet have been identified in an adolescent, healthy, and substance naive sample. This study further suggests possible neural substrates of an important risk factor for various psychiatric disorders like substance abuse and potentially relevant for defining a neural endophenotype for such disorders. For instance, novelty seeking, a personality dimension comprising impulsiveness, measured during childhood predicted alcohol abuse in young adults [Cloninger et al., 1988]. We have also shown that predictive studies might benefit from focusing on possible interactions of impulsiveness with aspects of cognitive functioning.

REFERENCES

- Andersson M, Ystad M, Lundervold A, Lundervold AJ (2009): Correlations between measures of executive attention and cortical thickness of left posterior middle frontal gyrus—A dichotic listening study. *Behav Brain Funct* 5:41.
- Antonucci AS, Gansler DA, Tan S, Bhadelia R, Patz S, Fulwiler C (2006): Orbitofrontal correlates of aggression and impulsivity in psychiatric patients. *Psychiatry Res* 147:213–220.
- Antshel KM, Faraone SV, Stallone K, Nave A, Kaufmann FA, Doyle A, Fried R, Seidman L, Biederman J (2007): Is attention deficit hyperactivity disorder a valid diagnosis in the presence of high IQ? Results from the MGH longitudinal family studies of ADHD. *J Child Psychol Psychiatry* 48:687–694.
- Aron AR, Poldrack RA (2005): The cognitive neuroscience of response inhibition: Relevance for genetic research in attention-deficit/hyperactivity disorder. *Biol Psychiatry* 57:1285–1292.
- Aron AR, Robbins TW, Poldrack RA (2004): Inhibition and the right inferior frontal cortex. *Trends Cogn Sci* 8:170–177.
- Ashburner J, Friston KJ (2000): Voxel-based morphometry—The methods. *Neuroimage* 11(6, Pt 1):805–821.
- Ashton MC, Lee K, Vernon PA, Jang KL (2000): Fluid intelligence, crystallized intelligence, and the openness/intellect factor. *J Res Personal* 34:198–207.
- Austin EJ, Deary IJ, Whiteman MC, Fowkes FGR, Pedersen L, Rabbitt P, Bent N, McInnes L (2002): Relationship between ability and personality: Does intelligence contribute positively to personal and social adjustment? *Personal Indiv Differ* 32:1391–1411.
- Avila C, Cuenca I, Felix V, Parcet MA, Miranda A (2004): Measuring impulsivity in school-aged boys and examining its relationship with ADHD and ODD ratings. *J Abnorm Child Psychol* 32:295–304.
- Bagge C, Nickell A, Stepp S, Durrett C, Jackson K, Trull TJ (2004): Borderline personality disorder features predict negative outcomes 2 years later. *J Abnorm Psychol* 113:279–288.
- Baron. 1982. Personality and intelligence. In: Sternberg RJ, editor. *Handbook of Human Intelligence*. New York: Cambridge University Press.
- Barratt E. 1994. Impulsiveness and aggression. In: Monahan JSH, editor. *Violence and Mental Disorder: Developments in Risk Assessments*. Chicago, IL: University of Chicago Press. pp 61–79.
- Barratt ES (1967): Perceptual-motor performance related to impulsiveness and anxiety. *Percept Mot Skills* 25:485–492.
- Berlin HA, Rolls ET, Iversen SD (2005): Borderline personality disorder, impulsivity, and the orbitofrontal cortex. *Am J Psychiatry* 162:2360–2373.
- Bizot JC, Thiebot MH (1996): Impulsivity as a confounding factor in certain animal tests of cognitive function. *Brain Res Cogn Brain Res* 3:243–250.
- Boehler CN, Appelbaum LG, Krebs RM, Hopf JM, Woldorff MG. (2010): Pinning down response inhibition in the brain—Conjunction analyses of the Stop-signal task. *Neuroimage* 52:1621–1632.
- Boes AD, Bechara A, Tranel D, Anderson SW, Richman L, Nopoulos P (2009): Right ventromedial prefrontal cortex: A neuroanatomical correlate of impulse control in boys. *Soc Cogn Affect Neurosci* 4:1–9.
- Brebner S. 1995. Theoretical and empirical relationships between personality and intelligence. In: Saklofske MZDH, editor. *International Handbook of Personality and Intelligence (Language of Science)*. Berlin: Springer. pp 322–348.
- Cale EM, Lilienfeld SO (2006): Psychopathy factors and risk for aggressive behavior: A test of the “threatened egotism” hypothesis. *Law Hum Behav* 30:51–74.
- Carmona S, Vilarroya O, Bielsa A, Tremols V, Soliva JC, Rovira M, Tomas J, Raheb C, Gispert JD, Batlle S, et al. (2005): Global and regional gray matter reductions in ADHD: A voxel-based morphometric study. *Neurosci Lett* 389:88–93.
- Clark L, Blackwell AD, Aron AR, Turner DC, Dowson J, Robbins TW, Sahakian BJ (2007): Association between response inhibition and working memory in adult ADHD: A link to right frontal cortex pathology? *Biol Psychiatry* 61:1395–1401.
- Cloninger CR (1986): A unified biosocial theory of personality and its role in the development of anxiety states. *Psychiatr Dev* 4:167–226.
- Cloninger CR (1999): *The Temperament and Character Inventory-Revised*. St. Louis, MO: Center for Psychobiology of Personality.
- Cloninger CR. (2008): The psychobiological theory of temperament and character: Comment on Farmer and Goldberg (2008). *Psychol Assess* 20:292–299; discussion 300–304.
- Cloninger CR, Sigvardsson S, Bohman M (1988): Childhood personality predicts alcohol abuse in young adults. *Alcohol Clin Exp Res* 12:494–505.
- Cloninger CR, Svrakic DM, Przybeck TR (1993): A psychobiological model of temperament and character. *Arch Gen Psychiatry* 50:975–990.
- Congdon E, Canli T (2008): A neurogenetic approach to impulsivity. *J Pers* 76:1447–1484.
- Conrod PJ, Pihl RO, Stewart SH, Dongier M (2000): Validation of a system of classifying female substance abusers on the basis of personality and motivational risk factors for substance abuse. *Psychol Addict Behav* 14:243–256.
- Corr PJ, Kumari V (1998): Impulsivity, time of day and stress: Effects on intelligence test performance. *J Res Personal* 32:1–12.
- de Wit H, Flory JD, Acheson A, McCloskey M, Manuck SB (2007): IQ and nonplanning impulsivity are independently associated with delay discounting in middle-aged adults. *Personal Indiv Differ* 42:111–121.

- Dickmann S (1993): Impulsivity and information processing. In: MCCown WG JJ, Shure MB, editors. *The Impulsive Client: Theory and Treatment*. Washington D.C.: American Psychological Association. pp 151–184.
- Dickman S (2000): Impulsivity, arousal, and attention. *Personal Indiv Differ* 28:563–582.
- Dougherty DM, Mathias CW, Marsh DM, Moeller FG, Swann AC (2004): Suicidal behaviors and drug abuse: Impulsivity and its assessment. *Drug Alcohol Depend* 76 (Suppl):S93–S105.
- Driemeyer J, Boyke J, Gaser C, Buchel C, May A (2008): Changes in gray matter induced by learning–revisited. *PLoS One* 3:e2669.
- Ellermann JM, Siegal JD, Strupp JP, Ebner TJ, Ugurbil K (1998): Activation of visuomotor systems during visually guided movements: A functional MRI study. *J Magn Reson* 131:272–285.
- Elovainio M, Kivimäki M, Puttonen S, Heponiemi T, Pulkki L, Keltikangas-Järvinen L (2004): Temperament and depressive symptoms: A population-based longitudinal study on Cloninger's psychobiological temperament model. *J Affect Disord* 83:227–232.
- Esposito C, Spirito A, Boergers J, Donaldson D (2003): Affective, behavioral, and cognitive functioning in adolescents with multiple suicide attempts. *Suicide Life Threat Behav* 33:389–399.
- Flory K, Molina BS, Pelham WE, Jr., Gnagy E, Smith B (2006): Childhood ADHD predicts risky sexual behavior in young adulthood. *J Clin Child Adolesc Psychol* 35:571–577.
- Fossati A, Barratt ES, Carretta I, Leonardi B, Grazioli F, Maffei C (2004): Predicting borderline and antisocial personality disorder features in nonclinical subjects using measures of impulsivity and aggressiveness. *Psychiatry Res* 125:161–170.
- Frangou S, Chitins X, Williams SC (2004): Mapping IQ and gray matter density in healthy young people. *Neuroimage* 23:800–805.
- Gallinat J, Meisenzahl E, Jacobsen LK, Kalus P, Bierbrauer J, Kienast T, Witthaus H, Leopold K, Seifert F, Schubert F, et al. (2006): Smoking and structural brain deficits: A volumetric MR investigation. *Eur J Neurosci* 24:1744–1750.
- Gardini S, Cloninger CR, Venneri A (2009): Individual differences in personality traits reflect structural variance in specific brain regions. *Brain Res Bull* 79:265–270.
- Gelnar PA, Krauss BR, Sheeha PR, Szeverenyi NM, Apkarian AV (1999): A comparative fMRI study of cortical representations for thermal painful, vibrotactile, and motor performance tasks. *Neuroimage* 10:460–482.
- Gerfen CR (2000): Molecular effects of dopamine on striatal-projection pathways. *Trends Neurosci* 23(10 Suppl):S64–S70.
- Goel V (2007): Anatomy of deductive reasoning. *Trends Cogn Sci* 11:435–441.
- Golestani N, Paus T, Zatorre RJ (2002): Anatomical correlates of learning novel speech sounds. *Neuron* 35:997–1010.
- Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS (2001): A voxel-based morphometric study of ageing in 465 normal adult human brains. *Neuroimage* 14(1, Pt 1):21–36.
- Goodman R, Ford T, Richards H, Gatward R, Meltzer H (2000): The development and well-being assessment: Description and initial validation of an integrated assessment of child and adolescent psychopathology. *J Child Psychol Psychiatry* 41:645–655.
- Greening SG, Finger EE, Mitchell DG (2011): Parsing decision making processes in prefrontal cortex: Response inhibition, overcoming learned avoidance, and reversal learning. *Neuroimage* 54:1432–1441.
- Harmon-Jones E, Barratt ES, Wigg C (1997): Impulsiveness, aggression, reading, and the P300 of the event-related potential. *Personal Indiv Differ* 22:439–445.
- Hesslinger B, Tebartz van Elst L, Thiel T, Haegele K, Hennig J, Ebert D (2002): Fronto-orbital volume reductions in adult patients with attention deficit hyperactivity disorder. *Neurosci Lett* 328:319–321.
- Horn NR, Dolan M, Elliott R, Deakin JF, Woodruff PW (2003): Response inhibition and impulsivity: An fMRI study. *Neuropsychologia* 41:1959–1966.
- Hur YM, Bouchard TJ Jr (1997): The genetic correlation between impulsivity and sensation seeking traits. *Behav Genet* 27:455–463.
- Kazantseva AV, Gaisina DA, Malykh SB, Khusnutdinova EK (2009): [Role of dopamine transporter gene (DAT1) polymorphisms in personality traits variation]. *Genetika* 45:1110–1117.
- Kubicka L, Matejcek Z, Dytrych Z, Roth Z (2001): IQ and personality traits assessed in childhood as predictors of drinking and smoking behaviour in middle-aged adults: A 24-year follow-up study. *Addiction* 96:1615–1628.
- Kuhn S, Haggard P, Brass M (2009): Intentional inhibition: How the “veto-area” exerts control. *Hum Brain Mapp* 30:2834–2843.
- Kühn S, Schubert F, Gallinat J (2010): Reduced thickness of the orbitofrontal cortex in smokers. *Biological Psychiatry* 68:1061–1065.
- Kumari V, Barkataki I, Goswami S, Flora S, Das M, Taylor P (2009): Dysfunctional, but not functional, impulsivity is associated with a history of seriously violent behaviour and reduced orbitofrontal and hippocampal volumes in schizophrenia. *Psychiatry Res* 173:39–44.
- Lijffijt M, Kenemans JL, Verbaten MN, van Engeland H (2005): A meta-analytic review of stopping performance in attention-deficit/hyperactivity disorder: Deficient inhibitory motor control? *J Abnorm Psychol* 114:216–222.
- Links PS, Heslegrave R, van Reekum R (1999): Impulsivity: Core aspect of borderline personality disorder. *J Pers Disord* 13:1–9.
- Lotfipour S, Ferguson E, Leonard G, Perron M, Pike B, Richer L, Seguin JR, Toro R, Veillette S, Pausova Z, et al. (2009): Orbitofrontal cortex and drug use during adolescence: Role of prenatal exposure to maternal smoking and BDNF genotype. *Arch Gen Psychiatry* 66:1244–1252.
- Luby JL, Svrakic DM, McCallum K, Przybeck TR, Cloninger CR (1999): The Junior Temperament and Character Inventory: Preliminary validation of a child self-report measure. *Psychol Rep* 84(3, Pt 2):1127–1138.
- Maguire EA, Gadian DG, Johnsrude IS, Good CD, Ashburner J, Frackowiak RS, Frith CD (2000): Navigation-related structural change in the hippocampi of taxi drivers. *Proc Natl Acad Sci USA* 97:4398–4403.
- Marsh DM, Dougherty DM, Mathias CW, Moeller FG, Hicks LR (2002): Comparison of women with high and low trait impulsivity using behavioral models of response-disinhibition and reward-choice. *Personal Indiv Differ* 33:1291–1310.
- Matsuo K, Nicoletti M, Nemoto K, Hatch JP, Peluso MA, Nery FG, Soares JC (2009): A voxel-based morphometry study of frontal gray matter correlates of impulsivity. *Hum Brain Mapp* 30:1188–1195.
- Mechelli AP, Price CJ, Friston KJ, Ashburner J (2005): Voxel-based morphometry of the human brain: Methods and applications. *Curr Med Imaging Rev* 1:105–113.
- Mesulam MM (1981): A cortical network for directed attention and unilateral neglect. *Ann Neurol* 10:309–325.
- Miller E, Stephen J, Tudway J (2004): Assessing the component structure of four self-report measures of impulsivity. *Personal Indiv Differ* 37:349–358.
- Mink JW (1996): The basal ganglia: Focused selection and inhibition of competing motor programs. *Prog Neurobiol* 50:381–425.

- Mohanty A, Gitelman DR, Small DM, Mesulam MM (2008): The spatial attention network interacts with limbic and monoaminergic systems to modulate motivation-induced attention shifts. *Cereb Cortex* 18:2604–2613.
- Monkul ES, Hatch JP, Nicoletti MA, Spence S, Brambilla P, Lacerda AL, Sassi RB, Mallinger AG, Keshavan MS, Soares JC (2007): Fronto-limbic brain structures in suicidal and non-suicidal female patients with major depressive disorder. *Mol Psychiatry* 12:360–366.
- Munafò MR, Yalcin B, Willis-Owen SA, Flint J (2008): Association of the dopamine D4 receptor (DRD4) gene and approach-related personality traits: Meta-analysis and new data. *Biol Psychiatry* 63:197–206.
- Nambu A, Tokuno H, Takada M (2002): Functional significance of the cortico-subthalamo-pallidal ‘hyperdirect’ pathway. *Neurosci Res* 43:111–117.
- Ogura K, Fujii T, Abe N, Hosokai Y, Shinohara M, Takahashi S, Mori E (2011): Small gray matter volume in orbitofrontal cortex in Prader-Willi syndrome: A voxel-based MRI study. *Hum Brain Mapp* 32:1059–1066.
- Patton JH, Stanford MS, Barratt ES (1995): Factor structure of the Barratt impulsiveness scale. *J Clin Psychol* 51:768–774.
- Pedersen NL, Plomin R, McClearn GE, Friberg L (1988): Neuroticism, extraversion, and related traits in adult twins reared apart and reared together. *J Pers Soc Psychol* 55:950–957.
- Perales JC, Verdejo-García A, Moya M, Lozano O, Perez-García M (2009): Bright and dark sides of impulsivity: Performance of women with high and low trait impulsivity on neuropsychological tasks. *J Clin Exp Neuropsychol* 31:927–944.
- Petermann FP, Ulrike. 2007. Hamburg-Wechsler-Intelligenztest für Kinder-IV Manual Übersetzung und Adaptation der WISC-IV von David Wechsler. Bern: Verlag Hans Huber, Hogrefe AG, Testverlag.
- Prado J, Noveck IA (2007): Overcoming perceptual features in logical reasoning: A parametric functional magnetic resonance imaging study. *J Cogn Neurosci* 19:642–657.
- Richter J, Eisemann M, Richter G, Cloninger CR. 1999. Das Temperament- und Charakterinventar. Frankfurt/M.: Swets Test Services.
- Romer D, Betancourt L, Giannetta JM, Brodsky NL, Farah M, Hurt H (2009): Executive cognitive functions and impulsivity as correlates of risk taking and problem behavior in preadolescents. *Neuropsychologia* 47:2916–2926.
- Russo PM, De Pascalis V, Varriale V, Barratt ES (2008): Impulsivity, intelligence and P300 wave: An empirical study. *Int J Psychophysiol* 69:112–118.
- Sabbagh MA (2004): Understanding orbitofrontal contributions to theory-of-mind reasoning: Implications for autism. *Brain Cogn* 55:209–219.
- Schneider MF, Krick CM, Retz W, Hengesbach G, Retz-Junginger P, Reith W, Rosler M (2010): Impairment of fronto-striatal and parietal cerebral networks correlates with attention deficit hyperactivity disorder (ADHD) psychopathology in adults—a functional magnetic resonance imaging (fMRI) study. *Psychiatry Res* 183:75–84.
- Schoenbaum G, Roesch MR, Stalnaker TA (2006): Orbitofrontal cortex, decision-making and drug addiction. *Trends Neurosci* 29:116–124.
- Schumann G, Loth E, Banaschewski T, Barbot A, Barker G, Büchel C, Conrod PJ, Dalley JF, H, Gallinat J, Garavan H, Heinz A, Itterman B, Lathrop M, Mallik C, Mann K, Martinot J-L, Paus T, Poline J-B, Robbins TW, Rietschel M, Reed L, Smolka M, Spanagel RC, Speiser C, Stephens DN, Strohle A, Struve M; the IMAGEN consortium. (2010): The IMAGEN study: Reinforcement-related behaviour in normal brain function and psychopathology. *Mol Psychiatry* 1–12.
- Schwartz DL, Mitchell AD, Lahna DL, Luber HS, Huckans MS, Mitchell SH, Hoffman WF (2010): Global and local morphometric differences in recently abstinent methamphetamine-dependent individuals. *Neuroimage* 50:1392–1401.
- Schweizer K (2002): Does impulsivity influence performance in reasoning. *Personal Individ Differ* 33:1031–1043.
- Seroczynski AD, Bergeman CS, Coccaro EF (1999): Etiology of the impulsivity/aggression relationship: Genes or environment? *Psychiatry Res* 86:41–57.
- Sharp DJ, Bonnelle V, De Boissezon X, Beckmann CF, James SG, Patel MC, Mehta MA (2010): Distinct frontal systems for response inhibition, attentional capture, and error processing. *Proc Natl Acad Sci USA* 107:6106–6111.
- Sher KJ, Bartholow BD, Wood MD (2000): Personality and substance use disorders: A prospective study. *J Consult Clin Psychol* 68:818–829.
- Solanto MV, Schulz KP, Fan J, Tang CY, Newcorn JH (2009): Event-related fMRI of inhibitory control in the predominantly inattentive and combined subtypes of ADHD. *J Neuroimaging* 19:205–212.
- Swann AC, Dougherty DM, Pazzaglia PJ, Pham M, Steinberg JL, Moeller FG (2005): Increased impulsivity associated with severity of suicide attempt history in patients with bipolar disorder. *Am J Psychiatry* 162:1680–1687.
- Szeremi A, Tarnok Z, Farkas L, Dotzi J, Gadoros J (2005): [Neurocognitive symptoms of childhood attention deficit hyperactivity disorder]. *Psychiatr Hung* 20:299–311.
- Tranel D, Manzel K, Anderson SW (2008): Is the prefrontal cortex important for fluid intelligence? A neuropsychological study using Matrix Reasoning. *Clin Neuropsychol* 22:242–261.
- Wallis JD (2007): Orbitofrontal cortex and its contribution to decision-making. *Annu Rev Neurosci* 30:31–56.
- Wechsler A. 2003. WISC-IV. Technical and Interpretive Manual. San Antonio, TX: Psychological Association.
- Westlye LT, Grydeland H, Walhovd KB, Fjell AM. (2011): Associations between regional cortical thickness and attentional networks as measured by the attention network test. *Cereb Cortex* 21:345–356.
- Whitney P, Jameson T, Hinson JM (2004): Impulsiveness and executive control of working memory. *Personal Individ Differ* 37:417–428.
- Yen S, Shea MT, Sanislow CA, Grilo CM, Skodol AE, Gunderson JG, McGlashan TH, Zanarini MC, Morey LC (2004): Borderline personality disorder criteria associated with prospectively observed suicidal behavior. *Am J Psychiatry* 161:1296–1298.
- Zimmerman G (2010): Risk perception, emotion regulation and impulsivity as predictors of risk behaviours among adolescents in Switzerland. *J Youth Stud* 13:83–99.