Distributed Cell Assemblies for General Lexical and Category-Specific Semantic Processing as Revealed by fMRI Cluster Analysis

Friedemann Pulvermüller,* Ferath Kherif, Olaf Hauk,
Bettina Mohr, and Ian Nimmo-Smith

Medical Research Council, Cognition and Brain Sciences Unit, Cambridge, United Kingdom

Abstract: Here, we ask whether frontotemporal cortex is functionally dissociated into distributed lexical and category-specific semantic networks. To this end, fMRI activation patterns elicited during the processing of words from different semantic categories were categorized using *k*-means cluster algorithms. Results showed a distributed pattern of inferiorfrontal, superior temporal, and fusiform activation shared by different word categories. This shared activation contrasted with patterns of category-specific semantic activation in widely distributed neural systems. Clustering revealed congruent functional specificity of focal area activations in frontal and temporal cortex; thus suggesting a correspondence between functional partitionings of frontocentral mirror neuron systems and those of inferior temporal lexical and semantic circuits. Action words related to the face, arms, and legs specifically activated the motor system in a somatotopic manner, whereas form-related words activated prefrontal areas. Similar functional specificity was evident in temporal cortex, where a different semantic topography emerged for form- and action-related words. Results were replicated in a separate data set, therefore recommending fMRI cluster analysis as a reliable method for scrutinizing the brain basis of lexical, semantic, and conceptual systems in humans. As focal modules do not explain the distributed character of functionally specific clusters and their distinct topographies are at variance with general distributed processing accounts, the functionally-homogenous distributed clusters specific to semantic types are best explained by specifically-distributed cortical circuits which, similar to Hebbian cell assemblies, represent functional units with specific roles in cognitive processing, especially in lexical and semantic access and memory. *Hum Brain Mapp* 30:3837–3850, 2009. © 2009 Wiley-Liss, Inc.

Key words: action-perception cycle; category-specific activation; cell assembly; concept; *k*-means clustering; language; neuroscience; semantics

INTRODUCTION

The quest for the ultimate processes and representations that form the basis of conceptual thought has been a main

motivation for the advent of neurocognitive imaging. Active areas, however, only allow for limited conclusions on the differential orchestration of brain networks in cognitive tasks. Especially, neuroimaging work so far has not

Contract grant sponsor: Medical Research Council (UK); Contract grant number: U1055.04.003.00001.01; Contract grant sponsor: European Community (New and Emerging Science and Technologies Programme, NEST-2005-PATH-HUM, NESTCOM); Contract grant number: 043374.

*Correspondence to: Friedemann Pulvermüller, MRC Cognition and Brain Sciences Unit, 15 Chaucer Road, Cambridge CB2 7EF, United Kingdom. E-mail: friedemann.pulvermuller@mrc-cbu.cam.ac.uk

Received for publication 4 June 2008; Revised 24 March 2009; Accepted 27 March 2009

DOI: 10.1002/hbm.20811

Published online 24 June 2009 in Wiley InterScience (www.interscience.wiley.com).

yet provided a definitive answer to the fundamental question of what kinds of representations and processing units in the brain form the critical building blocks of thought. Two main proposals had been formulated: the local modular position, according to which specific focal brain centers, single cardinal or “grandmother” cells in the extreme, represent an object, person, word, or concept [Barlow, 1972; Geschwind, 1970; Hubel, 1995], and the distributed holistic view positing that learning in a homogenous network leads to interference patterns and fully distributed representations of cognitive entities [Elman et al., 1996; Lashley, 1950; McClelland and Rumelhart, 1985]. Because cognitive neuroimaging provided evidence for both local specificity and wide distributedness of brain processes of thought, this discussion is still ongoing.

Discussions of global versus locally-specific processing systems have recently been focusing on cortical regions relevant for sensorimotor and conceptual binding [Barsalou, 2008; Patterson et al., 2007; Pulvermüller, 1999]. In frontocentral cortex, the mirror neuron (MN) system is critical for mental processes ranging from binding of information about action execution and perception of the same actions to the recognition of goals and intentions and the processing of words, sentences, and their meaning [Gallese and Lakoff, 2005; Olivier et al., 2007; Pulvermüller, 2005; Rizzolatti and Craighero, 2004; Rizzolatti and Wolpert, 2005]. In humans, this binding is attributable to a great extent to the inferior-frontal premotor cortex homologue to macaque area F5, but more recently, additional MN systems in dorsolateral premotor cortex, prefrontal and parietal cortex, possibly with more specific function, have been discovered [Fogassi et al., 2005; Hauk et al., 2004; Nelissen et al., 2005]. To what degree these different sections of the MN system carry the same or dissociable functions, especially in cognitive domains such as language and conceptual processing, is largely open. The question about functional specificity or more general responsiveness also arises for cortical areas that contribute to the binding of semantic information to words, signs, and symbols. A unitary semantic binding area was proposed in inferior-frontal cortex [Bookheimer, 2002], temporal pole [Patterson et al., 2006], anterior-superiortemporal gyrus [Scott and Johnsrude, 2003], and more posterior inferiortemporal cortex [Price, 2000]. As these areas may contribute to category-specific semantic and conceptual processes [Martin and Chao, 2001], it once again appears crucial to clarify whether one or all of them have a homogeneous function in linguistic, semantic, and conceptual processing or whether their contribution is specific to certain categories of knowledge. A further possibility is that some of these frontal and temporal areas carry a general role in symbol processing while other areas engage in category-specific semantic processing.

We here address the question of functional specificity versus more general responsiveness in local or widely distributed systems using cluster analysis of the activation signatures of small brain loci as they emerged during processing of different types of semantic and conceptual information. We focus on *conceptual* information systemati-

cally linked to words. We call this specific type of conceptual knowledge *semantic* as it is critical for defining word meaning [see also Lambon Ralph and Patterson, 2008; Patterson et al., 2007]. To address functional specificity and segregation one can, in principle, use classic methods such as statistical parametric mapping. However, as the number of relevant parametric contrasts increases rapidly with the number m of conditions, this approach is cumbersome. Cluster analysis offers an alternative, which does not require any a priori grouping of data into subcategories, and which can separate specific from global brain activity. In a data driven manner, cluster analysis creates natural groupings according to the specificity or more general activation of voxels [Baune et al., 1999; Golland et al., 2008; Simon et al., 2004]. The main principle of the k -means clustering algorithm applied in this study is that it minimizes the variance within clusters while at the same time maximizing variance between them. Therefore, the method classifies voxels into the same cluster if they exhibit homogeneous function and are maximally segregated functionally from those in other partitions. As the voxels belonging to the same cluster therefore show the same functional activation pattern across conditions, we will say they have the same functional signature. Critically, the clustering algorithm is not influenced by spatial proximity in the brain and can, therefore, reveal local and distributed neuronal systems with similar function in the very same way.

To monitor conceptual and linguistic processes, words of different semantic types were presented and subjects were asked to read them to understand their meaning. An advantage of word stimuli lies in the possibility to match them for perceptual properties (for example their length), familiarity, and other important psychological features, so that these factors do not act as confounds in the investigation of category-specific processes. Some words included in this study were semantically related to visual information (color words, e.g. “green,” and form/shape words, e.g. “square”) and others related to actions (face-, arm-, and leg-related words, e.g. “lick,” “pick,” “kick”). Results of a range of studies had provided evidence for both globally distributed and locally specific cortical activation to words and concepts. Written language generally activated a distributed frontotemporal network spread out over fusiform [the so-called “visual word form area”, McCandliss et al., 2003] and inferiorfrontal gyrus [containing the human homologue of mirror neuron area F5, Rizzolatti and Craighero, 2004]. Action words semantically related to the face, arms, and legs specifically sparked single foci in motor and premotor areas [e.g., Hauk et al., 2004] and words related to visual information ignited different parts of the temporal lobe [e.g., Martin and Chao, 2001]. For identifying brain voxels with the same functional characteristics, we used k -means cluster analysis of fMRI data [Simon et al., 2004]. The underlying assumption was that voxels within one functionally coherent cluster include neuronal elements from the same type of cortical circuit. To test the reliability of the functional clustering, the experiment was performed twice, in different subjects.

MATERIALS AND METHODS

Subjects

Twenty-one young monolingual right-handed healthy native speakers of English (11 males) participated in the experiments. They were, on average, 24.5 years old ($SD = 5.3$), with an average laterality quotient of 87.1 [$SD = 15.4$, Oldfield, 1971]. Twelve subjects participated in Experiment 1 and nine in Experiment 2; subgroups did not differ in age or laterality. All participants were paid for their participation and had signed an informed consent form prior to testing. Ethics approval had been obtained from the Local Research Ethics Committee, Cambridge.

Stimuli and Experimental Design

Two hundred and fifty words semantically related to actions and visual features of objects were used as stimuli along with 250 strings of hash marks matched pair-wise in length to the words. In addition, 100 null events during which a fixation cross was displayed were included. The word stimuli were submitted to extensive testing for psycholinguistic properties before the experiment [psycholinguistic evaluation procedures as in Hauk and Pulvermüller, 2004; Pulvermüller et al., 1999]. Meticulous investigation of stimulus features revealed that 50 words were semantically related to arm or hand actions, 50 to foot or leg actions, and 50 to actions to which the muscles of the mouth, face, or articulators mainly contribute. Fifty words referred to color and 50 to form or shape information. The action word groups were precisely matched to each other in word length, frequency, imageability, semantic relationship to visual and action information, and number of ambiguous items. The visually related word groups were matched for the same variables. Visually related words were more common than action words [word form frequency 508 vs. 265 occurrences in the CELEX database, Baayen et al., 1995]. Stimuli from the five semantic word categories, arm, color, face, form, and leg, and control items were each presented for 100 ms with a stimulus onset asynchrony between two consecutive items of 2.5 s, so that stimulus presentation and scanner trigger were out of phase by ~ 500 ms. Each stimulus word was presented once to each subject. Two pseudorandomized stimulus sequences were alternated between subjects. Subjects were instructed to attend to all stimuli flashed on the screen and to silently read the words.

Imaging Methods

Subjects were scanned in a 3T Bruker MR system using a head coil. Echo planar imaging (EPI) sequence parameters were $TR = 3.02$ s, $TE = 27$ ms, flip angle = 90° . The functional images consisted of 21 slices covering the whole brain (slice thickness 4 mm, interslice distance 1 mm, field-of-view 25 cm, in plane resolution 128×128). Imag-

ing data were processed using SPM99 software (Wellcome Department of Cognitive Neurology, London, UK).

Images were corrected for slice timing, and then realigned to the first image using sinc interpolation. Phase maps were used to correct for inaccuracies resulting from inhomogeneities in the magnetic field [Cusack et al., 2003; Jezzard and Balaban, 1995]. Any nonbrain parts were removed from the T1-weighted structural images using a surface model approach ("skull-stripping") [Smith, 2002]. The EPI images were co-registered to these skull-stripped structural T1-images using a mutual information co-registration procedure [Maes et al., 1997]. The structural MRI was normalized to the 152-subject T1 template of the Montreal Neurological Institute (MNI). The resulting transformation parameters were applied to the co-registered EPI images. During the spatial normalization process, images were resampled with a spatial resolution of $2 \times 2 \times 2$ mm³. Finally, all normalized images were spatially smoothed with a 12 mm full-width half-maximum Gaussian kernel, globally normalized, and single-subject statistical contrasts were computed using the general linear model [Friston et al., 1998]. The choice of the smoothing parameter was motivated by earlier work on category-specific brain activation [e.g., Hauk et al., 2004; Pulvermüller et al., 2006] and aimed at reducing the effect of intersubject variability. Low-frequency noise was removed applying a high-pass filter with time constant 60 s. Group data were analyzed with random-effects analyses. To control for low-level visual activation, all word category-induced activation was considered relative to the visual control condition, where hash mark strings matched in lengths to the written words were presented. For statistical analysis, the SPM99 canonical hemodynamic response function (HRF) was used to model the activation time course to each event type. Stereotaxic coordinates for voxels with maximal z -values within activation clusters are reported in MNI coordinates.

Data Analysis

Random-effects t -tests were performed for each voxel to test the hypothesis that activation was present at a given brain locus in a given condition. Activation effects given as t -values for all brain voxels found active at a criterion of $P < 0.005$ (uncorrected) in one or more of the conditions were analyzed using k -means clustering. This relatively lenient threshold was chosen to minimize the likelihood of misses. It was expected that any random result of clustering would fail the replication test (see Reliability Check below). Activation of each voxel was plotted in five-dimensional space, each dimension being defined by one of the five conditions.

The two phase algorithm started with random seeds for k cluster centroids. Data points were first partitioned by assigning them to the cluster with the nearest centroid. After all data points had been assigned to clusters,

centroids were updated. These phases were iterated until no data points were reassigned to different clusters in consecutive runs. As clustering is sensitive to starting conditions, the entire procedure was repeated 100 times using different random seeds to calculate centroids. From the 100 solutions, the best clustering was selected using a criterion value calculated as the between-cluster (explained) variance divided by the within-cluster (unexplained) variance.

As the a priori hypothesis immanent to this study was that there is at least one type of distributed cell assembly for each semantic type plus at least one additional type for the processing of words as signs or spoken/written “word forms” [cf. de Saussure, 1916], the number k of clusters was set to 6. To ascertain that this choice of k was in the range appropriate for explaining the maximum explainable variance in the present data set, we calculated the explained effect size (t values) for increasing numbers of k

(starting at $k = 1$). The average and maximal t values obtained for each k are plotted in Figure 1 (top). It can be seen that there was an increase of explained effect size up to a number of clusters of 6. This further endorses the choice of $k = 6$ for the main analyses.

The specificity of clusters was evaluated using the measure of normalized cluster entropy (NCE). Intuitively, cluster entropy is the specificity of a cluster to one stimulus condition. NCE is derived for each cluster i through the proportion p_{ij} of the effect size for the j th of m conditions,

$$\text{NCE}_i = - \sum_{j=1}^m p_{ij} \ln p_{ij} / \ln m$$

Large entropy values imply nonspecific activation, i.e. similar activation values to different semantic categories. Small entropy values imply order and thus category-specific activation. Specific clusters were so selective that their

Figure 1.

(a) Effect size explained as a function of the number k of clusters. Maximal and average effect size explained by all clusters are given as t values. Note the plateau at six clusters indicating that further adding clusters will not increase the variance explained. (b,c) Effect size (t values) explained by five different semantic word categories for each of six clusters. Two nonspe-

cific clusters (1, 2; in orange and brown) can be distinguished from specific clusters, which predominantly explain variance from one semantic word category (3 in cyan, face-related words; 4 in dark purple, arm related words; 5 in yellow, leg-related words; 6 in blue, form/shape words). (b) and (c) present data from Experiments 1 and 2, respectively.

normalized effect size was always greater than 50% for their “best activator” condition, so that less than 50% of their aggregated effect was accounted for by all other conditions taken together. Nonspecific clusters showed more balanced activation over conditions.

Reliability Check

To check the reliability of the method, we determined whether the clustering of data obtained from Experiments 1 and 2 led to similar clustering. To this end, the Rand index [Rand, 1971] was calculated. The Rand index provides a measure of agreement between two pairs of cluster analyses CA1 and CA2. It is defined as

$$\text{Rand index} = (a + d) / (a + b + c + d)$$

where a is the number of voxel pairs clustered in the same partition in both CA1 and CA2, b is the number of voxel pairs clustered in the same partition in CA1 but in different ones in CA2, c is the voxel pair number in different partitions in CA1 but in the same in CA2, and d is the number of voxel pairs assigned to different partitions in both analyses. The Rand index ranges between 0 and 1; a Rand index > 0.5 indicates good correspondence between two classifications. Rand indexes were calculated for the two optimal cluster analyses for Experiment 1 and 2 and were also calculated for nine additional pairs of clusterings obtained after random assignment of subjects to experiments.

RESULTS

Cluster analysis revealed six clusters of voxels, each characterized by a specific activation signature to semantic word categories (Fig. 1, Table I). Importantly, all of these clusters were distributed, consisting of 5–43 local cluster components in different lobes of the cortex. These “foci” each included between 1 and 1,147 voxels; foci ≥ 50 voxels were considered in detail. Two of the six distributed clusters were active to different degrees to a range of semantic word categories (nonspecific clusters 1 and 2, see Fig. 1B,C). In contrast, four distributed clusters had a preferred semantic-category condition activating them, as over 50% of their aggregate effect could be explained by the effect of one semantic category alone, the best activator category (specific clusters 3–6, Fig. 1B,C, Table I, columns H and J). In fact, best activator categories accounted for 75–90% of the aggregate effect captured by three out of the four specific clusters (Table I, column H). Most of the aggregate effect obtained in the two experiments (rank numbers 1–3, Table I, column F) was explained by two semantic-category specific clusters and one nonspecific cluster. In Experiment 1, specific (nonspecific) clusters accounted for 64% (34%) of the metabolic changes and the respective numbers for Experiment 2 were 63% (33%, Table I, column E).

It was surprising that all distributed clusters included local clusters in different cortical lobes, especially in frontal and temporal cortex (Table II). These foci in different lobes but belonging to the same distributed cluster thus showed the same selectivity to word categories, i.e. the

TABLE I. Summary of cluster analysis performed on data from Experiments 1 and 2

No. of Clust.	No. of local clust.	No. of voxel	Avg. effect size [t]	Overall variance explained [%]	Rank	Norm. cluster entropy	% Effect size, best activator condition	% Effect size, 2nd best	Specificity/best activator condition
Experiment 1 ($n = 12$)									
1	6	916	2.44	20	2	0.86	31.01	30.92	Nonspecific
2	9	1,060	1.54	14	4	0.62	45.99	43.06	Nonspecific
3	16	2,957	0.82	22	1	0.32	85.01	9.75	Face
4	14	1,830	0.72	12	5	0.30	87.24	8.34	Arm
5	20	1,437	0.80	10	6	0.57	65.63	25.41	Leg
6	19	3,016	0.73	20	2	0.25	90.94	3.32	Form
Experiment 2 ($n = 9$)									
1	5	1,053	2.416	17	3	0.93	31.01	28.28	Nonspecific
2	15	1,633	1.479	16	4	0.75	41.80	37.06	Nonspecific
3	17	4,156	0.710	20	1	0.49	76.97	7.92	Face
4	21	2,110	0.630	9	6	0.54	75.25	10.86	Arm
5	43	3,685	0.635	16	4	0.71	53.45	30.88	Leg
6	25	4,127	0.653	18	2	0.46	79.10	11.48	Form

For each of the distributed partitions are listed its ID number, the number of local clusters it consisted of, the number of its voxels, the effect size averaged over stimulus categories, and the percentage of the haemodynamic variance explained; a rank index for the amount of explained variance (“rank 1” means “cluster explains most of the variance,” “rank 2” means “second most”) is followed by normalized cluster entropy, percentage of the aggregated effect size of each cluster attributable to the most active and second most active semantic category, specificity/nonspecificity of the cluster and, for specific clusters, the best activator condition. Clusters for which maximally active categories explained more than 50% of their aggregated effect sizes were considered specific. Two specific and one nonspecific clusters explain most of the variation in the data.

TABLE II. Major foci of distributed clusters and their replication

Experiment 1			Experiment 2		
Local clusters	<i>n</i>	<i>x/y/z</i>	<i>n</i>	<i>x/y/z</i>	Local clusters
Cluster 1 (nonspecific-frontofusiform)					
L-F, triangular, opercular	595	-47/22/15	581	-49/22/16	L-F, opercular, precentral
L-T, inf., fusiform	207	-42/-44/-19	441	-45/-46/-14	L-T, inf., mid., fusiform
Cluster 2 (nonspecific-perisylvian)					
L-F, triangular, orbital	873	-45/20/8	1,226	-46/18/13	L-F, triangular, orbital
L-T, mid.	127	-53/-49/-1	288	-50/-50/-6	L-T, mid., inf.
Cluster 3 (face)					
L-F, opercular, precentral	809	-49/11/16	912	-50/9/10	L-F, opercular, precentral
L-P, sup., postcentral	142	-32/-38/58	659	-38/-34/51	L-FP, pre-, postcentral
L-P, inf., postcentral	243	-44/-29/40			
L-T, mid., inf., fusiform	712	-51/-50/-3	826	-49/-50/-4	L-T, mid., inf., fusiform
R-F, triangular, opercular	710	48/22/10	866	49/21/8	R-F, triangular, opercular
Cluster 4 (arm)					
L-F, precentral	383	-32/-5/52	574	-34/-10/50	L-FP, pre-, postcentral
L-T, inf., mid.	89	-49/-51/-9	20*	-52/-52/-9	L-T, inf.
R-FP, pre-, postcentral	730	27/-17/56	181	36/-8/48	R-F, precentral
			813	7/-8/58	R-FP, pre-, postcentral
R-F, triangular	93	37/21/13	124	38/19/12	R-F, triangular, insular
Cluster 5 (leg)					
L-FP, pre-, postcentral	417	-19/-29/61	1,017	-20/-26/57	L-FP, pre-, postcentral1
LR-F, suppl. motor	417	0/3/56	180	0/2/58	LR-F, suppl. Motor
			167	-4/20/48	LR-F, suppl. Motor
L-T, hippocampus, parah.	237	-29/-7/-15	980	-35/-8/-11	L-T, sup., hippocampal
L-TO, lingual	69	-12/-43/-6	65	-11/-38/-10	L-TO, lingual, parah.
R-F, precentral	57	28/-19/60	181	38/-17/53	R-FP, pre-, postcentral
Cluster 6 (form)					
L-F, triangular, insular	1,147	-36/21/8	1,051	-39/26/2	L-F, triangular, insular
L-FP, pre-, postcentral	261	-46/6/42	352	-50/-3/38	L-FP, pre-, postcentral
LR-F, suppl. motor	328	-4/20/52	166	-2/20/56	LR-F, suppl. Motor
LR-F, suppl. motor	82	1/-10/61	144	0/-14/61	LR-F, suppl. Motor
L-T, inf., fusiform	418	-32/-28/-17	987	-28/-19/-16	L-T, inf., fusiform
L-TOP	247	-33/-67/22	540	-28/-66/38	L-TOP
R-FP, pre-, postcentral	420	32/-26/53	198	38/-31/52	R-P, postcentral, mid

For both experiments, the six distributed clusters are decomposed into their largest local clusters (50 voxel cut-off; *exception). For each local cluster, its cortical gyrus/sulcus, the number of active voxels and the MNI coordinate of its voxel with maximal effect size are listed. Replicated local clusters are shown.

L, R, left, right hemisphere; F, frontal; P, parietal; T, temporal; O, occipital; inf., inferior; med., medial; mid., middle; parah., parahippocampus; sup., superior; suppl., supplementary.

same functional signature. Nonspecific cluster 1 consisted of strong left-hemispheric inferiorfrontal triangular-opercular and inferiortemporal fusiform, along with some right-inferiorfrontal activation, whereas nonspecific cluster 2 was distributed over perisylvian left-inferior-prefrontal and middle-temporal areas. Cluster 3 specifically active to face action concepts also showed a left-inferiorfrontal distribution extending into postcentral gyrus along with left-inferiortemporal and right-inferiorfrontal areas. Cluster 4 specifically active to arm words had a strong bihemispheric component with local clusters in bilateral precentral cortex, where arm motor and premotor cortex are situated, extending into postcentral gyrus on the right.

Left-inferiortemporal and right-inferiorfrontal activation were also seen. Leg words sparked an assembly of local clusters distributed over left and right dorsal pre- and postcentral cortex, where leg motor and premotor cortex is situated, left-frontodorsal supplementary motor cortex, left-middletemporal gyrus, hippocampus, parahippocampus, and temporooccipital lingual gyrus (cluster 5). The most widely distributed activation was seen for cluster 6 specific to words related to form concepts. They activated a network of local clusters in bilateral inferiorfrontal and lateral-pre- and -postcentral cortex along with left-dorso-lateral prefrontal, supplementary-motor, inferiortemporal, and parietotemporooccipital areas.

Cluster analysis results from Experiment 1 agreed well with those from Experiment 2, as demonstrated by a Rand index of 0.8696. Nine additional pairs of cluster analyses (performed on the same data sets with different random assignments of subjects to groups) yielded Rand indexes ranging between 0.6174 and 0.8696 (average = 0.7888). This proves replicability of cluster analysis results.

Most of the local clusters making up a distributed cluster in one analysis had a homologue in the analysis of the other experiment (Table II). Twenty-seven of the larger foci (≥ 50 voxels) in each analysis directly matched with a focus in the other analysis (there were sometimes two focal homologues to one larger focus, see Table II). In addition to the 27 replicated focal clusters, Experiment 1 revealed three “idiosyncratic” foci not matched by analyses of Experiment 2. Experiment 2 gave rise to 13 idiosyncratic foci; some of these additional foci may be related to the smaller sample size ($n = 9$) of that study, leading to increased opportunity to reflect inter-individual variation. Among the overall 70 larger foci (≥ 50 voxels) in both studies taken together, there were 54 matches and 16 mismatches. The number of voxels in these matching foci was 23,823 (90.8%) and that in idiosyncratic ones was 2,415 (9.2%). These numbers confirm good replicability of results. Replicability is further proven by the visual similarity of the cortical distributions of the six distributed clusters (see Fig. 2).

DISCUSSION

Cluster analysis of fMRI activation identified six functionally distinct distributed sets of cortical foci. Four of these distributed clusters were specifically activated by one semantic word category, suggesting that, in these cases, the foci comprised by the distributed clusters related to word-induced category-specific semantic or conceptual processes. Two clusters were activated in a nonspecific manner, to all or most word kinds under study suggesting a relationship to general mechanisms of word processing. Remarkably, all clusters were widely dispersed and included activation foci in both left frontocentral and temporal cortex. As all clusters showed specific cortical distribution and most of them signified one semantic word type, these data indicate topographical and functional specificity of distributed cortical networks involved in semantic processing.

Focal modules do not explain the distributed character of functionally specific clusters and general distributed processing accounts face difficulties accounting for the topographical specificity of the individual clusters. The distributed clusters are best explained by distributed cortical circuits which, similar to Hebb’s construct of cell assemblies [Braitenberg, 1978; Hebb, 1949; Milner, 1996], form functional units with a specific role in cognitive processing, especially in semantic access and memory. Similar to the observed clusters, these neuronal circuits had been proposed to be distributed but also specific in their distribution

and function [Pulvermüller, 1992, 1999]. As distributed circuits appear to connect neurons in frontocentral areas controlling actions with neurons in temporal areas receiving auditory and visual information, this organization provides a mechanism for multimodal response characteristics as they are found in memory cells [Fuster, 1997; Fuster and Alexander, 1971] and action-related mirror neurons [Gallese et al., 1996; Rizzolatti and Craighero, 2004].

We will now turn to a detailed discussion of the cortical organization of functional clusters revealed and confirmed by our experiments and their possible roles in cognitive processing. Results will be related to theories of semantic and conceptual brain systems and the question whether the distributed clusters and their local foci might index functional contributions or possibly epiphenomenal activation will be addressed in the light of neuropsychological and TMS studies. A final paragraph offers conclusions on the nature of cognitive representations in the brain.

Clusters Signify General Linguistic and Specific Conceptual Processes

Two of the distributed clusters, clusters 1 and 2, were nonspecific as voxels became active for a range of word types. These clusters’ local activations in left inferiorfrontal, middle-temporal, and fusiform gyrus, may reflect links between motor, acoustic, and visual representations of spoken and written words and possibly modality-nonspecific processing of such abstract “word forms.” We here use the term “word form” in the sense of “words as written or spoken signs” and distinguish word form from meaning adopting a distinction well-established in language science [de Saussure, 1916]. Although both nonspecific clusters included superior- and inferiortemporal along with inferiorfrontal foci, the relative size of the temporal foci was different. The superiortemporal foci were pronounced in cluster 2 (>100 adjacent voxels), whereas the inferiortemporal ones were smaller. The substantial foci included posterior superiortemporal sulcus, which is known to respond specifically to vowels [coordinates $(-64/-22/-2)$, Uppenkamp et al., 2006], individual speech sounds [$(-60/-20/-2)$, Pulvermüller et al., 2006], and spoken words [$(-48/-27/-5)$, Shtyrov et al., 2008]. In contrast, cluster 1 was characterized by substantial foci (including >200 voxels) in left inferiortemporal and fusiform cortex, which included a locus possibly contributing to visual word form processing [$(-42/-57/-15)$, McCandliss et al., 2003]. These distributional differences suggest that cluster 1 is more closely related to visual word form processing, whereas cluster 2 plays a greater role in phonological and spoken word processing. As only visual stimuli were used in the current experiments, these suggestions must however await further experimental confirmation.

Critically, the majority of clusters were specific and responded primarily to one semantic word type. Four

Figure 2.

Specifically distributed functionally coherent sets of foci as revealed by the distribution of voxels belonging to six distributed clusters obtained from Experiments 1 and 2. Neuronal assemblies revealed by two nonspecific clusters (top, in orange and brown) include foci in inferiorfrontal, middle-, and inferiortemporal cortex. Clusters specific to semantic categories shown in cyan, purple, yellow, and blue have an even wider specific distribution. Similarity of clusters on the left and right proves excellent replicability.

Figure 2 (Continued).

Figure 3.

All six distributed clusters obtained from Experiments 1 (left) and 2 (right) reveal functional segregation of general lexical and/or semantic processes on the one hand (brown and orange clusters) and category-specific semantic processes in frontocentral mirror neuron systems and in temporal cortex (all other colors).

clusters showed most of their effect size in response to one semantic category of words. Words referring to face (Cluster 3), arm (4), leg (5), and form (6) each activated a replicable and specific set of foci. The foci in the motor/premotor system indexing semantic relationship of action words to different parts of the body are each part of a distributed cluster also including specific temporal areas. Temporal foci may index circuits activated by the visual perception of actions to which the words refer and/or target areas of specific temporal projections from the sensorimotor cortex.

Although action word categories were matched for a range of psycholinguistic variables, visually-related words had higher word frequencies than action words. The absence of a separate cluster for color words in the present results may reflect the relatively high frequency of this word group, which may be the reason for the reduced brain response [Chee et al., 2003; Fiebach et al., 2002; Hauk et al., 2008b]. A further reason for the absence of a separate cluster for color words may be the spatial vicinity of relevant circuits to cluster 5. Previous studies indicated that color words activate inferior temporal fusiform and parahippocampal gyri [Pulvermüller and Hauk, 2006; Simmons et al., 2007], which are close to the temporal foci of cluster 5. As common words elicit less brain activity than rare ones [Chee et al., 2003; Fiebach et al., 2002; Hauk et al., 2008b], the more widespread activation compared with all other clusters elicited by the form-word cluster 6 cannot be due to word frequency. It appears more plausible that form word cluster 6 reflects sensorimotor semantic aspects of form words and concepts. Note that words semantically related to visual shapes ("square," "round") are related to the motor system, too, as observing shapes is usually accompanied by specific eye movement patterns tracing the shape. The frontoparietal and temporal circuits included in the form word cluster may provide a basis for such cross-modal binding of action and perception-related semantic information. Similar mechanisms for cross-modal information link-

age may underpin the processing of abstract words [Binder et al., 2005; Pexman et al., 2007a,b; Pulvermüller, 2008]

In inferior frontal cortex, foci that indexed general word-form processing were sandwiched in-between areas that responded specifically to semantic categories (Figs. 2 and 3), a precentral focus of the cluster specific to shape words ($-36/21/8$) and a precentral focus of that to face words ($-49/11/16$). Similarly, inferior and middle temporal foci contributing to word form processing ($-42/-44/-19$, $-53/-49/-1$) were surrounded by category-specific activations. The motor-premotor region was dominated by semantic somatotopy [Aziz-Zadeh et al., 2006; Boulenger et al., 2008a; Buccino et al., 2001; Hauk et al., 2004; Pulvermüller et al., 2001] with face, arm, and leg word activation foci at inferior lateral ($-49/11/16$), dorsolateral ($-32/-5/52$), and dorsal sites ($-19/-21/61$). A different semantic map was seen in the temporal cortex and the hippocampal formation where an anterior-to-posterior order emerged for foci included in specific clusters for leg words ($-29/-7/-15$), shape words ($-32/-28/-17$), and face and arm words ($-51/-50/3$ and $-49/-51/-9$).

Distributed Neuronal Circuits for Different Types of Cognitive Processes

These results offer answers to critical questions about the specificity of cortical area activations in linguistic, semantic, and conceptual processing. The observed pattern of local functional alteration between general word processing circuits and category-specific ones adds to the evidence about the functional parcellation of the left inferior frontal cortex [Bookheimer, 2002]. However, while confirming a role of anterior inferior frontal cortex to category-specific semantic processing, they also suggest a similar role, but for different categories, of posterior inferior frontal and motor/premotor cortex. In between category-specific semantic activations, the data suggest a "layer" for lexical or more general semantic computations (see Fig. 3).

As general language regions and category-specific conceptual-semantic foci reside side-by-side in inferiorfrontal cortex, this area can be seen to unite a general language processor operating on MNs [Rizzolatti and Arbib, 1998] and more specific MN circuits involved in understanding subtypes of action words and concepts [Pulvermüller, 2005]. The semantic map uncovered in temporal cortex adds to previous evidence about category-specificity in this region [Martin and Chao, 2001], as a wider variety of words and concepts, including face-, arm-, leg-, and form-related ones, elicited specific activations in this brain part. The general word-related activations together with the category-specific semantic ones in temporal areas now offer a solution of the controversy about the posterior inferior temporal cortex and its relationship to processing of either visual word form or semantics [e.g., McCandliss et al., 2003; Price and Devlin, 2003]. Both possibilities seem to be correct, but, however, for different closely adjacent areas.

Category-Specificity Explains Most of the Neurometabolic Effect

Table I shows that about two thirds of the aggregated effect in neurometabolic activation was accounted for by specific clusters, therefore demonstrating the predominance of category-specific brain activation over nonspecific responsiveness to lexical items. The clear and reproducible category differences obtained from concept groups preselected for specific semantic features might provide a hint why earlier neuroimaging studies failed to find category-specificity in brain activation [Devlin et al., 2002]. This may be due to the natural heterogeneity of the stimulus categories and the lack of behavioral control of semantic variables. In our view, control of variance in physical and psychological features of the stimulus set is important for obtaining category effects [Penolazzi et al., 2007; Pulvermüller, 1999]. The present data suggest that the lion's share of brain activation in word-elicited conceptual/semantic thought is category-specific. Therefore, studies of nonspecific brain activation to words averaging over semantic categories may miss out on these critical effects.

Category-Specific Semantic Activation Clusters and the Effect of Lesions and TMS

The present results from a metabolic imaging study show that topographically specific cortical clusters become active when words with different meaning are being processed. They also suggest that circuits with different distributions contribute to, and form the basis of, semantic processing. Although it is well-established in cognitive neuroscience that action-perception systems in the brain play a critical role in semantic and conceptual processing [e.g., Barsalou, 2008; Hoenig et al., 2008; Kemmerer and Gonzalez-Castillo, 2008; Martin, 2007; Olivier et al., 2007;

Pulvermüller, 1996, 2005; Rizzolatti and Craighero, 2004; Warrington and Shallice, 1984], it is important to point to the limits of any functional imaging study, which can show correlated activation but not a causal role of such activation. Some effort has been spent to extend the range of conclusions from fMRI work, for example by looking at priming and category-specificity of word frequency effects [Gold et al., 2006; Hauk et al., 2008a; Wheatley et al., 2005]. However, for finding out whether the activated distributed circuits represent a *necessary* brain basis of semantic processing, or rather a basis for epiphenomenal processes not related to semantics [see, e.g., Oliveri et al., 2004], it is necessary to consider complementary methods, especially TMS and lesion studies.

Previous patient work revealed double dissociations in support of category-specific semantic processing [Daniele et al., 1994; Warrington and Shallice, 1984]. Especially the double dissociation in the processing of animal and tool knowledge [e.g., Buxbaum and Saffran, 2002] and in processing object-nouns and action verbs [e.g., Neininger and Pulvermüller, 2003] provide strong evidence that the cortical systems necessary for processing visually based conceptual-perceptual knowledge and action knowledge differ. In addition, recent studies showed that motor cortex degeneration in motor neuron disease leads to a specific impairment of action words and concepts [Bak et al., 2001, 2006], frontoparietal lesion associated with apraxia degrades object utilization knowledge more strongly than other conceptual aspects of object understanding [Buxbaum and Saffran, 2002], focal lesion in the motor cortex selectively impairs action word processing [e.g., Neininger and Pulvermüller, 2003], and Parkinson's disease is also accompanied by a specific deficit in processing action verbs [Boulenger et al., 2008]. Although these studies are consistent with the notion of distributed semantic circuits with distinct topographies [for discussion, see Neininger and Pulvermüller, 2003], they address relatively large semantic and conceptual domains (animal/tool, object/action, noun/verb), whereas this study focused on more fine-grained distinctions (e.g., action subtypes). However, neuropsychological work also showed that the areas found active for the more fine-grained semantic subcategories play a critical role in their processing. TMS to hand and foot motor cortex respectively influenced the processing of hand- and leg-related words, yielding a double dissociation in processing these categories [Pulvermüller et al., 2005] and, vice versa, action language processing affected motor system activation and behavior [Boulenger et al., 2006; Buccino et al., 2005]. Also object color and form knowledge was found to dissociate after lesion in anterior-temporal areas, consistent with differential functional activation [Miceli et al., 2001; Pulvermüller and Hauk, 2006]. These results further support a critical functional role of category-specific circuits in semantic processing and in part validate the functional clusters. Still, as this study revealed for the first time that category-specific distributed clusters are each composed of several foci with similar

functionality, it is evident that lesion and TMS evidence is not available for the full set of foci. Future neuropsychological work is necessary to clarify whether the different foci included in the distributed clusters each play a critical role in semantic processing.

Specific Distribution of Semantic Circuits

The present results neither support a nonspecific holistic network approach according to which cognitive processes freely use distributed cortical resources, nor does it bolster a modular localist framework according to which semantics or concepts are bound to a single local site. Instead, the present results support what one may want to call “specific distributionalism”: semantic and conceptual processes specific to categories of knowledge are carried by scattered neuronal circuits with specific cortical distributions in the brain. Accordingly, the cortical binding devices for word-related conceptual information are distributed semantic circuits spread out over specific areas in different cortical lobes. It is possible that these conceptual representations are linked into word processing circuits at one critical location, a unitary centre for semantics. However, the present data indicate that the entirety of the frontotemporal perisylvian cortex, the distributed cluster array for (spoken and visual) word form circuits (i.e., Clusters 1 and 2), is available for semantic linkage between words and concepts. This conclusion is consistent with neuroimaging work arguing for a role in semantic linkage for both temporal [Price, 2000] and inferior-prefrontal cortex [Bookheimer, 2002]. In addition, studies of degenerative brain diseases characterized by semantic deficits for action and object processing and caused by degradation of either inferiorfrontal [Bak et al., 2001] or temporal systems [Patterson et al., 2007] further suggest the possibility of multiple semantic links between words and concepts. Our present results indicate that the areas of general lexical and/or semantic processing are, in both inferiorfrontal and temporal cortex, localized in-between areas specifically processing semantic categories.

SUMMARY

We here report evidence for scattered cortical circuits with specific distribution, which are either selectively activated by individual semantic word categories or show nonselective activation to words of different kinds. These results refute strong versions of both local modular and distributed holistic proposals of lexical and semantic processing and enforce proposals according to which distributed neuronal assemblies with specific and partly overlapping topographies carry language, concepts, and thought.

ACKNOWLEDGMENTS

We thank two anonymous referees for their comments on a previous version of this article and Clare Dine and Yury Shtyrov for their help at different stages of this work.

REFERENCES

- Aziz-Zadeh L, Wilson SM, Rizzolatti G, Iacoboni M (2006): Congruent embodied representations for visually presented actions and linguistic phrases describing actions. *Curr Biol* 16:1818–1823.
- Baayen RH, Piepenbrock R, Gulikers L (1995): The CELEX Lexical Database (Release 2). Philadelphia, PA: Linguistic Data Consortium, University of Pennsylvania.
- Bak TH, O'Donovan DG, Xuereb JH, Boniface S, Hodges JR (2001): Selective impairment of verb processing associated with pathological changes in brodmann areas 44 and 45 in the motor neurone disease-dementia-aphasia syndrome. *Brain* 124:103–120.
- Bak TH, Yancopoulou D, Nestor PJ, Xuereb JH, Spillantini MG, Pulvermüller F, Hodges JR (2006): Clinical, imaging and pathological correlates of a hereditary deficit in verb and action processing. *Brain* 129 (Part 2):321–332.
- Barlow H (1972): Single units and cognition: A neurone doctrine for perceptual psychology. *Perception* 1:371–394.
- Barsalou LW (2008): Grounded cognition. *Annu Rev Psychol* 59:617–645.
- Baune A, Sommer FT, Erb M, Wildgruber D, Kardatzki B, Palm G, Grodd W (1999): Dynamical cluster analysis of cortical fMRI activation. *Neuroimage* 9:477–489.
- Binder JR, Westbury CF, McKiernan KA, Possing ET, Medler DA (2005): Distinct brain systems for processing concrete and abstract concepts. *J Cognit Neurosci* 17:905–917.
- Bookheimer S (2002): Functional MRI of language: New approaches to understanding the cortical organization of semantic processing. *Annu Rev Neurosci* 25:151–188.
- Boulenger V, Roy AC, Paulignan Y, Deprez V, Jeannerod M, Nazir TA (2006): Cross-talk between language processes and overt motor behavior in the first 200 msec of processing. *J Cognit Neurosci* 18:1607–1615.
- Boulenger V, Mechtouff L, Thobois S, Broussolle E, Jeannerod M, Nazir TA (2008): Word processing in Parkinson's disease is impaired for action verbs but not for concrete nouns. *Neuropsychologia* 46:743–756.
- Boulenger V, Hauk O, Pulvermüller F (2008a): Grasping ideas with the motor system: Semantic somatotopy in idiom comprehension. *Cereb Cortex* doi: 10.1093/cercor/bhn217.
- Braitenberg V (1978): Cell assemblies in the cerebral cortex. In: Heim R, Palm G, editors. *Lecture Notes in Biomathematics*, Vol. 21: Theoretical Approaches to Complex Systems. Berlin: Springer. pp 171–188.
- Buccino G, Binkofski F, Fink GR, Fadiga L, Fogassi L, Gallese V, Seitz RJ, Zilles K, Rizzolatti G, Freund HJ (2001): Action observation activates premotor and parietal areas in a somatotopic manner: An fMRI study. *Eur J Neurosci* 13:400–404.
- Buccino G, Riggio L, Melli G, Binkofski F, Gallese V, Rizzolatti G (2005): Listening to action-related sentences modulates the activity of the motor system: A combined TMS and behavioral study. *Brain Res Cognit Brain Res* 24:355–363.

- Buxbaum LJ, Saffran EM (2002): Knowledge of object manipulation and object function: Dissociations in apraxic and nonapraxic subjects. *Brain Lang* 82:179–199.
- Chee MW, Westphal C, Goh J, Graham S, Song AW (2003): Word frequency and subsequent memory effects studied using event-related fMRI. *Neuroimage* 20:1042–1051.
- Cusack R, Brett M, Osswald K (2003): An evaluation of the use of magnetic field maps to undistort echo-planar images. *Neuroimage* 18:127–142.
- Daniele A, Giustolisi L, Silveri MC, Colosimo C, Gainotti G (1994): Evidence for a possible neuroanatomical basis for lexical processing of nouns and verbs. *Neuropsychologia* 32:1325–1341.
- de Saussure F (1916): *Cours de Linguistique Generale*. Paris: Payot.
- Devlin JT, Moore CJ, Mummery CJ, Gorno-Tempini ML, Phillips JA, Noppeney U, Frackowiak RS, Friston KJ, Price CJ (2002): Anatomic constraints on cognitive theories of category specificity. *Neuroimage* 15:675–685.
- Elman JL, Bates L, Johnson M, Karmiloff-Smith A, Parisi D, Plunkett K (1996): *Rethinking Innateness: A Connectionist Perspective on Development*. Cambridge, MA: MIT Press.
- Fiebach CJ, Friederici AD, Muller K, von Cramon DY (2002): fMRI evidence for dual routes to the mental lexicon in visual word recognition. *J Cognit Neurosci* 14:11–23.
- Fogassi L, Ferrari PF, Gesierich B, Rozzi S, Chersi F, Rizzolatti G (2005): Parietal lobe: From action organization to intention understanding. *Science* 308:662–667.
- Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg MD, Turner R (1998): Event-related fMRI: Characterizing differential responses. *Neuroimage* 7:30–40.
- Fuster JM (1997): Network memory. *Trends Neurosci* 20:451–459.
- Fuster JM, Alexander GE (1971): Neuron activity related to short-term memory. *Science* 173:652–654.
- Gallese V, Fadiga L, Fogassi L, Rizzolatti G (1996): Action recognition in the premotor cortex. *Brain* 119 (Pt 2):593–609.
- Gallese V, Lakoff G (2005): The brain's concepts: The role of the sensory-motor system in conceptual knowledge. *Cognit Neuropsychol* 22:455–479.
- Geschwind N (1970): The organization of language and the brain. *Science* 170:940–944.
- Gold BT, Balota DA, Jones SJ, Powell DK, Smith CD, Andersen AH (2006): Dissociation of automatic and strategic lexical-semantics: Functional magnetic resonance imaging evidence for differing roles of multiple frontotemporal regions. *J Neurosci* 26:6523–6532.
- Golland Y, Golland P, Bentin S, Malach R (2008): Data-driven clustering reveals a fundamental subdivision of the human cortex into two global systems. *Neuropsychologia* 46:540–553.
- Hauk O, Pulvermüller F (2004): Neurophysiological distinction of action words in the fronto-central cortex. *Hum Brain Mapp* 21:191–201.
- Hauk O, Johnsrude I, Pulvermüller F (2004): Somatotopic representation of action words in the motor and premotor cortex. *Neuron* 41:301–307.
- Hauk O, Davis MH, Kherif F, Pulvermüller F (2008a): Imagery or meaning? Evidence for a semantic origin of category-specific brain activity in metabolic imaging. *Eur J Neurosci* 27:1856–1866.
- Hauk O, Davis MH, Pulvermüller F (2008b): Modulation of brain activity by multiple lexical and word form variables in visual word recognition: A parametric fMRI study. *Neuroimage* 42:1185–1195.
- Hebb DO (1949): *The Organization of Behavior. A Neuropsychological Theory*. New York: Wiley.
- Hoenig K, Sim EJ, Bochev V, Herrnberger B, Kiefer M (2008): Conceptual flexibility in the human brain: Dynamic recruitment of semantic maps from visual, motor, and motion-related areas. *J Cognit Neurosci* 20:1799–1814.
- Hubel D (1995): *Eye, Brain, and Vision*. New York: Scientific American Library.
- Jezzard P, Balaban RS (1995): Correction for geometric distortion in echo planar images from B0 field variations. *Magn Reson Med* 34:65–73.
- Kemmerer D, Gonzalez-Castillo J (2008): The two-level theory of verb meaning: An approach to integrating the semantics of action with the mirror neuron system. *Brain Lang*.
- Lambon Ralph MA, Patterson K (2008): Generalization and differentiation in semantic memory: Insights from semantic dementia. *Ann N Y Acad Sci* 1124:61–76.
- Lashley KS (1950): In search of the engram. *Symp Soc Exp Biol* 4:454–482.
- Maes F, Collignon A, Vandermeulen D, Marchal G, Suetens P (1997): Multimodality image registration by maximization of mutual information. *IEEE Trans Med Imaging* 16:187–198.
- Martin A (2007): The representation of object concepts in the brain. *Annu Rev Psychol* 58:25–45.
- Martin A, Chao LL (2001): Semantic memory and the brain: Structure and processes. *Curr Opin Neurobiol* 11:194–201.
- McCandliss BD, Cohen L, Dehaene S (2003): The visual word form area: Expertise for reading in the fusiform gyrus. *Trends Cognit Sci* 7:293–299.
- McClelland JL, Rumelhart DE (1985): Distributed memory and the representation of general and specific information. *J Exp Psychol Gen* 114:159–188.
- Miceli G, Fouch E, Capasso R, Shelton JR, Tomaiuolo F, Caramazza A (2001): The dissociation of color from form and function knowledge. *Nat Neurosci* 4:662–667.
- Milner PM (1996): Neural representation: Some old problems revisited. *J Cognit Neurosci* 8:69–77.
- Neininger B, Pulvermüller F (2003): Word-category specific deficits after lesions in the right hemisphere. *Neuropsychologia* 41:53–70.
- Nelissen K, Luppino G, Vanduffel W, Rizzolatti G, Orban GA (2005): Observing others: Multiple action representation in the frontal lobe. *Science* 310:332–336.
- Oldfield RC (1971): The assessment and analysis of handedness: The Edinburgh Inventory. *Neuropsychologia* 9:97–113.
- Oliveri M, Finocchiaro C, Shapiro K, Gangitano M, Caramazza A, Pascual-Leone A (2004): All talk and no action: A transcranial magnetic stimulation study of motor cortex activation during action word production. *J Cognit Neurosci* 16:374–381.
- Olivier E, Davare M, Andres M, Fadiga L (2007): Precision grasping in humans: From motor control to cognition. *Curr Opin Neurobiol* 17:644–648.
- Patterson K, Lambon Ralph MA, Jefferies E, Woollams A, Jones R, Hodges JR, Rogers TT (2006): “Presemantic” cognition in semantic dementia: Six deficits in search of an explanation. *J Cognit Neurosci* 18:169–183.
- Patterson K, Nestor PJ, Rogers TT (2007): Where do you know what you know? The representation of semantic knowledge in the human brain. *Nat Rev Neurosci* 8:976–987.
- Penolazzi B, Hauk O, Pulvermüller F (2007): Early semantic context integration and lexical access as revealed by event-related brain potentials. *Biol Psychol* 74:374–388.
- Pexman PM, Hargreaves IS, Edwards JD, Henry LC, Goodyear BG (2007a): The neural consequences of semantic richness:

- When more comes to mind, less activation is observed. *Psychol Sci* 18:401–406.
- Pexman PM, Hargreaves IS, Edwards JD, Henry LC, Goodyear BG (2007b): Neural correlates of concreteness in semantic categorization. *J Cognit Neurosci* 19:1407–1419.
- Price CJ (2000): The anatomy of language: Contributions from functional neuroimaging. *J Anat* 197 (Part 3):335–359.
- Price CJ, Devlin JT (2003): The myth of the visual word form area. *Neuroimage* 19:473–481.
- Pulvermüller F (1992): Constituents of a neurological theory of language. *Concepts Neurosci* 3:157–200.
- Pulvermüller F (1996): Hebb's concept of cell assemblies and the psychophysiology of word processing. *Psychophysiology* 33:317–333.
- Pulvermüller F (1999): Words in the brain's language. *Behav Brain Sci* 22:253–336.
- Pulvermüller F (2005): Brain mechanisms linking language and action. *Nat Rev Neurosci* 6:576–582.
- Pulvermüller F (2008): Grounding language in the brain. In: de Vega M, Graesser A, Glenberg AM, editors. *Symbols, Embodiment, and Meaning*. Oxford: Oxford University Press. pp 85–116.
- Pulvermüller F, Hauk O (2006): Category-specific processing of color and form words in left fronto-temporal cortex. *Cereb Cortex* 16:1193–1201.
- Pulvermüller F, Lutzenberger W, Preissl H (1999): Nouns and verbs in the intact brain: Evidence from event-related potentials and high-frequency cortical responses. *Cereb Cortex* 9:498–508.
- Pulvermüller F, Hummel F, Härle M (2001): Walking or talking? Behavioral and neurophysiological correlates of action verb processing. *Brain Lang* 78:143–168.
- Pulvermüller F, Hauk O, Nikulin VV, Ilmoniemi RJ (2005): Functional links between motor and language systems. *Eur J Neurosci* 21:793–797.
- Pulvermüller F, Huss M, Kherif F, Moscoso del Prado Martin F, Hauk O, Shtyrov Y (2006): Motor cortex maps articulatory features of speech sounds. *Proc Natl Acad Sci USA* 103:7865–7870.
- Rand WM (1971): Objective criteria for the evaluation of clustering methods. *J Am Stat Assoc* 66:846–850.
- Rizzolatti G, Arbib MA (1998): Language within our grasp. *Trends Neurosci* 21:188–194.
- Rizzolatti G, Craighero L (2004): The mirror-neuron system. *Annu Rev Neurosci* 27:169–192.
- Rizzolatti G, Wolpert DM (2005): Motor systems. *Curr Opin Neurobiol* 15:623–625.
- Scott SK, Johnsrude IS (2003): The neuroanatomical and functional organization of speech perception. *Trends Neurosci* 26:100–107.
- Shtyrov Y, Osswald K, Pulvermüller F (2008): Memory traces for spoken words in the brain as revealed by the hemodynamic correlate of the mismatch negativity. *Cereb Cortex* 18:29–37.
- Simmons WK, Ramjee V, Beauchamp MS, McRae K, Martin A, Barsalou LW (2007): A common neural substrate for perceiving and knowing about color. *Neuropsychologia* 45:2802–2810.
- Simon O, Kherif F, Flandin G, Poline JB, Riviere D, Mangin JF, Le Bihan D, Dehaene S (2004): Automatized clustering and functional geometry of human parietofrontal networks for language, space, and number. *Neuroimage* 23:1192–1202.
- Smith SM (2002): Fast robust automated brain extraction. *Hum Brain Mapp* 17:143–155.
- Uppenkamp S, Johnsrude IS, Norris D, Marslen-Wilson W, Patterson RD (2006): Locating the initial stages of speech-sound processing in human temporal cortex. *Neuroimage* 31:1284–1296.
- Warrington EK, Shallice T (1984): Category specific semantic impairments. *Brain* 107:829–854.
- Wheatley T, Weisberg J, Beauchamp MS, Martin A (2005): Automatic priming of semantically related words reduces activity in the fusiform gyrus. *J Cognit Neurosci* 17:1871–1885.