Published in final edited form as:

J Am Chem Soc. 2019 May 22; 141(20): 8104–8109. doi:10.1021/jacs.9b04189.

Transition-Metal- and Light-Free Directed Amination of Remote Unactivated C(sp³)–H Bonds of Alcohols

Daria Kurandina[‡], Dongari Yadagiri[‡], Mónica Rivas[‡], Aleksei Kavun, Padon Chuentragool, Keiichi Hayama, Vladimir Gevorgyan^{*}

Department of Chemistry, University of Illinois at Chicago, 845 W. Taylor Street, Rm. 4500, Chicago, Illinois 60607, United States

Abstract

Due to the great value of amino alcohols, new methods for their synthesis are in high demand. Abundant aliphatic alcohols represent the ideal feedstock for the method development toward this important motif. To date, transition-metal-catalyzed approaches for the directed remote amination of alcohols have been well established. Yet, they have certain disadvantages such as the use of expensive catalysts and limited scope. Very recently, transition-metal-free visible-light-induced radical approaches have emerged as new powerful tools for directed remote amination of alcohols. Relying on 1,5-HAT reactivity, these methods are limited to β - or δ -amination only. Herein, we report a novel transition-metal- and visible-light-free room-temperature radical approach for remote β -, γ -, and δ -C(sp³)–N bond formation in aliphatic alcohols using mild basic conditions and readily available diazonium salt reagents.

A diverse range of natural products, pharmaceuticals, and catalyst designs feature amino alcohols as privileged motifs.¹ One of the most powerful strategies toward their synthesis is the directed site-selective C(sp³)–H amination of aliphatic alcohols,² which are ubiquitous in complex molecules and are inexpensive and sustainable starting materials. The seminal strategy for directed amination of alcohols developed by Du Bois, White, and others relies on insertion of Rh-,³ Ru-,⁴ Ag-,⁵ Fe-,⁶ Mn-,⁷ Cu-,⁸ or Co⁹-based transition-metal-nitrene species A into distal unactivated C(sp³)–H bonds¹⁰ (Scheme 1, a). However, the innate metallacyclic intermediates restrict these metal-based methods to γ - and β -site functionalization. Recently, radical approaches for remote functionalization of alcohols¹¹ have gained broad attention due to the discoveries of new mild methods for generation of radicals and their versatile hydrogen atom transfer (HAT) reactivity.¹² To date, only two methods on remote site-selective C(sp³)–H amination of alcohols have been developed relying on radical intermediates. In 2017, the Nagib group introduced an elegant metal-free visible-light-induced radical relay for synthesis of β -amino alcohols from their alcohol analogues (Scheme 1, b).¹³

^{*}Corresponding Author: vlad@uic.edu.

[‡]Author Contributions: D.K., D.Y., and M.R. contributed equally.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.9b04189.

Experimental procedures and compound characterization data (PDF)

The authors declare no competing financial interest.

This method utilizes a novel, easily installable/removable imidate-based chaperone capable of generating N-centered radical species **B**, which undergoes a 1,5-HAT process producing **C**. Subsequent 5-*endo*-trig radical cyclization of **3** and *in situ* hydrolysis delivers the amination products with exclusive β -selectivity. Later, the Zuo group developed another visible-light-induced protocol to access δ -amino alcohols (Scheme 1, c).¹⁴ In this case, Ce-catalyzed generation of alkoxy radicals **D** directly from alcohols followed by 1,5-HAT and radical addition of **E** to di-*tert*-butyl azodiformate leads to δ -selective C–N bond formation. Overall, these radical approaches eliminated the necessity for use of expensive transition metal catalysts and conventional heating for remote C(sp³)–H amination of alcohols.

However, due to the preference of heteroatom-centered radicals for 1,5-HAT,^{12b} the scope of these radical transformations remains limited and γ -amination toward valuable 1,3-amino alcohols has not yet been achieved. Therefore, development of more general metal-free methods featuring benign conditions is highly desired.¹⁵ Herein, we report the first transition-metal- and visible-light-free room-temperature protocol for remote β -, γ -, and δ -C(sp³)–N bond formation in aliphatic alcohols using mild basic conditions and readily available diazonium salts (Scheme 1, d). This method employs the ability of the latter to initiate radical reactions in the presence of bases or other reducing agents,¹⁶ as well as to trap nucleophilic radical intermediates to form a new C–N bond.¹⁷ Thus, previously unknown iodine atom abstraction from the silyl methyl iodide moiety of a tethered alcohol by an aryl radical, formed from the diazonium salt in the presence of base, leads to silyl methyl radical species **F**. This electrophilic radical,¹⁸ due to electronic mismatch, is not predisposed to coupling with the diazonium reagent and instead undergoes a selective 1,6-, 1,5-, or 1,7-HAT process to produce transposed nucleophilic radical species **G** at the remote C(sp³)–H site of the alcohol.¹⁹ The latter would be ultimately trapped by excess diazonium salt, which is known to be a facile process ($k = 10^6 \text{ M}^{-1} \text{ s}^{-1}$ for primary alkyl radicals; $k = 10^8 \text{ M}^{-1} \text{ s}^{-1}$ for tertiary alkyl radicals),^{17b} and reduced by base to produce β -, γ -, or δ -diazenylated alcohols, accordingly. Thus, the base plays a double duty by reducing both the diazonium cation and its radical adduct.

Continuing our previous work on visible-light-induced Pd-catalyzed remote functionalization of aliphatic alcohols with the employment of Si-based auxiliaries,^{19a} we began our studies on remote amination by subjecting standard substrate **1a** to the reactions with different electrophilic reagents. It turned out that when substrate **1a** and diazonium salt **2a** were mixed in methanol under our blue light/Pd-based conditions, in 1 h the γ -diazenylated Si-protected alcohol **3aa** was obtained in 22% GC yield (Table 1, entry 1). *To our surprise, the control experiments revealed that this reaction does not require employment of Pd catalyst or visible light to proceed with the same efficiency toward 3aa!* (entries 2, 3). The control experiment proved the necessity of base for the success of this transformation (entry 4). Excited by the obtained results, we began optimizing this novel transition-metal- and visible-light-free base-promoted remote amination by screening different solvent systems (see SI for full optimization). We realized that poor solubility of nonpolar silyl ether **1a** in methanol was a key problem; therefore a cosolvent was required for better efficiency and reproducibility. The mixture of *N,N*-dimethylacetamide (DMA) and methanol (1:1) was found to be an optimal solvent system, producing **3aa** in 60% yield (entry 5). Testing

different bases revealed that inexpensive and decently soluble in methanol $\text{HCO}_2\text{Li}\cdot\text{H}_2\text{O}$ was superior to other inorganic bases (entry 6). *Finally, using more diluted solution and decreasing the temperature to 0 °C allowed the reaction to deliver 3aa in 76% GC yield after 2 h* (entry 7).

With the optimized conditions in hand, we investigated the scope of diazonium salts (Table 2). Both electron-rich (**2b–g**) and electron-deficient (**2h–l**) diazonium salts were reactive toward remote diazenylation of **1a**, leading to the desired products **3** in good to excellent yields. We noticed that the yields were generally slightly higher in the case of more electron-rich diazonium salts. Interestingly, *ortho*-substitution at diazonium salts did not have much effect on the yield of this transformation (entries 3, 5).

The products possessing *p*-H (**4aa**), *p*-Me (**4ba**), *o*-Me (**4ca**), *p*-MeO (**4ga**), *p*-CF₃ (**4ha**), and *p*-CN (**4ia**) substituents were isolated as the γ -diazenylated alcohols **4** after *in situ* deprotection of silicon auxiliary in good yields. Based on these results, *p*-tolylN₂BF₄ (**2b**) was selected as the preferred diazenylating reagent for the remote amination of aliphatic alcohols.

The remote diazenylation of Si-tethered alcohols was found to be quite general (Scheme 2). In contrast to our recently developed Heck relay process,^{19b} the product of premature coupling was never observed in this transformation, due to the electronic mismatch of the silyl methyl radical and diazonium cation.¹⁸ First, using the optimized conditions and *p*-tolylN₂BF₄ (**2b**), the scope of γ -diazenylation was examined (Scheme 2, a). Similarly to the standard substrate **1a**, its symmetric analogue **1b** furnished the corresponding product **3b** in excellent yield. Substrates **1c** and **1d**, which possess competitive H β and H δ sites of abstraction, reacted preferentially at the γ -C–H sites, producing **2c** and **2d** in high yields. This result is in agreement with the previously observed reactivity preference of the silicon tether toward 1,6-HAT over 1,5-HAT and 1,7-HAT for tertiary sites with similar bond dissociation energies.^{19a} Alcohols possessing an unsubstituted long chain (**1e**) and –CO₂Me and –Ph-substituted chains (**1f**, **1g**, respectively) underwent γ -3° C–H diazenylation reaction in a highly efficient and regioselective manner. Moreover, functional groups such as 3-pyridyl- (**3h**), chloride- (**3i**), cyanide- (**3j**), phenylsulfide- (**4k**), *p*-methoxyphenoxide- (**4l**), *N*-tosylamide- (**3m**), *N*-phthalimide- (**3n**), carboxylic acid (**3o**), tertiary alcohol (**3s**), and alkene (**3t**) were all tolerated under these conditions (see robustness screening in the SI).²⁰ However, trace or no product was observed with substrates possessing a primary alcohol, amine, or bromide moiety (**3p**, **3q**, and **3s**, respectively). Interestingly, this method afforded the double γ -diazenylation of diol **1u**: product **3u** was isolated together with the monodiazenylated adduct **3u'**. (–)-Menthol derivative **1v** was selectively diazenylated at the isopropyl group in 82% yield. Reaction of its five-membered analogue **1w** was less selective, giving a mixture of γ - and β -diazenylated alcohols **3w** in 85% yield (4.5:1 ratio). γ -Diazenylation of C–H bonds of cyclic alcohols **1x–1z**, even in the complex setting of androsterone **1z**, proceeded uneventfully, delivering products **3k–3m** in reasonable yields. Furthermore, cyclic (**1ab–ad**) and acyclic (**1aa'**) tertiary alcohols were also compatible, giving rise to products **3aa'–ad**. Primary alcohol **1ae** furnished the desired product **4ae** in 47% yield in the γ -diazenylation reaction followed by one-pot Si-auxiliary deprotection.

Next, we investigated the scope of the kinetically less favorable β -diazenylation (Scheme 2, b). To our delight, secondary alcohols **1af** and **1ag** reacted quite efficiently under slightly modified conditions (see SI for details), producing **3af** and **3ag** in 81% and 53% yields, respectively. Diazenylation of **1ah** possessing accessible β -3° C–H, γ -2° C–H, and δ -3° C–H sites preferably occurred at the β -position, thus supporting our earlier conclusion that, for this silicon tether, the 1,5-HAT is more favorable than the 1,7-HAT.^{19a} In addition, γ - and δ -diazenylation products were also observed in this reaction as minor regioisomers. Cyclic (**1ai**, **1aj**) and bicyclic (**1ak**) alcohols underwent diazenylation reaction at the β -tertiary C–H sites, as well (products **3ai**, **3aj**, **3ak**). Moreover, this protocol was compatible with tertiary (**1al**) and primary alcohols (**1am**, **1an**), leading to the desired products **3al**, **3am**, and **3an**, respectively. Challenging δ -diazenylation was targeted as well (Scheme 2, c). Gratifyingly, reactions of secondary (**1ao**, **1ap**) and primary (**1aq**) alcohols led to the desired products as mixtures of δ - and γ -diazenylation products in good yields.

Finally, we applied this methodology for 2° C–H and even 1° C–H site functionalization (Scheme 2, d). *Both γ - and β -diazenylation reactions were accomplished for secondary (**1ar**, **1as**) and tertiary (**1at**, **1au**, **1av**) alcohols in good yields. Remarkably, this diazenylation protocol was also applicable for unactivated 1° C–H site functionalization, yielding primary diazene **3aw** in 28% yield!*

Lastly, we examined the synthetic utility of the obtained β -, γ -, and δ -diazenylated alcohols toward their highly valuable amino alcohol derivatives. Thus, the synthesized substrates were subjected to hydrogenation conditions²¹ followed by acylation to access the corresponding protected amino alcohols **5** (Scheme 3). *To our delight, the diverse range of secondary and tertiary cyclic and acyclic 1,2- 1,3- and 1,4-amino alcohols were obtained in reasonable yields via a three-step procedure starting from Si-tethered alcohols **1**.* Moreover, it was found that secondary diazenes, which are prone to isomerize into hydrazones, could efficiently be oxidized in the presence of oxygen into the corresponding ketones.²² Thus, this approach can be further used for the remote oxygenation of aliphatic alcohols²³ with secondary sites into γ -hydroxy ketones **6** (Scheme 4). This method potentially can be used to generate aldol equivalent products, which are difficult to access by other methods (**6ay**).

In summary, we have developed the first *classical-radical-initiator-, transition-metal- and visible-light-free*, auxiliary-enabled remote C–H functionalization protocol for selective γ -, β -, and δ -C(sp³)–N bond formation in aliphatic alcohols. This method features mild radical conditions and employs readily available diazonium salts to serve a double duty as the I atom abstracting agent and a coupling partner. The reaction is promoted by a weak base, lithium formate, and does not require use of transition metals or radical initiators. The scope of obtained diazenylated alcohols is very broad, and their transformations toward amino alcohols and hydroxy ketones were accomplished with good efficiency. Overall, this is a highly accessible, inexpensive, and practical method for functionalization of remote β -, γ -, and δ -unactivated C–H sites of alcohols, expected to find broad applications in synthesis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank the National Institutes of Health (GM120281) and National Science Foundation (CHE-1663779) for the financial support of this work. We also thank visiting researchers Michael Buck and Mirela Sparano and undergraduate researcher Yana McAuliff for their help with preparation of starting materials for this work.

REFERENCES

- (1). (a)Ager DJ; Prakash I; Schaad DR 1,2-Amino Alcohols and Their Heterocyclic Derivatives as Chiral Auxiliaries in Asymmetric Synthesis. *Chem. Rev* 1996, 96, 835. [PubMed: 11848773] (b)Bergmeier SC The Synthesis of Vicinal Amino Alcohols. *Tetrahedron* 2000, 56, 2561.(c)Lait SM; Rankic DA; Keay BA 1,3-Aminoalcohols and Their Derivatives in Asymmetric Organic Synthesis. *Chem. Rev* 2007, 107, 767. [PubMed: 17319729] (d)Gomes CRB; Moreth M; Cardinot D; Kopke V; Cunico W; Da Silva Lourenio MC; De Souza MVN Synthesis and Antimycobacterial Activity of Novel Amino Alcohols Containing Central Core of the Anti-HIV Drugs Lopinavir and Ritonavir. *Chem. Biol. Drug Des* 2011, 78, 1031. [PubMed: 21933353] (e)Quiliano M; Mendoza A; Fong KY; Pabón A; Goldfarb NE; Fabing I; Vettorazzi A; López de Ceraín A; Dunn BM; Garavito G; Wright DW; Deharo E; Pérez-Silanes S; Aldana I; Galiano S Exploring the scope of new arylamino alcohol derivatives: Synthesis, antimalarial evaluation, toxicological studies, and target exploration. *Int. J. Parasitol.: Drugs Drug Resist* 2016, 6, 184. [PubMed: 27718413]
- (2). (a)Davies HML; Manning JR Catalytic C-H functionalization by metal carbenoid and nitrenoid insertion. *Nature* 2008, 451, 417. [PubMed: 18216847] (b)Du Bois J Rhodium-Catalyzed C-H Amination. An Enabling Method for Chemical Synthesis. *Org. Process Res. Dev* 2011, 15, 758. [PubMed: 21804756] (c)Roizen JL; Harvey ME; Du Bois J Metal-Catalyzed Nitrogen-Atom Transfer Methods for the Oxidation of Aliphatic C-H Bonds. *Acc. Chem. Res* 2012, 45, 911. [PubMed: 22546004] (d)Jeffrey JL; Sarpong R Intramolecular C(sp³)-H amination. *Chem. Sci* 2013, 4, 4092.(e)Darses B; Rodrigues R; Neuville L; Mazurais M; Dauban P Transition metal-catalyzed iodine(III)-mediated nitrene transfer reactions: efficient tools for challenging syntheses. *Chem. Commun* 2017, 53, 493.(f)Park Y; Kim Y; Chang S Transition Metal-Catalyzed C-H Amination: Scope, Mechanism, and Applications. *Chem. Rev* 2017, 117, 9247. [PubMed: 28051855]
- (3). (a)Espino CG; Wehn PM; Chow J; Du Bois J Synthesis of 1,3-Difunctionalized Amine Derivatives through Selective C-H Bond Oxidation. *J. Am. Chem. Soc* 2001, 123, 6935.(b)Espino CG; Fiori KW; Kim M; Du Bois J Expanding the Scope of C-H Amination through Catalyst Design. *J. Am. Chem. Soc* 2004, 126, 15378. [PubMed: 15563154] (c)Espino CG; Du Bois J A Rh-Catalyzed C-H Insertion Reaction for the Oxidative Conversion of Carbamates to Oxazolidinones. *Angew. Chem., Int. Ed* 2001, 40, 598.(d)Grelrier G; Rey-Rodriguez R; Darses B; Retailleau P; Dauban P Catalytic Intramolecular C(sp³)-H Amination of Carbamimidates. *Eur. J. Org. Chem* 2017, 82, 1880.(e)Lebel H; Mamani Laparra L; Khalifa M; Trudel C; Audubert C; Szponarski M; Dicaire Leduc C; Azek E; Ernzerhof M Synthesis of Oxazolidinones: Rhodium-catalyzed C-H Amination of N-Mesyloxycarbamates. *Org. Biomol. Chem* 2017, 15, 4144. [PubMed: 28422263]
- (4). (a)Milczek E; Boudet N; Blakey S Enantioselective C-H Amination Using Cationic Ruthenium(II)-pybox Catalysts. *Angew. Chem., Int. Ed* 2008, 47, 6825.(b)Liang J-L; Yuan S-X; Huang J-S; Yu W-Y; Che C-M Highly Diastereo- and Enantioselective Intramolecular Amidation of Saturated C-H Bonds Catalyzed by Ruthenium Porphyrins. *Angew. Chem., Int. Ed* 2002, 41, 3465.(c)Zhang J-L; Huang J-S; Che C-M Oxidation Chemistry of Poly(ethylene glycol)-Supported Carbonylruthenium(II) and Dioxoruthenium(VI) *meso*-Tetrakis(pentafluorophenyl)porphyrin. *Chem. - Eur. J* 2006, 12, 3020. [PubMed: 16491497]
- (5). Alderson JM; Phelps AM; Scamp RJ; Dolan NS; Schomaker JM Ligand-Controlled, Tunable Silver-Catalyzed C-H Amination. *J. Am. Chem. Soc* 2014, 136, 16720. [PubMed: 25386769]
- (6). (a)Liu Y; Guan X; Wong EL-M; Liu P; Huang J-S; Che C-M Nonheme Iron-Mediated Amination of C(sp³)-H Bonds. Quinquepyridine-Supported Iron-Imide/Nitrene Intermediates by Experimental Studies and DFT Calculations. *J. Am. Chem. Soc* 2013, 135, 7194. [PubMed: 23634746] (b)Paradine SM; White MC Iron-Catalyzed Intramolecular Allylic C-H Amination. *J. Am. Chem. Soc* 2012, 134, 2036. [PubMed: 22260649]

- (7). Paradine SM; Griffin JR; Zhao J; Petronico AL; Miller SM; Christina White M A manganese catalyst for highly reactive yet chemoselective intramolecular C(sp³)-H amination. *Nat. Chem* 2015, 7, 987. [PubMed: 26587714]
- (8). Barman DN; Nicholas KM Copper-Catalyzed Intramolecular C-H Amination. *Eur. J. Org. Chem* 2011, 2011, 908.
- (9). Lu H; Tao J; Jones JE; Wojtas L; Zhang XP Cobalt(II)-Catalyzed Intramolecular C-H Amination with Phosphoryl Azides: Formation of 6- and 7-Membered Cyclophosphoramidates. *Org. Lett* 2010, 12, 1248. [PubMed: 20184343]
- (10). (a) Fraunhofer KJ; White MC *syn*-1,2-Amino Alcohols via Diastereoselective Allylic C-H Amination. *J. Am. Chem. Soc* 2007, 129, 7274. [PubMed: 17516648] (b) Rice GT; White MC Allylic C-H Amination for the Preparation of *syn*-1,3-Amino Alcohol Motifs. *J. Am. Chem. Soc* 2009, 131, 11707. [PubMed: 19645489] (c) Qi X; Rice GT; Lall MS; Plummer MS; White MC Diversification of a β -Lactam Pharmacophore via Allylic C-H Amination: Accelerating Effect of Lewis Acid co-catalyst. *Tetrahedron* 2010, 66, 4816. [PubMed: 21379377]
- (11). (a) Wu X; Zhang H; Tang N; Wu Z; Wang D; Ji M; Xu Y; Wang M; Zhu C Metal-free alcohol-directed regioselective heteroarylation of remote unactivated C(sp³)-H bonds. *Nat. Commun* 2018, 9, 3343. [PubMed: 30131495] (b) Friese FW; Mück-Lichtenfeld C; Studer A Remote C-H functionalization using radical translocating arylating groups. *Nat. Commun* 2018, 9, 2808. [PubMed: 30022072] (c) Zhu Y; Huang K; Pan J; Qiu X; Luo X; Qin Q; Wei J; Wen X; Zhang L; Jiao N Silver-catalyzed remote Csp³-H functionalization of aliphatic alcohols. *Nat. Commun* 2018, 9, 2625. [PubMed: 29980674] (d) Chen K; Richter JM; Baran PS 1,3-Diol Synthesis via Controlled, Radical-Mediated C-H Functionalization. *J. Am. Chem. Soc* 2008, 130, 7247. [PubMed: 18481847] (e) Short MA; Blackburn JM; Roizen JL Sulfamate Esters Guide Selective Radical-Mediated Chlorination of Aliphatic C-H Bonds. *Angew. Chem* 2018, 130, 302. (f) Sathyamoorthi S; Banerjee S; Du Bois J; Burns NZ; Zare RN Site-selective bromination of sp³ C-H bonds. *Chem. Sci* 2018, 9, 100. [PubMed: 29629078]
- (12). (a) Robertson J; Pillai J; Lush RK Radical translocation reactions in synthesis. *Chem. Soc. Rev* 2001, 30, 94. (b) Chiba S; Chen H sp³ C-H oxidation by remote H-radical shift with oxygen- and nitrogen-radicals: a recent update. *Org. Biomol. Chem* 2014, 12, 4051. [PubMed: 24781782] (c) Yan M; Lo JC; Edwards JT; Baran PS Radicals: Reactive Intermediates with Translational Potential. *J. Am. Chem. Soc* 2016, 138, 12692. [PubMed: 27631602] (d) Hu X-Q; Chen J-R; Xiao W-J Controllable Remote C-H Bond Functionalization by Visible-Light Photocatalysis. *Angew. Chem., Int. Ed* 2017, 56, 1960. (e) Green SA; Crossley SWM; Matos JLM; Vásquez-Céspedes S; Shevick SL; Shenvi RA The High Chemofidelity of Metal-Catalyzed Hydrogen Atom Transfer. *Acc. Chem. Res* 2018, 51, 2628. [PubMed: 30406655]
- (13). Wappes EA; Nakafuku KM; Nagib DA Directed β C-H Amination of Alcohols via Radical Relay Chaperones. *J. Am. Chem. Soc* 2017, 139, 10204. [PubMed: 28741940] While this manuscript was under revision, a report on light-free β C-H amination was published: Stateman LM; Wappes EA; Nakafuku KM; Edwards KM; Nagib DA Catalytic β C-H amination via an imidate radical relay. *Chem. Sci* 2019, 10, 2693. [PubMed: 30996986]
- (14). Hu A; Guo J-J; Pan H; Tang H; Gao Z; Zuo Z δ -Selective Functionalization of Alkanols Enabled by Visible-Light-Induced Ligand-to-Metal Charge Transfer. *J. Am. Chem. Soc* 2018, 140, 1612. [PubMed: 29381061]
- (15). Samanta R; Matcha K; Antonchick AP Metal-Free Oxidative Carbon-Heteroatom Bond Formation Through C-H Bond Functionalization. *Eur. J. Org. Chem* 2013, 2013, 5769.
- (16). (a) Abrams R; Lefebvre Q; Clayden J Transition Metal Free Cycloamination of Prenyl Carbamates and Ureas Promoted by Aryldiazonium Salts. *Angew. Chem., Int. Ed* 2018, 57, 13587. (b) Hartmann M; Li Y; Studer A Transition-metal-free oxyarylation of alkenes with aryl diazonium salts and TEMPO. *J. Am. Chem. Soc* 2012, 134, 16516. [PubMed: 23009829] (c) Naveen N; Sengupta S; Chandrasekaran S Metal-Free *S*-Arylation of Cysteine Using Arenediazonium Salts. *J. Org. Chem* 2018, 83, 3562. [PubMed: 29480006] (d) Koziakov D; Jacobi von Wangelin A Metal-free radical aromatic carbonylations mediated by weak bases. *Org. Biomol. Chem* 2017, 15, 6715. [PubMed: 28770941] (e) Kindt S; Wicht K; Heinrich MR Base-Induced Radical Carboamination of Nonactivated Alkenes with Aryldiazonium Salts. *Org. Lett* 2015, 17, 6122. [PubMed: 26636470]

- (17). (a)Blank O; Heinrich MR Carbodiazenylation of Olefins by Radical Iodine Transfer and Addition to Arenediazonium Salts. *Eur. J. Org. Chem* 2006, 2006, 4331. (b)Citterio A; Minisci F Free-radical diazocoupling. A new general reaction of diazonium salts. *J. Org. Chem* 1982, 47, 1759. (c)Cannella R; Clerici A; Pastori N; Regolini E; Porta O One-Pot Four-Component Reaction: Aqueous $\text{TiCl}_3/\text{PhN}^+2$ -Mediated Alkyl Radical Addition to Imines Generated in Situ. *Org. Lett* 2005, 7, 645. [PubMed: 15704915] (d)Cao L; Li C $p\text{-MeOC}_6\text{H}_4\text{N}^+ + 2\text{BF}_4^-/\text{TiCl}_3$: a novel initiator for halogen atom-transfer radical reactions in aqueous media. *Tetrahedron Lett.* 2008, 49, 7380. (e)Matcha K; Antonchick AP Cascade Multicomponent Synthesis of Indoles, Pyrazoles, and Pyridazinones by Functionalization of Alkenes. *Angew. Chem., Int. Ed* 2014, 53, 11960.
- (18). Kurandina D; Parasram M; Gevorgyan V Visible Light-Induced Room-Temperature Heck Reaction of Functionalized Alkyl Halides with Vinyl Arenes/Heteroarenes. *Angew. Chem., Int. Ed* 2017, 56, 14212.
- (19). (a)Parasram M; Chuentragool P; Wang Y; Shi Y; Gevorgyan V General, Auxiliary-Enabled Photoinduced Pd-Catalyzed Remote Desaturation of Aliphatic Alcohols. *J. Am. Chem. Soc* 2017, 139, 14857. [PubMed: 28992686] (b)Chuentragool P; Yadagiri D; Morita T; Sarkar S; Parasram M; Wang Y; Gevorgyan V Aliphatic Radical Relay Heck Reaction at Unactivated $\text{C}(\text{sp}^3)\text{-H}$ Sites of Alcohols. *Angew. Chem., Int. Ed* 2019, 58, 1794.
- (20). (a)Collins KD; Rühling A; Glorius F Application of a robustness screen for the evaluation of synthetic organic methodology. *Nat. Protoc* 2014, 9, 1348. [PubMed: 24833173] (b)Collins KD; Glorius F A robustness screen for the rapid assessment of chemical reactions. *Nat. Chem* 2013, 5, 597. [PubMed: 23787750]
- (21). Nelson HM; Patel JS; Shunatona HP; Toste FD Enantioselective α -amination enabled by a BINAM-derived phase-transfer catalyst. *Chem. Sci* 2015, 6, 170. [PubMed: 25485073]
- (22). (a)Harej M; Dolenc D Autoxidation of Hydrazones. Some New Insights. *J. Org. Chem* 2007, 72, 7214. [PubMed: 17696476] (b)Ohkatsu Y; Yoshino K; Tsuruta T Autoxidation of α -Methylbenzylhydrazones. *Sekiyu Gakkaishi* 1983, 26, 272.
- (23). Wappes EA; Vanitcha A; Nagib DA β C-H dihalogenation via iterative hydrogen atom transfer. *Chem. Sci* 2018, 9, 4500. [PubMed: 29896392]

Transition metal (TM)-catalyzed approach (well established) *Visible-light-induced approach (at initial stage of development)* **This work: Novel transition metal- and visible light-free approach**

•Room temperature reaction

•Broad scope of alcohols

•Selective β -, γ -, and δ -amination of inert $\text{C}(\text{sp}^3)\text{-H}$ bonds**Scheme 1.**

Directed Remote C–H Amination of Alcohols

Scheme 2. Scope of Transition-Metal-Free Amination of Remote Unactivated C(sp³)-H Sites of Aliphatic Alcohols:^a (a) γ -Diazenylation; (b) β -Diazenylation; (c) δ -Diazenylation; (d) Diazenylation at Secondary and Primary C-H Sites

^a0.3 or 0.4 mmol scale reactions. Isolated yield,%. ^bNMR yield, % (standard: CH₂Br₂).

^cWhen R⁵ = OH (**3p**): 20% NMR yield; R⁵ = NH₂ (**3q**): decomp; R⁵ = Br (**3r**): decomp.

^dProducts **3** were deprotected and isolated as free alcohols **4**. ^eCalculated yield, %. ^fMixture of regio- or stereoisomers: NMR yield of major isomer is represented.

Scheme 3. Transformations of Diazenylated Alcohols: Hydrogenation toward Amino Alcohols^a

^a0.3–0.4 mmol scale reactions. Isolated yield, %.

Scheme 4. Transformations of Diazenylated Alcohols: Oxygenation toward Hydroxy Ketones^a

^a0.2–0.4 mmol scale reactions. Isolated yield, %. ^bCompound synthesized via a semi-one-pot procedure from the corresponding Si-auxiliary protected alcohol **1ay**. Isolated yield over two steps, %.

Table 1.

Optimization Table

no.	cat.	base	solvent	cond	yield (3aa), ^a %	
1 ^b	[Pd] ^c	Cs ₂ CO ₃	MeOH	rt, 1 h	22	
2	[Pd] ^c	Cs ₂ CO ₃	MeOH	rt, 1 h	20	
3		Cs ₂ CO ₃	MeOH	rt, 1 h	23	
4			MeOH	rt, 1 h	0	
5		Cs ₂ CO ₃	DMA/MeOH	rt, 1 h	60	
6		HCO ₂ Li·H ₂ O	DMA/MeOH	rt, 30 min	64	
7		HCO ₂ Li·H ₂ O	DMA/MeOH ^d	0 °C to rt, 2 h	76	

^a 0.1 mmol scale reaction; GC-MS yield (standard: pentadecane).^b Reaction mixture was irradiated with visible light.^c 5 mol % Pd(OAc)₂/10 mol % Xantphos.^d 0.05 M concentration.

Table 2.

Screening of Diazonium Salts

no.	Ar	yield of 3 or 4, %
1	C ₆ H ₅ (2a)	84% ^a (3aa), 68% ^b (4aa)
2	<i>p</i> -MeC ₆ H ₄ (2b)	86% ^a (3ba), 81% ^b (4ba)
3	<i>o</i> -MeC ₆ H ₄ (2c)	69% ^b (4ca)
4	<i>m</i> -MeC ₆ H ₄ (2d)	83% ^a (3da)
5	<i>p,o</i> -diMeC ₆ H ₃ (2e)	87% ^a (3ea)
6	<i>p,m</i> -diMeC ₆ H ₃ (2f)	80% ^a (3fa)
7	<i>p</i> -MeOC ₆ H ₄ (2g)	81% ^a (3ga), 64% ^b (4ga)
8	<i>p</i> -CF ₃ C ₆ H ₄ (2h)	54% ^b (4ha)
9	<i>p</i> -CNC ₆ H ₄ (2i)	52% ^b (4ia)
10	<i>m</i> -ClC ₆ H ₄ (2j)	67% ^a (3ja)
11	<i>m</i> -FC ₆ H ₄ (2k)	73% ^a (3ka)
12	<i>p</i> -ClC ₆ H ₄ (2l)	76% ^a (3la)

^a0.1 mmol scale reaction; NMR yield (standard: CH₂Br₂).

^b0.5 mmol scale reaction, followed by deprotection with aqueous HCl (5 equiv); isolated yield.