Published in final edited form as:

Curr Opin Physiol. 2019 April ; 8: 161–169. doi:10.1016/j.cophys.2019.03.001.

Diversity of reticulospinal systems in mammals

Marie-Claude Perreault, Andrea Giorgi

Department of Physiology, Emory University School of Medicine, Atlanta, Georgia, USA

Abstract

Reticulospinal (RS) neurons provide the spinal cord with the executive signals for a large repertoire of motor and autonomic functions, ensuring at the same time that these functions are adapted to the different behavioral contexts. This requires the coordinated action of many RS neurons. In this mini-review, we examine how the RS neurons that carry out specific functions distribute across the three parts of the brain stem. Extensive overlap between populations suggests a need to explore multi-functionality at the single cell-level. We next contrast functional diversity and homogeneity in transmitter phenotype. Then, we examine the molecular genetic mechanisms that specify brain stem development and likely contribute to RS neurons identities. We advocate that a better knowledge of the developmental lineage of the RS neurons and a better knowledge of RS neuron activity across multiple behaviors will help uncover the fundamental principles behind the diversity of RS systems in mammals.

Keywords

Brain Stem; Reticular Formation; Reticulospinal; Control of Movement and Bodily Functions

Introduction

Reticulospinal (RS) neurons mediate diverse specific motor and autonomic functions including escape, micturition, reaching and grasping, sleep, respiration, vomiting, and locomotion. Referring to these RS neurons as one RS system fails to capture this functional diversity. Admittedly, the cellular and circuit organization that lies behind the diversity of the RS systems is still not well understood. Multi-functionality at the single cell-level has been little explored and so has the extent to which the different RS systems are linked to each other. For instance, we have yet to fully understand how the RS neurons that mediate motor functions are linked to the RS neurons that mediate autonomic functions so that our motor actions align to our constantly changing behavioral needs.

Correspondance : Marie-Claude Perreault, m-c.perreault@emory.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declarations of interest
None

The aim of this mini-review is to bring into light the diversity of the RS systems in term of spatial distributions, transmitter phenotypes, and molecular profiles. We focus on the role of RS neurons in motor and autonomic control but it should be kept in mind that another important role of RS neurons is sensory modulation, so that peripheral sensory inputs can ensure the smooth progression of our motor behaviors (McCrea and Perreault, 1998; Perreault et al., 1999; Rudomin, 2009). The mini-review should be placed in the context of the recently renewed interest in the RS systems, partly triggered by technological advances, which are now being applied to the brain stem. Excitingly, some of these advances promise to capture the molecular diversity of the RS neurons with new opportunities for their specific manipulation.

We give predominance to studies that have identified reticular neurons as spinally projecting neurons, either anatomically with retrograde tracing, or electrophysiologically with antidromic activation and collision test. In keeping with the editorial limits given in term of length and number of citations to older literature, many of the original studies that have shaped our current understanding of the RS systems are also left out. Several of these studies can be found in the recent review by Brownstone and Chopek [1], which has called attention to the lack of information regarding the internal organization of the reticular formation (RF) nuclei and how this information is necessary to understand how RS systems tune descending motor commands.

Spatial distribution throughout the brain stem

Nomenclature

We refer to RS neurons as spinally projecting neurons that reside within the RF of the brain stem (mesencephalon, pons, and medulla), excluding the spinally projecting neurons that have received dedicated names because of their residency in well-delineated RF nuclei (e.g., interstitial nucleus of Cajal, red nucleus, tectum, locus coeruleus, raphe nuclei). These RS neurons have axons that enter the spinal cord in the ventral funiculus, the ventral portion of the lateral funiculus, and, to a lesser extent, the dorsal portion of the lateral funiculus. RS axons in the different spinal funiculi have been grouped differently into medial and lateral reticulospinal tracts by classical anatomists and electrophysiologists, which has led to some confusion [2]. For this reason, we will have opted not to employ this terminology. Finally, with regard to the cytoarchitecture in the RF, we follow closely the nomenclature developed by Paxinos and Watson [3] which is the most commonly used for the mouse brain. Occasionally, we will also give the alternate nomenclature used for the cat brain. A direct correspondence between the two nomenclatures is, however, sometimes difficult to establish. For instance, the gigantocellular reticular nucleus pars alpha and ventralis (GiA and GiV) in the mouse most likely correspond to the nucleus reticularis magnocellularis in the cat.

Mesencephalon

The mesencephalon is not considered a major source of RS neurons but mesencephalic RS neurons can be found in the ventrolateral periaqueductal gray (PAG), the adjacent mesencephalic reticular tegmentum or pre-cuneiform/cuneiform nucleus (prCnF/CnF), and the pedunculopontine nucleus (PPTg or PPN)[4,5]. RS neurons in the prCnF/CnF project

predominantly or exclusively to the cervical and upper thoracic segments whereas the RS neurons in the PAG and PPTg/PPN project also to the lumbar segments [6,7]. Mesencephalic RS neurons in the PAG and the prCnF/CnF have not been recorded yet but mesencephalic RS neurons in the PPTg/PPN have been found to discharge during locomotor activity and also, together with non-RS neurons, during REM sleep [8]. Stimulation of the PAG and prCnF/CnF produces various defensive motor behaviors (e.g., freezing, micturition, fight, and running) [9–14] and cardiovascular and respiratory changes [15]. Stimulation of the PPTg/PPN produces different sleep/wake states [16] and low-speed locomotion [14,17]. All these functions, however, are believed to be mediated by non-RS mesencephalic neurons and to involve the activation of the RS neurons in the pons and the medulla (see below).

Pons

The pons is the second largest source of RS neurons [4,5,18]. RS neurons are found dorsally in the Barrington's nucleus (Bar) and the subcoeruleus nucleus (SubC or sublateralodorsal nucleus, SLD), and more ventrally, in the nucleus pontis oralis (PnO) and caudalis (PnC).

The pontine RS neurons in the Bar nucleus project to the sacral cord and have been implicated in the control of micturition and other pelvic motor functions [19,20,9,21]. Through their axon collaterals in the thoracolumbar segments, however, Bar RS neurons may also mediate the changes in posture and sympathetic tone that occur during these functions [21]. The pontine RS neurons in the SubC/SLD have been implicated in sleep atonia [22], a task they share with non-RS SubC/SLD neurons that project to the spinal cord via medullary RS neurons [23–28].

The pontine RS neurons in the PnO have not been attributed specific motor functions yet, whilst the pontine RS neurons in the PnC have been implicated in locomotion, reaching, and protective motor reactions [29–31]. Stimulation of the PnO nucleus decreases postural muscle tone whereas stimulation of the PnC increases it [32,33]. Despite the fact that both PnO and PnC RS neurons have axons that branch extensively in the spinal cord [34,35], PnO effect on muscle tone are thought to be mediated mostly through the activation of the RS neurons in the medulla [36]. Independently of the precise type of motor behavior they mediate, pontine RS neurons thus appear to have the capability to produce multi-segmental coordinated actions on motoneurons, a necessary competency for matching postural muscle tone to ongoing motor activity.

Medulla

Most RS neurons reside in the medulla [4,5,18] which rostral border we define operationally as the level where raphespinal neurons first appear [18]. Dorsally located medullary RS neurons are found in the dorsal paragigantocellular nucleus (DPGi), parvicellular reticular nucleus (PCRT), and the intermediate reticular nucleus (IRt). Ventrally located medullary RS neurons are found in the gigantocellular reticular nucleus (Gi including its pars alpha (GiA) and ventralis (GiV)), the parapyramidal nucleus (PPy or PPr), the Botzinger (Bo) and pre-Botzinger (PrBo) nuclei, the ventral respiratory group (VRG), and the lateral paragigantocellularis (LPGi). At the level of the pyramidal decussation, which marks the border between the medulla and the spinal cord, RS neurons are found in the medullary

reticular nucleus (or nucleus centralis), a nucleus that is divided in two parts by the IRt, the dorsal (MdD or Cd) and the ventral part (MdV or Cv).

The dorsally located medullary RS neurons in the DPGi, PCRt, IRt and MdD have been very little studied and their roles in motor control remain obscure, even though some of the nuclei in which they reside have received much attention. The DPGi, for instance, has been implicated in the regulation of sleep/wake cycle through its reticulo-reticular connections [37,24–28] but the participation of the DPGi RS neurons in this context has not been clarified. Similarly, the RS neurons in the MdD, a key region for the transmission and modulation of pain often referred to as subnucleus reticularis dorsalis or SRD, may produce pain-related motor reactions by sending axon collaterals to the RS neurons in the Gi [38]. RS neurons in PCRt and IRt have direct connections with triceps motoneurons [39] and could therefore be mediate forelimb extension but it remains unclear in which behavioral contexts.

In contrast, the ventrally located medullary RS neurons have been implicated in a wide variety of behaviors. The medullary RS in the Gi and its pars GiA and GiV have been implicated in reaching [40,41,30] and locomotion [42,43,36,29,44–46,30] with subsets of RS neurons controlling specific aspects of movement such as termination [47], muscle atonia [36,48,24,49,33] or adjustments to unexpected perturbations [50]. The medullary RS neurons in the MdV [39] may also participate in forelimb movements and those in the LPGi [51,52] in locomotor activity. Their specific contributions, however, awaits electrophysiologically documentation of ongoing discharge pattern since these RS neurons intermingle with MdV and LPGi RS neurons that control cardiovascular functions [53–55] and Bo, PrBo, and VRG RS neurons that control respiratory functions [56,57,53,58–60] (Figure 1A-C). The ventralmost RS neurons may in fact be more concerned with providing the metabolic and homeostatic support required at the onset and during motor activity than the movement itself, especially when this activity is prolonged or becomes intense (e.g., standing up from lying down, start walking, carrying a load, running for escape) [61–63].

Amino acid neurotransmitter phenotypes

As other bulbospinal neurons, RS neurons are by and large glutamatergic neurons and do not exhibit great diversity with regard to amino acid neurotransmitter phenotypes [23,64–67,39,49,47,20,52,21,28,14]. Still, a small proportion of ventrally located medullary RS neurons [68,52,69] and MLF-descending pontine RS neurons ([70], however, see [18]) are gabaergic or glycinergic. This presence of inhibitory neurons is not unique to the RS system. The vestibulospinal system also holds inhibitory spinally projecting neurons [71].

The fact that RS neurons involved in different functions are overwhelmingly glutamatergic and have overlapping spatial distributions (cf. previous section) represents a challenge for optogenetic experiments that use the VGLUT2 promoter, which may lead to apparently conflicting results [14,17]. Another difficulty is that all descending neurons that release glutamate, corticospinal neurons excepted, express VGLUT2, resulting in the activation of *en passant* glutamatergic fibers of non-RS sources. Although novel tools for intersectional approaches [72] and efficient viral receptor complementation strategies [73] are being

developed to address these issues, achieving neurotransmitter-independent specificity would be greatly beneficial. This, however, will likely require determining the molecular identity of the different groups of RS neurons.

Molecular profiles

The embryonic brain stem, under the action of Hox genes, transiently divides into a series of anteroposterior compartments called mesomeres (m1-m2) and rhombomeres (r0-r7 and pr8-pr11) which in the adult give rise to the mesencephalon and the rhombencephalon (cerebellum, pons and medulla) (Figure 1D) [74,75]. These compartments have common dorsoventral divisions (Table 1) whose precise architecture is controlled by the expression of compartment-specific developmental genes [76]. The compartment-specific interactions between transcriptional programs along the anteroposterior and the dorsoventral axes, which govern the production of specific neuronal phenotypes during development, likely give rise to the clustering of the RS neurons into defined anteroposterior and dorsoventral domains [77,78]. Transcription factors might therefore uniquely identify subsets of RS neurons.

An initial study of the molecular control underlying the development of RS neurons has evidenced RS subsets that differentially express LIM homeodomain transcription factors Lhx1/5 and Lhx3/4, and Chx10 [79]. In two different studies, it was found that about 50% of the Chx10+ RS neurons project to the cervical spinal cord [80] and 50% project to the lumbar cord [47]. It is unclear whether the two studies investigated the same contingent of cells, as the proportion of Chx10+ RS neurons projecting to cervical versus the lumbar cord has not been established. Chx10+ RS are primarily located in the medullary Gi (incl. GiV and GiA) [80,47] although a fair number of Chx10+ RS neurons are present in the medullary IRt and MdV [80] and pontine PnC [47]. Many of the Chx10+ RS neurons, especially in the Gi incl. GiV and GiA, are activated during MLR-evoked locomotion [80] whereas optogenetic activation of RS and non-RS Chx10 + neurons in the PnC neurons and rostral Gi halts ongoing locomotor activity [47].

Other major groups of RS neurons also express specific transcription factors. For instance, the RS neurons in the Bar involved in micturition express the atonal bHLH transcription factor 1 (Atoh1) [21] whereas the RS neurons in the VRG involved in breathing express either the developing brain homeobox 1 (Dbx1), but not En1 nor Sim1, or the ladybird homeobox 1 (Lbx1), but not Phox2b nor Tlx3 [69].

While determining the unique molecular signature of all the different subsets of RS neurons will clearly require more work, a molecular basis of their diversity is slowly emerging.

Concluding remarks

As we move towards a clearer understanding of the extent to which the various RS neurons assemble in distinct functional units, it will be important to understand how these functional units work together and coordinate since disruption of this coordination may itself results in pathological conditions [81]. The question as to whether RS neurons are dedicated to specific functions or whether most RS neurons can be recruited during most behaviors [82] thus remains open. In that context, determining the discharge patterns of individual RS

neurons in relation to different autonomic and motor behaviors will continue to be essential to gaining insights into how the different RS populations regulate these functions. Molecular genetic approaches that interrogate the developmental origins of RS neurons show great promise although these approaches have not quite reached the level of specificity needed to uniquely define a given RS subset. In this respect, the development of mesomere- or rhombomere-restricted reporter mice might be useful for fate-mapping analysis [83], enabling gene over-expression or knockdown for transgenic manipulation and physiological investigation that would otherwise not be possible.

The diversity of RS systems underlines their importance for nearly all our essential functions. Understanding how this diversity emerges will not only help understand how we normally perform these essential functions but also how these functions may be recovered after injury or disease. RS neurons are known to be plastic [84–89] and a number of studies have demonstrated that their plasticity can be harnessed to enhance functional recovery [90–96].

Acknowledgment:

We would like to thank all present and past collaborators who have shared our interest in the reticulospinal systems and the National Institute of Health (NIH), the Craig H Nielsen Foundation, and the International Research for Paraplegia for their support throughout the years. Both authors are currently supported from a NIH grant to M-C. P (NS085387).

Abbreviations

5N	Motor trigeminal nucleus
7N	Facial nucleus
Bar	Barrington's nucleus
Bo	Botzinger nucleus
Cb	Cerebellum
CnF/PreCnF	Cuneiform/precuneiform nucleus
Cu	Cuneate nucleus
DPGi	Dorsal paragigantocellular nucleus
Gi	Gigantocellular reticular nucleus
GiA	Gigantocellular reticular nucleus, pars alpha
GiV	Gigantocellular reticular nucleus, pars ventralis
Hox	Homeobox
IC	Inferior colliculus
IRt	Intermediate reticular nucleus

LPGi	Lateral paragigantocellularis
m1, m2...	Mesomere 1, 2...
MdD	Medullary reticular nucleus, dorsal part
MdV	Medullary reticular nucleus, ventral part
MVe	Medial vestibular nucleus
PAG	Periaqueductal gray
PCRt	Parvicellular reticular nucleus
Pn	Pontine nucleus
PnC	Nucleus pontis caudalis
PnO	Nucleus pontis oralis
PPTg or PPN	Pedunculopontine nucleus
Ppy/PPr	Parapyramidal nucleus
PrBo	Pre-Botzinger Nucleus
r1, r2...	Rhombomere 1, 2...
R	Red nucleus
REM	Rapid-eye movement
RF	Reticular formation
RS	Reticulospinal
SC	Superior colliculus
SLD	Sublaterodorsal nucleus
SO	Superior olivary complex
spVe	Spinal vestibular nucleus
SubC	Subcoeruleus nucleus
VRG	Ventral respiratory group

References

- [1]. Brownstone RM, Chopek JW, Reticulospinal Systems for Tuning Motor Commands, *Front. Neural Circuits* 12 (2018). doi:10.3389/fncir.2018.00030.
- [2]. Kuypers HGJM, Anatomy of the Descending Pathways, in: Brookhart J, Mountcastle V, Brooks V, Geiger S (Eds.), *Handb. Physiol. Nerv. Syst. Vol II Mot. Control Part I*, American Physiological Society, Bethesda, Maryland, USA, 1981. doi:10.1002/cphy.cp010213.
- [3]. Paxinos G, Watson C, *The rat brain in stereotaxic coordinates*, Academic Pr, Sydney, 1982.

- [4]. VanderHorst VGJM, Ulfhake B, The organization of the brainstem and spinal cord of the mouse: Relationships between monoaminergic, cholinergic, and spinal projection systems, *J. Chem. Neuroanat* 31 (2006) 2–36. doi:10.1016/j.jchemneu.2005.08.003. [PubMed: 16183250]
- [5]. Liang H, Paxinos G, Watson C, Projections from the brain to the spinal cord in the mouse, *Brain Struct. Funct* 215 (2011) 159–186. doi:10.1007/s00429-010-0281-x. [PubMed: 20936329]
- [6]. Castiglioni AJ, Gallaway MC, Coulter JD, Spinal projections from the midbrain in monkey, *J. Comp. Neurol* 178 (1978) 329–345. doi:10.1002/cne.901780208. [PubMed: 415074]
- [7]. Liang H, Paxinos G, Watson C, Spinal projections from the presumptive midbrain locomotor region in the mouse, *Brain Struct. Funct* 217 (2012) 211–219. doi: 10.1007/s00429-011-0337-6. [PubMed: 21735296]
- [8]. Thankachan S, Fuller PM, Lu J, Movement- and behavioral state-dependent activity of pontine reticulospinal neurons, *Neuroscience*. 221 (2012) 125–139. doi: 10.1016/j.neuroscience.2012.06.069. [PubMed: 22796072]
- [9]. Holstege G, How the Emotional Motor System Controls the Pelvic Organs, *Sex. Med. Rev* 4 (2016) 303–328. doi: 10.1016/j.sxmr.2016.04.002. [PubMed: 27872027]
- [10]. Tovote P, Esposito MS, Botta P, Chaudun F, Fadok JP, Markovic M, Wolff SBE, Ramakrishnan C, Fenno L, Deisseroth K, Herry C, Arber S, Luthi A, Midbrain circuits for defensive behaviour, *Nature*. 534 (2016) 206–212. doi: 10.1038/nature17996. [PubMed: 27279213]
- [11]. Noga BR, Opris I, The Hierarchical Circuit for Executive Control of Movement, in: Opris I, Casanova MF (Eds.), *Phys. Mind Brain Disord*, Springer International Publishing, Cham, 2017: pp. 95–127. doi: 10.1007/978-3-319-29674-6_5.
- [12]. Kim LH, Sharma S, Sharpies SA, Mayr KA, Kwok CHT, Whelan PJ, Integration of Descending Command Systems for the Generation of Context-Specific Locomotor Behaviors, *Front. Neurosci* 11 (2017). doi: 10.3389/fnins.2017.00581.
- [13]. Ferreira-Pinto MJ, Ruder L, Capelli P, Arber S, Connecting Circuits for Supraspinal Control of Locomotion, *Neuron*. 100 (2018) 361–374. doi: 10.1016/j.neuron.2018.09.015. [PubMed: 30359602]
- [14]. Caggiano V, Leiras R, Goñi-Erro H, Masini D, Bellardita C, Bouvier J, Caldeira V, Fisone G, Kiehn O, Midbrain circuits that set locomotor speed and gait selection, *Nature*. 553 (2018) 455–460. doi:10.1038/nature25448. [PubMed: 29342142]
- [15]. Subramanian HH, Holstege G, The midbrain periaqueductal gray changes the eupneic respiratory rhythm into a breathing pattern necessary for survival of the individual and of the species, in: *Prog. Brain Res*, Elsevier, 2014: pp. 351–384. doi:10.1016/B978-0-444-63488-7.00017-3.
- [16]. Kroeger D, Ferrari LL, Petit G, Mahoney CE, Fuller PM, Arrigoni E, Scammell TE, Cholinergic, Glutamatergic, and GABAergic Neurons of the Pedunculopontine Tegmental Nucleus Have Distinct Effects on Sleep/Wake Behavior in Mice, *J. Neurosci* 37 (2017) 1352–1366. doi: 10.1523/JNEUROSCI.1405-16.2016. [PubMed: 28039375] * A chemogenetic analysis of the precise role of cholinergic, glutamatergic, and GABAergic pedunculopontine neurons in driving sleep/wake behavior using designer receptors exclusively activated by designer drugs (DREADDs).
- [17]. Josset N, Roussel M, Lemieux M, Lafrance-Zoubga D, Rastqar A, Bretzner F, Distinct Contributions of Mesencephalic Locomotor Region Nuclei to Locomotor Control in the Freely Behaving Mouse, *Curr. Biol* 28 (2018) 884–901.e3. doi:10.1016/j.cub.2018.02.007. [PubMed: 29526593]
- [18]. Sivertsen MS, Perreault M-C, Glover JC, Pontine reticulospinal projections in the neonatal mouse: Internal organization and axon trajectories: Pontine Reticulospinal Projections in Newborn Mouse, *J. Comp. Neurol* 524 (2016) 1270–1291. doi: 10.1002/cne.23904. [PubMed: 26400815]
- [19]. Sasaki M, Feed-forward and feedback regulation of bladder contractility by Barrington's nucleus in cats: Feed-forward and feedback regulation of bladder contractility, *J. Physiol* 557 (2004) 287–305. doi: 10.1113/jphysiol.2003.059352. [PubMed: 15047771]
- [20]. Hou XH, Hyun M, Taranda J, Huang KW, Todd E, Feng D, Atwater E, Croney D, Zeidel ML, Osten P, Sabatini BL, Central Control Circuit for Context-Dependent Micturition, *Cell*. 167 (2016) 73–86.e12. doi:10.1016/j.cell.2016.08.073. [PubMed: 27662084] * This study identifies

glutamatergic RS neurons in the pontine micturition center (Bar), which express corticotropin-releasing hormone and are essential for context-dependent urination.

- [21]. Verstegen AMJ, Vanderhorst V, Gray PA, Zeidel ML, Geerling JC, Barrington's nucleus: Neuroanatomic landscape of the mouse "pontine micturition center": VERSTEGEN et al., J. Comp. Neurol 525 (2017) 2287–2309. doi:10.1002/cne.24215. [PubMed: 28340519]
- [22]. Weng FJ, Williams RH, Hawryluk JM, Lu J, Scammell TE, Saper CB, Arrigoni E, Carbachol excites sublaterodorsal nucleus neurons projecting to the spinal cord: Muscarinic activation of pontine atonia neurons, J. Physiol 592 (2014) 1601–1617. doi: 10.1113/jphysiol.2013.261800. [PubMed: 24344163]
- [23]. Lu J, Sherman D, Devor M, Saper CB, A putative flip–flop switch for control of REM sleep, Nature. 441 (2006) 589–594. doi:10.1038/nature04767. [PubMed: 16688184]
- [24]. Chase MH, Motor control during sleep and wakefulness: Clarifying controversies and resolving paradoxes, Sleep Med. Rev 17 (2013) 299–312. doi:10.1016/j.smrv.2012.09.003. [PubMed: 23499211]
- [25]. Arrigoni E, Chen MC, Fuller PM, The anatomical, cellular and synaptic basis of motor atonia during rapid eye movement sleep: Neural circuitry regulating REM atonia, J. Physiol 594 (2016) 5391–5414. doi:10.1113/JP271324. [PubMed: 27060683]
- [26]. Saper CB, Fuller PM, Wake–sleep circuitry: an overview, Curr. Opin. Neurobiol 44 (2017) 186–192. doi:10.1016/j.conb.2017.03.021. [PubMed: 28577468]
- [27]. Scammell TE, Arrigoni E, Upton JO, Neural Circuitry of Wakefulness and Sleep, Neuron. 93 (2017) 747–765. doi:10.1016/j.neuron.2017.01.014. [PubMed: 28231463]
- [28]. Valencia Garcia S, Brischoux F, Clément O, Libourel P-A, Arthaud S, Lazarus M, Luppi P-H, Fort P, Ventromedial medulla inhibitory neuron inactivation induces REM sleep without atonia and REM sleep behavior disorder, Nat. Commun 9 (2018). doi: 10.1038/s41467-017-02761-0.
- [29]. Prentice SD, Drew T, Contributions of the Reticulospinal System to the Postural Adjustments Occurring During Voluntary Gait Modifications, J. Neurophysiol 85 (2001) 679–698. doi: 10.1152/jn.2001.85.2.679. [PubMed: 11160503]
- [30]. Schepens B, Stapley P, Drew T, Neurons in the Pontomedullary Reticular Formation Signal Posture and Movement Both as an Integrated Behavior and Independently, J. Neurophysiol 100 (2008) 2235–2253. doi: 10.1152/jn.01381.2007. [PubMed: 18632892]
- [31]. Hormigo S, López DE, Cardoso A, Zapata G, Sepúlveda J, Castellano O, Direct and indirect nigrothalamic projections to the nucleus reticularis pontis caudalis mediate in the motor execution of the acoustic startle reflex, Brain Struct. Funct 223 (2018) 2733–2751. doi: 10.1007/s00429-018-1654-9. [PubMed: 29574585]
- [32]. Takakusaki K, Chiba R, Nozu T, Okumura T, Brainstem control of locomotion and muscle tone with special reference to the role of the mesopontine tegmentum and medullary reticulospinal systems, J. Neural Transm 123 (2016) 695–729. doi:10.1007/s00702-015-1475-4. [PubMed: 26497023]
- [33]. Takakusaki K, A hypothesis for understanding mechanisms of normal and abnormal behavior states based on operation hypothesis, (2018) 8.
- [34]. Matsuyama K, Kobayashi Y, Takakusaki K, Mori S, Kimura H, Termination mode and branching patterns of reticuloreticular and reticulospinal fibers of the nucleus reticularis pontis oralis in the cat: an anterograde PHA-L tracing study, Neurosci. Res 17 (1993) 9–21. doi: 10.1016/0168-0102(93)90024-k. [PubMed: 8414221]
- [35]. Matsuyama K, Mori F, Kuze B, Mori S, Morphology of single pontine reticulospinal axons in the lumbar enlargement of the cat: A study using the anterograde tracer PHA-L, J. Comp. Neurol 410 (1999) 413130. doi: 10.1002/(SICI)1096-9861(19990802)410:3<413::AID-CNE5>3.0.CO;2-Q.
- [36]. Iwakiri H, Oka T, Takakusaki K, Mori S, Stimulus effects of the medial pontine reticular formation and the mesencephalic locomotor region upon medullary reticulospinal neurons in acute decerebrate cats, Neurosci. Res 23 (1995) 47–53. [PubMed: 7501300]
- [37]. Sapin E, Lapray D, Béréd A, Goutagny R, Léger L, Ravassard P, Clément O, Hanriot L, Fort P, Luppi P-H, Localization of the Brainstem GABAergic Neurons Controlling Paradoxical (REM) Sleep, PLoS ONE. 4 (2009) e4272. doi:10.1371/journal.pone.0004272. [PubMed: 19169414]

- [38]. Leiras R, Martín-Cora F, Velo P, Liste T, Canedo A, Cat's medullary reticulospinal and subnucleus reticularis dorsalis noxious neurons form a coupled neural circuit through collaterals of descending axons, *J. Neurophysiol* 115 (2016) 324–344. doi:10.1152/jn.00603.2015. [PubMed: 26581870]
- [39]. Esposito MS, Capelli P, Arber S, Brainstem nucleus MdV mediates skilled forelimb motor tasks, *Nature*. 508 (2014) 351–356. doi: 10.1038/nature13023. [PubMed: 24487621]
- [40]. Schepens B, Drew T, Independent and Convergent Signals From the Pontomedullary Reticular Formation Contribute to the Control of Posture and Movement During Reaching in the Cat, *J. Neurophysiol* 92 (2004) 2217–2238. doi: 10.1152/jn.01189.2003. [PubMed: 15175364]
- [41]. Schepens B, Drew T, Descending Signals From the Pontomedullary Reticular Formation Are Bilateral, Asymmetric, and Gated During Reaching Movements in the Cat, *J. Neurophysiol* 96 (2006) 2229–2252. doi: 10.1152/jn.00342.2006. [PubMed: 16837662]
- [42]. Drew T, Dubuc R, Rossignol S, Discharge Patterns of Reticulospinal and Other Reticular Neurons in Chronic, Unrestrained Cats Waking on a Treadmill, (1986) 27.
- [43]. Perreault MC, Drew T, Rossignol S, Activity of medullary reticulospinal neurons during fictive locomotion, *J. Neurophysiol* 69 (1993) 2232–2247. [PubMed: 8350141]
- [44]. Matsuyama K, Drew T, Vestibulospinal and Reticulospinal Neuronal Activity During Locomotion in the Intact Cat. I. Walking on a Level Surface, *J. Neurophysiol* 84 (2000) 2237–2256. doi: 10.1152/jn.2000.84.5.2237. [PubMed: 11067969]
- [45]. Matsuyama K, Drew T, Vestibulospinal and Reticulospinal Neuronal Activity During Locomotion in the Intact Cat. II. Walking on an Inclined Plane, *J. Neurophysiol* 84 (2000) 2257–2276. doi: 10.1152/jn.2000.84.5.2257. [PubMed: 11067970]
- [46]. Matsuyama K, Mori F, Nakajima K, Drew T, Aoki M, Mori S, Locomotor role of the corticoreticular–reticulospinal–spinal interneuronal system, in: *Prog. Brain Res*, Elsevier, 2004: pp. 239–249. doi:10.1016/S0079-6123(03)43024-0.
- [47]. Bouvier J, Caggiano V, Leiras R, Caldeira V, Ballardita C, Balueva K, Fuchs A, Kiehn O, Descending Command Neurons in the Brainstem that Halt Locomotion, *Cell*. 163 (2015) 1191–1203. doi: 10.1016/j.cell.2015.10.074. [PubMed: 26590422]
- [48]. Brown RE, Basheer R, McKenna JT, Strecker RE, McCarley RW, Control of Sleep and Wakefulness, *Physiol. Rev* 92 (2012) 1087–1187. doi: 10.1152/physrev.00032.2011. [PubMed: 22811426]
- [49]. Weber F, Chung S, Beier KT, Xu M, Luo L, Dan Y, Control of REM sleep by ventral medulla GABAergic neurons, *Nature*. 526 (2015) 435–438. doi: 10.1038/nature14979. [PubMed: 26444238]
- [50]. Stapley PJ, Drew T, The Pontomedullary Reticular Formation Contributes to the Compensatory Postural Responses Observed Following Removal of the Support Surface in the Standing Cat, *J. Neurophysiol* 101 (2009) 1334–1350. doi:10.1152/jn.91013.2008. [PubMed: 19118108]
- [51]. Liu J, Jordan LM, Stimulation of the Parapyramidal Region of the Neonatal Rat Brain Stem Produces Locomotor-Like Activity Involving Spinal 5-HT₇ and 5-HT_{2A} Receptors, *J. Neurophysiol* 94 (2005) 1392–1404. doi:10.1152/jn.00136.2005. [PubMed: 15872068]
- [52]. Capelli P, Pivetta C, Soledad Esposito M, Arber S, Locomotor speed control circuits in the caudal brainstem, *Nature*. (2017). doi:10.1038/nature24064.
- [53]. Goodchild AK, Moon EA, Maps of cardiovascular and respiratory regions of rat ventral medulla: Focus on the caudal medulla, *J. Chem. Neuroanat* 38 (2009) 209–221. doi: 10.1016/j.jchemneu.2009.06.002. [PubMed: 19549567]
- [54]. Sclocco R, Beissner F, Desbordes G, Polimeni JR, Wald LL, Kettner NW, Kim J, Garcia RG, Renvall V, Bianchi AM, Cerutti S, Napadow V, Barbieri R, Neuroimaging brainstem circuitry supporting cardiovagal response to pain: a combined heart rate variability/ultrahigh-field (7 T) functional magnetic resonance imaging study, *Philos. Trans. R. Soc. Math. Phys. Eng. Sci* 374 (2016) 20150189. doi:10.1098/rsta.2015.0189.
- [55]. Dempsey B, Le S, Turner A, Bokiniec P, Ramadas R, Bjaalie JG, Menuet C, Neve R, Allen AM, Goodchild AK, McMullan S, Mapping and Analysis of the Connectome of Sympathetic Premotor Neurons in the Rostral Ventrolateral Medulla of the Rat Using a Volumetric Brain Atlas, *Front. Neural Circuits* 11 (2017). doi: 10.3389/fncir.2017.00009.

- [56]. Bianchi AL, Denavit-Saubie M, Champagnat J, Central control of breathing in mammals: neuronal circuitry, membrane properties, and neurotransmitters, *Physiol. Rev* 75 (1995) 1–45. doi: 10.1152/physrev.1995.75.1.1. [PubMed: 7831394]
- [57]. Iscoe S, Control of abdominal muscles, *Prog. Neurobiol* 56 (1998) 433–506. doi: 10.1016/S0301-0082(98)00046-X. [PubMed: 9775401]
- [58]. Rice CD, Weber SA, Waggoner AL, Jessell ME, Yates BJ, Mapping of neural pathways that influence diaphragm activity and project to the lumbar spinal cord in cats, *Exp. Brain Res* 203 (2010) 205–211. doi:10.1007/s00221-010-2197-3. [PubMed: 20186399]
- [59]. Road JD, Ford TW, Kirkwood PA, Connections between expiratory bulbospinal neurons and expiratory motoneurons in thoracic and upper lumbar segments of the spinal cord, *J. Neurophysiol* 109 (2013) 1837–1851. doi: 10.1152/jn.01008.2012. [PubMed: 23324322]
- [60]. Le Gal J-P, Juvin L, Cardoit L, Thoby-Brisson M, Morin D, Remote Control of Respiratory Neural Network by Spinal Locomotor Generators, *PLoS ONE*. 9 (2014) e89670. doi: 10.1371/journal.pone.0089670. [PubMed: 24586951]
- [61]. Waldrop TG, Iwamoto GA, Point:Counterpoint: Supraspinal locomotor centers do/do not contribute significantly to the hyperpnea of dynamic exercise, *J. Appl. Physiol* 100 (2006) 1077–1083. doi: 10.1152/japplphysiol.01528.2005. [PubMed: 16467394]
- [62]. Stornetta RL, Guyenet PG, C1 neurons: a nodal point for stress?, *Exp. Physiol* 103 (2018) 332–336. doi: 10.1113/EP086435. [PubMed: 29080216] * Catecholamine-synthesizing glutamatergic RS neurons in the area of the LPGi called the rostral ventrolateral medulla (RVL) are best known for their homeostatic contribution to the control of arterial pressure. This article describes how these neurons may also mediate adaptive non-homeostatic responses to acute stressors (e.g., pain, hypoxia, infection, hemorrhage).
- [63]. Cowley KC, A new conceptual framework for the integrated neural control of locomotor and sympathetic function: implications for exercise after spinal cord injury, *Appl. Physiol. Nutr. Metab* 43 (2018) 1140–1150. doi: 10.1139/apnm-2018-0310. [PubMed: 30071179]
- [64]. Stornetta RL, Identification of neurotransmitters and co-localization of transmitters in brainstem respiratory neurons, *Respir. Physiol. Neurobiol* 164 (2008) 18–27. doi: 10.1016/j.resp.2008.07.024. [PubMed: 18722563]
- [65]. Stornetta RL, Neurochemistry of bulbospinal presympathetic neurons of the medulla oblongata, *J. Chem. Neuroanat* 38 (2009) 222–230. doi:10.1016/j.jchemneu.2009.07.005. [PubMed: 19665549]
- [66]. Häggglund M, Borgius L, Dougherty KJ, Kiehn O, Activation of groups of excitatory neurons in the mammalian spinal cord or hindbrain evokes locomotion, *Nat. Neurosci* 13 (2010) 246–252. doi:10.1038/nn.2482. [PubMed: 20081850]
- [67]. Martin EM, Devidze N, Shelley DN, Westberg L, Fontaine C, Pfaff DW, Molecular and neuroanatomical characterization of single neurons in the mouse medullary gigantocellular reticular nucleus, *J. Comp. Neurol* 519 (2011) 2574–2593. doi: 10.1002/cne.22639. [PubMed: 21456014]
- [68]. Hossaini M, Goos JAC, Kohli SK, Holstege JC, Distribution of Glycine/GABA Neurons in the Ventromedial Medulla with Descending Spinal Projections and Evidence for an Ascending Glycine/GABA Projection, *PLoS ONE*. 7 (2012) e35293. doi: 10.1371/journal.pone.0035293. [PubMed: 22558137]
- [69]. Wu J, Capelli P, Bouvier J, Goulding M, Arber S, Fortin G, A V0 core neuronal circuit for inspiration, *Nat. Commun* 8 (2017). doi: 10.1038/s41467-017-00589-2.* This study shows that Dbx1 expressing RS neurons in the prBo and rostral VRG are driving phrenic motoneurons at early postnatal age.
- [70]. Du Beau A, Shakya Shrestha S, Bannatyne BA, Jalicy SM, Linnen S, Maxwell DJ, Neurotransmitter phenotypes of descending systems in the rat lumbar spinal cord, *Neuroscience*. 227 (2012) 67–79. doi: 10.1016/j.neuroscience.2012.09.037. [PubMed: 23018001]
- [71]. Wilson V, Yoshida M, Monosynaptic inhibition of neck motoneurons by the medial vestibular nucleus, *Exp. Brain Res* 9 (1969). doi:10.1007/BF00235245.
- [72]. Madisen L, Garner AR, Shimaoka D, Chuong AS, Klapoetke NC, Li L, van der Bourg A, Niino Y, Egolf L, Monetti C, Gu H, Mills M, Cheng A, Tasic B, Nguyen TN, Sunkin SM, Benucci A,

Nagy A, Miyawaki A, Helmchen F, Empson RM, Knöpfel T, Boyden ES, Reid RC, Carandini M, Zeng H, Transgenic Mice for Intersectional Targeting of Neural Sensors and Effectors with High Specificity and Performance, *Neuron*. 85 (2015) 942–958. doi:10.1016/j.neuron.2015.02.022. [PubMed: 25741722]

- [73]. Li S-J, Vaughan A, Sturgill JF, Kepecs A, A Viral Receptor Complementation Strategy to Overcome CAV-2 Tropism for Efficient Retrograde Targeting of Neurons, *Neuron*. 98 (2018) 905–917.e5. doi:10.1016/j.neuron.2018.05.028. [PubMed: 29879392]
- [74]. Marín F, Aroca P, Puelles L, Hox gene colinear expression in the avian medulla oblongata is correlated with pseudorhombomeric domains, *Dev. Biol* 323 (2008) 230–247. doi: 10.1016/j.ydbio.2008.08.017. [PubMed: 18786526]
- [75]. Tomás-Roca L, Corral-San-Miguel R, Aroca P, Puelles L, Marín F, Crypto-rhombomeres of the mouse medulla oblongata, defined by molecular and morphological features, *Brain Struct. Funct* 221 (2016) 815–838. doi:10.1007/s00429-014-0938-y. [PubMed: 25381007]
- [76]. Puelles L, Plan of the Developing Vertebrate Nervous System, in: *Patterning Cell Type Specif. Dev. CNS PNS*, Elsevier, 2013: pp. 187–209. doi:10.1016/B978-0-12-397265-1.00118-0.
- [77]. Auclair F, Marchand R, Glover JC, Regional patterning of reticulospinal and vestibulospinal neurons in the hindbrain of mouse and rat embryos, *J. Comp. Neurol* 411 (1999) 288–300. doi: 10.1002/(SICI)1096-9861(19990823)411:2<288::AID-CNE9>3.0.CO;2-U. [PubMed: 10404254]
- [78]. Perreault M-C, Glover JC, Glutamatergic reticulospinal neurons in the mouse: developmental origins, axon projections, and functional connectivity: Glutamatergic reticulospinal neurons in the mouse, *Ann. N. Y. Acad. Sci* 1279 (2013) 80–89. doi:10.1111/nyas.12054. [PubMed: 23531005]
- [79]. Cepeda-Nieto AC, Pfaff SL, Varela-Echavarria A, Homeodomain transcription factors in the development of subsets of hindbrain reticulospinal neurons, *Mol. Cell. Neurosci* 28 (2005) 30–41. doi: 10.1016/j.men.2004.06.016. [PubMed: 15607939]
- [80]. Bretzner F, Brownstone RM, Lhx3-Chx10 Reticulospinal Neurons in Locomotor Circuits, *J. Neurosci* 33 (2013) 14681–14692. doi:10.1523/JNEUROSCI.5231-12.2013. [PubMed: 24027269]
- [81]. Mileykovskiy BY, Kiyashchenko L.I., Kodama T, Lai Y-Y, Siegel JM, Activation of Pontine and Medullary Motor Inhibitory Regions Reduces Discharge in Neurons Located in the Locus Coeruleus and the Anatomical Equivalent of the Midbrain Locomotor Region, *J. Neurosci* 20 (2000) 8551–8558. doi:10.1523/JNEUROSCI.20-22-08551.2000.
- [82]. Siegel J, Behavioral functions of the reticular formation, *Brain Res. Rev* 1 (1979) 69–105. doi: 10.1016/0165-0173(79)90017-1.
- [83]. Di Bonito M, Boulland J-L, Krezel W, Setti E, Studer M, Glover JC, Loss of Projections, Functional Compensation, and Residual Deficits in the Mammalian Vestibulospinal System of Hoxb1-Deficient Mice, *ENeuro*. 2 (2015). doi:10.1523/ENEUR.0096-15.2015.* This study describes how a knockout of Hoxb1 prevents the development of the lateral vestibulospinal tract neurons and subpopulation of ipsilaterally projecting RS neurons, resulting in behavioral deficits. Some behavioral deficits in proprioception and interlimb coordination are maintained into adulthood.
- [84]. Hill CE, Beattie MS, Bresnahan JC, Degeneration and Sprouting of Identified Descending Supraspinal Axons after Contusive Spinal Cord Injury in the Rat, *Exp. Neurol* 171 (2001) 153–169. doi: 10.1006/exnr.2001.7734. [PubMed: 11520130]
- [85]. Jin Y, Transplants of Fibroblasts Genetically Modified to Express BDNF Promote Axonal Regeneration from Supraspinal Neurons Following Chronic Spinal Cord Injury, *Exp. Neurol* 177 (2002) 265–275. doi:10.1006/exnr.2002.7980. [PubMed: 12429228]
- [86]. Blesch A, Yang H, Weidner N, Hoang A, Otero D, Axonal responses to cellularly delivered NT-4/5 after spinal cord injury, *Mol. Cell. Neurosci* 27 (2004) 190–201. doi: 10.1016/j.mcn.2004.06.007. [PubMed: 15485774]
- [87]. Lee Y-S, Lin C-Y, Robertson RT, Hsiao I, Lin VW, Motor Recovery and Anatomical Evidence of Axonal Regrowth in Spinal Cord-Repaired Adult Rats, *J. Neuropathol. Exp. Neurol* 63 (2004) 233–245. doi:10.1093/jnen/63.3.223-a. [PubMed: 15055447]

- [88]. Schaal SM, Kitay BM, Cho KS, Lo TP, Barakat DJ, Marcillo AE, Sanchez AR, Andrade CM, Pearse DD, Schwann Cell Transplantation Improves Reticulospinal Axon Growth and Forelimb Strength after Severe Cervical Spinal Cord Contusion, *Cell Transplant*. 16 (2007) 207–228. doi: 10.3727/000000007783464768. [PubMed: 17503734]
- [89]. Vavrek R, Pearse DD, Fouad K, Neuronal Populations Capable of Regeneration following a Combined Treatment in Rats with Spinal Cord Transection, *J. Neurotrauma* 24 (2007) 1667–1673. doi: 10.1089/neu.2007.0290. [PubMed: 17970629]
- [90]. Ballermann M, Fouad K, Spontaneous locomotor recovery in spinal cord injured rats is accompanied by anatomical plasticity of reticulospinal fibers, *Eur. J. Neurosci* 23 (2006) 1988–1996. doi: 10.1111/j.1460-9568.2006.04726.X. [PubMed: 16630047]
- [91]. Filli L, Engmann AK, Zorner B, Weinmann O, Moraitis T, Gullo M, Kasper H, Schneider R, Schwab ME, Bridging the Gap: A Reticulo-Propriospinal Detour Bypassing an Incomplete Spinal Cord Injury, *J. Neurosci* 34 (2014) 13399–13410. doi:10.1523/JNEUROSCI.0701-14.2014. [PubMed: 25274818]
- [92]. García-Álías G, Truong K, Shah PK, Roy RR, Edgerton VR, Plasticity of subcortical pathways promote recovery of skilled hand function in rats after corticospinal and rubrospinal tract injuries, *Exp. Neurol* 266 (2015) 112–119. doi:10.1016/j.expneurol.2015.01.009. [PubMed: 25666586]
- [93]. May Z, Fenrich KK, Dahlby J, Batty NJ, Torres-Espín A, Fouad K, Following Spinal Cord Injury Transected Reticulospinal Tract Axons Develop New Collateral Inputs to Spinal Interneurons in Parallel with Locomotor Recovery, *Neural Plast.* 2017 (2017) 1–15. doi: 10.1155/2017/1932875.
- [94]. Baker SN, Perez MA, Reticulospinal Contributions to Gross Hand Function after Human Spinal Cord Injury, *J. Neurosci* 37 (2017) 9778–9784. doi:10.1523/JNEUROSCI.3368-16.2017. [PubMed: 28871033]
- [95]. Asboth L, Friedli L, Beauparlant J, Martinez-Gonzalez C, Anil S, Rey E, Baud L, Pidpruzhnykova G, Anderson MA, Shkorbatova P, Batti L, Pagès S, Kreider J, Schneider BL, Barraud O, Courtine G, Cortico–reticulo–spinal circuit reorganization enables functional recovery after severe spinal cord contusion, *Nat. Neurosci* 21 (2018) 576–588. doi: 10.1038/s41593-018-0093-5. [PubMed: 29556028]
- [96]. Darling WG, Ge J, Stilwell-Morecraft KS, Rotella D, Pizzimenti MA, Morecraft RJ, Hand Motor Recovery following Extensive Frontoparietal Cortical Injury is accompanied by Upregulated Corticoreticular Projections in the Monkey, *J. Neurosci* (2018) 0403–18. doi:10.1523/JNEUROSCI.0403-18.2018.

Highlights

- Review of reticulospinal systems implicated in control of motor and autonomic functions
- Advocate the need for single-cell recording to decipher reticulospinal activity across behaviors
- Draw attention on studies that exploit developmental identity of reticulospinal neurons to study their physiological roles

Figure 1. Schematic of parasagittal section through the brain stem at the level of the medulla. Depiction of spatial overlap in the ventral medulla of RF regions containing RS neurons implicated in **A)** movement-related, **B)** cardiovascular, and **C)** respiratory functions together with **D)** rhombomeric organization (B-C adapted from Goodchild and Moon 2009 and D adapted Toma-Rocas et al., 2016). Data from different animal species (c.f., literature cited in text) are mapped on a reference parasagittal section from an adult mouse brain stem (lateral

1 mm, Franklin and Paxinos 2007). Similar functional overlap has been reported in the other regions of the brain stem where RS neurons reside.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.
Transcription factors expressed in progenitors and post-mitotic neurons during brain stem development.

Transcription factors set up genetic diversity of brain stem neurons. Progenitor domains dP1-dP6 and p0-p3 give rise to different early- and late (L)-born cell groups. Partial list of transcription factors expressed within post-mitotic dorsal groups (dA1–dB4) and ventral groups (vO–v3L) (adapted from Gray 2013). Abbreviations: **Olig3**, Oligodendrocyte Transcription Factor 3; **Atoh1**, Atonal BHLH Transcription Factor 1; **Pax3/6/7**, Paired Box 3/6/7; **Msx1**, Msh Homeobox 1; **Ngn1/2**, Neurogenin 1/2; **Ascl1**, Achaete-Scute Family BHLH Transcription Factor 1; **Gsx2**, GS Homeobox 2; **Ptf1a**, Pancreas Associated Transcription Factor 1a; **Lbx1**, Ladybird Homeobox 1; **Dbx1/2**, Developing Brain Homeobox 1/2; **En1**, Engrailed 1; **GATA3**, GATA Binding Protein 3; **NKx2.2**, NK2 Homeobox 2; **Pou4f1**, POU Class 4 Homeobox 1; **Barh1**, BarH-like 1 Homeobox protein; **Lhx1/1b/2/5/9**, LIM Homeobox 1/1b/2/5/9; **Evx1**, Even-skipped Homeobox 1; **FoxP2/D3**, Forkhead Box P2/D3; **Phox2b**, Paired-like Homeobox 2b; **Tlx3**, T-cell leukemia homeobox 3; **Prrxl1**, Paired related Homeobox protein-like 1; **bHLHb5**, Basic Helix-loop-helix family, member b5; **Wt1**, Wilms Tumor 1; **Dmrt3**, Double sex and Mab-3 related Transcription Factor 3; **Pitx2**, Paired-like Homeodomain 2; **Chx10**, Visual System Homeobox 2.

Domains	TFs in progenitors neurons	TFs in post-mitotic neurons
dA1	Olig3, Atoh1, Pax3, Msx1	Pou4f1, Barh1, Lhx2, Lhx9, Evx1
dA2	Olig3, Ngn1, Ngn2, Pax3, Pax7, Msx1	Pou4f1, Lhx1, Lhx5, FoxP2
dA3	Olig3, Ascl1, Ngn2, Pax3, Pax6, Pax7, Gsx2	Pou4f1, Tlx3, Prrxl1, Phox2b, Lmx1b
dA4	Olig3, Ascl1, Ptf1a, Ngn2	FoxP2, Foxd3
dB1	Ascl1, Ptf1a, Ngn2, Pax3, Pax6, Pax7, Gsx1/2	Lbx1, Pax2, Lhx1, Lhx5
dB2	Lbx1	Phox2b, Atoh1
dB3	Ascl1, Pax3, Pax6, Pax7, Gsx1/2, Dbx2	Lbx1, Tlx3, Prrxl1, Pou4f1, FoxP2, Lmx1b
dB4	Ngn1, Ngn2, Pax3, Pax6, Pax7, Dbx2	Lbx1, bHLHb5, Pax2, Lhx1, Lhx5, Wt1, Dmrt3
v0d	Dbx1	Pax2
v0v	Dbx1	Evx1
v0c	Dbx1	Pitx2
v1	En1	Pax2
v2a	GATA3	Chx10
v2b	GATA3	
pMN	NKx2.2	Olig1
v3L	NKx2.2	Lmx1b, Pet1