Published in final edited form as:

ACS Catal. 2019 January 4; 9(1): 216–223. doi:10.1021/acscatal.8b04669.

Dative Directing Group Effects in Ti-Catalyzed [2+2+1] Pyrrole Synthesis: Chemo- and Regioselective Alkyne Heterocoupling

Hsin-Chun Chiu, Xin Yi See, Ian A. Tonks*

Department of Chemistry, University of Minnesota–Twin Cities, 207 Pleasant Street SE, Minneapolis, Minnesota 55455, United States

Abstract

Transient dative substrate–Ti interactions have been found to play a key role in controlling the regioselectivity of alkyne insertion and [2+2] cycloaddition in Ti-catalyzed [2+2+1] pyrrole synthesis and Ti-catalyzed alkyne hydroamination. TMS-protected alkynes with pendent Lewis basic groups can invert the regioselectivity of TMS-protected alkyne insertion, leading to the selective formation of highly substituted 3-TMS pyrroles. The competency of various potential directing groups was investigated, and it was found that the directing-group effect can be tuned by modifying the catalyst Lewis acidity, the directing-group basicity, or the directing-group tether length. Dative directing-group effects are unexplored with Ti catalysts, and this study demonstrates the potential power of dative substrate–Ti interactions in tuning selectivity.

Graphical Abstract

Keywords

directing-group effects; multicomponent reactions; pyrroles; titanium; alkynes

INTRODUCTION

Highly substituted pyrroles are important components of many bioactive natural products, drug candidates, and FDA-approved drugs.^{1–4} Recently, we have reported several multicomponent formal [2+2+1] pyrrole syntheses from alkynes and diazenes or azides

*Corresponding Author itonks@umn.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscatal.8b04669.

Full experimental procedures, characterization data, and NMR spectra (PDF)

Notes

The authors declare no competing financial interest.

(Figure 1).^{5–9} These methods provide facile, modular access to highly substituted pyrrole cores that are otherwise challenging to synthesize.

Attaining regio- and chemoselectivity can be inherently difficult [2+2+1] cycloaddition reactions such as our Ti-catalyzed pyrrole synthesis or the Pauson–Khand reaction.^{10,11} Previous studies demonstrated that the selectivity of Ti-catalyzed alkyne coupling is driven by intrinsic alkyne stereoelectronic properties, and this substrate-driven selectivity results in a statistical mixture of regioisomers.⁵ More recently, we discovered that the heterocoupling of TMS-protected alkynes can occur with exceptional regioselectivity, yielding 2-TMS pyrroles (Figure 1, top).⁸ These reactions are highly selective because of the electronic properties of the TMS-protected alkynes: they do not easily undergo the [2+2] cycloadditions with Ti=NR imidos, but they rapidly (because of their electron-richness) and regioselectively (because of the thermodynamic α -Si effect^{8,12–19}) insert into azatitanacyclobutenes. This protocol can access all possible penta- and tetrasubstituted pyrrole substitution patterns except for 1,2,3,5-tetrasubstitution.

An alternative strategy for regiocontrol is to use heteroatom-based directing groups to enforce selectivity. For example, sulfur and nitrogen-substituted alkenes have been used in the cobalt catalyzed [2+2+1] Pauson–Khand reaction to direct the product regioselectivity.^{20–22} This method has been applied further to the formal synthesis of Prostaglandin A2.²³ Removable directing groups have also been used in the Pauson–Khand reaction, notably by using silicon-tethered pyridinyl vinyl silanes which could go through facile hydrolytic desilylation.^{24,25}

Pendent alkoxide directing groups have been used extensively in Ti-catalyzed and mediated reactions to engender regio-, chemo-, and enantioselective transformations²⁶. For example, Sharpless' catalytic asymmetric alkene epoxidation relies on alkoxide directing groups to engender enantiofacial selectivity,²⁷ while more recently Micalizio has used alkoxides to direct alkyne insertion regioselectivity in Ti-mediated reductive coupling reactions (Figure 1, middle),^{18,19,28–31} and Burns has used alkoxides to direct stereoselective alkene dihalogenation and haloazidation reactions.^{32–35} Alkoxide-directed coupling reactions have also served as a platform for heterocycle syntheses, such as pyridines and indolizidines.^{36,37}

To exert complete regiocontrol over multisubstituted pyrrole synthesis, we have undertaken a study of pendent directing-group effects in Ti-catalyzed [2+2+1] pyrrole synthesis (Figure 1, bottom), envisioning that it would be possible to reverse the regiochemical course of the reaction. In contrast to anionic directing groups, the use of dative directing groups in titanium catalysis is unexplored. However, we envisioned that transient dative donor interactions could promote chemo- and regioselective reactions while also undergoing the facile ligand exchange needed for productive redox catalysis. We herein report that TMS-protected alkynes with pendent Lewis basic groups can invert the regioselectivity of TMS-protected alkyne insertion in Ti-catalyzed [2+2+1] pyrrole synthesis. This demonstrates that a dative directing-group effect can override the substrate's intrinsic selectivity. These reactions lead to the formation of 3-TMS-substituted pyrroles, which can be used in complex molecule synthesis.

RESULTS AND DISCUSSION

Unlike other TMS-protected alkynes, the [2+2+1] heterocoupling of ((2-methoxyphenyl)ethynyl)trimethylsilane (**1a**) with phenylpropyne (**2a**) was remarkably unselective for **4a-TMS**, giving a 1.1:1 ratio of **3a-TMS**:**4a-TMS** (Table 1, entry 1).⁸ We speculated that the increased yield of **3a-TMS** was a result of the *o*-OMe group precoordinating to Ti and directing second alkyne insertion in a manner opposite the electronic properties of the alkyne (Figure 2).

Thus, we sought to find conditions that favored the directed 3-TMS product **3a-TMS**. First, [py₂TiCl₂(NPh)]₂ was replaced by the more Lewis acidic catalyst THF₃TiI₂(NPh)^{38,39} to promote stronger coordination of the *o*-OMe group. This increased the **3a-TMS**:**4a-TMS** ratio to 5.5:1 (Table 1, entry 2). Lowering the reaction temperature further improved the selectivity, giving regioisomeric ratios >10.5:1 (Table 1, entries 2–6). At lower temperatures, dissociation of the *o*-OMe is likely slowed, resulting in formation of the kinetically preferred metallacycle leading to **3a-TMS** over **4a-TMS**.⁸ In our previous studies of TMS-alkyne heterocouplings, chemoselectivities were good at a 1:1 ratio of the two alkynes, but from a practical perspective, we found it better to use an excess of the TMS-alkyne in order to make product separations more simple. However, here reducing the equivalents of **1a** had little effect on yield or selectivity, indicating that the directing-group effect can more effectively outcompete **2a** homocoupling (entries 5 and 7). Thus, the conditions in entry 6 were deemed optimized.

The directing effect was explored on a range of TMS-protected alkynes with pendent Lewis basic functionality (Table 2). The optimized conditions from Table 1 were used (Condition A), along with the prior conditions for TMS-alkyne heterocoupling catalysis⁸ (Condition B).

First, substrates containing O atom donors (**1a–1h**) were tested. Substrates that can form five-membered chelates upon second insertion such as **1a**, vinyl methyl ether **1b**, and homopropargylic ethers **1c** and **1d** are excellent directing groups with the more Lewis acidic catalyst THF₃TiI₂(NPh) (Condition A), giving high regioselectivity and yields of pyrroles **3a–3d**. Directing-group effects are observed whether the 2-C linker is sp² (**1a**, **1b**) or sp³ (**1c**, **1d**) hybridized. Substrates **1b–1d** are significantly less selective with the less Lewis acidic catalyst [py₂TiCl₂(NPh)]₂, although in the case of **1d**, acceptable regioselectivity for **3d** is still obtained. Phenyl ether **1h**, which also contains a 2-C linker, manifests no directing-group effect—presumably due to a combination of the weaker aryl ether donor O and increased rotational degrees of freedom with respect to **1a**.

In contrast to 2-C linkers, shorter (1-C) linkers such as propargylic ethers (**1e**) and acetals (**1f**) do not yield productive reactivity due to C–O bond decomposition.⁴⁰ Longer (3-C) linkers (**1g**) display no directing-group effect, but are still competent for heterocoupling to the 2-unsubstituted pyrrole (**4g**).

Next, other Lewis basic heteroatom donors were examined. Arenes with an *ortho*-SMe (**1i**) or -NMe₂ (**1j**) couple with high regioselectivity to the directed products **3i** and **3j**. The less basic and softer -SMe group (**1i**) directed to higher regioselectivity (20:1) than harder -OMe

(**1a**, 9.3:1) and -NMe_2 (**1j**, 8.0:1), which is unexpected due to the preference of Ti^{IV} for harder ligands. Given the sensitivity to chelate size observed with O atom donors, the increased regioselectivity in **1i** may be a function of the S atomic size and resulting chelate angles of the S-containing ring.

Tertiary aryl amine **1j** displays good directing ability, although aliphatic pendent tertiary amines (**1l**, **1n**) do not. This may be due to an increase in tether flexibility coupled with the sterics of the 3° amine. Nonetheless, these substrates do not inhibit catalysis and are selective for heterocoupling to products **4l** and **4n**. Aryl fluorides (**1k**), alkyl thioethers (**1m**), and thioacetals (**1o**) give high yields of the nondirected products **4k**, **4m**, and **4o** with $[\text{py}_2\text{TiCl}_2(\text{NPh})]_2$, which is more functional-group tolerant than $\text{THF}_3\text{TiI}_2(\text{NPh})$ owing to reduced Lewis acidity.

Heterocyclic substituent reactivity is strongly dependent on the heterocycle basicity.⁴¹ *N*-Methylimidazole (**1s**) is too basic and inhibits catalysis for both $\text{THF}_3\text{TiI}_2(\text{NPh})$ and $[\text{py}_2\text{TiCl}_2(\text{NPh})]_2$ reactions through coordination to Ti. Pyridine (**1p**), which is a reasonably strong heterocyclic base but less basic than **1s**, inhibits catalysis with the more Lewis acidic $\text{THF}_3\text{TiI}_2(\text{NPh})$, but acts as a directing group and yields high regioselectivity (19:1) with $[\text{py}_2\text{TiCl}_2(\text{NPh})]_2$. Thiazole (**1q**), which is less basic than pyridine, acts as a selective directing group (5.2:1) with $\text{THF}_3\text{TiI}_2(\text{NPh})$, but not with the less Lewis-acidic $[\text{py}_2\text{TiCl}_2(\text{NPh})]_2$ (0.5:1). Benzoxazole (**1r**), which is similar to thiazole **1q** in basicity, yields similar selectivities but poor yields due to C–O bond cleavage, similar to other propargylic O-functional groups (**1e**, **1f**). Very weakly basic furan (**1t**) and thiophene (**1u**) pendants are not strong enough to serve as directing groups with either catalyst but still yield normal heterocoupled products **4t** and **4u**.

Internal alkyne **1w** is also suitable for directed heterocoupling, giving **3w** in moderate yield (Figure 3). This takes advantage of the extreme sensitivity of [2+2] cycloaddition to substrate sterics as well as the directing-group effect in second alkyne insertion. However, the window for success is narrow: less sterically encumbered alkynes (**1v**) predominantly homocouple due to their higher reactivity, while more sterically encumbered internal alkynes (**1x**) are too hindered even for second insertion and thus yield mainly pyrrole products of PhCCMe homocoupling.

Other internal alkynes can also serve as [2+2] partners along with directing-group-tethered TMS alkynes. For example, reaction of 3-hexyne (**2b**) maintains good chemo- and regioselectivity to form **3ab**, albeit with a longer reaction time (Figure 4, top). Additionally, internal alkynes with appended directing groups such as **1v**—which typically would undergo rapid homocoupling (vide supra)—can be heterocoupled with TMS alkynes with pendent strong directing groups such as **1p** to form products such as **3pv** (Figure 4, bottom). Unfortunately, all heterocoupling attempts with terminal alkynes have failed; these alkynes appear to be too reactive and instead undergo either alkyne trimerization or pyrrole formation via homocoupling.

[2+2+1] directed heterocoupling can enable facile synthesis of pyrrole-containing EP_1 antagonist derivatives (Figure 5).^{3,42–44} These moieties are typically synthesized through

classical Paal-Knorr syntheses, and require multistep pyrrole construction followed by postfunctionalization of the pyrrole core to introduce more intricate substitution patterns^{42,43} such as the 2-methyl-1,3,5-triaryl core of **5y**. Benzyl aryl ethers like that in **5y** do not serve as directing groups in the [2+2+1] reaction, but *o*-Me derivative **1y** is a deprotectable surrogate. Directed [2+2+1] heterocoupling of **1y** with **2a** and (*p*-BrPh)₂N₂ gives methoxy-protected **3y**, which can be deprotected to phenol **4y** prior to benzylation to **5y** in 22% overall yield (Figure 5). The advantage of this protocol is that the [2+2+1] reaction is widely tolerant of substitution on both azobenzene⁷ and alkyne⁸ fragments, allowing access to diverse structures with relative ease.

In addition to alkyne insertion, directing-group effects can also be seen during the [2+2] cycloaddition step (Figure 6). For example, reaction of **1z** under catalytic conditions results in exclusive formation of **6z**. In contrast, the reaction of 1-phenylpropyne gives a mixture of all 3 possible regioisomers.^{5,7,45–47} Since **1z** has little steric or electronic bias for [2+2] cycloaddition or insertion, it is likely that the –OMe group directs the [2+2] cycloaddition. Similarly, reaction of **1v** results exclusively in formation of **6v**.

Having established that directing groups can influence the regiochemistry of [2+2] cycloaddition, the possibility of performing substrate-directed Ti-catalyzed alkyne hydroamination was explored.⁴⁸ Reaction of electronically unbiased substrate **1z** with aniline resulted in selective formation of **7z**, after directed [2+2] cycloaddition and protonolysis of the metallacycle by aniline (Figure 7). This reaction represents the first example of Ti-catalyzed directed alkyne hydroamination, and it yields electronically unbiased dissymmetric ketones that are difficult to synthesize selectively via Wacker oxidation of internal alkenes.^{49–54} It is notable that weakly Lewis-basic dialkyl ether of **1z** is capable of directing selective alkyne [2+2] cycloaddition despite stoichiometric quantities of aniline.

CONCLUSIONS

In summary, we have shown that TMS-protected alkynes with pendent Lewis basic groups can invert the regioselectivity of TMS-protected alkyne insertion, leading to selective formation of highly substituted 3-TMS pyrroles. This method provides a complementary approach to regioselective [2+2+1] pyrrole formation and also demonstrates how dative substrate–Ti interactions can influence—and, importantly, tune—regioselectivity through directing-group effects, which was previously unexplored in Ti catalysis.

The inverse regioselectivity highly depends on the coordinating strength of the functional groups, the tether length, and the Lewis acidity of the Ti catalyst. These effects work in concert, providing a high degree of tunability of the pyrrole regiochemistry. Balancing the directing-group Lewis basicity with the catalyst Lewis acidity is critical for catalysis: too strong of an acid–base pairing inhibits catalysis, while too weak of a pairing results in formation of 2-TMS pyrrole **4** instead of the 3-TMS pyrrole **3**. Further, we have shown that dative directing-group effects can also play a role in other Ti-catalyzed reactions such as alkyne hydroamination, which leads to the selective formation of electronically unbiased imines/ketones via directed [2+2] alkyne/Ti=NR cycloaddition. This work will provide a

basis for the development of other, selective Ti-catalyzed reactions that can exploit transient dative interactions, expanding directed Ti catalysis beyond classical alkoxide directing groups.

EXPERIMENTAL SECTION

General Considerations.

All air- and moisture-sensitive compounds were manipulated in a glovebox under a nitrogen atmosphere. The solvent, PhCF₃, was predried in a Pure Process Technology Solvent System and passed through activated basic alumina before being used. All high-boiling liquid reagents were freeze pump-thawed three times, brought into the glovebox, diluted in hexanes or diethyl ether, passed through activated basic alumina, pumped dry *in vacuo* and checked by ¹H NMR in CDCl₃ to ensure the dryness. [(py)₂TiCl₂NPh]₂,⁵⁵ (THF)₃TiI₂(NPh)⁸, ((2-methoxyphenyl)ethynyl)trimethylsilane (**1a**), 2-((trimethylsilyl)ethynyl)pyridine (**1p**)⁸, trimethyl(thiophen-2-ylethynyl)silane (**1u**)⁸, 1-methoxy-2-(prop-1-yn-1-yl)benzene (**1v**)⁸, and (*E*)-1,2-bis(4-bromophenyl)diazene⁵ were prepared according to reported procedure. 1-Phenyl-1-propyne (**2a**) and 3-hexyne (**2b**) were purchased from Sigma-Aldrich. Azobenzene was purchased from TCI Chemicals and purified by dissolving it in hexanes and washing with water. The organic layer was collected, dried over Na₂SO₄, and the solvent was removed *in vacuo*.

¹H, ¹³C, ¹H-¹³C and ¹H-¹⁵N HMBC, NOESY, and No-D ¹H NMR spectra were collected on Bruker Avance III HD NanoBay 400 MHz, Bruker Avance III HD 500 MHz, or Varian Inova 500 MHz spectrometers. High-resolution mass spectra were collected on Agilent 7200 GC/QTOF-MS, Bruker BioTOF II ESI/TOF-MS, or AB-Sciex 4800 MALDI/TOF-MS. Chemical shifts are reported with references of residual protio-solvent impurity: ¹H (s, 7.16 ppm for C₆D₅H; s, 7.27 for ppm of CHCl₃), ¹³C (t, 128.39 ppm for C₆D₆; t, 77.23 ppm for CDCl₃). No-D NMR spectrum were referenced to the proton signal of the internal standard triphenylmethane (Ph₃CH, ppm = 5.0) at a delay time = 30 s and acquisition time = 5 s.

Example Reaction Condition Optimization (Table 1).

Example for Entry 2.—In a glovebox, to an NMR tube was added PhNNPh (8.2 mg, 0.045 mmol, 0.5 equiv), (THF)₃I₂TiNPh (6.1 mg, 0.01 mmol, 0.11 equiv), ((2-methoxyphenyl)ethynyl)trimethylsilane (**1a**) (20.4 mg, 0.1 mmol, 1.1 equiv), phenylpropyne (**2a**) (11.6 mg, 0.1 mmol, 1.1 equiv), triphenylmethane (4.9 mg, 0.02 mmol, 0.22 equiv) as the internal standard, and 0.5 mL of PhCF₃. The NMR tube was sealed and brought out of the glovebox, and a *t* = 0 h No-D ¹H NMR spectrum was taken. The NMR tube was heated at 115 °C in an oil bath. No-D ¹H NMR spectra were taken every 15 min until the full consumption of PhNNPh. Yields are reported with respect to **2a**.

Example Synthesis and Isolation of Multisubstituted Pyrroles via Heterocoupling Reactions (Table 2, Condition A).

Example for ((2-Methoxyphenyl)ethynyl)- trimethylsilane (1a).—In a glovebox, to a 7 mL vial was added PhNNPh (41.0 mg, 0.225 mmol, 0.5 equiv), (THF)₃I₂TiNPh (30.5 mg, 0.05 mmol, 0.11 equiv), ((2-methoxyphenyl)-ethynyl)trimethylsilane (**1a**) (102 mg, 0.5

mmol, 1.1 equiv), phenylpropyne (**2a**) (58.0 mg, 0.5 mmol, 1.1 equiv), and 2.5 mL of PhCF₃. The reaction was then sealed with a Teflon screw cap and heated at 60 °C outside of the glovebox in an oil bath for 2 h. The reaction was diluted by 2 mL of ethyl acetate, quenched by 2 mL of 2 M HCl in methanol for 1 h, washed with DI water, dried over Na₂SO₄, and concentrated prior to purification. The product **3a** was isolated by silica gel chromatography using 5% EA/Hex eluent to give a paleyellow oil. (120 mg, 64% corrected yield). Yields are reported with respect to **2a**.

Example Synthesis and Isolation of Multisubstituted Pyrroles via Heterocoupling Reactions (Table 2, Condition B).

Example for (5-(Benzyloxy)pent-1-yn-1-yl)- trimethylsilane (1g).—In a glovebox, to a 7 mL vial was added PhNNPh (41.0 mg, 0.225 mmol, 0.5 equiv), [(py)₂Cl₂TiNPh]₂ (18.4 mg, 0.025 mmol, 0.055 equiv), (5-(benzyloxy)pent-1-yn-1-yl)trimethylsilane (**1g**) (246 mg, 0.1 mmol, 2.2 equiv), phenylpropyne (**2a**) (58.0 mg, 0.5 mmol, 1.1 equiv), and 2.5 mL of PhCF₃. The reaction was then sealed with a Teflon screw cap and heated at 115 °C outside of the glovebox in an oil bath for 3 h. The reaction was diluted by 2 mL of ethyl acetate, quenched by 2 mL of 2 M HCl in methanol for 1 h, washed with DI water, dried over Na₂SO₄, and concentrated prior to the purification. The product **4g** was isolated by silica gel chromatography using 2% EA/Hex eluent to give a colorless oil (166 mg, 87% yield). Yields are reported with respect to **2a**.

Example No-D ¹H NMR Determination of Pyrrole Regioisomeric Ratios (Table 2, Condition A).

Example for ((2-Methoxyphenyl)ethynyl)trimethylsilane (1a).—In a glovebox, to an NMR tube was added PhNNPh (8.2 mg, 0.045 mmol, 0.5 equiv), (THF)₃I₂TiNPh (6.1 mg, 0.01 mmol, 0.11 equiv), ((2-methoxyphenyl)ethynyl)trimethylsilane (**1a**) (20.4 mg, 0.1 mmol, 1.1 equiv), phenylpropyne (**2a**) (11.6 mg, 0.1 mmol, 1.1 equiv), triphenylmethane (4.9 mg, 0.02 mmol, 0.22 equiv) as the internal standard, and 0.5 mL of PhCF₃. The NMR tube was sealed and brought out of the glovebox, and a *t* = 0 h No-D ¹H NMR spectrum was taken. The NMR tube was heated at 60 °C in an oil bath for 2 h. The NMR tube was then cooled to ambient temperature, and a *t* = 2 h No-D ¹H NMR spectrum was taken. Yields are reported with respect to **2a**.

Example No-D ¹H NMR Determination of Pyrrole Regioisomeric Ratios (Table 2, Condition B).

Example for (5-(Benzyloxy)pent-1-yn-1-yl)trimethylsilane (1g).—In a glovebox, to an NMR tube was added PhNNPh (8.2 mg, 0.045 mmol, 0.5 equiv), [(py)₂Cl₂TiNPh]₂ (6.1 mg, 0.005 mmol, 0.055 equiv), (5-(benzyloxy)pent-1-yn-1-yl)trimethylsilane (**1g**) (49.2 mg, 0.2 mmol, 2.2 equiv), phenylpropyne (**2a**) (11.6 mg, 0.1 mmol, 1.1 equiv), triphenylmethane (4.9 mg, 0.02 mmol, 0.22 equiv) as the internal standard, and 0.5 mL of PhCF₃. The NMR tube was sealed and brought out of the glovebox, and a *t* = 0 h No-D ¹H NMR spectrum was taken. The NMR tube was heated at 115 °C in an oil bath for 3 h. The NMR tube was then cooled to ambient temperature, and a *t* = 3 h No-D ¹H NMR spectrum was taken. Yields are reported with respect to **2a**.

Synthesis of 5-(5-Bromo-2-((2,3-difluorobenzyl)oxy)-phenyl)-1-(4-bromophenyl)-2-methyl-3-phenyl-1*H*-pyrrole (**5y**).

Reaction of **1y with **2a** to form **3y**.**—In a glovebox, to a 20 mL of vial was added 283 mg of **1y** (1 eq., 1 mmol), 116 mg of **2a** (1 eq., 1 mmol), 166 mg of (*E*)-1,2-bis(4-bromophenyl)diazene (0.5 eq., 0.5 mmol), 15 mg of (THF)₃I₂TiNPh (2.5%, 0.025 mmol), and 5 mL of PhCF₃. The reaction was then sealed with a Teflon screw cap and heated in the glovebox for 30 h at 60 °C. After being cooled down to ambient temperature, the reaction was removed from the glovebox, 10 mL of hexanes was added to the reaction, and the mixture was filtered through a fine frit. The filtrate was concentrated in vacuo and mixed with 10 mL of 2 M HCl in methanol. After the mixture was stirred for 2 h, it was diluted by 20 mL of DI water and extracted by hexanes (15 mL × 3). The organic layer was washed with brine, dried by MgSO₄, and concentrated in vacuo to get a yellow solid, **3y**, that was carried on in the reaction as-is.

Deprotection of **3y to **4y**.**—In a glovebox, the entirety of **3y** was dissolved in 2 mL of 2,4,6-collidine and transferred to a 7 mL vial, followed by the addition of 266 mg LiI (2 eq., 2 mmol). The vial was sealed and heated outside of the glovebox at 175 °C for 30 h. The reaction was diluted by 20 mL of methanol and 20 mL of DI water and then acidified by 20 mL of 2 M HCl in methanol. The mixture was extracted by hexanes (15 mL × 6), and the organic layer was washed with brine and dried by MgSO₄. Removal of the volatiles yielded the crude oily product **4y**, which was directly used in the next step.

Benylation of **4y to **5y**.**—Next, 0.2 g of K₂CO₃ (1.44 equiv, 1.44 mmol), 0.1 mL 2,3-difluorobenzyl bromide (0.78 equiv, 0.78 mmol) and 3 mL MeCN were added. The mixture was stirred at reflux for 3 h. The mixture was diluted by 10 mL of DI water and extracted by hexanes (20 mL × 3). The collected organic layer was washed with brine, dried by MgSO₄, and concentrated. The only purification of the final product was carried out via silica gel chromatography using 12% DCM/Hex eluent to give a white solid **5y** (135 mg, 22% overall yield with respect to **1y**). ¹H NMR (400 MHz, C₆D₆): δ 7.62–7.60 (m, 2H), 7.35 (t, ³J_{HH} = 7.8 Hz, 2H), 7.24 (dd, ³J_{HH} = 7.5 Hz, ⁴J_{HH} = 1.7 Hz, 1H), 7.18–7.15 (m, 1H), 6.97–6.93 (m, 2H), 6.90 (td, ³J_{HH} = 7.9 Hz, ⁴J_{HH} = 1.7 Hz, 1H), 6.77–6.71 (m, 3H), 6.65–6.58 (m, 3H), 6.54 (tdd, ³J_{HH} = 7.9, 5.0 Hz, ⁴J_{HH} = 1.4 Hz, 1H), 6.39 (d, ³J_{HH} = 8.0 Hz, 1H), 4.57 (s, 2H, –O–CH₂–Ar), 2.08 (s, 3H, 2-NC₄–CH₃) ppm. ¹³C NMR (101 MHz, C₆D₆): δ 156.0, 150.7 (dd, ¹J_{CF} = 249 Hz, ²J_{CF} = 12.4 Hz), 148.3 (dd, ¹J_{CF} = 249 Hz, ²J_{CF} = 13.2 Hz), 138.9, 137.8, 132.3, 131.9, 130.3, 129.8, 128.9, 128.9, 128.6, 127.5 (d, ³J_{CF} = 10.9 Hz), 126.6, 125.9, 124.4 (dd, ²J_{CF} = 6.7 Hz, ³J_{CF} = 4.7 Hz), 124.0 (app t, ^{2,3}J_{CF} = 3.1 Hz), 123.8 (d, ³J_{CF} = 1.1 Hz), 121.5, 120.8, 116.6, 116.4, 113.2, 111.8, 63.6 (dd, ³J_{CF} = 4.5 Hz, ⁴J_{CF} = 3.0 Hz), 12.7 ppm. MALDI-HRMS: calcd for C₃₀H₂₁NOBr₂F₂ [M–Br⁺], 528.0775; found, 528.0305.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

Financial support was provided by the National Institutes of Health (1R35GM119457) and the Alfred P. Sloan Foundation (I.A.T. is a 2017 Sloan Fellow). Equipment for the Chemistry Department NMR facility was supported through a grant from the National Institutes of Health (S10OD011952) with matching funds from the University of Minnesota.

REFERENCES

- (1). Fan H; Peng J; Hamann MT; Hu J-F Lamellarins and Related Pyrrole-Derived Alkaloids from Marine Organisms. *Chem. Rev* 2008, 108, 264–287. [PubMed: 18095718]
- (2). Baumann M; Baxendale IR; Ley SV; Nikbin N An Overview of the Key Routes to the Best Selling 5-Membered Ring Heterocyclic Pharmaceuticals. *Beilstein J. Org. Chem* 2011, 7, 442–495. [PubMed: 21647262]
- (3). Bhardwaj V; Gumber D; Abbot V; Dhiman S; Sharma P Pyrrole: A Resourceful Small Molecule in Key Medicinal Hetero-Aromatics. *RSC Adv* 2015, 5, 15233–15266.
- (4). Khajuria R; Dham S; Kapoor KK Active Methylenes in the Synthesis of a Pyrrole Motif: An Imperative Structural Unit of Pharmaceuticals, Natural Products and Optoelectronic Materials. *RSC Adv*. 2016, 6, 37039–37066.
- (5). Gilbert ZW; Hue RJ; Tonks IA Catalytic Formal [2+2+1] Synthesis of Pyrroles from Alkynes and Diazenes via $\text{Ti}^{\text{II}}/\text{Ti}^{\text{IV}}$ Redox Catalysis. *Nat. Chem* 2016, 8, 63–68. [PubMed: 26673265]
- (6). Davis-Gilbert ZW; Tonks IA Titanium Redox Catalysis: Insights and Applications of an Earth-abundant Base Metal. *Dalton Trans.* 2017, 46, 11522–11528. [PubMed: 28795719]
- (7). Davis-Gilbert ZW; Wen X; Goodpaster JD; Tonks IA Mechanism of Ti-Catalyzed Oxidative Nitrene Transfer in [2+2+1] Pyrrole Synthesis from Alkynes and Azobenzene. *J. Am. Chem. Soc* 2018, 140, 7267–7281. [PubMed: 29763560]
- (8). Chiu H-C; Tonks IA Trimethylsilyl-Protected Alkynes as Selective Cross-Coupling Partners in Titanium-Catalyzed [2+2+1] Pyrrole Synthesis. *Angew. Chem. Int. Ed* 2018, 57, 6090–6094.
- (9). Pearce AJ; See XY; Tonks IA Oxidative Nitrene Transfer from Azides to Alkynes via $\text{Ti}(\text{ii})/\text{Ti}(\text{iv})$ Redox Catalysis: Formal [2+2+1] Synthesis of Pyrroles. *Chem. Commun* 2018, 54, 6891–6894.
- (10). Fager-Jokela E; Muuronen M; Patzschke M; Helaja J Electronic Regioselectivity of Diarylalkynes in Cobalt-Mediated Pauson–Khand Reaction: An Experimental and Computational Study with Para- and Meta-Substituted Diarylalkynes and Norbornene. *J. Org. Chem* 2012, 77, 9134–9147. [PubMed: 22985418]
- (11). Fager-Jokela E; Muuronen M; Khaizourane H; Vázquez-Romero A; Verdaguer X; Riera A; Helaja J Regioselectivity of Intermolecular Pauson–Khand Reaction of Aliphatic Alkynes: Experimental and Theoretical Study of the Effect of Alkyne Polarization. *J. Org. Chem* 2014, 79, 10999–11010. [PubMed: 25321602]
- (12). Stockis A; Hoffmann R Metallacyclopentanes and Bisolefin Complexes. *J. Am. Chem. Soc* 1980, 102, 2952–2962.
- (13). Skibbe V; Erker G Zur Bildung Metallacyclischer Fünfringverbindungen: Regioselektivität der Addition von Alkinen an Zirconocenkomplexe. *J. Organomet. Chem* 1983, 241, 15–26.
- (14). Negishi E; Holmes SJ; Tour JM; Miller JA; Cederbaum FE; Swanson DR; Takahashi T Novel Bicyclization of Enynes and Diynes Promoted by Zirconocene Derivatives and Conversion of Zirconabicycles into Bicyclic Enones via Carbonylation. *J. Am. Chem. Soc* 1989, 111, 3336–3346.
- (15). Lefebvre C; Ohff A; Tillack A; Baumann W; Kempe R; Burlakov VV; Rosenthal U Darstellung und Regioselektive Reaktionen des Phosphinfreien Zirconocen-Alkin-Komplexes $\text{Cp}_2\text{Zr}(\text{THF})(\text{BuC}_2\text{SiMe}_3)$. *J. Organomet. Chem* 1995, 501, 189–194.
- (16). List AK; Koo K; Rheingold AL; Hillhouse GL Preparation of Trimethylsilyl Alkyne Complexes of Bis-(pentamethylcyclopentadienyl)zirconium, $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zr}(\text{RC}\equiv\text{CSiMe}_3)$, and Their Regioselective Reactions with Nitrous Oxide. *Inorg. Chim. Acta* 1998, 270, 399–404.
- (17). Gao Y; Shirai M; Sato F Synthesis of Substituted Pyrroles from an Alkyne, an Imine and Carbon Monoxide via an Organotitanium Intermediate. *Tetrahedron Lett.* 1996, 37, 7787–7790.

- (18). Mizoguchi H; Micalizio GC Synthesis of Angularly Substituted trans-Fused Decalins through a Metallacycle-Mediated Annulative Cross-Coupling Cascade. *Angew. Chem. Int. Ed* 2016, 55, 13099–13103.
- (19). Jeso V; Aquino C; Cheng X; Mizoguchi H; Nakashige M; Micalizio GC Synthesis of Angularly Substituted Trans-Fused Hydroindanes by Convergent Coupling of Acyclic Precursors. *J. Am. Chem. Soc* 2014, 136, 8209–8212. [PubMed: 24856045]
- (20). Krafft ME Regiocontrol in the Intermolecular Cobalt-Catalyzed Olefin-Acetylene Cycloaddition. *J. Am. Chem. Soc* 1988, 110, 968–970.
- (21). Krafft ME; Juliano CA; Scott IL; Wright C; McEachin MD The Directed Pauson-Khand Reaction. *J. Am. Chem. Soc* 1991, 113, 1693–1703.
- (22). Krafft ME; Juliano CA Studies on the Use of Bidentate Ligands in the Directed Pauson-Khand Reaction. *J. Org. Chem* 1992, 57, 5106–5115.
- (23). Krafft ME; Wright C A Formal Total Synthesis of PGA2 Using the Directed Pauson-Khand Reaction. *Tetrahedron Lett.* 1992, 33, 151–152.
- (24). Itami K; Mitsudo K; Yoshida J. i. A Pyridylsilyl Group Expands the Scope of Catalytic Intermolecular Pauson–Khand Reactions. *Angew. Chem., Int. Ed* 2002, 41, 3481–3484.
- (25). Itami K; Mitsudo K; Fujita K; Ohashi Y; Yoshida J.-i. Catalytic Intermolecular Pauson–Khand-Type Reaction: Strong Directing Effect of Pyridylsilyl and Pyrimidylsilyl Groups and Isolation of Ru Complexes Relevant to Catalytic Reaction. *J. Am. Chem. Soc* 2004, 126, 11058–11066. [PubMed: 15339192]
- (26). Hoveyda AH; Evans DA; Fu GC Substrate-Directable Chemical Reactions. *Chem. Rev* 1993, 93, 1307–1370.
- (27). Gao Y; Klunder JM; Hanson RM; Masamune H; Ko SY; Sharpless KB Catalytic Asymmetric Epoxidation and Kinetic Resolution: Modified Procedures Including in situ Derivatization. *J. Am. Chem. Soc* 1987, 109, 5765–5780.
- (28). Ryan J; Micalizio GC An Alkoxide-Directed Carbometalation of Internal Alkynes. *J. Am. Chem. Soc* 2006, 128, 2764–2765. [PubMed: 16506731]
- (29). Micalizio GC; Mizoguchi H The Development of Alkoxide-Directed Metallacycle-Mediated Annulative Cross-Coupling Chemistry. *Isr. J. Chem* 2017, 57, 228–238. [PubMed: 28652644]
- (30). Umemura S; McLaughlin M; Micalizio GC Convergent Synthesis of Stereodefined exo-Alkylidene γ -Lactams from β -Halo Allylic Alcohols. *Org. Lett* 2009, 11, 5402–5405. [PubMed: 19877599]
- (31). McLaughlin M; Takahashi M; Micalizio GC An Alkoxide-Directed Intermolecular [2+2+1] Annulation: A Three-Component Coupling Reaction for the Synthesis of Tetrasubstituted $\alpha\beta$ -Unsaturated γ -Lactams. *Angew. Chem. Int. Ed* 2007, 46, 3912–3914.
- (32). Hu DX; Seidl FJ; Bucher C; Burns NZ Catalytic Chemo-, Regio-, and Enantioselective Bromochlorination of Allylic Alcohols. *J. Am. Chem. Soc* 2015, 137, 3795–3798. [PubMed: 25738419]
- (33). Landry ML; Hu DX; McKenna GM; Burns NZ Catalytic Enantioselective Dihalogenation and the Selective Synthesis of (–)-Deschloromylipin A and (–)-Danicalipin A. *J. Am. Chem. Soc* 2016, 138, 5150–5158. [PubMed: 27018981]
- (34). Landry ML; Burns NZ Catalytic Enantioselective Dihalogenation in Total Synthesis. *Acc. Chem. Res* 2018, 51, 1260–1271. [PubMed: 29664281]
- (35). Seidl FJ; Min C; Lopez JA; Burns NZ Catalytic Regio- and Enantioselective Haloazidation of Allylic Alcohols. *J. Am. Chem. Soc* 2018, 140, 15646–15650. [PubMed: 30403852]
- (36). Yang D; Micalizio GC A Convergent Stereoselective Synthesis of Quinolizidines and Indolizidines: Chemoselective Coupling of 2-Hydroxymethyl-Substituted Allylic Silanes with Imines. *J. Am. Chem. Soc* 2009, 131, 17548–17549. [PubMed: 19950989]
- (37). Chen MZ; Micalizio GC Three-Component Coupling Sequence for the Regiospecific Synthesis of Substituted Pyridines. *J. Am. Chem. Soc* 2012, 134, 1352–1356. [PubMed: 22103772]
- (38). Billow BS; McDaniel TJ; Odom AL Quantifying Ligand Effects in High-Oxidation-State Metal Catalysis. *Nat. Chem* 2017, 9, 837. [PubMed: 28837168]

- (39). DiFranco SA; Maciulis NA; Staples RJ; Batrice RJ; Odom AL Evaluation of Donor and Steric Properties of Anionic Ligands on High Valent Transition Metals. *Inorg. Chem* 2012, 51, 1187–1200. [PubMed: 22200335]
- (40). Okamoto S Synthetic Reactions Using Low-valent Titanium Reagents Derived from Ti(OR)_4 or CpTiX_3 (X = O-i-Pr or Cl) in the Presence of Me_3SiCl and Mg. *Chem. Rec* 2016, 16, 857–872. [PubMed: 26969349]
- (41). Lide DR *CRC Handbook of Chemistry and Physics*; CRC Press: Boca Raton, FL, 2005.
- (42). Hall A; Atkinson S; Brown SH; Chessell IP; Chowdhury A; Clayton NM; Coleman T; Giblin GMP; Gleave RJ; Hammond B; Healy MP; Johnson MR; Michel AD; Naylor A; Novelli R; Spalding DJ; Tang SP Structure–Activity Relationships of 1,5-Biaryl Pyrroles as EP1 Receptor Antagonists. *Bioorg. Med. Chem. Lett* 2006, 16, 3657–3662. [PubMed: 16697196]
- (43). Hall A; Atkinson S; Brown SH; Chessell IP; Chowdhury A; Giblin GMP; Goldsmith P; Healy MP; Jandu KS; Johnson MR; Michel AD; Naylor A; Sweeting JA Discovery of Novel, Non-Acidic 1,5-Biaryl Pyrrole EP1 Receptor Antagonists. *Bioorg. Med. Chem. Lett* 2007, 17, 1200–1205. [PubMed: 17196385]
- (44). Hall A; Brown SH; Chessell IP; Chowdhury A; Clayton NM; Coleman T; Giblin GMP; Hammond B; Healy MP; Johnson MR; Metcalf A; Michel AD; Naylor A; Novelli R; Spalding DJ; Sweeting J 1,5-Biaryl Pyrrole Derivatives as EP1 Receptor Antagonists: Structure–Activity Relationships of 4- and 5-Substituted Benzoic Acid Derivatives. *Bioorg. Med. Chem. Lett* 2007, 17, 732–735. [PubMed: 17098427]
- (45). Haak E; Bytschkov I; Doye S Intermolecular Hydroamination of Alkynes Catalyzed by Dimethyltitanocene. *Angew. Chem., Int. Ed* 1999, 38, 3389–3391.
- (46). Khedkar V; Tillack A; Beller M A Dramatic Effect of Aryloxo Ligands on the Titanium-Catalyzed Hydroamination of Alkynes. *Org. Lett* 2003, 5, 4767–4770. [PubMed: 14653669]
- (47). Ochi N; Nakao Y; Sato H; Sakaki S {2+2} Cycloaddition of Alkyne with Titanium–Imido Complex: Theoretical Study of Determining Factor of Reactivity and Regioselectivity. *J. Phys. Chem. A* 2010, 114, 659–665. [PubMed: 19883057]
- (48). Müller TE; Hultsch KC; Yus M; Foubelo F; Tada M Hydroamination: Direct Addition of Amines to Alkenes and Alkynes. *Chem. Rev* 2008, 108, 3795–3892. [PubMed: 18729420]
- (49). Zhu C; Cook SP A Concise Synthesis of (+)-Artemisinin. *J. Am. Chem. Soc* 2012, 134, 13577–13579. [PubMed: 22866604]
- (50). DeLuca RJ; Edwards JL; Steffens LD; Michel BW; Qiao X; Zhu C; Cook SP; Sigman MS Wacker-Type Oxidation of Internal Alkenes using Pd(Quinox) and TBHP. *J. Org. Chem* 2013, 78, 1682–1686. [PubMed: 23363387]
- (51). Morandi B; Wickens ZK; Grubbs RH Practical and General Palladium-Catalyzed Synthesis of Ketones from Internal Olefins. *Angew. Chem., Int. Ed* 2013, 52, 2944–2948.
- (52). Morandi B; Wickens ZK; Grubbs RH Regioselective Wacker Oxidation of Internal Alkenes: Rapid Access to Functionalized Ketones Facilitated by Cross-Metathesis. *Angew. Chem., Int. Ed* 2013, 52, 9751–9754.
- (53). Lerch MM; Morandi B; Wickens ZK; Grubbs RH Rapid Access to β -Trifluoromethyl-Substituted Ketones: Harnessing Inductive Effects in Wacker-Type Oxidations of Internal Alkenes. *Angew. Chem., Int. Ed* 2014, 53, 8654–8658.
- (54). Carlson AS; Calcanas C; Brunner RM; Topczewski JJ Regiocontrolled Wacker Oxidation of Cinnamyl Azides. *Org. Lett* 2018, 20, 1604–1607. [PubMed: 29498865]
- (55). See XY; Beaumier EP; Davis-Gilbert ZW; Dunn PL; Larsen JA; Pearce AJ; Wheeler TA; Tonks IA Generation of Ti^{II} Alkyne Trimerization Catalysts in the Absence of Strong Metal Reductants. *Organometallics* 2017, 36, 1383–1390. [PubMed: 28690352]

Previous work: electronically-controlled [2+2+1] cross coupling:

Prior art: anionic directing groups

This work: inverted regioselectivity through directing effects

Figure 1.

Top: general overview of directing groups in Ti catalyzed/mediated chemistry; Bottom: Ti-catalyzed [2+2+1] pyrrole synthesis and methods for chemo- and regioselective alkyne heterocoupling.

Figure 2.

Initial observation of a directing-group effect: *o*-Me substituted aryl alkynes (**1a**) have poor selectivity for electronically preferred 2nd insertion product **4a-TMS**.

Figure 3.
Directing-group effects on internal alkyne heterocoupling.

Figure 4.
Directed heterocoupling with other internal alkyne [2+2] partners.

Figure 5.
Synthesis of an EP_1 antagonist derivative **5y**.

Figure 6.
Substrate-directed regioselective pyrrole formation via alkyne homocoupling.

Figure 7.
Substrate-directed regioselective alkyne hydroamination.

Table 1.

Optimization of Reaction Conditions to Promote Directed Alkyne Insertion^a

entry	T (°C)	X (equiv)	time (h)	Y (%)	yield of 3a ^b (%)	ratio of 3a/4a ^b
1 ^c	115	2	2	10	51	1.1
2	115	2	1	10	76	5.5
3	80	2	1	10	81	7.2
4	60	2	2	10	79	8.8
5	60	1	2	10	67	9.3
6	45	2	24	10	74	10.5
7	45	1	24	10	74	10.3

^aA mixture of 0.1–0.2 mmol **1a**, 0.1 mmol **2a**, 0.225 mmol PhNNPh , 0.0025–0.01 mmol $(\text{THF})_3\text{TiI}_2(\text{NPh})$, and 0.01 Ph_3CH (internal standard) in 0.5 mL of CF_3Ph were heated at desired temperature and time.

^bYield and selectivity are determined by ^1H NMR and reported with respect to **2a**.

^cReported in ref 8 with $[\text{py}_2\text{TiCl}_2(\text{NPh})]_2$ as catalyst.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

Scope of Directing-Group Effects in Ti-Catalyzed [2+2+1] Pyrrole Synthesis

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^a Condition A: A mixture of 0.5 mmol **1** (1.1 equiv), 0.5 mmol **2** (1.1 equiv), 0.225 mmol PhNNPh (0.5 equiv) and 0.05 mmol THF₃[₂Ti(NPh)]₂ (0.1 equiv) in 2.5 mL CF₃Ph was heated for 2 h. Reactions were quenched with 2 M HCl in MeOH;

^b Condition B: A mixture of 1.0 mmol **1** (2.2 equiv), 0.5 mmol **2** (1.1 equiv), 0.225 mmol PhNNPh (0.5 equiv) and 0.025 mmol [py₂Cl₂TiNPh]₂ (0.05 equiv) in 2.5 mL CF₃Ph was heated for 3 h. Reactions were quenched with 2 M HCl in MeOH;

^c Isolated and/or NMR yield of major regioisomer drawn and are reported with respect to **2a**.

^d Reaction run at 80 °C;

^e Reaction run at 115 °C.