Published in final edited form as:

Tetrahedron. 2013 November 11; 69(45): 9406–9416. doi:10.1016/j.tet.2013.08.081.

Lithium Perchlorate-, Acetic Anhydride-, and Triphenylphosphine-assisted Multicomponent Syntheses of 4-Unsubstituted 2,5-Dioxooctahydroquinoline-3-carboxylates and 3-carbonitriles

Xingxian Gu^{a,b}, Gunda I. Georg^{b,*}

^aDepartment of Medicinal Chemistry, University of Kansas, Lawrence, KS 66045

^bDepartment of Medicinal Chemistry and Institute for Therapeutics Discovery and Development, University of Minnesota, 717 Delaware Street, SE, Minneapolis, Minnesota 55414, United States

Abstract

Lithium perchlorate and acetic anhydride were the key additives for the multi-component reaction between 3-aminocyclohex-2-enones, formaldehyde, and malonates yielding adducts that were annulated under acidic conditions to afford bicyclic 2,5-dioxooctahydroquinoline-3-carboxylates. When methyl cyanoacetate was subjected to the same reaction conditions in the presence of a catalytic amount of triphenylphosphine, the bicyclic 2,5-dioxooctahydroquinoline-3-carbonitriles were obtained in a one-flask reaction.

Graphical abstract

*Corresponding Author Tel: (612) 626-6320. Fax: (612) 626-6318. georg@umn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary data

¹H NMR and ¹³C NMR spectra for all new compounds are available.

Notes

The authors declare no competing financial interest.

Keywords

Multicomponent reaction; catalysis; lithium perchlorate; acetic anhydride; triphenylphosphine; 2,5-dioxooctahydroquinolines

1. Introduction

1,4-Dihydropyridines, 1,4-dihydropyridine-2-pyridones, pyridines, and piperidines are closely related structural motifs found in many natural products and bioactive molecules. 1,4-Dihydropyridines are considered to be privileged structures, because they bind to multiple ion channels and receptors.¹ For example, the 4-aryl-1,4-dihydropyridine moiety is a component of the cardiovascular drug nifedipine (**1**, Figure 1), which is a calcium channel antagonist.^{2, 3} The annulated 1,4-dihydropyridine analogue **2** (Figure 1) initiates cardiac differentiation in mouse embryonic stem cells by inhibiting TGF β signaling – a mechanism unrelated to calcium channel blockade.⁴ These analogues represent a new class of agents for potential use in cardiac regenerative medicine.⁵ The structurally related 1,4-dihydropyridine-2-pyridone scaffold is also of medicinal interest. The 2-pyridone moiety is a component of clinically used phosphodiesterase-3 (PDE3) inhibitors such as milrinone (**3**, Figure 1) and related derivatives,^{6, 7} including the annulated PDE3 inhibitor **4** (Figure 1).⁸ 2-Pyridones are also found as structural elements in a variety of other bioactive compounds including natural products.^{7, 9} Octahydro-2-pyridones such as LY191704 and the related tetrahydro analogues were found to be specific inhibitors of human type I steroid 5- α -reductase.¹⁰⁻¹² Despite the fact that these structural types have displayed interesting bioactivities, 3,4-dihydropyridinones and their annulated analogues, 2,5-dioxooctahydroquinoline-3-carboxylates **5** (target scaffold **5**, Figure 1) have not been examined extensively yet in terms of synthesis, biological activity, and structure-activity relationships.

It is well known that 1,3-diketones, including cyclic ketones,¹³ react via enamine formation (Scheme 1, eq 1) to produce substituted 1,4-dihydropyridines (Hantzsch reaction). Similar to the Hantzsch-type reaction that is used to synthesize annulated 1,4-dihydropyridines (Scheme 1, eq 1), formal [3+3] cycloaddition reactions were employed to construct dihydro-2-pyridones including annulated core structures.^{9, 14-17} In these multi-component reactions 1,3-cyclohexanediones, malonates or Meldrum's acid, aldehydes, and amines or ammonia are reacted in one flask, as is depicted in Scheme 1, eq 2. In the example shown in eq 2, ZnO serves as an efficient catalyst to effect the reaction with aromatic aldehydes in excellent yields, but apparently does not work with formaldehyde or aliphatic aldehydes.¹⁸ When Meldrum's acid is used in the multicomponent reaction, formaldehyde and aliphatic aldehydes can be employed, but the reaction products lack the 3-carboxylic acid moiety (Scheme 1, eq 3), which is lost by decarboxylation.¹⁷ In related chemistry (Scheme 1, eq 4), hydrazinyl enamines of 1,3-cyclohexanediones, and the arylidenemalonates are preformed and then reacted in the presence of butyl lithium to form the Michael addition products that are subsequently cyclized to yield 4-aryl-2,5-dioxooctahydroquinoline-3-carboxylates.¹⁹

2. Results and discussion

In our efforts to prepare structurally diverse and novel nitrogen-containing heterocyclic compounds with privileged structures for biological screening, we targeted the synthesis of bicyclic 1,4-dihydropyridine-2-pyridones **5** (Figure 1) that do not contain a substituent at C4 but an ester moiety at C3 (2,5-dioxooctahydroquinoline-3-carboxylates). While aromatic aldehydes have been employed in the multicomponent reaction shown in eq 2 in Scheme 1, the reaction apparently did not take place when formaldehyde was used.¹⁸ The availability of a method that provides the target compounds **5** without a substituent at C4 is of importance to expand the scope of this chemistry. Once an aryl or aliphatic group is present at C4 it will require multiple steps for removal if unsubstituted analogues are desired for biological investigations.

We started our investigations by carrying out a one-flask reaction similar to the one shown in Scheme 1, eq 3, by following a reported procedure¹⁷ using 1,3-cyclohexanedione, formaldehyde, dimethyl malonate and benzylamine, but failed to obtain the desired bicyclic reaction product. We then decided to explore the possibility of synthesizing the target molecule in a step-wise manner in order to simplify the reaction system. 3-(Benzylamino)cyclohex-2-enone (**6a**), easily prepared by condensation of 1,3-cyclohexanedione and benzylamine,²⁰ paraformaldehyde, and dimethyl malonate, were envisaged to form an adduct, followed by a subsequent cyclization to yield the target molecule. Yet, when a mixture of the enaminone, formaldehyde, and dimethyl malonate were heated in a reaction vessel, only little of adduct **7a** formed and instead *bis*-addition of the enaminone to formaldehyde took place and **8a** was isolated as the major product in 74% (Scheme 2, eq 1). Clearly, this side reaction needed to be suppressed in order to allow the desired multi-component reaction to become the major reaction pathway and yield the desired adduct **7a**. Since LiClO₄ has been introduced as a useful reagent to facilitate Mannich-type reactions,²¹⁻²⁵ we tested this Lewis acid as an additive in the reaction (Scheme 2, eq 2). The reaction was heated to 90 °C in acetonitrile in the presence of one equivalent of LiClO₄, and the desired adduct **7a** was isolated as the major product in 77%, along with the *bis*-addition side product **8** in 13% as the minor product.

Optimization of the reaction conditions (Table 1) showed that an equimolar amount of LiClO₄ was crucial to obtain good yields (Table 1, entries 1-3) and that excess LiClO₄ was unnecessary (entries 4 and 5). In an attempt to annulate the adduct in the same flask, acetic anhydride was added as an additive. Whereas no annulated product was detected, the yield for the adduct formation of **7a** improved to 93% (Table 1, entry 6). When acetic anhydride was added, the reaction could be completed at a lower temperature (60 °C instead of 90 °C) in a comparable amount of time (0.5 - 3.0 h). Several other mild Lewis acids (Table 1, entries 7-9) were also tested, but the results were not satisfactory.

In order to explore the scope of the reaction we subjected a variety of *N*-substituted 3-aminocyclohex-2-enones and malonates to the reaction under the optimized conditions (Table 2). Most of the reactions yielded the corresponding adducts in moderate to excellent yields (66%-97%). Attempts to replace formaldehyde with benzaldehyde were not successful. No product formation was detected when a sterically hindered malonate *tert*-

butyl ester (Table 2, entry 16) or dimethyl methylmalonate (Table 2, entry 17) was employed in the reaction. Methyl acetoacetate, methyl cyanoacetate, and triethyl phosphonoacetate were tested as surrogates of malonates, but no product was detected under the standard conditions. We propose that methyl acetoacetate forms an intermediate similar to **9a** through Knoevenagel reaction,²⁶ however such an intermediate would be a more reactive Michael acceptor than **9a**, due to the higher electrophilicity provided by the ketone carbonyl versus the ester carbonyl. Thus, this intermediate would get consumed rapidly via self-polymerization,²⁶ without reacting with **6a**. We have provided the ¹H NMR of **9a** (see supplementary material), showing its rapid self-polymerization. The Michael acceptor intermediate derived from methyl acetoacetate can be expected to polymerize even faster.^{26, 27}

Similar to what has been proposed in the literature¹⁷ and is shown in Scheme 1, eq 4,¹⁹ we assume that product **7a** formed via a conjugate addition of the enaminone to an intermediate generated from an *in situ* Knoevenagel reaction between dimethyl malonate and formaldehyde, followed by an imine-enamine tautomerism (Scheme 3). To investigate the proposed pathway, dimethyl methylenemalonate (**9a**) was prepared following a literature procedure,²⁸ and the conjugate addition by the enaminone successfully yielded **7a** in 90% without the assistance of LiClO₄ and acetic anhydride. Thus, LiClO₄ and acetic anhydride facilitate the formation of intermediate **9a** and thereby, favoring the Michael addition of the enamine **6a** to intermediate **9a** over the formation of the bis-addition product **8**, although the detailed mechanism is not clear.

No product formation was detected when benzaldehyde was used in the LiClO₄-assisted reaction (Table 2, entry 20). Based on the proposed mechanism, the difficulty in the corresponding conjugate addition to the sterically more demanding intermediate and/or a difference in electronic properties could be the cause of the lack of reactivity. To confirm that hypothesis, dimethyl benzylidenemalonate (**9b**)²⁹ was prepared and subjected to the reaction with the enaminone. Indeed, the enaminone did not add to dimethyl 2-benzylidenemalonate (**9b**) in the presence of LiClO₄ or without it.

The observed rate enhancement after the addition of acetic anhydride most likely facilitates the LiClO₄-catalyzed Knoevenagel reaction by *in situ* formation of the more reactive geminal formaldehyde diacetate,³⁰ similar to what has been described for the iodine-catalyzed Knoevenagel reaction in the presence of acetic anhydride (Scheme 4).³¹

Intermediates **7** were next treated with trifluoroacetic acid at 90 °C in dichloroethane or toluene in a sealed reaction vessel to yield the desired 2,5-dioxooctahydroquinoline-3-carboxylate esters **5** in mostly moderate yields as shown in Table 3.³²⁻³⁴ The cyclization of dimethylmalonate adducts was carried out in dichloroethane, while toluene was a better choice of solvent for the dibenzylmalonate adducts, with an approximately 10% improvement in yields on average. Adducts containing a 1-morpholino group (**7h** and **7i**, Table 2) decomposed under the acidic conditions, furnishing none of the annulated products.

We next subjected methyl cyanoacetate (instead of a malonate ester) to the same reaction conditions but no reaction product was obtained. However, when a catalytic amount of

triphenylphosphine was added to the reaction mixture, the annulated products **10a-e** (Table 4) were formed directly.

The triphenylphosphine catalyst presumably plays multiple roles in this reaction. It is well known that triphenylphosphine catalyzes the Knoevenagel condensation between aldehydes and cyanoacetates to form α -cyanoacrylonitriles, but not with malonates.³⁵ To probe whether triphenylphosphine catalyzed this reaction in our case, we carried out the reaction in the presence of triphenylphosphine but without LiClO₄. We did not obtain any of the desired product and only bis-addition product **8** was found. We therefore concluded that LiClO₄ is responsible for the formation of the presumed methyl 2-cyanoacrylate intermediate, rather than triphenylphosphine.

Triphenylphosphine is a strong nucleophile that adds to electron-deficient double and triple bonds.³⁶⁻³⁹ Scheme 5 shows a possible mechanism for the formation of compounds **10**, which is similar to the mechanism reported by Steward *et al.* for the triphenylphosphine catalyzed conjugate addition of alcohols and water to enones.⁴⁰ Triphenylphosphine adds to methyl 2-cyanoacrylate and forms phosphonium enolate **11**, which abstracts a proton from **6a** forming ion pair **12**, which then adds to an equivalent of methyl 2-cyanoacrylate. The new ion pair **13** abstracts a proton from another molecule of **6a** to form intermediate **14** that cyclizes to yield product **10a**. Deprotonated **6a** and regenerated **11** reform ion pair **12**. (Steward *et al.* were able to rule out a direct proton abstraction by triphenylphosphine.⁴⁰)

The cyclization reaction is most likely catalyzed by triphenylphosphine as well since it has been reported that triphenylphosphine can be a catalyst for transesterifications,⁴¹ although an influence of steric and/or electronic effects of the nitrile group on the cyclization cannot be ruled out. We propose that triphenylphosphine adds to the ester **14** and forms acylphosphonium intermediate **15** that would cyclize to form compound **10a** and regenerate PPh₃.

Formaldehyde was the only viable aldehyde in this reaction. Malonates were also subjected to such reaction conditions in order to obtain **5** in one step, but no cyclized product was detected after 12 h. Instead the bis-addition product **8** was the major product, and only a small amount of the conjugated addition adduct **7a** was detected.

The syntheses of structurally related monocyclic 4-aryl-2-oxo-tetrahydropyridine-3-carbonitriles from 4*H*-pyrans has recently been reported.^{42, 43} The 4*H*-pyrans were prepared from aromatic aldehydes, malononitrile, and ethyl acetoacetate in the presence of ZnO as the catalyst.⁴³ Another recent publication describes a three-component reaction that employed aromatic aldehydes (and one aliphatic aldehyde) with either malononitrile or 2-cyanoacetamide in the presence of PEG-SO₃H or vitamin B1, respectively, to provide bicyclic 4-substituted-2,5-dioxooctahydroquinolone-3-carbonitriles.⁴⁴ Thus, our technique complements existing methods for the synthesis of these types of scaffolds in that we are employing formaldehyde rather than aromatic or aliphatic aldehydes in the multi-component reaction and thereby providing a method for the synthesis of 4-unsubstituted analogues.

The compounds were submitted to the NIH Molecular Libraries Program and screened in several assays that are reported in PubChem.⁴⁵

3. Conclusion

In summary, a LiClO₄-assisted multi-component reaction of 3-aminocyclohex-2-enones **6** with malonates or cyanoacetates, and formaldehyde in the presence of acetic anhydride is described, providing access to 4-unsubstituted 2,5-dioxooctahydroquinoline-3-carboxylates and 3-carbonitriles.

4. Experimental section

4.1 General

Proton nuclear magnetic resonance spectra were recorded at 400 MHz and carbon nuclear magnetic resonance spectra were recorded at 100 MHz. All chemical shifts were recorded as parts per million (ppm), and all samples were dissolved in CDCl₃ using the residual solvent peak as internal standard. Mass spectra were obtained from a ZAB HS mass spectrometer equipped with an 11/250 data system. Fast-atom bombardment mass spectrometry (FAB-MS) experiments were performed with a Xenon gun operated at 8 Kev energy and 0.8 mA emission at the Institute for Therapeutic Discovery and Development at University of Minnesota. Fast-atom bombardment high resolution mass spectra (FAB-HRMS) were recorded at 1:10,000 resolution using linear voltage scans under data system control and collected in a multi-channel analyzer mode (MCA). A Recording Infrared Spectrophotometer or a FT-IR was used to record infrared spectra. Melting points are uncorrected. Solvents and reagents that are commercially available were used without further purification. Acetonitrile, dichloroethane, and toluene were distilled freshly from calcium hydride under argon. Silica gel (230 - 400 mesh) was used for flash column chromatography. All compounds were concentrated using a standard rotary evaporator and high-vacuum techniques.

General Method for LiClO₄-assisted Reaction of Enaminones with Malonates and Formaldehyde to Form Adducts **7 (Table 2).**—A suspension of the enaminone, LiClO₄ (1.0 equiv), the malonate diester (1.5 equiv), and acetic anhydride (1.0 equiv) in acetonitrile (10 mL, ~35 mL/mmol) was stirred and pre-heated in a sealed screw-cap vial until the reaction mixture turned into a clear solution after approximately 5 min. To the reaction medium was then added paraformaldehyde (2.0 equiv). The reaction was then sealed, stirred at 60 °C, and monitored by TLC. In general, the reaction reached completion within 0.5 to 3 h, and was quenched subsequently with H₂O (20 mL, ~70 mL/mmol). The aqueous phase was extracted with EtOAc (3 × 10 mL, ~35 mL/mmol), and then, the organic layers were combined, and washed with brine, and dried over MgSO₄. The organic solvent was then removed under reduced pressure and the resulting residue was purified by flash column chromatography on silica gel, using 50% acetone/hexanes as the eluent.

Dimethyl 2-((2-(Benzylamino)-6-oxocyclohex-1-enyl)methyl)malonate (7a): The *N*-benzyl-enaminone (50 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024

mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (80 mg, 0.23 mmol, 93%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.36 - 7.34 (m, 2H), 7.30 - 7.26 (m, 3H), 6.35 (m, 1H), 4.43 (d, *J* = 6.1 Hz, 2H), 3.80 (t, *J* = 7.2 Hz, 1H), 3.69 (s, 6H), 2.83 (d, *J* = 7.2 Hz, 2H), 2.39 (t, *J* = 6.2 Hz, 2H), 2.26 (t, *J* = 6.2 Hz, 2H), 1.84 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 194.9, 171.1, 162.8, 138.4, 129.0, 127.8, 127.0, 105.8, 52.7, 50.6, 47.2, 36.3, 25.8, 23.6, 21.3; ESI-HRMS: calc'd m/e for [M+Na⁺] C₁₉H₂₃NNaO₅: 368.1474, found 368.1484; IR (neat, NaCl, cm⁻¹): 2951, 1733, 1575, 1436, 1409, 1357, 1299, 1246, 1202, 1158, 1118, 1031.

Dimethyl 2-((6-Oxo-2-(phenylamino)cyclohex-1-enyl)methyl)malonate (7b): The *N*-phenyl-enaminone (47 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions and yielded a viscous colorless oil (74 mg, 0.22 mmol, 90%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 8.03 (s, 1H), 7.33 - 7.35 (m, 2H), 7.16 (m, 1H), 7.06 - 7.09 (m, 2H), 3.82 (t, *J* = 7.2 Hz, 1H), 3.74 (s, 6H), 2.91 (d, *J* = 7.2 Hz, 2H), 2.50 (t, *J* = 6.0 Hz, 2H), 2.37 (t, *J* = 6.0 Hz, 2H), 1.86 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.9, 171.3, 160.6, 139.3, 129.3, 125.0, 124.1, 109.0, 52.9, 50.9, 36.9, 27.4, 23.8, 22.2; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₈H₂₂NO₅: 332.1498, found 332.1505; IR (neat, NaCl, cm⁻¹): 2593, 1733, 1567, 1499, 1435, 1398, 1306, 1245, 1204, 1080, 1038.

Dibenzyl 2-((6-Oxo-2-(phenylamino)cyclohex-1-enyl)methyl)malonate (7c): The *N*-phenyl-enaminone (47 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dibenzyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (120 mg, 0.24 mmol, 97%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.89 (s, 1H), 7.35-7.29 (m, 12H), 7.15 (m, 1H), 7.01 (m, 2H), 5.17 (q-like, *J* = 12.4 Hz, 4H), 3.95 (t, *J* = 7.4 Hz, 1H), 2.95 (d, *J* = 7.4 Hz, 2H), 2.39 (t, *J* = 6.0 Hz, 2H), 2.32 (t, *J* = 6.0 Hz, 2H), 1.76 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 196.0, 170.6, 160.8, 139.2, 135.7, 129.3, 128.7, 128.4, 128.2, 125.1, 124.2, 108.6, 67.4, 51.1, 36.8, 27.3, 23.7, 22.0; ESI-HRMS: calc'd m/e for [M+H⁺] C₃₀H₃₀NO₅: 484.2124, found 484.2118; IR (neat, NaCl, cm⁻¹): 2948, 1736, 1578, 1498, 1454, 1431, 1396, 1304, 1233, 1199, 1153, 1080, 1030, 1003.

1-Benzyl 3-Methyl 2-((2-(Benzylamino)-6-oxocyclohex-1-en-1-yl)methyl)malonate (7d): The *N*-benzyl-enaminone (50 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), benzyl methyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (77 mg, 0.18 mmol, 73%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.37 - 7.24 (m, 10H), 6.24 (t, *J* = 6.2 Hz, 1H), 5.17 (q-like, *J* = 12.4 Hz, 2H), 4.36 (d, *J* = 6.1 Hz, 2H), 3.89 (t, *J* = 7.6 Hz, 1H), 3.67 (s, 3H), 2.84 (m, 2H), 2.33 (m, 2H), 2.24 (m, 2H), 1.81 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 194.9, 170.9, 170.7, 162.7, 138.4, 129.0, 128.6, 128.4, 128.2, 127.0, 105.8, 67.2, 54.2, 52.7, 50.71, 47.2, 36.4, 25.8, 23.6, 21.3; ESI-HRMS: calc'd m/e for [M+H⁺] C₂₅H₂₈NO₅: 422.1967, found 422.1973; IR (neat, NaCl, cm⁻¹): 2949, 1732, 1575, 1497, 1455, 1436, 1409, 1378, 1356, 1297, 1242, 1202, 1154, 1119, 1079, 1029.

Dibenzyl 2-((2-(Benzylamino)-6-oxocyclohex-1-enyl)methyl)malonate (7e): The *N*-benzyl-enaminone (50 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dibenzyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (112 mg, 0.23 mmol, 90%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.36 - 7.22 (m, 15H), 6.20 (t, *J* = 6.1 Hz, 1H), 5.12 (q-like, *J* = 12.4 Hz, 4H), 4.30 (d, *J* = 6.1 Hz, 2H), 3.95 (t, *J* = 7.3 Hz, 1H), 2.87 (d, *J* = 7.3 Hz, 2H), 2.28 (t, *J* = 6.2 Hz, 2H), 2.22 (t, *J* = 6.3 Hz, 2H), 1.77 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): 194.9, 170.4, 162.7, 138.3, 135.7, 129.0, 128.6, 128.3, 128.1, 127.7, 127.0, 105.6, 67.2, 50.7, 47.1, 36.3, 25.7, 23.5, 21.1; ESI-HRMS: calc'd m/e for [M+H⁺] C₃₁H₃₂NO₅: 498.2280, found 498.2275; IR (neat, NaCl, cm⁻¹): 2945, 1731, 1577, 1497, 1456, 1434, 1372, 1356, 1295, 1200, 1149, 1028.

Dimethyl 2-((2-(Butylamino)-6-oxocyclohex-1-enyl)methyl)malonate (7f): The *N*-butyl-enaminone (42 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (52 mg, 0.17 mmol, 66%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 5.88 (s, 1H), 3.75 (t, *J* = 6.2 Hz, 1H), 3.69 (s, 6H), 3.18 (m, 2H), 2.76 (d, *J* = 7.1 Hz, 2H), 2.41 (t, *J* = 6.2 Hz, 2H), 2.26 (t, *J* = 6.3 Hz, 2H), 1.87 (m, 2H), 1.57 (m, 2H), 1.40 (m, 2H), 0.95 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 194.4, 171.2, 163.0, 104.9, 52.7, 50.6, 43.2, 36.3, 32.4, 25.8, 23.4, 21.3, 20.1, 13.9; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₆H₂₆NO₅: 312.1811, found 312.1801; IR (neat, NaCl, cm⁻¹): 2954, 1733, 1570, 1436, 1417, 1368, 1301, 1243, 1203, 1155, 1100.

Dibenzyl 2-((2-(Butylamino)-6-oxocyclohex-1-enyl)methyl)malonate (7g): The *N*-butyl-enaminone (42 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dibenzyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (92 mg, 0.20 mmol, 80%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.32 - 7.26 (m, 10H), 5.69 (t, *J* = 4.8 Hz, 1H), 5.13 (q-like, *J* = 12.4 Hz, 4H), 3.92 (t, *J* = 7.4 Hz, 1H), 3.04 (m, 2H), 2.81 (d, *J* = 7.3 Hz, 2H), 2.28 (t, *J* = 6.2 Hz, 2H), 2.22 (t, *J* = 6.1 Hz, 2H), 1.81 (m, 2H), 1.50 (m, 2H), 1.37 (m, 2H), 0.94 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 194.4, 170.6, 162.8, 135.8, 128.6, 128.3, 128.1, 104.7, 67.1, 50.7, 43.1, 36.3, 32.3, 25.8, 23.4, 21.2, 20.1, 13.79; ESI-HRMS: calc'd m/e for [M+H⁺] C₂₈H₃₄NO₅: 464.2437, found 464.2444; IR (neat, NaCl, cm⁻¹): 2955, 1731, 1575, 1438, 1413, 1377, 1198, 1152.

Dimethyl 2-((2-(Morpholinoamino)-6-oxocyclohex-1-enyl)methyl)malonate (7h): The *N*-morpholino-enaminone (49 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (58 mg, 0.17 mmol, 72%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 6.78 (s, 1H), 3.77 (t, *J* = 7.2 Hz, 1H), 3.76 (br, 4H), 3.71 (s, 6H), 2.76 (br, 4H), 2.74 (d, *J* = 7.2 Hz, 2H), 2.64 (t, *J* = 6.2 Hz, 2H), 2.27 (t, *J* = 6.2 Hz, 2H), 1.86 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.6, 171.1, 162.6, 103.6, 66.7, 56.9, 52.7, 50.6, 36.7, 25.2, 23.2, 21.4; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₆H₂₅N₂O₆: 341.1713, found 341.1699; IR

(neat, NaCl, cm^{-1}): 2953, 2857, 1733, 1583, 1438, 1403, 1303, 1267, 1243, 1201, 1157, 1111, 1074, 1042, 1028.

Dibenzyl 2-((2-(Morpholinoamino)-6-oxocyclohex-1-enyl)methyl)malonate (7i): The *N*-morpholino-enaminone (49 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dibenzyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (106 mg, 0.22 mmol, 86%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.33 - 7.25 (m, 10H), 6.64 (s, 1H), 5.14 (q-like, J = 12.5 Hz, 4H), 3.94 (t, J = 7.4 Hz, 1H), 3.90 - 3.40 (br, 4H), 2.78 (d, J = 7.3 Hz, 2H), 2.58 (s, 4H), 2.54 (t, J = 6.2 Hz, 2H), 2.25 (t, J = 6.2 Hz, 2H), 1.82 (m, 2H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 195.5, 170.4, 162.6, 135.8, 128.6, 128.3, 127.9, 103.5, 67.2, 66.6, 56.7, 50.8, 36.7, 25.2, 23.1, 21.3; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{28}\text{H}_{33}\text{N}_2\text{O}_6$: 493.2339, found 493.2342; IR (neat, NaCl, cm^{-1}): 2917, 2850, 1732, 1584, 1497, 1455, 1402, 1385, 1266, 1233, 1200, 1152, 1112, 1076.

Dimethyl 2-((4,4-Dimethyl-6-oxo-2-(phenylamino)cyclohex-1-enyl)methyl)-malonate (7j): The *N*-phenyl-enaminone (54 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (75 mg, 0.21 mmol, 83%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.94 (s, 1H), 7.35 (m, 2H), 7.16 (m, 1H), 7.06 (m, 2H), 3.86 (t, J = 7.4 Hz, 1H), 3.75 (s, 6H), 2.93 (d, J = 7.4 Hz, 2H), 2.36 (s, 2H), 2.23 (s, 2H), 0.97 (s, 6H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 195.5, 171.3, 158.4, 139.3, 129.4, 124.9, 123.9, 108.0, 52.9, 50.7, 50.5, 40.8, 32.8, 28.1, 23.8; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{20}\text{H}_{26}\text{NO}_5$: 360.1811, found 360.1801; IR (neat, NaCl, cm^{-1}): 2954, 1735, 1579, 1498, 1436, 1395, 1321, 1270, 1243, 1150.

Dibenzyl 2-((4,4-Dimethyl-6-oxo-2-(phenylamino)cyclohex-1-enyl)methyl)-malonate (7k): The *N*-phenyl-enaminone (54 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dibenzyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (111 mg, 0.22 mmol, 87%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.90 (s, 1H), 7.34 (m, 2H), 7.29 (m, 10H), 7.16 (m, 1H), 7.00 (m, 2H), 5.16 (dt, J = 3.2, 14.0 Hz, 4H), 3.99 (t, J = 7.5 Hz, 1H), 2.95 (d, J = 7.5 Hz, 2H), 2.27 (s, 2H), 2.18 (s, 2H), 0.91 (s, 6H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 195.5, 170.5, 158.4, 139.3, 135.5, 129.3, 128.7, 128.4, 128.2, 124.9, 124.0, 107.9, 67.4, 51.0, 50.5, 40.7, 32.7, 28.1, 23.7; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{32}\text{H}_{34}\text{NO}_5$: 512.2437, found 512.2431; IR (neat, NaCl, cm^{-1}): 2956, 1733, 1579, 1498, 1455, 1395, 1384, 1321, 1271, 1227, 1147, 1071.

Dimethyl 2-((2-(Benzylamino)-4,4-dimethyl-6-oxocyclohex-1-enyl)methyl)-malonate (7l): The *N*-benzyl-enaminone (57 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (81 mg, 0.24 mmol, 95%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.37 (m, 2H), 7.29 (m, 3H), 6.20 (m, 1H), 4.43 (d, J = 6.2 Hz, 2H), 3.82 (t, J = 7.4

Hz, 1H), 3.69 (s, 6H), 2.84 (d, $J = 7.3$ Hz, 2H), 2.24 (s, 2H), 2.16 (s, 2H), 0.96 (s, 6H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 194.4, 171.1, 161.0, 138.5, 129.0, 127.8, 126.9, 104.5, 52.7, 50.5, 50.0, 47.1, 39.3, 31.8, 28.5, 23.4; ESI-HRMS: ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{21}\text{H}_{28}\text{NO}_5$: 374.1967, found 374.1964; IR (neat, NaCl , cm^{-1}): 2953, 1732, 1576, 1436, 1407, 1357, 1325, 1271, 1245, 1150, 1118, 1036.

Dibenzyl 2-((2-(Benzylamino)-4,4-dimethyl-6-oxocyclohex-1-enyl)methyl)malonate (7m):

The *N*-benzyl-enaminone (57 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dibenzyl malonate (0.070 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous light yellow oil (115 mg, 0.22 mmol, 88%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.37 - 7.23 (m, 15H), 6.10 (t, $J = 6.5$ Hz, 1H), 5.11 (q-like, $J = 12.4$ Hz, 4H), 4.32 (d, $J = 6.3$ Hz, 2H), 3.95 (t, $J = 7.4$ Hz, 1H), 2.87 (d, $J = 7.4$ Hz, 2H), 2.13 (s, 2H), 2.10 (s, 2H), 0.92 (s, 6H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 194.4, 170.4, 160.9, 138.5, 135.6, 129.0, 128.6, 128.3, 128.2, 127.7, 126.9, 104.3, 67.2, 50.8, 49.9, 47.0, 39.2, 31.8, 28.5, 23.3; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{33}\text{H}_{36}\text{NO}_5$: 526.2593, found 526.2586; IR (neat, NaCl , cm^{-1}): 2955, 1731, 1577, 1497, 1455, 1407, 1356, 1324, 1271, 1233, 1147, 1028.

Dimethyl 2-((2-(Butylamino)-4,4-dimethyl-6-oxocyclohex-1-enyl)methyl)malonate (7n):

The *N*-butyl-enaminone (49 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (75 mg, 0.22 mmol, 88%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 5.73 (s, 1H), 3.77 (t, $J = 7.3$ Hz, 1H), 3.69 (s, 6H), 3.18 (m, 2H), 2.77 (d, $J = 7.3$ Hz, 2H), 2.25 (s, 2H), 2.16 (s, 2H), 1.57 (m, 2H), 1.40 (m, 2H), 1.02 (s, 6H), 0.96 (t, 7.3 Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 193.8, 171.1, 161.1, 103.5, 52.6, 50.5, 49.9, 43.2, 39.5, 32.7, 31.8, 28.6, 23.3, 20.1, 13.9; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{18}\text{H}_{30}\text{NO}_5$: 340.2124, found 340.2126; IR (neat, NaCl , cm^{-1}): 2953, 1733, 1576, 1436, 1407, 1357, 1325, 1271, 1245, 1150, 1118, 1036.

Dimethyl 2-((5-Oxo-2-(phenylamino)cyclopent-1-enyl)methyl)malonate (7o): The *N*-phenyl-enaminone (43 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024 mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (76 mg, 0.24 mmol, 96%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 8.20 (s, 1H), 7.34 (t, $J = 7.7$ Hz, 2H), 7.14 (m, 3H), 3.75 (s, 6H), 3.72 (t, $J = 7.2$ Hz, 1H), 2.78 (d, $J = 7.0$ Hz, 2H), 2.70 (s, 2H), 2.37 (s, 2H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 202.9, 171.5, 170.8, 139.3, 129.5, 124.9, 122.3, 111.6, 52.9, 50.3, 33.2, 26.3, 21.7; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{17}\text{H}_{20}\text{NO}_5$: 318.1341, found 318.1344; IR (neat, NaCl , cm^{-1}): 2953, 1734, 1667, 1581, 1499, 1435, 1410, 1340, 1293, 1240, 1158, 1079, 1063.

Dimethyl 2-((2-(Benzylamino)-5-oxocyclopent-1-enyl)methyl)malonate (7p): The *N*-benzyl-enaminone (47 mg, 0.25 mmol), LiClO_4 (27 mg, 0.25 mmol), paraformaldehyde (16 mg, 0.50 mmol), dimethyl malonate (0.042 mL, 0.37 mmol), and acetic anhydride (0.024

mL, 0.25 mmol) were subjected to the standard reaction conditions, and yielded a viscous colorless oil (78 mg, 0.24 mmol, 94%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.36 (m, 2H), 7.29 (m, 3H), 6.43 (s, 1H), 4.47 (d, *J* = 6.4 Hz, 2H), 3.71 (t, *J* = 6.9 Hz, 1H), 3.69 (s, 6H), 2.69 (d, *J* = 7.2 Hz, 2H), 2.54 (s, 2H), 2.33 (s, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 202.3, 174.1, 170.6, 138.0, 129.0, 127.9, 127.0, 108.6, 52.8, 50.3, 47.3, 32.9, 24.9, 21.8; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₈H₂₂NO₅: 332.1498, found 332.1492; IR (neat, NaCl, cm⁻¹): 2952, 1733, 1669, 1594, 1578, 1496, 1436, 1353, 1279, 1238, 1156, 1118, 1077, 1056, 1030.

2,2'-Methylenebis(3-(benzylamino)cyclohex-2-enone) (8): The *N*-benzyl enaminone (50 mg, 0.25 mmol) was dissolved in acetonitrile (10 mL), and paraformaldehyde (16 mg, 0.50 mmol), and dimethyl malonate (0.042 mL, 0.37 mmol) were added sequentially. The reaction was stirred at 60 °C for 2 h, the solvent was then removed under reduced pressure, and the crude material was purified by flash column chromatography on silica gel, using 50% acetone:hexanes (v/v) as the eluent. After purification, a light yellow oil was obtained (38 mg, 74%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 9.51 (s, 2H), 7.33 - 7.20 (m, 10H), 4.48 (d, *J* = 6.4 Hz, 4H), 3.77 (s, 2H), 2.36 (br, 4H), 2.27 (br, 4H), 1.82 (m, 4H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 194.9, 166.9, 139.1, 128.8, 127.3, 126.7, 108.9, 63.8, 46.8, 35.7, 25.6, 21.5; ESI-HRMS: calc'd m/e for [M+H⁺] C₂₇H₃₁N₂O₂: 415.2386, found 415.2390; IR (neat, NaCl, cm⁻¹): 3262, 3171, 3030, 2946, 1734, 1610, 1560, 1478, 1433, 1355, 1317, 1201, 1127, 1106, 1051, 1011.

Dimethyl 2-Methylenemalonate (9a): Dimethyl malonate (11.5 mL, 100.0 mmol) was dissolved in acetic acid (45 mL), Cu(OAc)₂ (2.0 g, 11.2 mmol) was added, and the reaction mixture was heated at 100 °C (external bath temperature) for 4 h. After cooling to ambient temperature, the salts were filtered off, and the acetic acid was removed using bulb-to-bulb distillation (bp 50-60 °C, 100 mmHg). Further bulb-to-bulb distillation (bp 120-130 °C, 20 mm Hg) gave a colorless oil (3.7 g, 25.7 mmol, 26%). The product tended to polymerize rapidly and was therefore, taken to the reaction with the enaminone immediately after isolation. ¹H NMR (CDCl₃, 400 MHz, ppm): δ 6.56 (s, 2H), 3.83 (s, 6H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 167.1, 135.2, 134.5, 52.7; ESI-HRMS: calc'd m/e for [M+H⁺] C₆H₉O₄: 145.0501, found 145.0500.

Dimethyl 2-Benzylidenemalonate (9b): To a solution of benzaldehyde (6.6 mL, 60 mmol) in DMSO (20 mL), proline (690.0 mg, 6.0 mmol) was added. The mixture was stirred for 5 min before the addition of dimethyl malonate (13.7 mL, 120 mmol). After stirring at ambient temperature for 24 h, the reaction was diluted with EtOAc (60 mL) and washed with water (60 mL × 2). The combined organic layer was dried over Na₂SO₄, and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography on silica gel using 50% acetone:hexanes (v/v) as the eluent to furnish a viscous oil (12.8 g, 98%): ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.78 (s, 1H), 7.42 - 7.40 (m, 5H), 3.86 (s, 3H), 3.85 (s, 3H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 167.3, 164.6, 143.1, 132.9, 130.8, 129.5, 129.0, 125.6, 52.87, 52.86; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₂H₁₃O₄: 221.0814, found 221.0805.

Four reactions were set up in parallel, utilizing the enaminone (3-(benzylamino)cyclohex-2-enone) to react with dimethyl 2-methylenemalonate **9a** and dimethyl 2-benzylidenemalonate **9b** synthesized above, respectively:

- a) A solution of enaminone (50.2 mg, 0.25 mmol), LiClO₄ (30 mg, 0.25 mmol), and dimethyl 2-methylenemalonate (54 mg, 0.38 mmol) was heated in a sealed screw-cap vial at 60 °C;
- b) A solution of enaminone (50.2 mg, 0.25 mmol) and dimethyl 2-methylenemalonate (54 mg, 0.38 mmol) was heated in a sealed screw-cap vial at 60 °C;
- c) A solution of enaminone (50.2 mg, 0.25 mmol), LiClO₄ (30 mg, 0.25 mmol), and dimethyl 2-benzylidenemalonate (83 mg, 0.38 mmol) was heated in a sealed screw-cap vial at 60 °C;
- d) A solution of enaminone (50.2 mg, 0.25 mmol) and dimethyl 2-benzylidenemalonate (83 mg, 0.38 mmol) was heated in a sealed screw-cap vial at 60 °C.

The reactions were monitored by TLC for product formation. After the starting material was completely consumed, the reaction medium was then allowed to cool to ambient temperature and was subsequently washed with H₂O (20 mL). The aqueous phase was extracted with EtOAc (3 × 10 mL). The organic layers were combined, washed with brine, and dried with MgSO₄. The organic solvent was then removed under reduced pressure, and the residue was purified by flash column chromatography on silica gel. Reactions a and b gave the desired adduct in a yield of 90%, while only starting material was recovered for reactions c) and d).

General Method for the Synthesis of 2,5-Dioxo-1,2,3,4,5,6,7,8-

octahydroquinoline-3-carboxylates 5 (Table 3).—The adduct **7** was dissolved in the corresponding solvent (30 mL/mmol) in a screw-cap vial. Dichloroethane was selected as the solvent for the adducts derived from dimethyl malonate, and toluene was used for those derived from dibenzyl malonate. To the solution was slowly added trifluoroacetic acid (30 equiv). The vial was sealed and stirred at 90 °C for 0.5 to 3 hr. The reaction medium was cooled to ambient temperature and washed with saturated NaHCO₃ solution (20 mL). The aqueous phase was extracted with CH₂Cl₂ or EtOAc (3 × 10 mL). The organic layers were combined, washed with brine, and dried over MgSO₄. The organic solvent was then removed under reduced pressure, and the residue was purified by flash column chromatography on silica gel, using 20% EtOAc:hexanes (v/v) as the eluent.

Methyl 1-Benzyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5a): The adduct (80 mg, 0.23 mmol) was dissolved in dichloroethane (7 mL); then TFA (0.54 mL, 7.0 mmol) was added, and the reaction was treated under the standard conditions to yield a light yellow oil (58 mg, 0.18 mmol, 79%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.27 (m, 2H), 7.20 (m, 1H), 7.11 (m, 2H), 4.96 (q-like, *J* = 16.4 Hz, 2H), 3.70 (s, 3H), 3.57 (t, *J* = 8.3 Hz, 1H), 3.00 (dd, *J* = 8.4, 16.9 Hz, 1H), 2.78 (dd, *J* = 8.5, 16.9 Hz, 1H), 2.42 (t, *J* = 6.3 Hz, 2H), 2.28 (t, *J* = 6.2 Hz, 2H), 1.88 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.8, 169.6, 167.2, 154.7, 136.7, 129.0, 127.7, 126.2, 115.0, 52.8, 47.4, 45.7, 36.0, 26.5, 21.8, 20.8; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₈H₂₀NO₄: 314.1392, found 314.1385; IR (neat, NaCl, cm

⁻¹): 2953, 1743, 1692, 1654, 1622, 1497, 1454, 1437, 1392, 1365, 1326, 1299, 1280, 1259, 1195, 1167, 1128.

Benzyl 1-Benzyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5b): The adduct (108 mg, 0.22 mmol) was dissolved in toluene (7 mL), TFA (0.50 mL, 6.5 mmol) was added, and the reaction mixture was then treated under the standard conditions to yield a light yellow oil (60 mg, 0.15 mmol, 70%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.37 -7.24 (m, 8H), 7.16 (m, 2H), 5.20 (q-like, *J* = 12.2 Hz, 2H), 5.00 (q-like, *J* = 16.5 Hz, 2H), 3.68 (t, *J* = 7.0 Hz, 1H), 3.15 (dd, *J* = 7.5, 16.8 Hz, 1H), 2.80 (dd, *J* = 6.6, 16.8 Hz, 1H), 2.41 (m, 2H), 2.26 (m, 2H), 1.84 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.6, 168.9, 167.1, 154.7, 136.6, 135.4, 129.0, 128.6, 128.4, 128.3, 127.5, 126.1, 115.0, 67.3, 47.5, 45.6, 35.9, 26.3, 21.6, 20.8; ESI-HRMS: calc'd m/e for [M+H⁺] C₂₄H₂₄NO₄: 390.1705, found 390.1700; IR (neat, NaCl, cm⁻¹): 2953, 1738, 1691, 1652, 1622, 1497, 1455, 1428, 1390, 1364, 1297, 1258, 1194, 1164, 1126.

Methyl 2,5-Dioxo-1-phenyl-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5c): The adduct (57 mg, 0.17 mmol) was dissolved in dichloroethane (5 mL), TFA (0.40 mL, 5.2 mmol) was added, and the reaction mixture was treated under the standard conditions to yield a white solid (43mg, 0.14 mmol, 84%, mp: 120 - 121 °C). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.48 - 7.40 (m, 3H), 7.17 (s, 2H), 3.76 (s, 3H), 3.68 (t, *J* = 7.6 Hz, 1H), 3.18 (dd, *J* = 8.4, 17.0 Hz, 1H), 2.90 (dd, *J* = 8.6, 17.0 Hz, 1H), 2.39 (t, *J* = 6.0 Hz, 2H), 2.09 (t, *J* = 6.0 Hz, 2H), 1.93 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 196.0, 169.6, 167.1, 154.8, 137.0, 129.7, 129.0, 128.8, 113.9, 52.8, 47.5, 36.3, 28.0, 21.9, 21.1; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₇H₁₈NO₄: 300.1236, found 300.1228; IR (neat, NaCl, cm⁻¹): 2953, 1742, 1702, 1652, 1625, 1594, 1493, 1380, 1271, 1244, 1154.

Benzyl 2,5-Dioxo-1-phenyl-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5d): The adduct (117 mg, 0.24 mmol) was dissolved in toluene (7 mL), TFA (0.55 mL, 7.2 mmol) was added, and the reaction mixture was treated under the standard conditions to yield a light yellow solid (44 mg, 0.12 mmol, 49%, mp: 117 - 118 °C). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.43 (m, 3H), 7.35 (m, 5H), 7.09 (m, 2H), 5.20 (q-like, *J* = 12.1 Hz, 2H), 3.73 (t, *J* = 6.5 Hz, 1H), 3.30 (dd, *J* = 6.5, 17.0 Hz, 1H), 2.85 (dd, *J* = 6.6, 16.9 Hz, 1H), 2.33 (m, 2H), 1.99 (m, 2H), 1.81 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.8, 168.8, 167.0, 154.8, 136.9, 135.3, 129.6, 128.9, 128.6, 128.5, 128.4, 113.8, 67.4, 47.6, 36.1, 27.9, 21.7, 21.2; ESI-HRMS: calc'd m/e for [M+H⁺] C₂₃H₂₂NO₄: 376.1549, found 376.1546; IR (neat, NaCl, cm⁻¹): 2951, 1740, 1704, 1652, 1624, 1595, 1491, 1455, 1379, 1350, 1311, 1271, 1244, 1197, 1155.

Methyl 1-Butyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5e): The adduct (42 mg, 0.14 mmol) was dissolved in dichloroethane (4 mL), TFA (0.31 mL, 3.8 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a viscous yellow oil (24 mg, 0.077 mmol, 57%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 3.74 (s, 3H), 3.69 (m, 2H), 3.48 (dd, *J* = 6.8, 9.3 Hz, 1H), 2.94 (dd, *J* = 9.3, 17.0 Hz, 1H), 2.79 (dd, *J* = 6.8, 16.6 Hz, 1H), 2.60 (m, 2H), 2.40 (t, *J* = 6.2 Hz, 2H), 2.07 (m, 2H), 1.54 (m, 2H), 1.35 (m, 2H), 0.94 (t, *J* = 7.3 Hz, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.7,

169.6, 166.8, 154.3, 114.7, 52.6, 47.3, 42.6, 36.0, 31.2, 26.2, 21.8, 20.6, 20.1, 13.8; ESI-HRMS: calc'd m/e for $[M+H]^+$ $C_{15}H_{22}NO_4$: 280.1549, found 280.1544; IR (neat, NaCl, cm^{-1}): 2957, 1744, 1690, 1652, 1618, 1435, 1393, 1279, 1245, 1201, 1121.

Benzyl 1-Butyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5f): The adduct (96 mg, 0.21 mmol) was dissolved in toluene (6 mL), TFA (0.48 mL, 6.2 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a viscous colorless oil (44 mg, 0.13 mmol, 60%). 1H NMR ($CDCl_3$, 400 MHz, ppm): δ 7.36 - 7.29 (m, 5H), 5.16 (q-like, J = 12.2 Hz, 2H), 3.68 (m, 2H), 3.52 (dd, J = 6.6, 8.0 Hz, 1H), 3.04 (dd, J = 8.0, 16.7 Hz, 1H), 2.73 (dd, J = 6.4, 17.2 Hz, 1H), 2.51 (m, 2H), 2.34 (m, 2H), 1.96 (m, 2H), 1.50 (m, 2H), 1.34 (m, 2H), 0.93 (t, J = 7.2 Hz, 3H); ^{13}C NMR ($CDCl_3$, 100 MHz, ppm): δ 195.6, 168.9, 166.7, 154.5, 135.4, 128.6, 128.4, 128.3, 114.7, 67.2, 47.5, 42.6, 35.9, 31.2, 26.1, 21.7, 20.8, 20.1, 13.7; ESI-HRMS: calc'd m/e for $[M+H]^+$ $C_{21}H_{26}NO_4$: 356.1862, found 356.1861; IR (neat, NaCl, cm^{-1}): 2958, 1741, 1691, 1652, 1618, 1498, 1456, 1394, 1278, 1245, 1197, 1172, 1121.

Methyl 1-Benzyl-7,7-dimethyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5g): The adduct (52 mg, 0.14 mmol) was dissolved in dichloroethane (5 mL), TFA (0.32 mL, 4.2 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a viscous colorless oil (32 mg, 0.093 mmol, 67%). 1H NMR ($CDCl_3$, 400 MHz, ppm): δ 7.34 (m, 2H), 7.26 (m, 1H), 7.17 (m, 2H), 5.09 (q-like, J = 16.4 Hz, 2H), 3.75 (s, 3H), 3.64 (t, J = 6.9 Hz, 1H), 3.14 (dd, J = 7.2, 16.8 Hz, 1H), 2.80 (dd, J = 6.6, 16.8 Hz, 1H), 2.34 (s, 2H), 2.22 (s, 2H), 0.95 (s, 3H), 0.92 (s, 3H); ^{13}C NMR ($CDCl_3$, 100 MHz, ppm): δ 195.7, 169.6, 167.4, 152.9, 136.8, 129.1, 127.7, 126.1, 114.1, 52.8, 49.8, 47.5, 45.6, 40.1, 33.2, 28.9, 27.5, 20.6; ESI-HRMS: calc'd m/e for $[M+H]^+$ $C_{20}H_{24}NO_4$: 342.1705, found 342.1698; IR (neat, NaCl, cm^{-1}): 2956, 1741, 1693, 1653, 1625, 1497, 1454, 1439, 1386, 1303, 1280, 1252, 1204, 1168.

Benzyl 1-Benzyl-7,7-dimethyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5h): The adduct (117 mg, 0.22 mmol) was dissolved in toluene (7 mL), TFA (0.46 mL, 6.7 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a viscous colorless oil (60 mg, 0.14 mmol, 64%). 1H NMR ($CDCl_3$, 400 MHz, ppm): δ 7.35 - 7.25 (m, 8H), 7.16 (m, 2H), 5.18 (q-like, J = 12.2 Hz, 2H), 5.00 (q-like, J = 16.6 Hz, 2H), 3.69 (t, J = 6.8 Hz, 1H), 3.19 (dd, J = 7.0, 16.8 Hz, 1H), 2.79 (dd, J = 4.7, 16.8 Hz, 1H), 2.27 (q-like, J = 10.2 Hz, 2H), 2.15 (q-like, J = 16.3 Hz, 2H), 0.90 (s, 3H), 0.84 (s, 3H); ^{13}C NMR ($CDCl_3$, 100 MHz, ppm): δ 195.6, 169.0, 167.3, 153.0, 136.7, 135.4, 129.0, 128.7, 128.5, 128.4, 127.6, 126.0, 114.0, 67.4, 49.7, 47.6, 45.5, 40.0, 33.0, 28.6, 27.7, 20.7; ESI-HRMS: calc'd m/e for $[M+H]^+$ $C_{26}H_{28}NO_4$: 418.2018, found 418.2021; IR (neat, NaCl, cm^{-1}): 2959, 1739, 1694, 1653, 1625, 1497, 1455, 1386, 1369, 1303, 1252, 1213, 1169, 1124.

Methyl 7,7-Dimethyl-2,5-dioxo-1-phenyl-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate (5i): The adduct (54 mg, 0.21 mmol) was dissolved in toluene (6 mL), TFA (0.48 mL, 6.2 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a white solid (44 mg, 0.12 mmol, 64%, mp: 123 - 124 °C). 1H NMR ($CDCl_3$, 400

MHz, ppm): δ 7.50 - 7.42 (m, 3H), 7.21 - 7.10 (m, 2H), 3.75 (s, 3H), 3.69 (t, J = 5.8 Hz, 1H), 3.25 (dd, J = 6.6, 16.9 Hz, 1H), 2.86 (dd, J = 6.6, 16.9 Hz, 1H), 2.26 (s, 2H), 1.95 (q-like, J = 17.6 Hz, 2H), 0.97 (s, 3H), 0.95 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 195.8, 169.5, 167.2, 153.0, 136.9, 129.7, 128.9, 128.5, 113.0, 52.7, 49.9, 47.5, 41.5, 33.2, 28.8, 27.4, 20.8; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{19}\text{H}_{22}\text{NO}_4$: 328.1549, found 328.1540; IR (neat, NaCl, cm^{-1}): 2956, 1740, 1704, 1654, 1628, 1596, 1492, 1437, 1380, 1317, 1292, 1263, 1239, 1144.

Benzyl 7,7-Dimethyl-2,5-dioxo-1-phenyl-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate

(5j): The adduct (111 mg, 0.22 mmol) was dissolved in toluene (7 mL), TFA (0.50 mL, 6.5 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a white solid (63 mg, 0.16 mmol, 72%, mp: 126 - 127 °C). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.43 (m, 3H), 7.34 (m, 5H), 7.07 (m, 2H), 5.18 (q-like, J = 12.2 Hz, 2H), 3.73 (t, J = 6.3 Hz, 1H), 3.33 (dd, J = 5.8, 16.9 Hz, 1H), 2.84 (dd, J = 6.3, 16.9 Hz, 1H), 2.19 (q-like, J = 16.4 Hz, 2H), 1.88 (q-like, J = 17.0 Hz, 2H), 0.94 (s, 3H), 0.86 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 195.7, 168.8, 167.2, 152.9, 136.9, 135.2, 129.62, 129.61, 128.9, 128.7, 128.52, 128.47, 112.8, 67.5, 49.8, 47.6, 41.4, 33.0, 28.6, 27.6, 20.9; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{25}\text{H}_{26}\text{NO}_4$: 404.1862, found 404.1864; IR (neat, NaCl, cm^{-1}): 2958, 1740, 1705, 1654, 1628, 1596, 1492, 1455, 1379, 1351, 1317, 1292, 1240, 1262, 1173, 1143.

Methyl 1-Butyl-7,7-dimethyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carboxylate

(5k): The adduct (75 mg, 0.22 mmol) was dissolved in dichloroethane (7 mL), TFA (0.51 mL, 6.6 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a viscous light yellow oil (42 mg, 0.14 mmol, 62%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 3.71 (s, 3H), 3.68 (m, 2H), 3.48 (dd, J = 6.6, 8.0 Hz, 1H), 3.00 (dd, J = 8.1, 16.7 Hz, 1H), 2.72 (dd, J = 6.6, 16.8 Hz, 1H), 2.42 (s, 2H), 2.27 (s, 2H), 1.53 (m, 2H), 1.35 (m, 2H), 1.10 (s, 3H), 1.07 (s, 3H), 0.94 (t, J = 7.3 Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 195.6, 169.7, 167.1, 152.6, 113.8, 52.7, 49.8, 47.4, 42.5, 40.1, 33.2, 31.4, 29.0, 27.9, 20.5, 20.2, 13.9; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{17}\text{H}_{26}\text{NO}_4$: 308.1862, found 308.1868; IR (neat, NaCl, cm^{-1}): 2958, 2872, 1745, 1692, 1653, 1623, 1439, 1391, 1310, 1279, 1248, 1206, 1173, 1118, 1032, 1009.

Methyl 1-Benzyl-2,5-dioxo-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridine-3-carboxylate

(5l): The adduct (51 mg, 0.15 mmol) was dissolved in dichloroethane (5 mL), TFA (0.36 mL, 4.6 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a white solid (29 mg, 0.095 mmol, 62%, mp: 134 - 135 °C). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.37 - 7.29 (m, 3H), 7.24 (m, 2H), 4.96 (q-like, J = 15.8 Hz, 2H), 3.77 (s, 3H), 3.72 (t, J = 7.6 Hz, 1H), 2.96 (dd, J = 7.3, 17.3 Hz, 1H), 2.75 (dd, J = 7.9, 17.0 Hz, 1H), 2.65 (m, 2H), 2.46 (m, 2H); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 201.5, 169.6, 168.4, 167.4, 136.3, 129.1, 128.0, 126.9, 116.9, 53.0, 47.5, 46.5, 34.3, 25.4, 19.8; ESI-HRMS: calc'd m/e for $[\text{M}+\text{H}^+]$ $\text{C}_{17}\text{H}_{18}\text{NO}_4$: 300.1236, found 300.1230; IR (neat, NaCl, cm^{-1}): 2953, 2928, 1742, 1687, 1640, 1496, 1437, 1412, 1363, 1322, 1281, 1253, 1213, 1167, 1137, 1123, 1027, 1083.

Methyl 2,5-Dioxo-1-phenyl-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridine-3-carboxylate (5m): The adduct (76 mg, 0.24 mmol) was dissolved in dichloroethane (7 mL), TFA (0.55 mL, 7.2 mmol) was added, and then, the reaction mixture was treated under the standard conditions to yield a white solid (34 mg, 0.12 mmol, 49%, mp: 100 - 101 °C). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.50 - 7.44 (m, 3H), 7.24 (m, 2H), 3.79 (t, *J* = 6.4 Hz, 1H), 3.78 (s, 3H), 3.06 (dd, *J* = 6.3, 17.1 Hz, 1H), 2.83 (dd, *J* = 7.8, 17.1 Hz, 1H), 2.45 (m, 2H), 2.33 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 201.9, 169.7, 168.6, 167.3, 136.1, 129.8, 129.4, 128.3, 116.1, 53.1, 47.6, 34.3, 26.0, 20.1; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₆H₁₆NO₄: 286.1079, found 286.1074; IR (neat, NaCl, cm⁻¹): 2954, 2929, 1743, 1690, 1641, 1595, 1564, 1492, 1439, 1395, 1352, 1317, 1293, 1257, 1140, 1024, 1000.

General Procedure for the Preparation of 2,5-Dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carbonitriles 10 (Table 4).—The enaminone **6** was dissolved in acetonitrile (100 mL/mmol) and stirred at ambient temperature. To the solution, LiClO₄ (1 equiv) was added, followed by the addition of PPh₃ (0.5 equiv), methyl cyanoacetate (1.5 equiv), and paraformaldehyde (1.0 equiv) in sequence. The reaction was heated to 60 °C in a sealed vessel for 14 h (monitored by TLC). The reaction was quenched with saturated NaHCO₃ solution, upon consumption of starting material. The mixture was then extracted with CH₂Cl₂ (10 mL × 2), and the organic layers were combined, dried over MgSO₄, and concentrated *in vacuo*. The crude product was purified by flash column chromatography on silica gel, using 25% EtOAc:hexanes (v/v) as the eluent.

1-Benzyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carbonitrile (10a): The *N*-benzyl enaminone (50 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (8 mg, 0.25 mmol), methyl cyanoacetate (0.033 mL, 0.37 mmol), and triphenylphosphine (33 mg, 0.13 mmol) in acetonitrile (25 mL) were subjected to the standard conditions and yielded a colorless oil (60 mg, 0.21 mmol, 85%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.37 - 7.27 (m, 3H), 7.13 (m, 2H), 5.12 (d, *J* = 16.3 Hz, 1H), 4.91 (d, 16.3 Hz, 1H), 3.75 (dd, *J* = 6.3, 11.3 Hz, 1H), 3.17 (ddd, *J* = 1.4, 4.8, 15.3 Hz, 1H), 2.87 (m, 1H), 2.53 (m, 2H), 2.38 (t, *J* = 6.2 Hz, 2H), 1.99 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.1, 163.0, 154.6, 135.8, 129.2, 127.9, 126.1, 115.6, 114.6, 46.0, 35.7, 34.4, 26.3, 21.8, 21.5; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₇H₁₇N₂O₂: 281.1290, found 281.1281; IR (neat, NaCl, cm⁻¹): 2959, 2890, 2256, 1756, 1700, 1655, 1627, 1497.

2,5-Dioxo-1-phenyl-1,2,3,4,5,6,7,8-octahydroquinoline-3-carbonitrile (10b): The *N*-phenyl enaminone (47 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (8 mg, 0.25 mmol), methyl cyanoacetate (0.033 mL, 0.37 mmol), and triphenylphosphine (33 mg, 0.13 mmol) in acetonitrile (25 mL) were subjected to the standard conditions and yielded a yellow solid (43 mg, 0.16 mmol, 65%, mp: 158 - 160 °C). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 7.57 - 7.46 (m, 3H), 7.15 (m, 2H), 3.83 (dd, *J* = 6.4, 11.1 Hz, 1H), 3.24 (dd, *J* = 6.3, 15.2 Hz, 1H), 2.97 (m, 1H), 2.42 (dt, *J* = 1.9, 6.4 Hz, 2H), 2.13 (m, 2H), 1.96 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.3, 162.7, 154.8, 136.1, 129.9, 129.4, 128.5, 115.5, 113.6, 36.0, 34.7, 27.9, 22.1, 21.6; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₆H₁₅N₂O₂: 267.1134, found 267.1134; IR (neat, NaCl, cm⁻¹): 2940, 2260, 1765, 1710, 1655, 1627, 1500.

1-Butyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carbonitrile (10c): The *N*-butyl enaminone (42 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (8 mg, 0.25 mmol), methyl cyanoacetate (0.033 mL, 0.37 mmol), and triphenylphosphine (33 mg, 0.13 mmol) in acetonitrile (25 mL) were subjected to the standard conditions and yielded a colorless oil (44 mg, 0.18 mmol, 73%). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 3.77 (m, 1H), 3.63 (m, 2H), 3.11 (dd, *J* = 6.3, 16.5 Hz, 1H), 2.74 (m, 1H), 2.61 (m, 2H), 2.43 (t, *J* = 7.1 Hz, 2H), 2.11 (m, 2H), 1.53 (m, 2H), 1.34 (m, 2H), 0.94 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.0, 162.5, 154.4, 115.7, 114.3, 43.1, 35.8, 34.4, 31.2, 26.2, 21.8, 21.7, 20.0, 13.7; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₄H₁₉N₂O₂: 247.1447, found 247.1449; IR (neat, NaCl, cm⁻¹): 2964, 2897, 2256, 1756, 1700, 1655, 1627.

7,7-Dimethyl-2,5-dioxo-1-propyl-1,2,3,4,5,6,7,8-octahydroquinoline-3-carbonitrile (10d): The *N*-propyl enaminone (45 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (8 mg, 0.25 mmol), methyl cyanoacetate (0.033 mL, 0.37 mmol), and triphenylphosphine (33 mg, 0.13 mmol) in acetonitrile (25 mL) were subjected to the standard conditions and yielded a white solid (49 mg, 0.19 mmol, 75%, mp: 136 - 137 °C). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 3.71 (m, 1H), 3.62 (m, 2H), 3.03 (dd, *J* = 6.3, 16.6 Hz, 1H), 2.81 (m, 1H), 2.43 (m, 2H), 2.29 (d, *J* = 16.9 Hz, 2H), 1.53 (m, 2H), 1.08 (s, 3H), 1.06 (s, 3H), 0.88 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 194.0, 161.7, 151.5, 114.8, 112.2, 48.5, 43.6, 38.9, 33.4, 32.1, 27.5, 27.3, 21.5, 20.5, 10.0; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₅H₂₁N₂O₂: 261.1603, found 261.1612; IR (neat, NaCl, cm⁻¹): 2962, 2903, 2252, 1754, 1702, 1652, 1628.

1-(2-Methoxyethyl)-7,7-dimethyl-2,5-dioxo-1,2,3,4,5,6,7,8-octahydroquinoline-3-carbonitrile (10e): The *N*-methoxyethyl enaminone (49 mg, 0.25 mmol), LiClO₄ (27 mg, 0.25 mmol), paraformaldehyde (8 mg, 0.25 mmol), methyl cyanoacetate (0.033 mL, 0.37 mmol), and triphenylphosphine (33 mg, 0.13 mmol) in acetonitrile (25 mL) were subjected to the standard conditions and yielded a white solid (47 mg, 0.17 mmol, 67%, mp: 77 - 79 °C). ¹H NMR (CDCl₃, 400 MHz, ppm): δ 3.91 (m, 2H), 3.63 (dd, *J* = 6.2, 11.1 Hz, 1H), 3.52 (m, 2H), 3.28 (s, 3H), 3.08 (dd, *J* = 6.2, 16.3 Hz, 1H), 2.80 (m, 1H), 2.61 (d, *J* = 17.2 Hz, 1H), 2.49 (d, *J* = 17.2 Hz, 1H), 2.28 (d, *J* = 3.0 Hz, 2H), 1.10 (s, 3H), 1.08 (s, 3H); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 195.4, 163.1, 153.6, 115.7, 112.6, 70.3, 59.1, 49.5, 43.3, 40.2, 34.4, 33.0, 28.4, 28.3, 21.5; ESI-HRMS: calc'd m/e for [M+H⁺] C₁₅H₂₁N₂O₃: 277.1552, found 277.1558; IR (neat, NaCl, cm⁻¹): 2957, 2889, 2252, 1770, 1715, 1655, 1627, 1249, 1207.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Triggle DJ Cell. Mol. Neurobiol 2003, 23, 293–303. [PubMed: 12825828]
2. Rose U Arch. Pharm 1990, 323, 281–286.
3. Takahashi D; Oyzunzul L; Onoue S; Ito Y; Uchida S; Simsek R; Gunduz MG; Safak C; Yamada S Biol. Pharm. Bull 2008, 31, 473–479. [PubMed: 18310913]

4. Willems E; Cabral-Teixeira J; Schade D; Cai W; Reeves P; Bushway PJ; Lanier M; Walsh C; Kirchhausen T; Izpisua BJC; Cashman J; Mercola M *Cell Stem Cell* 2012, 11, 242–252. [PubMed: 22862949]
5. Schade D; Lanier M; Willems E; Okolotowicz K; Bushway PJ; Wahlquist C; Gilley C; Mercola M; Cashman JR *J. Med. Chem* 2012, 55, 9946–9957. [PubMed: 23130626]
6. Movsesian MA; Kukreja RC, Phosphodiesterase Inhibition in Heart Failure In *Phosphodiesterases as Drug Targets*, Francis SH; Conti M; Houslay MD, Eds. Springer: Berlin, 2011; pp 237–250.
7. Gorobets NY; Yousefi BH; Belaj F; Kappe CO *Tetrahedron* 2004, 60, 8633–8644.
8. Fossa P; Menozzi G; Dorigo P; Floreani M; Mosti L *Bioorg. Med. Chem* 2003, 11, 4749–4759. [PubMed: 14556790]
9. Noguez MO; Marcelino V; Rodriguez H; Martin O; Martinez JO; Arroy GA; Perez FJ; Suarez M; Miranda R *Int. J. Mol. Sci* 2011, 12, 2641–2649. [PubMed: 21731463]
10. Hirsch KS; Jones CD Preparation of Hexahydrobenzo [f]quinolinones as 5- α -Reductase Inhibitors. 1992–307636 531026, 19920820., 1993.
11. Jones CD; Audia JE; Lawhorn DE; McQuaid LA; Neubauer BL; Pike AJ; Pennington PA; Stamm NB; Toomey RE; Hirsch KS *J. Med. Chem* 1993, 36, (3), 421–423. [PubMed: 8381185]
12. Astleford BA; Audia JE; Deeter J; Heath PC; Janisse SK; Kress TJ; Wepsiec JP; Weigel LO *J. Org. Chem* 1996, 61, (13), 4450–4454. [PubMed: 11667351]
13. Ko S; Sastry MNV; Lin C; Yao C *Tetrahedron Lett.* 2005, 46, 5771–5774.
14. Sun J; Xia E-Y; Wu Q; Yan C-G *ACS Comb. Sci* 2011, 13, 421–426. [PubMed: 21568339]
15. Fan XS; Li YZ; Zhang XY; Qu GR; Wang JJ; Hu XY *Heteroatom. Chem* 2006, 17, 382–388.
16. Tu S; Zhu X; Zhang J; Xu J; Zhang Y; Wang Q; Jia R; Jiang B; Zhang J; Yao C *Bioorg. Med. Chem. Lett* 2006, 16, 2925–2928. [PubMed: 16563758]
17. Suárez M; Ochoa E; Verdecia Y; Pita B; Morán L; Martín N; Quinteiro M; Seoane C; Soto J; Novoa H; Blaton N; Peters OM *Tetrahedron* 1999, 55, 875–884.
18. Azzam SHS; Siddekha A; Pasha MA *Tetrahedron Lett.* 2012, 53, (46), 6306–6309.
19. Enders D; Demir AS; Puff H; Franken S *Tetrahedron Lett.* 1987, 28, (33), 3795–3798.
20. Wang G; Miao C *Green Chem.* 2006, 8, 1080–1085.
21. Saidi MR; Azizi N; Zali-Boinee H *Tetrahedron* 2001, 57, 6829–6832.
22. Azizi N; Saidi MR *Tetrahedron* 2003, 59, 5329–5332.
23. Saidi MR; Azizi N; Naimi-Jamal MR *Tetrahedron Lett.* 2001, 42, 8111–8113.
24. Saidi MR; Azizi N *Tetrahedron: Asymmetry* 2002, 13, 2523–2527.
25. Aryanasab F; Saidi MR *Synthetic Commun.* 2008, 38, 4036–4044.
26. Hoye TR; Caruso AJ; Magee AS *J. Org. Chem* 1982, 47, (21), 4152–6.
27. Nenajdenko VG; Statsuk AV; Balenkova ES *Tetrahedron* 2000, 56, (35), 6549–6556.
28. Cannon JG *J. Am. Pharmaceut. Assoc* 1956, 45, 430.
29. Smith AB; Liu Z *Org. Lett* 2008, 10, 4363–4365. [PubMed: 18754624]
30. Deka N; Kalita DJ; Borah R; Sarma JC *J. Org. Chem* 1997, 62, (5), 1563–1564.
31. Li Z-X; Liu X-P; Qiu Z; Xu D; Yu X-J *J. Chem. Res* 2011, 35, (1), 35–36.
32. Hourcade S; Ferdenzi A; Retailleau P; Mons S; Marazano C *Eur. J. Org. Chem* 2005, 2005, 1302–1310.
33. Gray D; Gallagher T *Angew. Chem. Int. Ed* 2006, 45, 2419–2423.
34. Barnett CJ; Wilson TM; Evans DA; Somers TC *Tetrahedron Lett.* 1997, 38, 735–738.
35. Yadav JS; Reddy BVSR; Basak AK; Visali BN, A.V.; Nagagaiah K *Eur. J. Org. Chem* 2004, 546–551.
36. Lu X; Zhang C; Xu Z *Acc. Chem. Res* 2001, 34, 535–544. [PubMed: 11456471]
37. Nair V; Menon RS; Sreekanth AR; Abhilash N; Biju AT *Acc. Chem. Res* 2006, 39, 520–530. [PubMed: 16906748]
38. Methot JL; Roush WR *Adv. Synth. Catal* 2004, 346, 1035–1050.
39. Liu H; Zhang Q; Wang L; Tong X *Chem. Commun* 2010, 46, 312–314.

40. Stewart IC; Bergman RG; Toste FD J. Am. Chem. Soc 2003, 125 (29) 8696–8697 [PubMed: 12862443]
41. Yadav JS; Reddy BVS; Krishna AD; Reddy CS; Narsaiah AV J. Mol. Catal. A: Chem 2007, 261, (1), 93–97.
42. Dyachenko VD; Chernega AN Russ. J. Gen. Chem 2005, 75, 952–960.
43. Bhattacharyya P; Pradhan K; Paul S; Das AR Tetrahedron Lett. 2012, 53, (35), 4687–4691.
44. Pradhan K; Bhattacharyya P; Paul S; Das AR Tetrahedron Lett. 2012, 53, (44), 5840–5844.
45. <http://pubchem.ncbi.nlm.nih.gov>.

Figure 1.

Representative structures of cardiovascular drugs, 1,4-dihydropyridine nifedipine (1), a calcium channel blocker, TGFβ inhibitor 2, milrinone (3), annulated 2-pyridone phosphodiesterase-3 inhibitor 4, human type I steroid 5-α-reductase inhibitor LY191704, and target scaffold 5.

Scheme 1.

Multicomponent literature methods for the synthesis of 5-oxohexahydroquinoline-3-carboxylates (eq 1), 2,5-Dioxooctahydroquinoline-3-carboxylates (eq 2 and 3) and a stepwise reaction to form 2,5-dioxooctahydroquinoline-3-carboxylates (eq 4).

Scheme 2.

Multi-component reactions of 3-(benzylamino)cyclohex-2-enone (**6a**) with and without LiClO₄ as additive.

Scheme 3.
Reaction of enaminone **6a** with Knoevenagel intermediate **9a**.

Scheme 4.

Proposed mechanism for the LiClO_4 -catalyzed Knoevenagel reaction in the presence of acetic anhydride.

Scheme 5.

Proposed mechanism for the triphenylphosphine catalyzed synthesis of **10a**.

Table 1Optimization of the LiClO₄-assisted formation of **7a**

entry	temp (°C)	LiClO ₄ (equiv)	Ac ₂ O (equiv)	yield (%)
1	90	0	0	12
2	90	0.9	0	63
3	90	1.0	0	77
4	90	1.9	0	75
5	90	10.0	0	62
6	60	1.0	1.0	93 ^a
7	60	Li ⁺ TFA ⁻ 1.0	1.0	57
8	60	Ti(O <i>i</i> Pr) ₄ 1.0	1.0	0 ^b
9	60	MgBr ₂ •OEt ₂	1.0	0 ^b

^a used as the standard condition;^b *bis*-addition product observed,

none of the desired product was observed.

Table 2

Scope of the LiClO₄-assisted multi-component reaction

entry	compound	n	R ¹	R ²	R ³	R ⁴	yield (%)
1	7b	1	H	Ph	Me	Me	90
2	7c	1	H	Ph	Bn	Bn	97
3	7d	1	H	Bn	Me	Bn	73
3	7e	1	H	Bn	Bn	Bn	90
5	7f	1	H	<i>n</i> Bu	Me	Me	66
6	7g	1	H	<i>n</i> Bu	Bn	Bn	80
7	7h	1	H	morpholine	Me	Me	72
8	7i	1	H	morpholine	Bn	Bn	86
9	7j	1	Me	Ph	Me	Me	83
10	7k	1	Me	Ph	Bn	Bn	87
11	7l	1	Me	Bn	Me	Me	95
12	7m	1	Me	Bn	Bn	Bn	88
13	7n	1	Me	<i>n</i> Bu	Me	Me	88
14	7o	0	H	Ph	Me	Me	96
15	7p	0	H	Bn	Me	Me	94
16	7q	1	H	Bn	<i>t</i> Bu	Et	np
17	7r	1	H	Bn	dimethyl methylmalonate		np ^a
18	7s	1	H	Bn	methyl acetoacetate		np
19	7t	1	H	Bn	triethyl phosphonoacetate		np
20	7u	1	H	Bn	benzaldehyde, dimethyl malonate		np

^a np = no product.

Table 3

Synthesis of dioxooctahydroquinoline esters **5**

entry	compound	n	R ¹	R ²	^a R ³	yield (%)
1	5a	1	H	Bn	Me	79
2	5b	1	H	Bn	Bn	70
3	5c	1	H	Ph	Me	84
4	5d	1	H	Ph	Bn	49
5	5e	1	H	<i>n</i> Bu	Me	57
6	5f	1	H	<i>n</i> Bu	Bn	60
7	5g	1	Me	Bn	Me	67
8	5h	1	Me	Bn	Bn	64
9	5i	1	Me	Ph	Me	64
10	5j	1	Me	Ph	Bn	72
11	5k	1	Me	<i>n</i> Bu	Me	62
12	5l	0	H	Bn	Me	62
13	5m	0	H	Ph	Me	49

^a reaction performed in dichloroethanewhen R³ = Me, in toluene when R³ = Bn

Table 4Synthesis of dioxooctahydroquinoline-3-carbonitriles **10**

entry	compound	R ¹	R ²	yield (%)
1	10a	H	Bn	85
2	10b	H	Ph	65
3	10c	H	Bu	73
4	10d	Me	Pr	75
5	10e	Me	MeO(CH ₂) ₂ -	67