Published in final edited form as:

Gen Comp Endocrinol. 2019 January 01; 270: 75–81. doi:10.1016/j.ygcen.2018.10.008.

Exendin-4 antagonizes the metabolic action of acylated ghrelinergic signaling in the hypothalamic paraventricular nucleus

Shayan Abtahi, Erin Howell, Jack T. Salvucci, Joshua M.R. Bastacky, David P. Dunn, Paul J. Currie*

Department of Psychology, Reed College, 3203 SE Woodstock Blvd., Portland, OR 97202

Abstract

In the current study we investigated the interaction of hypothalamic paraventricular nucleus (PVN) glucagon-like peptide-1 (GLP-1) and ghrelin signaling in the control of metabolic function. We first demonstrated that acylated ghrelin injected directly into the PVN reliably altered the respiratory exchange ratio (RER) of adult male Sprague Dawley rats. All testing was carried out during the initial 2 h of the nocturnal cycle using an indirect open circuit calorimeter. Results indicated that acylated ghrelin induced a robust increase in RER representing a shift toward enhanced carbohydrate oxidation and reduced lipid utilization. In contrast, treatment with comparable dosing of des-acyl ghrelin failed to significantly impact metabolic activity. In separate groups of rats we subsequently investigated the ability of exendin-4 (Ex-4), a GLP-1 analogue, to alter acylated ghrelin's metabolic effects. Rodents were treated with either systemic or direct PVN Ex-4 followed by acyl ghrelin microinjection. While our results showed that both systemic and PVN administration of Ex-4 significantly reduced RER, importantly, Ex-4 pretreatment itself reliably inhibited the impact of ghrelin on RER. Overall, these findings provide increasingly compelling evidence that GLP-1 and ghrelin signaling interact in the neural control of metabolic function within the PVN.

Keywords

Acyl Ghrelin; Des-acyl Ghrelin; Energy substrate oxidation; Ghrelin; GLP-1; Hypothalamus; Neuropeptide Y; Respiratory exchange ratio

Introduction

Ghrelin, encoded by the *GHRL* gene, is an evolutionarily conserved neuroendocrine modulator critical for homeostatic energy balance and metabolism in a wide variety of species and first characterized in the rat (*Rattus norvegicus*) stomach by Kojima et al., (1999) (Abtahi et al., 2016, 2017; Blanco et al., 2017; Boswell and Dunn, 2017; Geelissen et

*Corresponding Author, Department of Psychology, Reed College, Portland, OR, 97202. Tel: 503-777-7267; pcurrie@reed.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

al., 2006; Nakazato et al., 2001; Velasco et al., 2015). The acylated peptide functions as the endogenous ligand for the peripherally and centrally expressed growth hormone secretagogue 1a receptor (GHSR-1a) (Hosoda et al., 2003; Kojima et al., 1999; Zigman et al., 2006). A particularly unique feature of ghrelin is the requirement of octanoylation at the serine-3 residue by the enzyme ghrelin-o-acyl-transferase (GOAT) in order to render it acylated and therefore active at the GHSR-1a (Gutierrez et al., 2008; Mohan and Unniappan, 2013; Romero et al., 2010; Yang et al., 2008). A number of studies have demonstrated that acylated ghrelin not only stimulates food intake but also increases respiratory exchange ratio (RER) in mammals and teleosts (Abtahi et al., 2016, 2017; Blanco et al., 2017; Currie et al., 2005; Currie et al., 2012; Lopez-Ferreras et al., 2017; Thomas et al., 2016). RER is an index of the nutrient source that animals metabolize to generate energy and is measured using indirect calorimetry as the volume of carbon dioxide produced divided by the volume of oxygen consumed (VCO_2/VO_2). RER values typically range from 0.7–1.0 with the lower bound indicating lipid oxidation and the higher bound representing carbohydrate utilization. Values in the 0.8 range indicate a combined reliance on carbohydrate and fat utilization (Bray et al., 1989; Coscina et al., 1998; Kleiber et al., 1987).

Although ghrelin is orexigenic in mammalian species, in avians, the peptide is found to be anorexigenic and is observed to decrease RER (Geelissen et al., 2006; Oclon and Pietras, 2010). However, one report did demonstrate that low doses of ghrelin stimulated food intake in the Japanese Quail (*Coturnix japonica*) (Shousha et al., 2005). Indeed, the majority of circulating ghrelin consists largely of des-acyl ghrelin, with less than 10% of the acylated molecule available to bind to the GHSR-1a (Delporte, 2013). While the function of des-acyl ghrelin remains unclear, recent evidence suggests that it may play a role in thermoregulation and stress activation in mammals (Inoue et al., 2016; Mahbod et al., 2018). Unlike the acylated peptide, des-acyl ghrelin does not bind to the GHSR-1a (Chen et al., 2017; Delporte, 2013). In both goldfish (*Carassius auratus*) and rodents, high doses of des-acyl ghrelin may antagonize the effect of acylated ghrelin on food intake by preventing the binding of acyl ghrelin to the 1a receptor (Matsuda et al., 2006). Therefore, although des-acyl ghrelin may not bind directly to the GHSR-1a, it does appear to play a role in regulating acyl ghrelin signaling in these species. Further, it remains possible that des-acyl ghrelin could bind and elicit its actions through a separate, yet unidentified, receptor protein (Chen et al., 2017).

As indicated above, the orexigenic and metabolic activity of acylated ghrelin are purported to be exerted centrally via GHSR-1a, which has wide expression throughout the hypothalamus in teleosts, mammals, and avians (Chen et al., 2007; Sanchez-Bretano et al., 2015; Zigman et al., 2006). For example, in teleosts and mammals, ghrelin stimulates food intake via neuropeptide Y (NPY)/agouti related peptide (AgRP) immunoreactive neurons in the arcuate nucleus (ArcN) (Chen et al., 2017; Delgado et al., 2017; Guan et al., 2008; Kohno et al., 2003; van den Top et al., 2007). In avians however, ghrelin's feeding effects appear to be largely mediated through corticotropin releasing hormone (CRH), given that intracerebroventricular (ICV) ghrelin elevates plasma corticosterone, whereas the CRH antagonist astressin reduces ghrelin's anorexigenic effect (Saito et al., 2005). Despite the greater ratio of circulating des-acyl to acyl ghrelin, robust evidence demonstrates that the acylated molecule is responsible for the ghrelinergic control of appetite, energy balance, and

metabolic function (Abtahi et al., 2016, 2017; Currie et al., 2005, 2011, 2012; Naleid et al., 2005; Nakazato et al., 2001; Tschöp et al., 2000). Other studies report that ghrelin injection into the ventral tegmental area (VTA) and nucleus accumbens (NAcc) of the mesolimbic reward pathway increases consumption of standard chow and palatable food, indicating that ghrelin influences both homeostatic and hedonic appetitive motivation (King et al., 2011; Naleid et al., 2005; Weinberg et al., 2011). Moreover, ghrelin administration into the CA1 of the hippocampus enhances social transmission of food preference, which is impaired by local viral knockdown of GHSR-1a (Hsu et al., 2018). These findings suggest a role for ghrelin in mediating food choice.

In addition to ghrelin, another critical homeostatic peptide involved in food intake and energy metabolism is glucagon-like peptide 1 (GLP-1), a peptide derived from the tissue specific post-translational processing of the proglucagon peptide (Alhadeff et al., 2012; Blanco et al., 2017; Hayes, 2012; Honda, 2016; Kanoski et al., 2011; Reiner et al., 2018). Encoded by the *GCG* gene, the predominant active form of GLP-1 in mammals upon cleaving is GLP-1(7–36) amide (Hayes et al., 2010). GLP-1 binds to the peripherally and centrally expressed GLP-1 receptor (GLP-1R) and is consistently anorexigenic across species (Kanoski et al., 2011; Silverstein et al., 2001; Tachibana et al., 2004). ICV treatment of exendin-4 (Ex-4), a GLP-1 analogue, enhances c-fos expression in ghrelin immunoreactive neurons in the arcuate and paraventricular (PVN) nuclei of the hypothalamus in mice (Dalvi et al., 2012). Systemic and ICV Ex-4 administration attenuates ghrelin plasma levels (Perez-Tilve et al., 2007). In gastrointestinal L-cells, ghrelin *in vitro* stimulates GLP-1 release (Gagnon et al., 2015), and circulating GLP-1 and proglucagon levels are higher in GHSR-1a null mice (Xu et al., 2015). Other work has shown that ghrelin decreases proglucagon mRNA in goldfish and that peripheral GLP-1 antagonizes the appetitive action of ghrelin (Blanco et al., 2017). Similarly, ghrelin's metabolic effects in rats are also suppressed by GLP-1 receptor antagonism (Abtahi et al., 2016).

GLP-1 has been well characterized with respect to its anorexigenic role in the nucleus of the solitary tract (NTS) in the hindbrain of mammals, however, emerging evidence now suggests that the peptide is integral to hypothalamic appetitive and metabolic circuitry (Alhadeff et al., 2012; Chen et al., 2017; Guan et al., 2008; Kohno et al., 2003; van den Top et al., 2007). For example, we have recently demonstrated that Ex-4 antagonizes the stimulatory effect of ArcN ghrelin on RER (Abtahi et al., 2016). In this same report, ArcN Ex-4 reliably suppressed the increase in RER found after co-infusion of ghrelin and NPY (Abtahi et al., 2016). As a result, and in order to further investigate the metabolic function of hypothalamic GLP-1, ghrelin, and NPY, in the present study we targeted the PVN, given the well-established role of this nucleus in regulating energy metabolism and substrate oxidation (Currie et al., 2010; Gao et al., 2018; Thomas et al., 2015; Vasselli et al., 2017). Specifically, the overall objective of the current report was to initially compare the impact of acyl and des-acyl ghrelin on energy substrate utilization. We then determined whether or not direct PVN treatment with the GLP-1 agonist Ex-4 would suppress the effect of PVN ghrelin or the combined effect of ghrelin and NPY on RER.

Materials and methods

Animals

Adult male Sprague-Dawley rats (N=38; Envigo Laboratories, Madison, WI, USA) were housed individually in polypropylene cages equipped with an automated watering system. Rats weighing between 275–300 g were 10 weeks of age at the time of surgery. All animals had free access to standard rodent chow (LabDiet, St, Louis, MO, USA) and water. The colony facility was maintained on a 12 h light/dark cycle (lights off at 1500 h) and at a temperature of $22 \pm 2^\circ\text{C}$. The Institutional Animal Care and Use Committee of Reed College approved all experimental protocols (#A4425–01).

Stereotaxic surgery

Unilateral guide cannulae were implanted into the PVN. Stereotaxic coordinates relative to bregma were 1.8 mm posterior, ± 0.3 mm lateral, and 3.9 mm ventral (Paxinos and Watson, 2014). Rats were anesthetized with ketamine (100 mg/kg IP, Henry Schein, Melville, NY, USA) co-administered with xylazine (5 mg/kg IP, Sigma, St. Louis, MO, USA) and placed in a Kopf stereotaxic frame (Kopf Instruments, Tujunga, CA, USA), with the incisor bar set at -3.5 mm. Guide cannulae (22-gauge; Plastics One, Roanoke, VA, USA) were implanted 4 mm dorsal to target in order to minimize damage to PVN tissue. Each implant was secured with acrylic cement and a 28-gauge stainless-steel inner stylet was used to maintain cannula patency. RER testing was initiated after a two-week postoperative recovery period. During this time animals were handled daily and habituated with mock injections.

Measurement of RER

Oxygen (O_2) consumption and carbon dioxide (CO_2) production were measured using an open-circuit indirect calorimeter (Columbus Instruments, Columbus, OH, USA). Gas concentrations were sequentially derived for each test chamber using separate O_2 and CO_2 sensors. The flow rate was set at 2000 ml/min. Gases were assessed in ml/kg body weight/minute. Body weights for each rat were uploaded into the program immediately prior to testing. RER was expressed as the volume of CO_2 produced (VCO_2) divided by the volume of O_2 (VO_2) consumed. The analyzers were calibrated prior to each test using primary gas standards of high purity (Praxair, Vancouver, WA, USA).

In an initial study, rats with guide cannulae targeting the PVN, were administered acyl ghrelin (50–200 pmol; n=8) or des-acyl ghrelin (50–200 pmol; n=7) and then immediately placed into the Oxyscan calorimeter for 2 hours. All rats were tested with each dose of the peptide and vehicle, with treatments presented on separate days and in randomized order. At least three non-injection days separated successive test days. In order to investigate the impact of Ex-4 on ghrelin-induced changes in RER, individual groups of rats were pretreated with systemic (0.1–1.0 $\mu\text{g/kg}$ IP; n=8) or PVN (0.01–0.05 μg ; n=8) Ex-4 and then immediately administered acyl ghrelin (50 pmol) into the PVN. We did not investigate the effects of Ex-4 paired with des-acyl ghrelin given that the preceding study demonstrated that des-acyl ghrelin alone failed to alter RER. In a final study (n=7), low doses of ghrelin and NPY were co-administered in order to determine if Ex-4 pretreatment would antagonize the combined effects of PVN ghrelin-NPY signaling. These very low doses of ghrelin and NPY

were expected to be subthreshold when injected individually. In this experiment rats were treated with PVN ghrelin (10 pmol), NPY (15 pmol), ghrelin paired with NPY, or Ex-4 (0.01 µg) combined with ghrelin and NPY. Again, a repeated measures design was used with rats subjected to all treatments in randomized order.

Acyl ghrelin, des-acyl ghrelin, NPY, and Ex-4 were dissolved in sterile physiological saline and were each purchased from Tocris (Minneapolis, MN, USA). Doses were selected based on past work examining RER, body weight, and appetite (Abtahi et al., 2016; Alhadeff et al., 2012; López-Ferreras et al., 2017; Mietlicke-Baase et al., 2014; Nakazato et al., 2001). PVN injections were administered via an injector cannula extending 4 mm beyond the indwelling guide cannula. The volume of each injection was 0.2 µl. Each microinjection was carried out over a 4-minute period. Intraperitoneal Ex-4 was delivered in a volume of 1 ml/kg.

Metabolic testing began immediately postinjection. In order to avoid changes in metabolism associated with energy intake, food was removed from the home cage 1 h prior to testing and was not available in the metabolic chamber during RER testing. All experiments were carried out during the initial 2 h of the nocturnal active cycle.

Histological confirmation

Cannula placements were confirmed via histological examination as described previously (Abtahi et al., 2016, 2017). Immediately following CO₂ induced euthanasia, rats were injected with 0.4 µl of black ink to facilitate the identification of microinjector placement. Tissue sections were cut at 20 µm on a coronal plane using a cryostat (CryoStar NX50, Thermo Fisher, Waltham, MA, USA), mounted on slides, processed, stained with Cresyl violet (Sigma, St. Louis, MO, USA), and then coverslipped. Slides were examined via light microscopy and assessed using the stereotaxic atlas of Paxinos and Watson (2014). All animals reported here were found to have accurate placements in the PVN. A representative coronal section of the PVN is shown in Figure 1.

Statistical analyses

Data were analyzed using one or two-way repeated measures analyses of variance (ANOVA) followed by post hoc Tukey tests of mean comparisons when justified. The criterion for statistical significance was $p < 0.05$. The statistical software used was TIBCO Statistica (Palo Alto, CA, USA).

Results

The RER effects of PVN acyl and des-acyl ghrelin are illustrated in Figure 2. One-way repeated measures ANOVA revealed that acyl ghrelin administration evoked reliable, dose-dependent increases in RER ($F(3,21)=66.91$, $p < 0.001$) whereas the des-acyl ghrelin molecule failed to alter RER at any of the doses tested ($F(3,18)=0.23$, $p > 0.05$). In an additional group of rats, Ex-4 pretreatment was used to assess whether the GLP-1 receptor agonist could modify the robust effect of acyl ghrelin on substrate utilization. Two-way repeated measures ANOVA indicated that intraperitoneal Ex-4 injection elicited a significant reduction in the stimulatory impact of acyl ghrelin on RER ($F(2,14)=4.22$, $p < 0.01$). The inhibitory effect of Ex-4 was dose-dependent and while both doses of Ex-4 suppressed

ghrelin's effect, only the higher dose of 1.0 µg/kg significantly lowered RER on its own. (See Figure 3). Comparable results were detected following direct hypothalamic PVN Ex-4 pretreatment ($F(2,14)=9.97$, $p<0.004$) with both subthreshold and threshold doses reliably inhibiting the stimulatory action of acyl ghrelin on RER (Figure 4). Again, while the low dose of 0.01 µg did not significantly alter RER on its own, this contrasted with the higher dose of 0.05 µg Ex-4 which was found to attenuate RER in comparison to vehicle control. Finally, as illustrated in Figure 5, PVN co-administration of a low, subthreshold dose of acyl ghrelin paired with a subthreshold dose of the orexigenic molecule NPY, reliably increased RER values. This combined metabolic action was effectively reversed by PVN Ex-4 pretreatment ($F(4,24)=7.35$, $p<0.001$).

Discussion

In the present report we first demonstrated that acyl ghrelin, but not des acyl ghrelin, elicited robust increases in RER when administered into the hypothalamic PVN. The stimulatory effect of acyl ghrelin on RER was blocked by either systemic or PVN treatment with the GLP-1 agonist Ex-4. PVN Ex-4 injection further attenuated the potentiation of RER elicited by co-administration of ghrelin and NPY. Previous reports have found that ghrelin and NPY not only stimulate food intake, but also shift energy substrate utilization in favor of increased carbohydrate oxidation (Currie et al., 2005, 2010, 2011, 2012; Nakazato et al., 2001; Thomas et al., 2016; Wren et al., 2000). We have recently shown that Ex-4 antagonizes the metabolic effects of arcuate ghrelin (Abtahi et al., 2016) suggesting that GLP-1 may play a critical role in the control of metabolic regulation within distinct regions of the mammalian hypothalamus. Specifically, our findings indicate that GLP-1 modulates the appetitive and metabolic actions of ghrelin and NPY via medial hypothalamic neurons localized to the PVN and ArcN in rodents.

Prior evidence suggests that hypothalamic GLP-1 may regulate energy balance within other species as well. For example, ICV GLP-1 treatment decreases food intake in both chicks and catfish (*Ictalurus punctatus*) (Tachibana et al., 2004; Silverstein et al., 2001). Similarly, ICV microinjection of GLP-1 attenuates the orexigenic effect of NPY in neonatal chicks (Furuse et al., 1997) and increases c-fos expression and RER when administered into the ventromedial nucleus (VMN) and lateral hypothalamic (LH) area in chicks (Tachibana et al., 2004, 2007). In addition to suppressing appetite (Tachibana et al., 2004), ICV GLP-1 increases plasma corticosterone and this effect is inhibited by the CRH antagonist astressin (Tachibana et al., 2006). GLP-1 administration into the PVN of rats also elevates plasma corticosterone (Kinzig et al., 2003). Therefore, despite reports of distinct species differences with respect to ghrelin's impact on food intake, ghrelin-GLP-1 neurohormonal interactions appear to be conserved in both avians and mammals, and this may be particularly evident via CRH signaling.

In the mammalian hypothalamus, a critical appetitive circuit incorporates the anorexigenic POMC/CART and orexigenic NPY/AgRP neurons, both of which express GHSR-1a, in the ArcN with projections to the PVN (Kohnno et al., 2008; Guan et al., 2008; van den Top et al., 2007). One report suggests that GLP-1R mRNA is coexpressed with 68% of POMC neurons in the ArcN of rats (Sandoval et al., 2008). The same study reported that GLP-1

administration elicits a decrease in glucose uptake in the ArcN, but not the PVN, and reduces food intake when microinjected into the PVN, but not the ArcN (Sandoval et al., 2008). In addition, ICV Ex-4 treatment has been shown to enhance c-fos expression in ghrelin immunoreactive neurons in the ArcN and PVN (Dalvi et al., 2012). Ventricular Ex-4 has also been reported to decrease plasma ghrelin levels (Perez-Tilve et al., 2007), and in diet-induced obese mice and C57BL/6J mice, third ventricular injection of Ex-4 decreases gastric ghrelin protein expression, hypothalamic GHSR-1a mRNA, GOAT mRNA expression, and circulating ghrelin (Hong et al., 2016). With respect to the appetitive effects, a number of studies suggest that the anorexigenic action of PVN GLP-1 is dependent on glutamatergic signaling (Liu et al., 2017; Mietlicke-Baase et al., 2014). Overall, therefore, emerging evidence now indicates that GLP-1 plays an active role within the ArcN to PVN circuitry, impacting both food intake and metabolic function through its interactions with endogenous ghrelin receptor mechanisms. Specifically, increased ghrelin receptor activation within the ArcN and PVN could in turn promote GLP-1 expression and subsequent receptor stimulation. Elevated GLP-1 signaling could, as a result, suppress not only eating elicited by ghrelin, but also associated metabolic changes, including enhanced carbohydrate oxidation.

Finally, recent research has demonstrated that Ex-4 attenuates ethanol consumption when administered into the nucleus accumbens shell, and inhibits appetite when injected into the accumbens core. Both of these effects are augmented when rats are co-administered the GHSR-1a antagonist, JMV2959 (Abtahi et al., 2017). While ghrelin microinjection into the VTA and NAcc stimulates voluntary ethanol consumption and conditioned place preference (CPP) (Cepko et al., 2014; Jerlhag et al., 2009; Suchankova et al., 2013), in contrast, GLP-1 suppresses ethanol intake and CPP (Shirazi et al., 2013). VTA and accumbal ghrelin injections also increase palatable food consumption (King et al., 2011; Skibicka et al., 2011), and potentiate the stimulatory effect of cocaine on alcohol reward (Cepko et al., 2014), whereas Ex-4 treatment attenuates food intake and operant responding for cocaine (Hernandez et al., 2018; Mietlicki-Baase et al., 2013, 2014; Schmidt et al., 2016).

Given the above work, therefore, it has become increasingly clear that ghrelin and GLP-1 influence both homeostatic and hedonic behavior within both the hypothalamus and the mesolimbic reward pathway. While the present report focused on homeostatic ghrelinergic and GLP-1 signaling within the PVN, future research could be directed at investigating how ghrelin and GLP-1 interact with other endogenous signals. As an example, CRH is well integrated within ghrelin signaling pathways in both avians and mammals, despite the divergent effects on appetite induced by ghrelin in different species (Azzam et al., 2017; Kinzig et al., 2003; Khan et al., 2014; Tachibana et al., 2006). Recently, it has been demonstrated that GOAT KO increases PVN c-fos expression in response to stress (Stark et al., 2016) and GLP-1R knockdown in the PVN reduces anxiety-like behavior (Ghosal et al., 2017), indicating that both ghrelin and GLP-1 impact stress activation via the PVN. Ghrelin similarly activates CRH immunoreactive neurons within the PVN via gamma-aminobutyric acid (GABA) inhibition independently of both NPY and the ArcN, suggesting that PVN ghrelin may be part of a unique CRH circuit (Cabral et al., 2012, 2016). Considering the relationship between ghrelin, stress, and appetite (Brockway et al., 2016; Spencer 2015), it remains possible that the mechanisms through which ghrelin and GLP-1 interact in order to regulate metabolism within the PVN and ArcN could be partially independent. Finally,

chronic peripheral Ex-4 treatment results in decreased hypothalamic NPY expression (Yang et al., 2014), suggesting that the interaction between GLP-1 and NPY should be further examined. These investigations would in turn elucidate how various regulatory peptides may interact in the control of appetitive motivation, metabolism, and glucose homeostasis.

Acknowledgments

Supported by a grant from the M.J. Murdock Charitable Trust (Life Sciences) and the National Institutes of Health (NIH 070496–01A1) to PJC. Additional support was provided by an Esther Hyatt Wender Research Fellowship to SA.

References

- Abtahi S, Howell E, Currie PJ, 2017 Accumbal ghrelin and glucagon-like peptide 1 modulation of ethanol reward and ingestive behavior in female rats. Program No.76.08. 2017 Neuroscience Meeting Planner. Washington, DC: Society for Neuroscience Online. [Presented at the 47th annual meeting of the Society for Neuroscience, Washington, DC, 11–15 November].
- Abtahi S, Mirza A, Howell E, Currie PJ, 2017 Ghrelin enhances food intake and carbohydrate oxidation in a nitric oxide dependent manner. *General and Comparative Endocrinology* 250, 9–14. [PubMed: 28552460]
- Abtahi S, VanderJagt HL, Currie PJ, 2016 The glucagon-like peptide-1 analog exendin-4 antagonizes the effect of acyl ghrelin on the respiratory exchange ratio: *NeuroReport* 27, 992–996. [PubMed: 27454242]
- Alhadeff AL, Rupprecht LE, Hayes MR, 2012 GLP-1 neurons in the nucleus of the solitary tract project directly to the ventral tegmental area and nucleus accumbens to control for food intake. *Endocrinology* 153, 647–658. [PubMed: 22128031]
- Azzam I, Gilad S, Limor R, Stern N, Greenman Y, 2017 Ghrelin stimulation by hypothalamic-pituitary-adrenal axis activation depends on increasing cortisol levels. *Endocr. Connect.* 6, 847–855. [PubMed: 29038331]
- Blanco AM, Bertucci JJ, Valenciano AI, Delgado MJ, Unniappan S, 2017 Ghrelin suppresses cholecystokinin (CCK), peptide YY (PYY) and glucagon-like peptide-1 (GLP-1) in the intestine, and attenuates the anorectic effects of CCK, PYY and GLP-1 in goldfish (*Carassius auratus*). *Horm Behav* 93, 62–71. [PubMed: 28506816]
- Boswell T, Dunn IC, 2017 Regulation of agouti-related protein and pro-opiomelanocortin Gene Expression in the avian arcuate nucleus. *Frontiers in Endocrinology (Lausanne)* 8.
- Bray GA, 1989 Nutrient balance and obesity: An approach to control of food intake in humans. *The Medical clinics of North America*, 73, 29–45. [PubMed: 2562997]
- Brockway ET, Krater KR, Selva JA, Wauson SER, & Currie PJ (2016). Impact of [d-Lys(3)]-GHRP-6 and feeding status on hypothalamic ghrelin-induced stress activation. *Peptides*, 79, 95–102. [PubMed: 27020248]
- Cabral A, Portiansky E, Sánchez-Jaramillo E, Zigman JM, Perello M. Ghrelin activates hypophysiotropic corticotropin-releasing factor neurons independently of the arcuate nucleus. *Psychoneuroendocrinology* 2016 67 27–39. [PubMed: 26874559]
- Cabral A, Suescun O, Zigman JM, Perello M. Ghrelin indirectly activates hypophysiotropic CRF neurons in rodents. *PLoS One*. 2012;7:e31462. [PubMed: 22363652]
- Cepko LCS, Selva JA, Merfeld EB, Fimmel AI, Goldberg SA, Currie PJ, 2014 Ghrelin alters the stimulatory effect of cocaine on ethanol intake following mesolimbic or systemic administration. *Neuropharmacology* 85, 224–231. [PubMed: 24880084]
- Chen L-L, Jiang Q-Y, Zhu X-T, Shu G, Bin Y-F, Wang X-Q, Gao P, Zhang Y-L, 2007 Ghrelin ligand-receptor mRNA expression in hypothalamus, proventriculus and liver of chicken (*Gallus gallus domesticus*): Studies on ontogeny and feeding condition. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 147, 893–902.

- Chen S-R, Chen H, Zhou J-J, Pradhan G, Sun Y, Pan H-L, Li D-P, 2017 Ghrelin receptors mediate ghrelin-induced excitation of agouti-related protein/neuropeptide Y but not pro-opiomelanocortin neurons. *J. Neurochem* 142, 512–520. [PubMed: 28547758]
- Coscina DV, Currie PJ, Szechtman H, 1998 Association of altered whole-body metabolism with locomotor sensitization induced by quinoprole. *Physiology & Behavior*, 63, 755–761. [PubMed: 9617996]
- Currie PJ, Coiro CD, Duenas R, Guss JL, Mirza A, Tal N, 2011 Urocortin I inhibits the effects of ghrelin and neuropeptide Y on feeding and energy substrate utilization. *Brain Res.* 1385, 127–134. [PubMed: 21303672]
- Currie PJ, Khelemsky R, Rigsbee EM, Dono LM, Coiro CD, Chapman CD, Hinchcliff K, 2012 Ghrelin is an orexigenic peptide and elicits anxiety-like behaviors following administration into discrete regions of the hypothalamus. *Behav. Brain Res.* 226, 96–105. [PubMed: 21907737]
- Currie PJ, John CS, Nicholson ML, Chapman CD, Loera KE, 2010 Hypothalamic paraventricular 5-hydroxytryptamine inhibits the effects of ghrelin on eating and energy substrate utilization. *Pharmacol Biochem Behav* 97, 152–155. [PubMed: 20573591]
- Currie PJ, Mirza A, Fuld R, Park D, Vasselli JR, 2005 Ghrelin is an orexigenic and metabolic signaling peptide in the arcuate and paraventricular nuclei. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 289, R353–R358. [PubMed: 15817841]
- Delvi PS, Nazarians-Armavil A, Purser MJ, Belsham DD, 2012 Glucagon-like peptide-1 receptor agonist, exendin-4, regulates feeding-associated neuropeptides in hypothalamic neurons *in Vivo* and *in Vitro*. *Endocrinology* 153, 2208–2222. [PubMed: 22334721]
- Delgado MJ, Cerda-Reverter JM, Soengas JL, 2017 Hypothalamic integration of metabolic, endocrine, and circadian signals in fish: Involvement in the control of food intake. *Frontiers in Neuroscience*. 11, 354. [PubMed: 28694769]
- Delporte C, 2013 Structure and Physiological Actions of Ghrelin. *Scientifica*. 2013, 1–25.
- Fernandez G, Cabral A, Cornejo MP, De Francesco PN, Garcia-Romero G, Reynaldo M, Perello M, 2016 Des-acyl ghrelin directly targets the arcuate nucleus in a ghrelin-receptor independent manner and impairs the orexigenic effect of ghrelin. *Journal of Neuroendocrinology* 28.
- Furuse M, Matsumoto M, Mori R, Sugahara K, Kano K, Hasegawa S, 1997 Influence of fasting and neuropeptide Y on the suppressive food intake induced by intracerebroventricular injection of glucagon-like peptide-1 in the neonatal chick. *Brain Research* 764, 289–292. [PubMed: 9295227]
- Gagnon J, Baggio LL, Drucker DJ, Brubaker PL, 2015 Ghrelin is a novel regulator of GLP-1 secretion. *Diabetes* 64, 1513–1521. [PubMed: 25412624]
- Gao Y, Vidal-Itriago A, Milanova I, Korpel NL, Kalsbeek MJ, Tom RZ, Kalsbeek A, Hofman SM, Yi C-X, 2017 Deficiency of leptin receptor in myeloid cells disrupts hypothalamic metabolic circuits and causes body weight increase. *Molecular Metabolism*. 7, 155–160. [PubMed: 29174000]
- Geelissen SME, Swennen Q, Geyten SV der, Kühn ER, Kaiya H, Kangawa K, Decuypere E, Buyse J, Darras VM, 2006 Peripheral ghrelin reduces food intake and respiratory quotient in chicken. *Domestic Animal Endocrinology* 30, 108–116. [PubMed: 16054797]
- Ghosal S, Packard AEB, Mahbod P, McKlveen JM, Seeley RJ, Myers B, Ulrich-Lai Y, Smith EP, D'Alesso DA & Herman JP (2017). Disruption of Glucagon-Like Peptide 1 Signaling in *Sim1* Neurons Reduces Physiological and Behavioral Reactivity to Acute and Chronic Stress. *The Journal of Neuroscience*, 37(1), 184–193. [PubMed: 28053040]
- Guan J-L, Okuda H, Takenoya F, Kintaka Y, Yagi M, Wang L, Seki M, Hori Y, Kageyama H, Shioda S, 2008 Synaptic relationships between proopiomelanocortin-and ghrelin-containing neurons in the rat arcuate nucleus. *Regulatory Peptides* 145, 128–132. [PubMed: 17936371]
- Gutierrez JA, Solenberg PJ, Perkins DR, Willency JA, Knierman MD, Jin Z, Witcher DR, Luo S, Onyia JE, Hale JE, 2008 Ghrelin octanoylation mediated by an orphan lipid transferase. *Proceedings of the National Academy of Sciences* 105, 6320–6325.
- Hayes MR, 2012 Neuronal and intracellular signaling pathways mediating GLP-1 energy balance and glycemic effects. *Physiology & Behavior* 106, 413–416. [PubMed: 22366059]
- Hayes MR, De Jonghe BC, Kanoski SE, 2010 Role of the glucagon-like-peptide-1 receptor in the control of energy balance. *Physiology & Behavior* 100, 503–510. [PubMed: 20226203]

- Hernandez NS, Ige KY, Mietlicki-Baase EG, Molina-Castro GC, Turner CA, Hayes MR, Schmidt HD, 2018 Glucagon-like peptide-1 receptor activation in the ventral tegmental area attenuates cocaine seeking in rats. *Neuropsychopharmacology*.
- Honda K, 2016 Glucagon-related peptides and the regulation of food intake in chickens. *Anim. Sci. J.* 87, 1090–1098. [PubMed: 27150835]
- Hong X, Zhang H, Liang H, Li D, Huang J, Li Z, Jiang S, Zhang W, Xu G, 2016 Exendin-4 decreases ghrelin levels through mTOR signaling. *Molecular and Cellular Endocrinology* 437, 201–212. [PubMed: 27569528]
- Hosoda H, Kojima M, Mizushima T, Shimizu S, Kangawa K, 2003 Structural divergence of human ghrelin. Identification of multiple ghrelin-derived molecules produced by post-translational processing. *J. Biol. Chem.* 278, 64–70. [PubMed: 12414809]
- Hsu TM, Noble EE, Reiner DJ, Liu CM, Suarez AN, Konanur VR, Hayes MR, Kanoski SE, 2018 Hippocampus ghrelin receptor signaling promotes socially-mediated learned food preference. *Neuropharmacology* 131, 487–496. [PubMed: 29191751]
- Inoue Y, Hayashi Y, Kangawa K, Suzuki Y, Murakami N, Nakahara K, 2016 Des-acyl ghrelin prevents heatstroke-like symptoms in rats exposed to high temperature and high humidity. *Neuroscience Letters* 615, 28–32. [PubMed: 26773867]
- Jerlhag E, Egencioglu E, Landgren S, Salomé N, Heilig M, Moechars D, Datta R, Perrissoud D, Dickson SL, Engel JA, 2009 Requirement of central ghrelin signaling for alcohol reward. *PNAS* 106, 11318–11323. [PubMed: 19564604]
- Kanoski SE, Fortin SM, Arnold M, Grill HJ, Hayes MR, 2011 Peripheral and central GLP-1 receptor populations mediate the anorectic effects of peripherally administered GLP-1 receptor agonists, liraglutide and exendin-4. *Endocrinology* 152, 3103–3112. [PubMed: 21693680]
- Khan MSI, Kaiya H, Tachibana T, 2014 Central injection of urocortin-3 but not corticotrophin-releasing hormone influences the ghrelin/GHS-R1a system of the proventriculus and brain in chicks. *Domest. Anim. Endocrinol.* 47, 27–34. [PubMed: 24484650]
- King SJ, Isaacs AM, O'Farrell E, Abizaid A, 2011 Motivation to obtain preferred foods is enhanced by ghrelin in the ventral tegmental area. *Hormones and Behavior* 60, 572–580. [PubMed: 21872601]
- Kinzig KP, D'Alessio DA, Herman JP, Sakai RR, Vahl TP, Figueredo HF, Murphy EK, Seeley RJ, 2003 CNS glucagon-like peptide-1 receptors mediate endocrine and anxiety responses to interoceptive and psychogenic stressors. *J. Neurosci.* 23, 6163–6170. [PubMed: 12867498]
- Kohno D, Gao H-Z, Muroya S, Kikuyama S, Yada T, 2003 Ghrelin directly interacts with neuropeptide-y-containing neurons in the rat arcuate nucleus Ca^{2+} signaling via protein kinase a and n-type channel-dependent mechanisms and cross-talk with leptin and orexin. *Diabetes* 52, 948–956. [PubMed: 12663466]
- Kleiber M The fire of life: An introduction to animal energetics. Malabar, FL: R.E. Krieger; 1987.
- Kohno D, Sone H, Minokoshi Y, Yada T, 2008 Ghrelin raises $[Ca^{2+}]_i$ via AMPK in hypothalamic arcuate nucleus NPY neurons. *Biochem. Biophys. Res. Commun.* 366, 388–392. [PubMed: 18068666]
- Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K, 1999 Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 402, 656–660. [PubMed: 10604470]
- López-Ferreras L, Richard JE, Anderberg RH, Nilsson FH, Olandersson K, Kanoski SE, Skibicka KP, 2017 Ghrelin's control of food reward and body weight in the lateral hypothalamic area is sexually dimorphic. *Physiol. Behav.* 176, 40–49. [PubMed: 28213203]
- Mahbod P, Smith EP, Fitzgerald ME, Morano RL, Packard BA, Ghosal S, Scheimann JR, Perez-Tilve D, Herman JP, Tong J, 2018 Desacyl Ghrelin Decreases Anxiety-like Behavior in Male Mice. *Endocrinology* 159, 388–399. [PubMed: 29155981]
- Matsuda K, Miura T, Kaiya H, Maruyama K, Himakura S-I, Uchiyama M, Kangawa K, Shioda S, 2006 Regulation of food intake by acyl and des-acyl ghrelins in the goldfish. *Peptides* 27, 2321–2325. [PubMed: 16687192]
- Mietlicki-Baase EG, Ortinski PI, Reiner DJ, Sinon CG, McCutcheon JE, Pierce RC, Roitman MF, Hayes MR, 2014 Glucagon-like peptide-1 Receptor Activation in the nucleus accumbens core suppresses feeding by increasing glutamatergic AMPA/kainate signaling. *Journal of Neuroscience* 34, 6985–6992. [PubMed: 24828651]

- Mietlicki-Baase EG, Ortinski PI, Rupprecht LE, Olivos DR, Alhadeff AL, Pierce RC, Hayes MR, 2013 The food intake-suppressive effects of glucagon-like peptide-1 receptor signaling in the ventral tegmental area are mediated by AMPA/kainate receptors. *AJP: Endocrinology and Metabolism* 305, E1367–E1374.
- Mohan H, Unniappan S, 2013 Discovery of Ghrelin O-Acyltransferase 25, 16–24.
- Nakazato M, Murakami N, Date Y, Kojima M, Matsuo H, Kangawa K, Matsukura S, 2001 A role for ghrelin in the central regulation of feeding : Article : Nature. *Nature* 409, 194–198. [PubMed: 11196643]
- Naleid AM, Grace MK, Cummings DE, Levine AS, 2005 Ghrelin induces feeding in the mesolimbic reward pathway between the ventral tegmental area and the nucleus accumbens. *Peptides* 26, 2274–2279. [PubMed: 16137788]
- Oclon E, Pietras M, 2011 Peripheral ghrelin inhibits feed intake through hypothalamo-pituitary-adrenal axis-dependent mechanism in chicken. *J. Anim. Feed Sci.* 20, 118–130.
- Paxinos G, Watson C, 2014 The rat brain in stereotaxic coordinates, 7th ed Academic Press, New York, NY.
- Perez-Tilve D, Gonzalez-Matias L, Alvarez-Crespo M, Leiras R, Tovar S, Dieguez C, Mallo F, 2007 Exendin-4 potentially decreases ghrelin levels in fasting rats. *Diabetes* 56, 143–151. [PubMed: 17192476]
- Reiner DJ, Leon RM, McGrath LE, Koch-Laskowski K, Hahn JD, Kanoski SE, Mietlicki-Baase EG, Hayes MR, 2018 Glucagon-like peptide-1 receptor signaling in the lateral dorsal tegmental nucleus regulates energy balance. *Neuropsychopharmacology* 43, 627–637. [PubMed: 28920591]
- Romero A, Kirchner H, Heppner K, Pfluger PT, Tschop MH, Nogueiras R, 2010 GOAT: the master switch for the ghrelin system? *European Journal of Endocrinology* 163, 1–8. [PubMed: 20421334]
- Saito ES, Kaiya H, Tachibana T, Tomonaga S, Denbow DM, Kangawa K, Furuse M, 2005 Inhibitory effect of ghrelin on food intake is mediated by the corticotropin-releasing factor system in neonatal chicks. *Regul. Pept.* 125, 201–208. [PubMed: 15582733]
- Sanchez-Bretano A, Blanco AM, Unniappan S, Kah O, Gueguen MM, Bertucci JI, Alonso-Gomez AL, Valenciano AI, Isorna E, Delgado MJ, 2015 In situ localization and rhythmic expression of ghrelin and ghs-r1 ghrelin receptor in the brain and gastrointestinal tract of goldfish (*Carassius auratus*). *PLoS One* 10, e0141043. [PubMed: 26506093]
- Sandoval DA, Bagnol D, Woods SC, D'Alessio DA, Seeley RJ, 2008 Arcuate glucagon-like peptide 1 receptors regulate glucose homeostasis but not food intake. *Diabetes* 57, 2046–2054. [PubMed: 18487451]
- Schmidt HD, Mietlicki-Baase EG, Ige KY, Maurer JJ, Reiner DJ, Zimmer DJ, Van Nest DS, Guercio LA, Wimmer ME, Olivos DR, De Jonghe BC, Hayes MR, 2016 Glucagon-like peptide-1 receptor activation in the ventral tegmental area decreases the reinforcing efficacy of cocaine. *Neuropsychopharmacology.* 41, 1917–1928. [PubMed: 26675243]
- Shirazi RH, Dickson SL, Skibicka KP, 2013 Gut peptide GLP-1 and its analogue, exendin-4, decrease alcohol intake and reward. *PLoS One* 8, e61965. [PubMed: 23613987]
- Shousha S, Nakahara K, Kojima M, Miyazato M, Hosoda H, Kangawa K, Murakami N, 2005 Different effects of peripheral and central ghrelin on regulation of food intake in the Japanese quail. *General and Comparative Endocrinology* 141, 178–183. [PubMed: 15748719]
- Silverstein JT, Bondareva VM, Leonard JBK, Plisetskaya EM, 2001 Neuropeptide regulation of feeding in catfish, *Ictalurus punctatus*: a role for glucagon-like peptide-1 (GLP-1)? *Comp. Biochem. Physiol. B-Biochem. Mol. Biol.* 129, 623–631. [PubMed: 11399498]
- Skibicka KP, Hansson C, Alvarez-Crespo M, Friberg PA, Dickson SL, 2011 Ghrelin directly targets the ventral tegmental area to increase food motivation. *Neuroscience* 180, 129–137. [PubMed: 21335062]
- Spencer SJ, Emmerzaal TL, Kozicz T, Andrews ZB Ghrelin's role in the hypothalamic-pituitary-adrenal axis stress response: Implications for mood disorders. *Biol. Psychiatry.* (2015);78:19–27. [PubMed: 25534754]
- Suchankova P, Steensland P, Fredriksson I, Engel JA, Jerlhag E, 2013 Ghrelin receptor (GHS-R1A) antagonism suppresses both alcohol consumption and the alcohol deprivation effect in rats following long-term voluntary alcohol consumption. *PLoS ONE* 8, e71284. [PubMed: 23977009]

- Tachibana T, Hirofujii K, Matsumoto M, Furuse M, Hasegawa S, Yoshizawa F, Sugahara K, 2004 The hypothalamus is involved in the anorexic effect of glucagon-like peptide-1 in chicks. *Comp. Biochem. Physiol. A-Mol. Integr. Physiol.* 137, 183–188. [PubMed: 14720603]
- Tachibana T, Oikawa D, Adachi N, Boswell T, Furuse M, 2007 Intracerebroventricular injection of glucagon-like peptide-1 changes lipid metabolism in chicks. *Comp. Biochem. Physiol. A-Mol. Integr. Physiol.* 147, 1104–1108. [PubMed: 17466552]
- Tachibana T, Sato M, Oikawa D, Furuse M, 2006 Involvement of CRF on the anorexic effect of GLP-1 in layer chicks. *Comp. Biochem. Physiol. A-Mol. Integr. Physiol.* 143, 112–117. [PubMed: 16377222]
- Thomas MA, Ryu V, Bartness TJ, 2016 Central ghrelin increases food foraging/hoarding that is blocked by GHSR antagonism and attenuates hypothalamic paraventricular nucleus neuronal activation. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* 310, R275–R285.
- Tschöp M, Smiley DL, Heiman ML 2000 Ghrelin induces adiposity in rodents. *Nature* 407, 908–913. [PubMed: 11057670]
- van den Top M, Lyons DJ, Lee K, Coderre E, Renaud LP, Spanswick D, 2007 Pharmacological and molecular characterization of ATP-sensitive K⁺ conductances in CART and NPY/AgRP expressing neurons of the hypothalamic arcuate nucleus. *Neuroscience* 144, 815–824. [PubMed: 17137725]
- Vasselli JR, Pi-Sunyer FX, Wall DG, John CS, Chapman CD, & Currie PJ, 2017 Central effects of insulin detemir on feeding, body weight, and metabolism in rats. *American Journal of Physiology. Endocrinology and Metabolism* 313, E613–E621. [PubMed: 28720583]
- Velasco C, Libran-Perez M, Otero-Rodino C, Lopez-Patino MA, Miguez JM, Cerda-Reverter JM, Soengas JL, 2015 Ghrelin modulates hypothalamic fatty acid-sensing and control of food intake in rainbow trout. *Journal of Endocrinology* 228, 25–37. [PubMed: 26459641]
- Weinberg ZY, Nicholson ML, Currie PJ, 2011 6-Hydroxydopamine lesions of the ventral tegmental area suppress ghrelin's ability to elicit food-reinforced behavior. *Neuroscience Letters* 499, 70–73. [PubMed: 21640160]
- Wren AM, Small CJ, Ward HL, Murphy KG, Dakin CL, Taheri S, Kennedy AR, Roberts GH, Morgan DG, Ghatei MA, Bloom SR, 2000 The novel hypothalamic peptide ghrelin stimulates food intake and growth hormone secretion. *Endocrinology* 141, 4325–4328. [PubMed: 11089570]
- Xu G, Hong X, Tang H, Jiang S, Liu F, Shen Z, Li Z, Zhang W, 2015 Ghrelin regulates GLP-1 production through mTOR signaling in L cells. *Molecular and Cellular Endocrinology* 416, 9–18. [PubMed: 26279396]
- Yang J, Brown MS, Liang G, Grishin NV, Goldstein JL 2008 Identification of the acyltransferase that octanoylates ghrelin, an appetite-stimulating peptide hormone. *Cell* 132, 387–396 [PubMed: 18267071]
- Yang Y, Moghadam AA, Corder ZA, Liang N-C, Moran TH, 2014 Long Term Exendin-4 Treatment Reduces Food Intake and Body Weight and Alters Expression of Brain Homeostatic and Reward Markers. *Endocrinology* 155, 3473–3483. [PubMed: 24949661]
- Zigman JM, Jones JE, Lee CE, Saper CB, Elmquist JK, 2006 Expression of ghrelin receptor mRNA in the rat and the mouse brain. *J Comp Neurol* 494, 528–548. [PubMed: 16320257]

Highlights:

- Peripheral and PVN Ex-4 attenuates ghrelin induced increases in the RER
- Ex-4 reverses the effect of subthreshold ghrelin paired with subthreshold NPY on RER
- Des-acyl ghrelin did not alter the respiratory exchange ratio

Fig. 1. Coronal section through the PVN of the hypothalamus illustrating the site of microinjection. One arrow targets blank ink residue within PVN, while the second arrow indicates PVN tissue as referenced by Paxinos & Watson (2014). Sections were cut at 20 μm , stained with Cresyl violet, and examined via light microscopy.

Fig. 2.

The effect of acyl (n=8) and des-acyl (n=7) ghrelin microinjection on RER. Only PVN acyl ghrelin elicited significant increases in RER indicative of enhanced carbohydrate utilization and reduced lipid oxidation. PVN des-acyl ghrelin treatment was ineffective in altering RER. Values represent mean $\pm$ S.E.M.; *P<0.05 compared to vehicle (Veh).

Fig. 3.

Intraperitoneal injections of the GLP-1 agonist Ex-4 reduced RER and attenuated the stimulatory effect of ghrelin on substrate oxidation. Values are mean $\pm$ S.E.M.; $n=8$. * $P < 0.05$ compared to vehicle (Veh). ** $P < 0.05$ compared to ghrelin paired with Veh.

Fig. 4. PVN treatment of Ex-4 decreased RER and attenuated the stimulatory effect of ghrelin on substrate oxidation. Values represent mean $\pm$ S.E.M.; $n=8$. * $P<0.05$ compared to vehicle (Veh). ** $P<0.05$ compared to ghrelin paired with Veh.

Fig. 5. PVN co-administration of subthreshold doses of ghrelin (10 pmol) and NPY (15 pmol) elicited significant increases in RER, an effect reversed by pretreatment with PVN Ex-4 (0.01 μ g). Values represent mean $\pm$ S.E.M.; n=6. *P<0.05 compared vehicle (Veh).