- the periphery on fat tissue and mediates stress-induced obesity and metabolic syndrome. *Nat Med* 2007;**13**:803–811.
22. Medzikovic L, Schumacher CA, Verkerk AO, van Deel ED, Wolswinkel R, van der Made I, Bleeker N, Kakici D, van den Hoogenhof MMG, Meggouh F, Creemers EE, Ann Remme C, Baartscheer A, de Winter RJ, de Vries CJM, de Arkenbout EK, de Waard V. Orphan nuclear receptor Nur77 affects cardiomyocyte calcium homeostasis and adverse cardiac remodelling. *Sci Rep* 2015;**5**:15404.
 23. Kurakula K, Wal E Van Der, Geerts D, Tiel CM Van, Vries CJM De. FHL2 protein is a novel co-repressor of nuclear receptor Nur77. *J Biol Chem* 2011;**286**:44336–44343.
 24. Sekiya T, Kashiwagi I, Yoshida R, Fukaya T, Morita R, Kimura A, Ichinose H, Metzger D, Chambon P, Yoshimura A. Nr4a receptors are essential for thymic regulatory T cell development and immune homeostasis. *Nat Immunol* 2013;**14**:230–237.
 25. Baartscheer A, Schumacher CA, Fiolet JW. SR calcium depletion following reversal of the Na⁺/Ca²⁺-exchanger in rat ventricular myocytes. *J Mol Cell Cardiol* 2000;**32**:1025–1037.
 26. Wieland HA, Engel W, Eberlein W, Rudolf K, Doods HN. Subtype selectivity of the novel nonpeptide neuropeptide Y Y1 receptor antagonist BIBO 3304 and its effect on feeding in rodents. *Br J Pharmacol* 1998;**125**:549–555.
 27. Pellieux C, Sauthier T, Domenighetti A, Marsh DJ, Palmiter RD, Brunner HR, Pedrazzini T. Neuropeptide Y (NPY) potentiates phenylephrine-induced mitogen-activated protein kinase activation in primary cardiomyocytes via NPY Y5 receptors. *Proc Natl Acad Sci USA* 2000;**97**:1595–1600.
 28. Dolmetsch RE, Lewis RS, Goodnow CC, Healy JL. Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration. *Nature* 1997;**386**:855–858.
 29. Gradin KA, Li J-Y, Andersson O, Simonsen U. Enhanced neuropeptide Y immunoreactivity and vasoconstriction in mesenteric small arteries from spontaneously hypertensive rats. *J Vasc Res* 2003;**40**:252–265.
 30. Wang R-H, He J-P, Su M-L, Luo J, Xu M, Du X-D, Chen H-Z, Wang W-J, Wang Y, Zhang N, Zhao B-X, Zhao W-X, Shan Z-G, Han J, Chang C, Wu Q. The orphan receptor TR3 participates in angiotensin II-induced cardiac hypertrophy by controlling mTOR signalling. *EMBO Mol Med* 2013;**5**:137–148.
 31. Cui M, Cai Z, Chu S, Sun Z, Wang X, Hu L, Yi J, Shen L, He B. Orphan nuclear receptor Nur77 inhibits angiotensin II-induced vascular remodeling via downregulation of beta-catenin. *Hypertens (Dallas, TX 1979)* 2016;**67**:153–162.
 32. Hirsch D, Zukowska Z. NPY and stress 30 years later: the peripheral view. *Cell Mol Neurobiol* 2012;**32**:645–659.
 33. Zukowska-Grojec Z, Konarska M, McCarty R. Differential plasma catecholamine and neuropeptide Y responses to acute stress in rats. *Life Sci* 1988;**42**:1615–1624.
 34. Myers S, Eriksson N, Burow R, Wang S-CM, Muscat GEO. Beta-adrenergic signaling regulates NR4A nuclear receptor and metabolic gene expression in multiple tissues. *Mol Cell Endocrinol* 2009;**309**:101–108.
 35. Hilgendorf I, Gerhardt LMS, Tan TC, Winter C, Holderried TAW, Chousterman BG, Iwamoto Y, Liao R, Zirlik A, Scherer-Crosbie M, Hedrick CC, Libby P, Nahrendorf M, Weissleder R, Swirski FK. Ly-6Chigh monocytes depend on Nr4a1 to balance both inflammatory and reparative phases in the infarcted myocardium. *Circ Res* 2014;**114**:1611–1622.
 36. Yan G, Zhu N, Huang S, Yi B, Shang X, Chen M, Wang N, Zhang G-X, Talarico JA, Tilley DG, Gao E, Sun J. Orphan nuclear receptor Nur77 inhibits cardiac hypertrophic response to beta-adrenergic stimulation. *Mol Cell Biol* 2015;**35**:3312–3323.
 37. Gicquiaux H, Lecat S, Gaire M, Dieterlen A, Mély Y, Takeda K, Bucher B, Galzi JL. Rapid internalization and recycling of the human neuropeptide Y Y1 receptor. *J Biol Chem* 2002;**277**:6645–6655.
 38. Lehmann LH, Jebessa ZH, Kreusser MM, Horsch A, He T, Kronlage M, Dewenter M, Sramek V, Oehl U, Krebs-Haupenthal J, von der Lieth AH, Schmidt A, Sun Q, Ritterhoff J, Finke D, Völkers M, Jungmann A, Sauer SW, Thiel C, Nickel A, Kohlhaas M, Schäfer M, Sticht C, Maack C, Gretz N, Wagner M, El-Armouche A, Maier LS, Londoño JEC, Meder B, Freichel M, Gröne H-J, Most P, Müller OJ, Herzig S, Furlong EEM, Katus HA, Backs J. A proteolytic fragment of histone deacetylase 4 protects the heart from failure by regulating the hexosamine biosynthetic pathway. *Nat Med* 2017;**24**:62–72.
 39. Liu JJ, Shi SG, Han QD. Evaluation of plasma neuropeptide Y levels in patients with congestive heart failure. *Zhonghua Nei Ke Za Zhi* 1994;**33**:687–689.
 40. Cuculi F, Herring N, Caterina AR, De Banning AP, Prendergast BD, Forfar JC, Choudhury RP, Channon KM, Kharbanda RK. Relationship of plasma neuropeptide Y with angiographic, electrocardiographic and coronary physiology indices of reperfusion during ST elevation myocardial infarction. *Heart* 2013;**99**:1198–1203.
 41. Kurakula K, Vos M, Logiantara A, Roelofs JJ, Nieuwenhuis MA, Koppelman GH, Postma DS, Rijt LS, van de Vries CJM. Nuclear receptor Nur77 attenuates airway inflammation in mice by suppressing NF-κB activity in lung epithelial cells. *J Immunol* 2015;**195**:1388–1398.
 42. Li S, Koziol-White C, Jude J, Jiang M, Zhao H, Cao G, Yoo E, Jester W, Morley MP, Zhou S, Wang Y, Lu MM, Panettieri RA, Morrissey EE. Epithelium-generated neuropeptide y induces smooth muscle contraction to promote airway hyperresponsiveness. *J Clin Invest* 2016;**126**:1978–1982.
 43. Cardell LO, Uddman R, Edvinsson L. Low plasma concentrations of VIP and elevated levels of other neuropeptides during exacerbations of asthma. *Eur Respir J* 1994;**7**:2169–2173.
 44. Duren RP, Boudreaux SP, Conneely OM. Genome wide mapping of NR4A binding reveals cooperativity with ETS factors to promote epigenetic activation of distal enhancers in acute myeloid leukemia cells. *PLoS One* 2016;**11**:e0150450–e0150420.

Corrigendum

doi:10.1093/cvr/cvy194
Online publish-ahead-of-print 20 August 2018

Corrigendum to: Loss of AZIN2 splice variant facilitates endogenous cardiac regeneration [*Cardiovasc Res* 2018;**114**:1642–1655]

The authors of the above paper wish to inform readers that they did not disclose to the Journal at a pre-publication stage that they had not secured the permission to use the abbreviation “ROCR” from the HUGO Gene Nomenclature Committee (HGNC). In cooperation with the HGNC, it was decided to remove the use of “ROCR” and replace with “non-coding splice variant of AZIN2” to avoid confusion in the field. The title and content of the article has been edited to reflect this change.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2018. For permissions, please email: journals.permissions@oup.com.