Published in final edited form as:

Chemistry. 2015 July 20; 21(30): 10681–10686. doi:10.1002/chem.201502132.

Total Synthesis and Structural Reassignment of Lyngbyaloside C Highlighted by Intermolecular Ketene Esterification

Chia-Fu Chang, Eric Stefan, Richard E. Taylor^a

^[a]Warren Family Research Center for Drug Discovery and Development, and the Department of Chemistry & Biochemistry, 250 Nieuwland Science Hall, University of Notre Dame, Notre Dame, IN 46556-5670 (USA)

Abstract

Lyngbyaloside C, a classic macrolide, isolated from *Lyngbya bouillonii*, has shown moderate anticancer activity against several cancer cell lines. Here, we report the first total synthesis and stereochemical configuration reassignment of lyngbyaloside C. The synthesis highlights a one-pot intermolecular ketene esterification reaction to form the crucial tertiary ester and tetrahydropyran. In addition, a novel and concise synthetic pathway towards the 1,3-*syn* secondary, tertiary diol fragment is described using a regio- and stereospecific electrophilic ether transfer reaction.

Graphical Abstract

taylor.61@nd.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/chem.201502132>.

Reassigned: The total synthesis of the nominal structure for the polyketide natural product lyngbyalloside C has been reported along with a correction of its originally assigned stereochemistry. Three diastereomerisomers of the compound were prepared using a highly convergent strategy.

Keywords

electrophilic ether transfer; ketene; lyngbyalloside C; natural products; polyketides; total synthesis

Naturally occurring, biologically active molecules that have evolved structurally over centuries, have long served as useful therapeutics agents.^[1] Polyketides, a unique class of natural products, possess a broad range of privileged scaffolds, structural diversity, and biological activities. Since 2007, a series of polyketide natural products have been isolated from the marine cyanobacteria,^[2] *Lyngbya bouillonii*, including lyngbyalloside C (*Z/E*) 1, lyngbouilloside, and lyngbyalloside B (Figure 1).^[3] These structures are characterized as 14-membered glycosylated macrolides, embedded with a 2,6-*cis* tetrahydropyran ring and an appended diene side chain. Notably, the tertiary ester moiety is not only interesting from a biosynthetic standpoint, but it also poses a unique synthetic challenge. Preliminary biological studies show that the lyngbyallosides exhibit moderate cytotoxicity against several cancer cell lines.^[3] As a result, the lyngbyallosides have garnered recent interest from the synthetic community.^[4] These efforts culminated in the first total synthesis of lyngbyalloside B by Fuwa et al. in 2014, in which the stereochemical configuration of lyngbyalloside B was reassigned.^[4f] However, previous synthetic endeavors all focused on the lyngbyallosides

which possess the (*E,E*) diene appendage (lyngbouilloside and lyngbyaloside B). Lyngbyaloside B also contains an additional methyl stereogenic center at C6. With that in mind, we also envisioned an intermolecular ketene esterification reaction that could generate the challenging tertiary ester from readily accessible fragments. Accordingly, lyngbyaloside C is the perfect target for the development of a general pathway to both isomers via intermolecular ketene esterification. Furthermore, the total synthesis of lyngbyaloside C would allow the confirmation of its stereochemical configuration assignment. Herein, we report the first total synthesis of lyngbyaloside C which warrants a structural revision, while providing a significant amount of material including two diastereoisomers, for more thorough biological investigation.

Our synthetic plan is outlined in Figure 2. We conceived that selective glycosylation of the C5 hydroxyl would allow late-stage installation of the sugar moiety. The precursory aglycone would be constructed via a highly convergent, sequential ketene esterification and ring-closing metathesis (RCM). Accordingly, the aglycone was broken into tertiary alcohol and ketene precursor fragments. Yadav et al. recently reported a similar strategy towards the synthesis of lyngbyaloside B.^[4d] For preparation of the tertiary alcohol, we envisioned the installation of the 1,3-*syn* diol through utilization of an electrophilic-induced ether transfer methodology recently developed in our labs.^[5] The alkene required for the subsequent RCM step provided us with an opportunity to explore a unique ether transfer substrate which contains potentially competing 1,1-disubstituted and terminal alkenes.

Synthesis of the ether transfer substrate commenced with the application of a Leighton crotylation^[6] of known aldehyde **2** (Scheme 1). Sequential protection of the resulting alcohol followed by a Wittig olefination furnished homoallylic alkoxy ether **6**. Gratifyingly, the ether transfer reaction, carried out using iodine monochloride (ICl, -95 °C), produced the desired 1,3-*syn* diol monoether **7** in 76% yield with reasonable diastereoselectivity (10:1 *syn:anti*). Interestingly, no reactivity was observed at the monosubstituted olefin, representing the first report of a regioselective ether transfer in the presence of two reactive olefins. The resulting diol monoether **7** was then converted to epoxide **8** via the three-step functional group manipulation, followed by ring opening with the corresponding Grignard reagent in the presence of copper(I) iodide to afford tertiary alcohol **9a**.

With tertiary alcohol **9a** in hand, we then turned our focus towards the synthesis of the ketene precursor fragment. The application of ketene chemistry for the synthesis of sterically encumbered esters has been investigated.^[7] On the basis of these prior studies, several ketene precursors, such as carboxylic acid **I**,^[7a-f] acid halides **II**, and alkynyl ether **IV**^[7g-I] were explored (Figure 3). Unfortunately, none of these ketene precursors proved to be efficient for generating the desired tertiary ester moiety. We believe that this is the result of the instability of the ketene intermediates and a preference for degradative over productive pathways. However, β -keto esters **V** have been shown to generate stabilized β -keto ketenes in the context of macrolactonization reactions.^[4d] In our preliminary investigation of these precursors, low yields were obtained when we tried to scale up (~ 100 mg), which was presumably caused by the competing ketene polymerization.^[7g] To address this issue, we explored the use of a thioester substrate **VI** activated by a thiophilic metal salt,^[8] allowing the corresponding ketene to be generated under comparatively mild conditions.

Synthesis of β -keto ester fragment **16** started from the Mukaiyama aldol reaction of dienol ether **10** with acrolein,^[9] followed by Sharpless kinetic resolution,^[10] to afford chiral, non-racemic allylic alcohol **12**. Desired ester **13** was accessed by trapping the acetylketene intermediate of **12** with methanol. Utilization of intramolecular hydride delivery^[11] provided the *1,3-anti* diol, which was subsequently silylated to afford ester **14**. Reduction of the aforementioned ester with DiBALH gave the corresponding aldehyde, which was then treated with *tert*-butylthiol diazoacetate **15**^[12] in the presence of tin(II) chloride to successfully afford β -keto- thioester **16** (Scheme 2).

With the two requisite building blocks of **1** in hand, we turned our attention to developing a robust and scalable ketene esterification reaction. After screening numerous conditions, silver trifluoroacetate was found to affect gram-scale one-pot intermolecular ketene esterification in which the tertiary ester and pyran are formed consecutively to afford **17** (Scheme 3). The formation of the methyl ketal was achieved by re-fluxing with dry MeOH in the presence of citric acid, and subsequent protection of the secondary alcohol provided diene **18a**. The ensuing RCM proceeded smoothly upon treatment with Hoveyda-Grubbs 2nd generation catalyst^[13] in the presence of benzoquinone^[14] to yield **19a** exclusively as the *E* isomer. Regioselective hydrogenation of the disubstituted olefin and deprotection of the benzyl ether was accomplished by using activated Raney nickel, which was followed by Grieco olefination conditions^[15] to provide **20a**. At this point, we attempted to prepare the diene side chain through the method of Fuwa et al.^[4f] Alkene **20a** was converted to the aldehyde via two-step oxidation cleavage. Unfortunately, Takai olefination^[16] conditions yielded vinyl iodide **21** but also resulted in an unexpected double elimination, a problem that could not be resolved.

The unexpected elimination process was successfully circumvented by use of a cross-metathesis homologation and a Wittig olefination. Deprotection of the PMB ether followed by treatment of the resultant free alcohol with the corresponding glycosylated trichloroimidate **23**, in the presence of catalytic La(OTf)₃^[17] yielded the desired glycosylated macrolide **24a** as a single β -isomer. Cross-metathesis of the terminal alkene with crotonaldehyde afforded **25a** as a single *E* isomer. The installation of bromodiene was completed by using Menche's optimized Wittig reaction conditions,^[18] which successfully inhibited the formation of the dibromide side product.^[18] Finally, global deprotection with hydrogen fluoride furnished nominal lyngbyaloside C *Z/E*-**1a** (*Z/E* 4.3/1) (Scheme 4). During preliminary studies on compounds earlier in the sequence, we found that *Z/E* mixtures of related bromodienes could be converted to exclusively *Z* by exposure to catalytic quantities of DDQ. Much to our delight, DDQ treatment of a *Z/E* mixture of **1a** provided isomerically pure *Z*-**1a** in 70% yield. At this point, the mechanism of the reaction remains unclear and the reaction may, in fact, involve selective degradation of the *E*-bromo-diene.

Comparing the ¹H and ¹³C NMR data for the synthetic nominal compound **1a** to those in the literature for natural lyngbyaloside C, we found there were appreciable disparities between them (Figure 4). In the ¹H spectrum, the doublet at δ = 2.80 ppm and the singlet at δ = 4.57 ppm are missing in the synthetic compound. In addition, several signals in both ¹H and ¹³C show broadening or are absent.^[19] These substantial disparities indicate one or more of the stereochemical configurations of natural lyngbyaloside C and nominal structure **1a** are

different. During our studies, Fuwa et al. reported the synthesis of nominal lyngbyaloside B, which resulted in the structural configuration reassignment at three stereogenic centers (C₁₀, C₁₁, C₁₃).^[4f] Therefore, we surmised the spectral disagreements in the current effort resulted from a misassignment of lyngbyaloside C at these same positions.

Based on NOE studies of the natural product, we considered two potential diastereomers as the likely stereochemical configuration for natural lyngbyaloside C. To access both tertiary alcohol fragments **9 b** and **9 c**, divergent routes from a common intermediate were developed highlighted by stereochemically complementary Ti-mediated aldol reactions (Scheme 5).^[20,21] Gratifyingly, fully diastereomeric targets *Z/E*-**1b** and *Z/E*-**1c** were then generated efficiently via the optimized synthetic pathway developed for the synthesis of *Z/E*-**1a** through ketene fragment **16**.^[22] Upon a comparison of NMR data, **1b** encountered similar irregularities with the originally proposed structure **1a**. In contrast, characterization data for **1c** was consistent with those of the natural product, indicating the stereochemical configuration of lyngbyaloside C is unambiguously assigned as structure **1c** (Figure 5).^[23] Furthermore, lyngbyaloside C shares the same absolute stereochemistry with lyngbyaloside B, and we propose that lyngbouilloside can be reassigned accordingly.

In summary, we have completed the first total synthesis of lyngbyaloside C **1c**, which has confirmed the need for reassignment of the stereochemical configuration of the natural product. Our synthetic pathway featured an intermolecular thioester transesterification through a ketene intermediate. In addition, we also developed a concise approach toward key tertiary alcohol **12a** by a regio-and-stereoselective electrophilic ether transfer reaction on a unique substrate. Lastly, in synthesizing the diene bromide appendage, we determined that a catalytic DDQ-mediated isomerization could be successfully applied at a late-stage to access exclusively the *Z*-bromodiene for each of the three stereoisomeric compounds. The mechanism of this potentially useful methodology along with the biological evaluation of the isomeric macrolides are currently topics of investigation in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Support from the NIH and the National Institute for General Medical Sciences is gratefully acknowledged (GM084922). C.-F. Chang would like to acknowledge the generous support and mentorship of Dr. Yean-Jang “Super” Lee.

References

- [1]. a)Firn RD, Jones CG, Nat. Prod. Rep. 2003, 20, 382; [PubMed: 12964834] b)William PG, Trends Biotechnol. 2008, 26, 45;c)Villa FA, Gerwick L, Immunopharm. Immunot. 2010, 32, 228;d)Mayer AMS, Glaser KB, Cuevas G, Jacobs RS, Kem W, Little RD, McIntosh JM, Newman DJ, Potts BC, Shusters DE, Trends Pharmacol. Sci. 2010, 31, 255; [PubMed: 20363514] e)Gerwick WH, Moore BS, Chem. Biol. 2012, 19, 85; [PubMed: 22284357] f)Newman DJ, Cragg GM. , J. Nat. Prod. 2012, 75, 311; [PubMed: 22316239] g)Gerwick WH, Fenner AM, Microb. Ecol. 2013, 65, 800. [PubMed: 23274881]

- [2]. a)Tan LT, Phytochemistry 2007, 68, 954; [PubMed: 17336349] b)Nunnery JK, Mevers E, Gerwick WH, Curr. Opin. Biotechnol. 2010, 21, 787. [PubMed: 21030245]
- [3]. a)Tan LT, Márquez BL, Gerwick WH, J. Nat. Prod. 2002, 65, 925; [PubMed: 12088441] b)Luesch H, Yoshida WY, Harrigan GG, Doom JP, Moore RE, Paul VJ, J. Nat. Prod. 2002, 65, 1945; [PubMed: 12502348] c)Matthew S, Salvador LA, Schupp PJ, Paul VJ, Luesch H, J. Nat. Prod. 2010, 73, 1544. [PubMed: 20704304]
- [4]. a)Methodologies toward pyran-embedded 14-members macrolactone:Hoye TR, Danielson ME, May AE, Zhao H, Angew. Chem. Int. Ed. 2008, 47, 9743; Angew. Chem. 2008, 120, 9889;b)Bai Y, Davis DC, Dai M, Angew. Chem.Int. Ed. 2014, 53, 6519; Angew. Chem. 2014, 126, 6637. Synthesis of aglycon and total synthesis:c)ElMarrouni A, Lebeuf R, Gebauer J, Heras M, Stellios A, Cossy J, Org. Lett. 2012, 14, 314; [PubMed: 22149647] d)Yadav JS, Haldar A, Maity T, Eur. J. Org. Chem. 2013, 3076;e)Fuwa H, Yamagata N, Saito A, Sasaki M, Org. Lett. 2013, 15, 1630; [PubMed: 23472842] f)Fuwa H, Okuaki Y, Yamagata N, Sasaki M, Angew. Chem. Int. Ed. 2015, 54, 868; Angew. Chem. 2015, 127, 882.
- [5]. a)Liu K, Taylor RE, Kartika R, Org. Lett. 2006, 8, 5393; [PubMed: 17078726] b)Kartika R, Taylor RE, Angew. Chem. Int. Ed. 2007, 46, 6874; Angew. Chem. 2007, 119, 6998;c)Kartika R, Frein JD, Taylor RE, J. Org. Chem. 2008, 73, 5592; [PubMed: 18570473] d)Kartika R, Gruffi TR, Taylor RE, Org. Lett. 2008, 10, 5047; [PubMed: 18855401] e)Liu K, Arico JW, Talyor RE, J. Org. Chem. 2010, 75, 3953; [PubMed: 20481506] f)Wilson MR, Taylor RE, Org. Lett. 2012, 14, 3408; [PubMed: 22720980] g)Stefan E, Taylor RE, Org. Lett. 2012, 14, 3490; [PubMed: 22716968] h)Stefan E, Nalin AP, Taylor RE, Tetrahedron 2013, 69, 7706. [PubMed: 24058215]
- [6]. a)Suen LM, Steigerwald ML, Leighton JL, Chem. Sci. 2013, 4, 2413; [PubMed: 25165504] b)Ho S, Bucher C, Leighton JL, Angew. Chem. Int. Ed. 2013, 52, 6757; Angew. Chem. 2013, 125, 6889.
- [7]. a)Funk RL, Abelman MM, Jellison KM, Synlett 1989, 36;<lit b>Shelkov R, Nahmany M, Melmann A, J. Org. Chem. 2002, 67, 8975; [PubMed: 12467417] c)Shelkov R, Nahmany M, Melmann A, Org. Biomol. Chem. 2004, 2, 397; [PubMed: 14747868] d)Paquette LA, Efremov I, J. Org. Chem. 2005, 70, 510; [PubMed: 15651794] e)Sano S, Ichikawa T, Nakao M, Nagao Y, Chem. Lett. 2012, 41, 68;f)Zhong Y-L, Gauthier DR Jr. , Shi Y-J, McLaughlin M, Chung JYL, Dagneau P, Marcune B, Krska SW, Ball RG, Reamer RA, J. Org. Chem. 2012, 77, 3297; [PubMed: 22423625] g)Peric\$ MA, Serratos F, Valentí E, Tetrahedron 1987, 43, 2311;h)Magriotis PA, Vourloumis D, Scott ME, Tarli A, Tetrahedron Lett. 1993, 34, 2071;i)Liang L, Ramaseshan M, MaGee DI, Tetrahedron 1993, 49, 2159;j)Moslin RM, Jamison TF, J. Am. Chem. Soc. 2006, 128, 15106;k)Hoye TR, Danielson ME, May AE, Zhao H, J. Org. Chem. 2010, 75, 7052; [PubMed: 20932014] l)Tran V, Minehan TG, Org. Lett. 2011, 13, 6588. [PubMed: 22103709]
- [8]. a)Booth PM, Fox CMJ, Ley SV, J. Chem. Soc. Perkin Trans. 1 1987, 121;b)Park PU, Broka CA, Johnson BF, Kishi Y, J. Am. Chem. Soc. 1987, 109, 6205;c)Clarke T, Ley SV, J. Chem. Soc. Perkin Trans. 1 1987, 1, 131–135;d)Ley SV, Woodward PR, Tetrahedron Lett. 1987, 28, 345–346;e)Fox CMJ, Ley SV, Org. Synth. 1988, 66, 108–115;f)Booth PM, Broughton HB, Ford MJ, Fox CMJ, Ley SV, Slawin AMZ, Williams DJ, Woodward PR, Tetrahedron 1989, 45, 7565–7580;g)Parenty A, Moreau X, Niel G, Campagne J-M, Chem. Rev. 2013, 113, PR1.
- [9]. a)Denmark SE, Beutner GL, Wynn T, Eastgate MD, J. Am. Chem. Soc. 2005, 127, 3774; [PubMed: 15771512] b)Navarro I, Basset J-F, Hebbe S, Major SM, Werner T, Howsham C, Bräckow J, Barrett AGM, J. Am. Chem. Soc. 2008, 130, 10293.
- [10]. Sugita Y, Sakaki J-I, Sato M, Kaneko C, J. Chem. Soc. Perkin Trans. 1 1992, 2855
- [11]. Evans DA, Chapman KT, Carreira EM, J. Am. Chem. Soc. 1988, 110, 3560.
- [12]. Danheiser RL, Miller RF, Brisbois RG, J. Org. Chem. 1990, 55, 1959.
- [13]. Gradillas A, Pérez-Castells J, Angew. Chem. Int. Ed. 2006, 45, 8086; Angew. Chem. 2006, 118, 8264.
- [14]. Hong SH, Sanders DP, Lee CW, Grubbs RH, J. Am. Chem. Soc. 2005, 127, 1716.
- [15]. a)Grieco PA, Gilman S, Nishizawa M, J. Org. Chem. 1976, 41, 1485;b)Sharpless KB, Young MW, J. Org. Chem. 1975, 40, 7.
- [16]. a)Takai K, Nitta K, Utimoto K, J. Am. Chem. Soc. 1986, 108, 7408;b)Evans DA, Black WC, J. Am. Chem. Soc. 1993, 115, 4497.

- [17]. Notably, during the screening of the glycosylation conditions, we found that the hemiketal pyran is a highly acid-sensitive moiety; acids like TrClO_4 , TBSOTf, TMSOTf, PPTS, and AgClO_4 , lead to either double eliminated product (similar to **21**), or decomposition of starting material. **21**
- [18]. Li P, Li J, Arikan F, Ahlbrecht W, Dieckmann M, Menche D, J. Org. Chem. 2010, 75, 2429. [PubMed: 20334428]
- [19]. See Supporting Information; Table S1; Figure S2.
- [20]. a)Crimmins MT, Chaudhary K, Org. Lett. 2000, 2, 775; [PubMed: 10754681] b)Crimmins MT, King BW, Tabet EA, Chaudhary K, J. Org. Chem. 2001, 66, 869.
- [21]. See Supporting Information; Scheme S3.
- [22]. See supporting information; Scheme S4.
- [23]. See Supporting Information;

Figure 1.
Proposed structures of polykeide natural products isolated from *Lyngbya bouilloni*.

Figure 4.
¹H NMR (CDCl₃) comparison between natural lyngbyaloside C(*Z*)^{3c} and *Z*-**1a** (nominal).

* Deviation in ^{13}C (ppm) between Natural lyngbyaloside (Z) and compound 1c-Z

Figure 5.
 ^1H NMR (CDCl_3) and ^{13}C NMR comparison between natural lyngbyaloside C(Z) 3c and Z-1c.

Scheme 1.

Synthesis of tertiary alcohol **9a** by electrophilic ether transfer reaction. Reagents and conditions: a) *Z*-crotyl-trichlorosilane, Leighton catalyst (*R,R*)-**3**, DBU, CH₂Cl₂, 0°C, 1 h; TBAF, HCl, 93%, 99% *ee*; b) chloromethyl-2-naphthylmethyl ether **5**, DIPEA, TBAI, CH₂Cl₂, 40°C, 12 h, 70%; c) CH₃PPh₃Br, *n*BuLi, THF, 0°C to RT, overnight, 71 %; d) ICl, PhCH₃, -95 °C, 1 min; DIPA/H₂O, 3 h, 76%, d.r. = 10:1; e) TBSOTf, 2,6-lut., CH₂Cl₂, 0°C, 30 min, 95%; f) DDQ, CH₂Cl₂, pH 7 phosphate buffer, RT, 1 h, 93%; g) NaH, THF, 0°C to RT, overnight, 86%; h) 3-benzoxylpropyl magnesium bromide, CuI, THF, -78 to -15°C, 2 h, 65%. DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene; TBAI: tetrabutylammonium iodide; DIPA: diisopropylamine; TBSOTf: trimethylsilyl trifluoromethanesulfonate; 2,6-lut.: 2,6-lutidine; DDQ: 2,3-dichloro-5,6-dicyano-*p*-benzoquinone; RT: room temperature.

Scheme 2.

Synthesis of β -keto thioester **16**. Reagents and conditions: a) acrolein, SiCl_4 , CH_2Cl_2 , -78°C , overnight, 80%; b) TBHP, L-(+)-DIPT, $\text{Ti}(\text{O}i\text{Pr})_4$, MS 4 Å, CH_2Cl_2 , -20°C , 40 h, 90%, 99% *ee*; c) CH_3OH , PhCH_3 , re-fluxing, 2 h, 90%; d) $\text{Me}_4\text{NBH}(\text{OAc})_3$, $\text{AcOH}/\text{CH}_3\text{CN}$, -35°C , 60 h, 92%; e) Et_3SiOTf , 2,6-lut., CH_2Cl_2 , 0°C , 2 h to RT, 87%; f) DiBALH, CH_2Cl_2 , -76°C , 30 min, 93%; g) *tert*-butylthiol diazoacetate **15**, SnCl_2 , CH_2Cl_2 , RT, 4 h, 80%. TBHP: *tert*-butylhydrogenperoxide; DiBALH: diisobutylaluminum hydride.

Scheme 3.

Synthesis of macrocyclic core **20a**. Reagents and conditions: a) AgTFA, Et₂O (0.08 M), RT, 10 min, 73%; b) citric acid, MeOH, 70 °C, overnight, 90%; c) PMBONHCCl₃, La(OTf)₃, PhCH₃, RT, 30 min, 80%; d) Hoyveda-Grubbs II, benzoquinone, CH₃Ph (0.003 M), refluxing, overnight, 80%; e) H₂ (1 atm), Raney Ni, MeOH, RT, overnight, 99%; f) 2-nitrophenyl seleocyanate, PBu₃, THF, RT, 15 min; H₂O₂, NaHCO₃, THF, 55 °C, 40 min, 85%; g) OsO₄, NMO, CH₂Cl₂, RT, overnight; NaIO₄, 0 °C, 1 h, 78%; h) CrCl₂, CHI₃, 1,4-dioxane/THF, 0 °C to RT, 72%. AgTFA: silver(I) trifluoroacetate; PMB: *p*-methoxylbenzyl; NMO: *N*-methylmorpholine-*N*-oxide.

Scheme 4.

Total synthesis of nominal lynybyaloside C. Reagents and conditions: a) DDQ, CH₂Cl₂, pH 7 phosphate buffer, RT, 1 h, 95%; b) **23**, La(OTf)₃, PhCH₃, RT, 15 min, 81 %; c) crotonaldehyde, Hoveyda-Grubbs II, CH₂Cl₂, re-fluxing, 90%; d) Ph₃PCH₂BrBr, NaHMDS, HMPA, THF, RT to -78°C, 15 min, 84%, Z/E=4.3/1; e) HF, CH₃CN, RT, 4 h 77%; f) DDQ, CH₂Cl₂, pH 7 phosphate buffer, RT, 10 min, 70%. NaHMDS: sodium bis(trimethylsilyl)amide; HMPA: hexamethylphosphoramide.

Scheme 5.

Total synthesis of diastereomers **1b** and **1c** and completion of the total synthesis of lyngbyaloside C.