Published in final edited form as:

Nat Chem Biol. 2019 February ; 15(2): 189–195. doi:10.1038/s41589-018-0192-3.

Metalloprotein switches that display chemical-dependent electron transfer in cells

Joshua T. Atkinson¹, Ian J. Campbell², Emily E. Thomas², Sheila C. Bonitatibus³, Sean J. Elliott³, George N. Bennett^{4,5}, Jonathan J. Silberg^{4,6,*}

¹Systems, Synthetic, and Physical Biology Graduate Program, Rice University, Houston, Texas, USA

²Biochemistry and Cell Biology Graduate Program, Rice University, Houston, Texas, USA

³Department of Chemistry, Boston University, Boston, Massachusetts, USA

⁴Department of BioSciences, Rice University, Houston, Texas, USA

⁵Department of Chemical and Biomolecular Engineering, Rice University, Houston, Texas, USA

⁶Department of Bioengineering, Rice University, Houston, Texas, USA

Abstract

Biological electron transfer is challenging to directly regulate using environmental conditions. To enable dynamic, protein-level control over energy flow in metabolic systems for synthetic biology and bioelectronics, we created ferredoxin logic gates that utilize transcriptional and post-translational inputs to control energy flow through a synthetic electron transfer pathway that is required for bacterial growth. These logic gates were created by subjecting a thermostable, plant-type ferredoxin to backbone fission and fusing the resulting fragments to a pair of proteins that self-associate, a pair of proteins whose association is stabilized by a small molecule, and to the termini of a ligand-binding domain. We show that the latter domain insertion design strategy yields an allosteric ferredoxin switch that acquires an oxygen-tolerant 2Fe-2S cluster and can use different chemicals, including a therapeutic drug and an environmental pollutant, to control the production of a reduced metabolite in *Escherichia coli* and cell lysates.

*Address correspondence to: Jonathan J. Silberg, Biosciences Department, 6100 Main Street, Houston TX 77005, Tel: 713-348-3849, joff@rice.edu.

AUTHOR CONTRIBUTIONS

J.T.A. designed and constructed all DNA vectors, performed the multiple sequence analysis, and did all cellular experiments. I.J.C. purified proteins and performed the lysate experiments. E.E.T. evaluated the switch substrate specificity profile. S.C.B and S.J.E. performed the voltammetry. J.T.A., J.J.S., and G.N.B. conceptualized the project. J.T.A. and J.J.S. wrote the manuscript.

COMPETING INTERESTS

J.J.S., J.T.A., G.N.B. and I.J.C. have submitted a patent application (No 62/583,770) covering the use of fragmented Fds as chemical-dependent electron carriers, entitled “Regulating electron flow using fragmented proteins.”

DATA AVAILABILITY

There are no restrictions on data availability. Accession codes and sequences used for multiple sequence alignments are provided in the supplement. Raw data for Figures 1–3 is available upon request.

INTRODUCTION

Achieving direct, dynamic control over biological electron transfer (ET) through synthetic biology is critical for creating living sensors and bioelectronics^{1–4}, controlling microbial electro-synthesis and -fermentation⁵, and building efficient metabolic pathways for chemical synthesis^{6,7}. At the hub of mediating the transfer of low potential electrons is the ferredoxin (Fd) family, ancient Fe-S proteins that are deeply rooted in the tree of life^{8,9}. These electron carriers are important for ancient metabolic processes, such as hydrogen-dependent CO₂ fixation by autotrophs (acetogens and methanogens) and light-dependent CO₂ fixation by photoautotrophs¹⁰. Additionally, Fds are widespread across all three domains of life. Some organisms contain over a dozen Fd paralogs, and Fds have been found to support ET to over eighty classes of oxidoreductases¹¹, suggesting that these electron carriers helped spur the evolution of diverse bioenergetics strategies¹². Recently, Fds have been used to construct synthetic ET chains. These pathways have been developed to evolve Fd-dependent enzymes with improved catalytic functions¹³ and to understand Fd-partner specificity¹⁴. These efforts have demonstrated the potential for manipulating protein electron carriers for synthetic biology. However, we still lack Fds whose ET can be switched “on” and “off” in response to specific environmental conditions.

Some organisms use Fds to dynamically control ET through post-translational modifications. Upon Fe-S cluster oxidation, nitrogenase-protecting Fds alter their conformation and binding affinity to nitrogenase to protect from oxidative damage^{15,16}. Phosphorylated and calcium-bound forms of Fds were recently discovered in cyanobacteria¹⁷ and rhizobia¹⁸, respectively. These discoveries suggest that Fd ET may also be dynamically regulated through signaling cascades, although the exact mechanisms by which these modifications affect Fd ET remain unclear. Additionally, protein design efforts have investigated whether Fd ET can be deliberately controlled. Fds have been rationally mutated using structural information¹⁹ and fused to acceptor proteins to tune the relative proportions of electrons transferred to different metabolic pathways^{20–22}. While these efforts have shown how protein evolution can redirect energy flow, they have yet to yield insight into the ways that Fds acquire new allosteric functions.

Previous studies have shown that proteins can be fragmented into polypeptides whose stable association and cooperative activities are dependent upon ligand-binding events²³. This has been achieved by fusing engineered fragments to pairs of proteins whose association is stabilized by ligand binding^{24,25}, and by fusing protein fragments to the termini of ligand binding domains^{25,26}. To investigate if a similar design approach can be used to create protein electron carriers that are regulated by environmental conditions, we rationally designed four split Fds (sFds) and evaluated their ability to support ET in cells. All four sFds supported ET in cells when they were fused to peptides that stabilize fragment association, and three of the sFds could be used as transcriptional AND gates in which a pair of conditional promoters regulate Fd-mediated ET. One of the sFds could also be used to create a three-hybrid system that only supports cellular ET transfer when both fragments are expressed and cells are exposed to the small-molecule rapamycin. Additionally, when the fragments of this sFd were fused to the termini of the estrogen receptor (ER) ligand-binding

domain (LBD), the resulting polypeptide presented ET that could be enhanced post-translationally by different estrogen-receptor modulators.

RESULTS

Monitoring protein electron transfer using a selection.

To rapidly assay ET mediated by engineered Fds, we utilized *Escherichia coli* EW11¹³, a strain that cannot grow in minimal medium containing sulfate as a sulfur source unless it expresses a synthetic ET chain consisting of a Fd-NADP⁺ reductase (FNR) electron donor, a Fd electron carrier, and a Fd-dependent sulfite reductase (SIR) electron acceptor (Fig. 1a). Endogenous oxidoreductases were systematically removed from the genome of this strain to insulate Fd-dependent redox pathways from native redox reactions¹³. To evaluate Fd ET in cells, we constitutively expressed *Zea mays* FNR and SIR and used a TetR-repressible promoter to control the expression of plant-type 2Fe-2S Fds. With this expression strategy, growth depended upon the level of anhydrotetracycline (aTc) (Fig. 1b), and thus, the amount of Fd available to mediate ET.

The effect of aTc on complementation by five 2Fe-2S Fds was analyzed, including plant, cyanobacterial, algal, and phage family members (Fig. 1c). Two of the Fds were unable to complement growth, including *Spinacia oleracea* and *Prochlorococcus* phage P-SSM2 Fds. The other three Fds presented half-maximal concentration at different aTc concentrations. *Mastigocladus laminosus* Fd (*MI*-Fd) yielded half-maximal complementation at the lowest aTc concentration. This thermostable Fd, which displays optimal ET at 65 °C²⁷, was targeted for all subsequent design efforts to minimize the destabilizing effects of mutations, since previous studies have shown that tolerance to fission correlates with thermostability²⁸.

Design of split protein electron carriers.

To identify backbone fission locations that are non-disruptive to *MI*-Fd structure, we generated a multiple sequence alignment of 2Fe-2S Fds (Supplementary Figure 1) and calculated the number of different amino acids at each native *MI*-Fd position (Fig. 1d). We targeted four sites with moderate sequence divergence for backbone fission, including the peptide bonds following residues 9, 35, 65, and 72. We chose these sites because they vary in their distances from the 2Fe-2S cluster, and they differ in the number of cofactor and partner-binding residues encoded distributed across the pair of fragments.

In the *MI*-Fd structure²⁷, the backbone cleavage sites occur 9.5 to 18.5 Å from the 2Fe-2S cluster (Supplementary Figure 2). For two of the split Fd (sFd) proteins, sFd-9 and sFd-35, all of the cysteines that coordinate iron reside on a single polypeptide (Fig. 1d). The other split proteins, sFd-65 and sFd-72, have their iron-coordinating cysteines dispersed across the fragments, potentially requiring 2Fe-2S cluster maturation following fragment complementation. Mapping the backbone fission sites onto the Fd-SIR complex structure²⁹ reveals that the SIR contacting residues in these sFds are divided among the different polypeptide fragments in sFd-35, sFd-65, and sFd-72 (Supplementary Figure 3). In contrast, the FNR contacting residues in the Fd-FNR complex³⁰ are only divided among the different fragments in sFd-65 and sFd-72.

Split ferredoxins support electron transfer in cells.

To assist with sFd ET, each pair of Fd fragments was initially expressed as fusions to SYNZIP-17 and SYNZIP-18 (Fig. 1e), synthetic peptides that stably associate to form a coiled coil³¹. When the sFds were expressed in *E. coli* EW11, complementation was observed with all four variants (Fig. 1f), albeit to varying extents. Removal of one or both SYNZIP peptides abolished growth in all cases, suggesting that these split proteins uniformly require assistance with fragment association for efficient ET. TetR- and LacI-repressible promoters were also used to control the expression of each Fd-fragment fusion (Fig. 1g), and complementation was analyzed over a range of aTc and isopropyl β -D-1-thiogalactopyranoside (IPTG) concentrations (Fig. 1h). sFd-35, sFd-65, and sFd-72 all functioned as two-input AND gates to control the production of a reduced sulfur metabolite with sFd-35 exhibiting the highest tunable range. In the case of sFd-35, a low level of complementation was observed when cells were exposed to aTc in the absence of IPTG. This finding is thought to arise as a consequence of leaky expression of the C-terminal Fd fragment.

To investigate whether sFd ET can be regulated post-translationally, we created vectors for expressing the three most active split proteins as fusions to the FKBP12 and the FKBP-rapamycin binding (FRB) domain of mTOR (Fig. 2a), proteins whose interaction is stabilized by rapamycin³², and we assessed their complementation of *E. coli* EW11 growth $\pm$ rapamycin. These sFds were expressed using TetR- and LacI-repressible promoters (Supplementary Figure 4a), and complementation was measured using inducer concentrations that yielded maximal growth when sFds were fused to SYNZIP peptides. In the absence of rapamycin, none of the sFds complemented growth. Addition of rapamycin enhanced the growth of cells expressing sFd-35 (Fig. 2b), but not sFd-65 and sFd-72 (Supplementary Figure 5). With sFd-35, growth was dose dependent and could be tuned to a similar maximal density as cells expressing *MI*-Fd.

To evaluate whether rapamycin affected sFd-35 expression, we examined rapamycin effects on protein accumulation using fragments fused to red fluorescent protein (RFP). With each sFd fragment, this analysis revealed similar RFP levels in the presence and absence of rapamycin (Fig. 2c). These findings provide evidence that ET is controlled post-translationally by rapamycin stabilization of the sFd fragment complex. Complementation measurements performed using different combinations of aTc, IPTG, and rapamycin revealed that sFd-35 can be used as a three-input AND gate to control ET for sulfide production and growth (Fig. 2d).

A ferredoxin switch created using domain insertion.

Allosteric protein switches have been engineered by fusing protein fragments to the termini of domains that undergo chemical-dependent conformational changes^{25,26}. To test whether this domain-insertion strategy can be used to diversify the chemical regulation of *MI*-Fd, we inserted the ligand-binding domain (LBD) of the human estrogen receptor (ER) after residue 35 to create sFd-35-ER (Fig. 3a). This LBD was chosen because it undergoes a conformational change upon 4-hydroxytamoxifen (4-HT) binding that alters the proximity

of the protein termini³³. In addition, we recently found that insertion of the ER-LBD into an amino acyl tRNA synthetase yields protein switches with 4-HT dependent activities²⁵.

We evaluated the activity of sFd-35-ER by expressing this protein using an aTc-inducible promoter (Supplementary Figure 4b). In medium containing aTc, sFd-35-ER did not significantly enhance *E. coli* EW11 growth following a 48 hour incubation (Fig. 3b). In contrast, complementation was enhanced to *MI*-Fd levels when cells were grown in the presence of aTc and 4-HT. We also expressed this protein as a fusion to RFP (sFd-35-ER-RFP) to evaluate how 4-HT affects protein expression. This analysis revealed similar expression across a range of 4-HT concentrations (Fig. 3c), providing evidence that the increased complementation arises from 4-HT effects on sFd-35-ER ET rather than increases in expression.

The ER-LBD binds two classes of estrogen receptor modulators, agonists and antagonists, which elicit distinct conformations (Supplementary Figure 6) and tissue-specific responses *in vivo*^{33,34}. To evaluate how these different modulators affect sFd-35-ER ET, we evaluated sFd-35-ER growth complementation in the presence of agonists, antagonists, and an endocrine disruptor (Fig. 3d). This analysis revealed that sFd-35-ER ET is chemically selective and activated by the ER antagonists (4-HT, raloxifene, lasoxifene) and endocrine disruptor (bisphenol-A). However, this switch is not activated by ER agonists (17 β -estradiol, diethylstilbestrol, hexestrol).

***In vitro* characterization of the ferredoxin switch.**

To understand how *MI*-Fd Fe-S cluster binding and ET are affected by domain insertion, sFd-35-ER was overexpressed in *E. coli* and purified under aerobic conditions. Recombinant sFd-35-ER was brown in color throughout the purification protocol (Supplementary Figure 7) and presented absorbance (Fig. 4a) and visible circular dichroism (CD) spectra (Fig. 4b) that were consistent with the presence of a 2Fe-2S metallocluster and similar to *MI*-Fd and other plant-type Fds^{35,36}.

To investigate if domain insertion affects the reduction potential of *MI*-Fd, we characterized *MI*-Fd and sFd-35-ER using protein film voltammetry³⁷. In the absence of 4-HT, we found that the midpoint potentials of both *MI*-Fd and sFd-35-ER (Fig. 4c) were -336 and -320 mV, respectively. These measurements indicate that sFd-35-ER retains a potential that is adequate for coupling NADPH oxidation by FNR ($\text{FAD } E^\circ = -337$ mV)³⁸ to sulfite reduction by SIR ($[4\text{Fe-4S}] E^\circ = -415$ mV, siroheme = -285 mV)³⁹. In the presence of 4-HT, *MI*-Fd presented a similar midpoint potential (-330 mV) as in the absence of ligand, while sFd-35-ER yielded a $+27$ mV shift in potential (-293 mV). These findings provide evidence that domain insertion can be used to create a Fd whose redox potential is tuned by chemical binding, and they suggest that this design approach could be applied to proteins with alternative 2Fe-2S cluster ligands to create switches capable of regulating ET over a range of potentials^{11,40}.

To directly demonstrate that purified sFd-35-ER functions as a switch *in vitro*, we examined whether this protein could support sulfide production in lysates derived from *E. coli* EW11 expressing only FNR and SIR (Fig. 4d). We monitored SIR activity in desalted cell lysates

using sulfidefluor-7 (SF7)⁴¹, a fluorescent probe for hydrogen sulfide. Cell extracts derived from *E. coli* EW11 expressing FNR and SIR presented sulfide production rates that depended upon NADPH concentration (Supplementary Figure 8). To minimize this background signal, we added a low level of NADPH (40 μ M) that did not yield large sulfide production rates over the time course of our measurements. With this *in vitro* assay, the rate of sulfide production is proportional to the concentration of *MI*-Fd added to cell lysates (Supplementary Figure 9). The sulfide production rate was not enhanced by adding 4-HT alone, demonstrating that this chemical does not support ET to SIR in the absence of a Fd. Addition of purified sFd-35-ER to cell lysates containing FNR and SIR increased the sulfide production rate (Fig. 4e), indicating that this protein can mediate ET to SIR in the absence of 4-HT. However, in the presence of 4-HT, the sulfide production rate was significantly enhanced. Taken together, these measurements provide direct evidence that 4-HT regulates sulfide production by altering the conformation of sFd-35-ER and the efficiency with which it transfers electrons from FNR to SIR.

DISCUSSION

Chemical-dependent protein electron carriers represent a new class of synthetic switches that will enable precision control over electron flow in cells. Herein, we show that these electrical switches can be used to control ET from FNR to SIR. These electron donor and acceptor proteins were used as partner proteins because they allowed for the rapid assessment of Fd-mediated ET using growth complementation of a nutritional requirement. Previous studies have shown that Fds can support ET in diverse non-native pathways, such as hydrogen^{13,19,20}, propane⁷, and antibiotic synthesis⁶. These observations suggest that our chemical-dependent Fds could be used to regulate diverse metabolic outputs by expressing them in parallel with FNR and alternative electron acceptors that are also dependent on 2Fe-2S Fds, such as aldehyde decarbonylase⁷, cytochrome P450s²², and glutamine oxoglutarate aminotransferase⁴².

A thermostable Fd was used as a starting point for switch design to maximize the likelihood of discovering Fd variants capable of ET. Among the different backbone fission sites tested, cleavage after residue 35 yielded the most robust fragments for different design strategies. With this sFd, ET was enhanced by fusion to SYNZIP-17/18 or fusion to FKBP/FRB; the other sFds only displayed function when fused to the SYNZIP peptides. Additionally, sFd-35 fragments could be fused to the end of the ER LBD to create a chemical-dependent Fd. These findings suggest that structurally-related fragments in other 2Fe-2S Fds could be used to create similar protein switches. By applying this design strategy to Fds with distinct partner specificities from *MI*-Fd, Fds could be created that extend the diversity of metabolic outputs that can be regulated using ER antagonists at the level of ET.

Fd switches having 2Fe-2S clusters should be useful for regulating ET across the range of midpoint potentials where native Fds function (−455 to −262 mV)¹¹. To create switches that support ET at more negative potentials, our design approach could be applied to clostridial-type Fds which possess multiple 4Fe-4S clusters^{11,40}. Modern clostridial Fds are thought to have arisen from duplication of smaller peptides that required association to function⁸. These smaller peptides have been synthesized *in vitro* and shown to coordinate iron-sulfur

clusters *in vitro*⁴³, suggesting that they may be capable of supporting ET from FNR to SIR when expressed in cells. Additionally, our design approach could be applied to families of protein electron carriers that support ET at more positive midpoint potentials, such as cytochromes, thioredoxins, rieske-type Fds, and cupredoxins¹¹. The major challenge with applying this design strategy to other protein electron carriers will be the development of high throughput assays that can be used to rapidly mine combinatorial libraries of engineered proteins for variants that display chemical-dependent ET.

Electrical protein switches will have a wide range of applications in bioelectronics, metabolic engineering, and synthetic biology. They can be used in place of native oxidoreductases to regulate the flow of electrons derived from oxidative metabolism and photosynthesis in response to specific metabolic or environmental cues. They can also be used to dynamically regulate electron flow between central metabolism and non-native metabolic pathways^{6,7,13,19,20}. Furthermore, electrical protein switches can be used to couple the exquisite sensing capabilities of proteins to electrical communication between cells and conductive materials outside of cells¹. To date, electron conduits have been described whose extracellular ET can be tuned using transcriptional regulation²⁻⁴. By coupling exoelectrogen current production to electrical protein switches, post-translational control over cellular current production may be possible that does not require slow transcriptional and translational processes. In the future, the ligand specificity of these switches can be diversified by fusing Fd fragments to different ligand-binding domains and by mutating the ligand-binding site of our prototype switches.

ONLINE METHODS

Materials.

Rapamycin was from TCI America, isopropyl β -D-1-thiogalactopyranoside (IPTG) was from RPI, and all other chemicals were purchased from Sigma-Aldrich.

Vector design.

Supplementary Table 1 lists all of the plasmids. These were constructed using Golden Gate DNA assembly⁴³ of PCR products amplified using Phusion DNA polymerase (Thermo-Fisher). The genes encoding *Zea mays* FNR, *Zea mays* SIR, and *Spinacia oleracea* Fd were a gift from Pam Silver (Harvard University). The genes encoding *Mastigocladus laminosus*, *Zea mays*, *Chlamydomonas reinhardtii*, and cyanomyophage PSSM-2 Fd were synthesized as G-blocks by Integrated DNA Technologies. The genes for *MI*-Fd and sFd-35-ER were cloned into pET-28b to create expression vectors for protein purification. All vectors were sequence verified.

Calculations.

A multiple sequence alignment (MSA) of 60 plant-type Fd sequences was generated using MUSCLE⁵. Positional amino acid sequence divergence was calculated as the number of unique amino acids observed at each *MI*-Fd native site. Any sites containing a gap in one or more Fd sequences was given a sequence divergence value of 20. The sequence divergence value for each Fd native site was calculated as the average for a sliding window of three

sites. The UniProt numbers for each Fd are provided in Supplementary Figure 1. The FASTA formatted MSA is included as Supplementary MSA.fasta and described in Supplementary Note 1.

Growth assay.

For all growth experiments, *E. coli* EW11 were freshly transformed with two plasmids using electroporation, one encoding the e-donor and acceptor pair (FNR and SIR) and the other encoding either a native Fd, a C42A mutant of *MI*-Fd that is unable to coordinate a 2Fe-2S cluster, or an engineered Fd. Starter cultures were inoculated using single colonies obtained by selecting for colonies containing both plasmids on LB-agar plates. These starter cultures were grown in deep-well 96-well plates for 18 hours at 37°C in 1 mL of a non-selective modified M9 medium (M9c), which contained sodium phosphate, dibasic (6.8 g/L), sodium phosphate, monobasic (3 g/L), sodium chloride (0.5 g/L), 2% glucose, ammonium chloride (1 g/L), calcium chloride (0.1 mM), magnesium sulfate (10 mM), ferric citrate (24 mg/L), p-aminobenzoic acid (2 mg/L), inositol (20 mg/L), adenine (5 mg/L), uracil (20 mg/L), tryptophan (40 mg/L), tyrosine (1.2 mg/L), and the remaining 18 amino acids (80 mg/L each). For complementation analysis, starter cultures that had been grown to stationary phase in M9c were diluted ~1:100 using a 96-well replicator pin into 100 µL of a selective modified M9 medium (M9sa), which is identical to M9c but lacks cysteine and methionine. Cells were grown in the presence of the indicated amount of inducers (aTc, IPTG, rapamycin, or 4-HT) in an Infinite m1000 Pro plate reader (Tecan) at 37°C with shaking at 250 rpm at an amplitude of 1.5 mm in double-orbital mode. Optical density (OD) measurements were taken every 10 minutes. To select for the Fd and e-donor/acceptor plasmids, all growth steps included chloramphenicol and streptomycin at 34 µg/mL and 100 µg/mL, respectively.

Fluorescence spectroscopy.

Whole cell RFP measurements were done as with the growth assay except starter cultures were diluted ~1:100 using a 96-well replicator pin into M9c medium (100 µL) and grown in an incubator at 37°C with shaking at 250 rpm. After 24 hours of growth, OD and fluorescence ($\lambda_{\text{ex}} = 560$ nm, $\lambda_{\text{em}} = 650$ nm) were measured using an Infinite m1000 Pro plate reader (Tecan). Fluorescence was normalized to OD and scaled relative to the condition without chemical inducer.

Protein purification.

E. coli Rosetta transformed with pET28b containing the *MI*-Fd or sFd-35-ER genes were grown at 37°C in LB medium containing 50 µg/mL kanamycin to mid-log phase, induced using 50 µM IPTG, and grown overnight at 30°C while shaking at 250 rpm. Cells harvested by centrifugation (4000g) were resuspended in 20 mL of lysis buffer (per L of culture), which contained 10 mM Tris pH 8, 5 mM dithiothreitol (DTT), 10 mg/L DNase I, and 0.5 mg/mL lysozyme. After freezing overnight at -80°C, cells were thawed and mixed with a cOmplete™ Mini, EDTA-Free protease inhibitor tablet (Sigma-Aldrich) at a ratio of one tablet per 400 mL of total cell lysate. Cell lysates were loaded onto a DE52 anion exchange column (Whatman) that had been equilibrated with TED buffer (25mM Tris pH 8, 1 mM EDTA, and 1 mM DTT), the column was washed with TED containing 200 mM NaCl, and

Fe-S proteins were eluted using TED containing 300 mM NaCl. The brown eluent was diluted with TED to bring NaCl below 100 mM and loaded onto HiTrap Q XL column (GE Healthcare) that had been equilibrated with TED using an AKTA Start FPLC system (GE Healthcare). This column was washed using TED buffer, a linear gradient was run from 0 to 375 mM NaCl TED, and an isocratic 500 mM NaCl TED solution was used to elute the Fe-S proteins. SDS-page was performed using NuPage 12% Bis-Tris Gels (Invitrogen) and the PageRuler Unstained Broad Range Protein Ladder (Thermo Scientific). Samples were concentrated 20-fold using an Amicon Ultra 10K MWCO spin column. Concentrated samples were flash frozen with liquid nitrogen.

Spectroscopy.

Purified *MI*-Fd and sFd-35-ER were dialyzed into TED buffer (25 mM Tris pH 8, 1 mM EDTA, 1 mM DTT). Each protein was incubated with 4-HT (100 μ M) or the carrier DMSO used to dissolve 4-HT (1% of final volume) for 30 min. A J-815 spectropolarimeter (Jasco) was used to measure the ellipticity of samples from 700 nm to 300 nm. UV/Vis absorbance of samples was read using a Cary 50 UV/Vis Spectrophotometer (Varian) from 600 nm to 250 nm. Measurements used quartz cuvettes with a 1 cm path length. Experiments performed twice using two distinct protein preparations yielded the same result.

Cell lysates.

Electrocompetent *E. coli* EW11 were transformed with the plasmid (pSAC01) that constitutively expresses *Zea mays* FNR and SIR. Cells were grown to stationary phase at 37°C and 250 rpm in 2xYTPG medium (16 g/L tryptone, 10 g/L yeast extract, 5 g/L NaCl, 3 g/L KH₂PO₄, 7 g/L K₂HPO₄, and 18 g/L glucose) containing 100 μ g/mL streptomycin. Cells from overnight cultures were diluted 1:100 in fresh 2xYTPG medium containing streptomycin, grown for 2 hours at 37°C while shaking at 250 rpm until mid-log phase, and then shifted to 30°C and grown for an additional 4 hours. Cell cultures were harvested by centrifugation (4000g), cells were resuspended in equal volume of S30 buffer (10 mM Tris-acetate, 14 mM magnesium acetate, 60 mM potassium acetate pH 8.2), cells were washed twice with S30 buffer, and the pellets were weighed and flash frozen with liquid nitrogen. Pellets were thawed, resuspended in 2 mL of S30 buffer per gram of cell pellet, and sonicated using a Q500 Sonicator (Qsonica) with the probe at 20 kHz and 40% maximal amplitude until samples had been exposed to ~0.5 Joules of sonication energy per μ L of cell slurry³⁶. To minimize sample overheating, sonication proceeded for 25 seconds followed by a 15 second rest period between sonication pulses. To remove native reduced cofactors like NADH and NADPH, lysates were applied to a Zeba Spin Desalting Column 7K MWCO (Thermo Scientific) that had been equilibrated with S30 buffer. Protein content in lysates was quantified using Bio-Rad Protein Assay Dye Reagent Concentrate (Bio-Rad), using bovine serum albumin (NEB) as a standard.

Sulfide production assay.

To visualize sulfide production, cells lysates were diluted 2-fold into S30 buffer to 4.8 mg/mL of total protein. Desalted lysates were diluted 1.25-fold with 5x LSR buffer, which contained 50 mM sodium sulfite, 200 μ M β -Nicotinamide adenine dinucleotide 2'-phosphate reduced tetrasodium salt (NADPH), 10 μ M Sulfidefluor 7 AM (Tocris)²⁶, 670

mM potassium acetate, 50 mM ammonium acetate, 40 mM magnesium acetate, and 50 mM potassium phosphate, dibasic pH 7.2. To analyze background sulfide production from lysates alone, additional NADPH was added to the lysate/LSR mixture to achieve higher concentrations. To analyze Fd-dependent sulfide production, purified *MI*-Fd or *sFd*-35-ER was added at the indicated concentrations to the lysate diluent S30 buffer. To limit disulfide bond formation in the ER-LBD *sFd*-35-ER, which has been observed in past studies⁴², this protein was first reduced with freshly made 1 mM DTT in S30 buffer for 20 minutes on ice and then desalted immediately before addition to the assay using a Zeba Spin Desalting Column 7K MWCO. Following NADPH or protein addition, reactions were arrayed in a Corning 96 well plate at room temperature (Costar #3603), transferred to an Infinite m1000 Pro plate reader (Tecan), and heated to 37°C and shaken at 250 rpm with fluorescence readings ($\lambda_{\text{ex}} = 495 \text{ nm}$, $\lambda_{\text{em}} = 520 \text{ nm}$) every 15 seconds. When analyzing chemical-dependent sulfide production, 12.5 μM 4-hydroxytamoxifen (4HT) or blank ethanol was injected into the plate following two minutes of incubation and the lysate diluent S30 buffer volume was reduced accordingly.

Midpoint potentials.

Electrochemical experiments were carried out anaerobically in a MBraun Labmaster glovebox using a PGSTAT12 potentiostat. A three-electrode configuration was used in a water-jacketed glass cell. A platinum wire was used as the counter electrode and a standard calomel electrode was used as the reference electrode. Reported potentials are relative to the standard hydrogen electrode. Baseline measurements were collected using an edge-plane graphite (EPG) electrode that was modified with a 100mM neomycin trisulfate solution, rinsed, and placed into a glass cell containing a 23.5 °C mixed buffer solution (5mM acetate/MES/MOPS/TAPS/CHES/CAPS) pH 7.0, with 100 mM NaCl. A 5 μL aliquot of 720 μM *MI*-Fd or 5mM *sFd*-35-ER was applied directly to the electrode surface with or without 1 μL of 50mM 4-HT, the protein was allowed to reduce in size for approximately one minute at room temperature before being placed into the buffer cell solution. Square wave voltammograms were collected at 23.5 °C with a frequency of 10Hz and electrochemical signals were analyzed using QSoas. Experiments performed twice using two distinct protein preparations yielded the same result.

Statistical analysis.

Growth assays and whole cell fluorescence data are reported as the mean and standard deviation of biologically independent samples ($n=3$). The sulfide production data are reported as the mean and standard deviation of independent experiments ($n=3$). All reported p-values were obtained using two-tailed, independent t-tests. Sample sizes were in accordance with community standards.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

E. coli EW11 and the genes encoding *Zea mays* FNR, *Zea mays* SIR, and *Spinacia oleracea* Fd were a gift from P. Silver (Harvard University). Cellular assay development was supported by DOE grant DE-SC0014462 (G.N.B. and J.J.S.), split Fd efforts were supported by NASA grant NNX15AL28G (J.J.S. and G.N.B), domain insertion efforts were supported by ONR grant N00014-17-1-2639 (J.J.S.), and electrochemistry was supported by DOE grant DE-SC0012598 (S.J.E.). J.T.A. was supported by NSF GRFP and DOE SGCSR fellowships.

REFERENCES

1. Jensen HM et al. Engineering of a synthetic electron conduit in living cells. *Proceedings of the National Academy of Sciences of the United States of America* 107, 19213–19218 (2010). [PubMed: 20956333]
2. Webster DP et al. An arsenic-specific biosensor with genetically engineered *Shewanella oneidensis* in a bioelectrochemical system. *Biosensors & Bioelectronics* 62, 320–324 (2014). [PubMed: 25038536]
3. Golitsch F, Bücking C. & Gescher J. Proof of principle for an engineered microbial biosensor based on *Shewanella oneidensis* outer membrane protein complexes. *Biosensors & Bioelectronics* 47, 285–291 (2013). [PubMed: 23584391]
4. West EA, Jain A. & Gralnick JA Engineering a native inducible expression system in *Shewanella oneidensis* to control extracellular electron transfer. *ACS Synthetic Biology* 6, 1627–1634 (2017). [PubMed: 28562022]
5. Kracke F, Vassilev I. & Krömer JO Microbial electron transport and energy conservation – the foundation for optimizing bioelectrochemical systems. *Frontiers in Microbiology* 6, 575 (2015). [PubMed: 26124754]
6. Shomar H. et al. Metabolic engineering of a carbapenem antibiotic synthesis pathway in *Escherichia coli*. *Nature Chemical Biology* (2018). doi:10.1038/s41589-018-0084-6
7. Kallio P, Pásztor A, Thiel K, Akhtar KM & Jones PR An engineered pathway for the biosynthesis of renewable propane. *Nature Communications* 5, 4731 (2014).
8. Eck RV & Dayhoff MO Evolution of the Structure of Ferredoxin Based on Living Relics of Primitive Amino Acid Sequences. *Science* 152, 363–366 (1966). [PubMed: 17775169]
9. Harel A, Bromberg Y, Falkowski PG & Bhattacharya D. Evolutionary history of redox metal-binding domains across the tree of life. *Proceedings of the National Academy of Sciences* 111, 7042–7047 (2014).
10. Sousa FL et al. Early bioenergetic evolution. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 368, 20130088 (2013).
11. Atkinson JT, Campbell I, Bennett GN & Silberg JJ Cellular Assays for Ferredoxins: A Strategy for Understanding Electron Flow through Protein Carriers That Link Metabolic Pathways. *Biochemistry* 55, 7047–7064 (2016). [PubMed: 27966889]
12. Hall D, Cammack R. & Rao K. Role for ferredoxins in the origin of life and biological evolution. *Nature* 233, 136–8 (1971). [PubMed: 12058758]
13. Barstow B. et al. A synthetic system links FeFe-hydrogenases to essential *E. coli* sulfur metabolism. *Journal of Biological Engineering* 5, 7 (2011). [PubMed: 21615937]
14. Yang J, Xie X, Yang M, Dixon R. & Wang Y-P Modular electron-transport chains from eukaryotic organelles function to support nitrogenase activity. *Proceedings of the National Academy of Sciences of the United States of America* 114, E2460–E2465 (2017). [PubMed: 28193863]
15. Schlesier J, Rohde M, Gerhardt S. & Einsle O. A Conformational Switch Triggers Nitrogenase Protection from Oxygen Damage by Shethna Protein II (FeSII). *Journal of the American Chemical Society* 138, 239–247 (2016). [PubMed: 26654855]
16. Milton RD et al. The In Vivo Potential-Regulated Protective Protein of Nitrogenase in *Azotobacter vinelandii* Supports Aerobic Bioelectrochemical Dinitrogen Reduction In Vitro. *Journal of the American Chemical Society* 139, 9044–9052 (2017). [PubMed: 28595003]

17. Angeleri M, Zorina A, Aro E-M & Battchikova N. Interplay of SpkG kinase and the Slr0151 protein in the phosphorylation of ferredoxin 5 in *Synechocystis* sp. strain PCC 6803. *FEBS letters* 592, 411–421 (2018). [PubMed: 29331018]
18. Moscattello R. et al. Identification of ferredoxin II as a major calcium binding protein in the nitrogen-fixing symbiotic bacterium *Mesorhizobium loti*. *BMC microbiology* 15, 16 (2015). [PubMed: 25648224]
19. Rumpel S. et al. Enhancing hydrogen production of microalgae by redirecting electrons from photosystem I to hydrogenase. *Energy Environ. Sci.* 7, 3296–3301 (2014).
20. Eilenberg H. et al. The dual effect of a ferredoxin-hydrogenase fusion protein in vivo: successful divergence of the photosynthetic electron flux towards hydrogen production and elevated oxygen tolerance. *Biotechnology for Biofuels* 9, 182 (2016). [PubMed: 27582874]
21. Yacoby I. et al. Photosynthetic electron partitioning between [FeFe]-hydrogenase and ferredoxin:NADP⁺-oxidoreductase (FNR) enzymes in vitro. *Proceedings of the National Academy of Sciences of the United States of America* 108, 9396–9401 (2011). [PubMed: 21606330]
22. Mellor S. et al. Fusion of Ferredoxin and Cytochrome P450 Enables Direct Light-Driven Biosynthesis. *ACS chemical biology* 11, 1862–1869 (2016). [PubMed: 27119279]
23. Stein V. & Alexandrov K. Synthetic protein switches: design principles and applications. *Trends in Biotechnology* 33, 101–110 (2015). [PubMed: 25535088]
24. Pelletier J, Campbell-Valois F. & Michnick S. Oligomerization domain-directed reassembly of active dihydrofolate reductase from rationally designed fragments. *Proceedings of the National Academy of Sciences of the United States of America* 95, 12141–12146 (1998). [PubMed: 9770453]
25. Thomas EE, Pandey N, Knudsen S, Ball ZT & Silberg JJ Programming Post-Translational Control over the Metabolic Labeling of Cellular Proteins with a Noncanonical Amino Acid. *ACS synthetic biology* 6, 1572–1583 (2017). [PubMed: 28419802]
26. Guntas G, Mansell TJ, Kim J. & Ostermeier M. Directed evolution of protein switches and their application to the creation of ligand-binding proteins. *Proceedings of the National Academy of Sciences of the United States of America* 102, 11224–11229 (2005). [PubMed: 16061816]
27. Fish A, Danieli T, Ohad I, Nechushtai R. & Livnah O. Structural basis for the thermostability of ferredoxin from the cyanobacterium *Mastigocladus laminosus*. *Journal of Molecular Biology* 350, 599–608 (2005). [PubMed: 15961101]
28. Segall-Shapiro TH et al. Mesophilic and hyperthermophilic adenylate kinases differ in their tolerance to random fragmentation. *Journal of Molecular Biology* 406, 135–148 (2011). [PubMed: 21145325]
29. Kim J, Nakayama M, Toyota H, Kurisu G. & Hase T. Structural and mutational studies of an electron transfer complex of maize sulfite reductase and ferredoxin. *Journal of Biochemistry* 160, 101–109 (2016). [PubMed: 26920048]
30. Kurisu G. et al. Structure of the electron transfer complex between ferredoxin and ferredoxin-NADP(+) reductase. *Nature Structural Biology* 8, 117–121 (2001). [PubMed: 11175898]
31. Reinke AW, Grant RA & Keating AE A synthetic coiled-coil interactome provides heterospecific modules for molecular engineering. *Journal of the American Chemical Society* 132, 6025–6031 (2010). [PubMed: 20387835]
32. Michnick S, Rosen M, Wandless T, Karplus M. & Schreiber S. Solution structure of FKBP. *Science* (New York, N.Y.) 252, 836–839 (1991).
33. Shiau A. et al. The structural basis of estrogen receptor/coactivator recognition and the antagonism of this interaction by tamoxifen. *Cell* 95, 927–937 (1998). [PubMed: 9875847]
34. Paige L. et al. Estrogen receptor (ER) modulators each induce distinct conformational changes in ER alpha and ER beta. *Proceedings of the National Academy of Sciences of the United States of America* 96, 3999–4004 (1999). [PubMed: 10097152]
35. Stephens P. et al. Circular dichroism and magnetic circular dichroism of iron-sulfur proteins. *Biochemistry* 17, 4770–8 (1978). [PubMed: 728385]
36. Ta D. & Vickery L. Cloning, sequencing, and overexpression of a [2Fe-2S] ferredoxin gene from *Escherichia coli*. *The Journal of Biological Chemistry* 267, 11120–11125

37. Bak DW, Zuris JA, Paddock ML, Jennings PA & Elliott SJ Redox Characterization of the FeS Protein MitoNEET and Impact of Thiazolidinedione Drug Binding. *Biochemistry* 48, 10193–10195 (2009). [PubMed: 19791753]
38. Aliverti A. et al. Biochemical and crystallographic characterization of ferredoxin-NADP(+) reductase from nonphotosynthetic tissues. *Biochemistry* 40, 14501–14508 (2001). [PubMed: 11724563]
39. Hirasawa M, Nakayama M, Hase T. & Knaff DB Oxidation-reduction properties of maize ferredoxin: sulfite oxidoreductase. *Biochimica Et Biophysica Acta* 1608, 140–148 (2004). [PubMed: 14871491]
40. Bak DW & Elliott SJ Alternative FeS cluster ligands: tuning redox potentials and chemistry. *Current Opinion in Chemical Biology* 19, 50–58 (2014). [PubMed: 24463764]
41. Lin VS, Lippert AR & Chang CJ Cell-trappable fluorescent probes for endogenous hydrogen sulfide signaling and imaging H₂O₂-dependent H₂S production. *Proceedings of the National Academy of Sciences of the United States of America* 110, 7131–7135 (2013). [PubMed: 23589874]
42. Navarro F, Martín-Figueroa E, Candau P. & Florencio F. Ferredoxin-dependent iron-sulfur flavoprotein glutamate synthase (GlsF) from the Cyanobacterium *synechocystis* sp. PCC 6803: expression and assembly in *Escherichia coli*. *Archives of biochemistry and biophysics* 379, 267–76 (2000). [PubMed: 10898944]
43. Mulholland SE, Gibney BR, Rabanal F. & Dutton LP Characterization of the Fundamental Protein Ligand Requirements of [4Fe-4S]₂^{+/+} Clusters with Sixteen Amino Acid Maquettes. *Journal of the American Chemical Society* 120, 10296–10302 (1998).
44. Salis HM, Mirsky EA & Voigt CA Automated design of synthetic ribosome binding sites to control protein expression. *Nature Biotechnology* 27, 946–950 (2009).

METHODS-ONLY REFERENCES

45. Engler C, Kandzia R. & Marillonnet S. A One Pot, One Step, Precision Cloning Method with High Throughput Capability. *PLoS ONE* 3, e3647 (2008).

Figure 1. Designing split Fds that transfer electrons within cells.

(a) A synthetic ET pathway that couples NADPH oxidation by FNR to sulfide production by SIR using Fd-mediated ET. (b) The effect of varying concentrations of aTc (0.4, 0.8, 1.6, 3.1, 6.3, 12.5, and 25 ng/mL) on the growth of cells expressing *MI*-Fd from an aTc-inducible P_{LtetO1} promoter and FNR and SIR from constitutive promoters. The red line represents the mean in the absence of aTc, black lines represent the mean in the presence of aTc, and the error bars represent the s.d. (n=3 biologically independent samples). (c) Growth complementation observed after 24 hours with *Mastigocladus laminosus* (*MI*; k_{1/2} = 1.8 ng/mL aTc), *Zea mays* root (*Zm*; k_{1/2} = 2.5 ng/mL aTc), *Chlamydomonas reinhardtii* (*Cr*; k_{1/2} = 14.2 ng/mL aTc), *Spinacia oleracea* (*So*; k_{1/2} > 250 ng/mL), and *Prochlorococcus*

phage (Pp ; $k_{1/2} > 250$ ng/mL) Fds. Symbols and error bars represent the mean and s.d., respectively ($n=3$ biologically independent samples). **(d)** The number of different amino acids observed at each residue position within a Fd multiple sequence alignment mapped onto *MI*-Fd primary structure (top). The backbone fission sites used to generate sFds (red lines) are compared with the number of intramolecular residue-residue contacts ($n = 0$ to 10 displayed as proportionally-sized black circles) within the structure of *MI*-Fd (PDB = 1RFX), as well as intermolecular binding residues, including the cysteines that coordinate the 2Fe-2S cluster (yellow) and the residues that contact SIR²⁹ (green) and FNR³⁰ (purple). **(e)** Each pair of sFd fragments, Fd_{f1} and Fd_{f2} (blue) was expressed as a fusion to SYNZIP-17 (SZ17; orange) and SYNZIP-18 (SZ18; red), respectively. **(f)** Complementation of *E. coli* EW11 growth by *MI*-Fd and a C42A mutant of *MI*-Fd that lacks a cysteine required for coordinating iron is compared with the complementation observed in cells expressing Fd_{f1} and Fd_{f2} alone, Fd_{f1} and the SZ18-Fd_{f2} fusion, and Fd_{f1} and Fd_{f2} fused to SZ17 and SZ18, respectively. Bars and error bars represent the average and s.d., respectively ($n=3$ biologically independent samples). * represents significant growth relative to the C42A mutant (p-values [sFd-9: 2×10^{-3} ; sFd-35: 8×10^{-4} ; sFd-65: 6×10^{-4} ; sFd-72: 1×10^{-4}] using a two-tailed, independent t-test). **(g)** Genetic circuit used to regulate Fd fragment expression. The Fd_{f1}-S17 fusion is expressed using an aTc-inducible promoter, the S18-Fd_{f2} fusion is expressed using an IPTG-inducible promoter, and the partner proteins are constitutively expressed. **(h)** Effects of sFd fragment concentration, controlled by aTc and IPTG, on *E. coli* EW11 complementation. Each box shows cell density from cultures with the growth rate (h^{-1}) indicated by the color. The white lines and shaded region within each box represent the cell density mean and s.d., respectively ($n=3$ biologically independent samples).

Figure 2. Post-translational control over Fd electron transfer in cells.

(a) sFd-35 fragments Fd_{f1} and Fd_{f2} (blue) were expressed as fusions to FKBP (orange) and FRB (red), respectively. (b) Effect of rapamycin concentration on *E. coli* EW11 growth in the presence of aTc (150 ng/mL) and IPTG (500 μM) concentrations that yielded strong complementation with SZ17/SZ18 fusions. Complementation by Fd_{f1}-FRB/FKBP-Fd_{f2} ($k_{1/2}$ = 9.95 μM rapamycin) is compared with *MI*-Fd and a C42A mutant of *MI*-Fd that lacks a cysteine required for coordinating iron. Growth of cells expressing Fd_{f1}-FRB/FKBP-Fd_{f2} is enhanced significantly above the C42A mutant when 25 μM rapamycin is added (p-values [25 μM- 1×10^{-4} , 50 μM- 6×10^{-5}] using a two-tailed, independent t-test). Symbols and error bars represent the mean and s.d., respectively (n=3 biologically independent samples). (c) The effect of rapamycin on the relative fluorescence of cells expressing Fd_{f1}-FKBP-RFP and FRB-Fd_{f2}-RFP. Upon addition of rapamycin in ethanol, there was no significant increase in fluorescence for either fragment compared to ethanol alone (p-value > 0.01 using a two-tailed, independent t-test). Symbols and error bars represent the mean and s.d., respectively (n=3 biologically independent samples). All fluorescent reporters were amended to the C-terminus of proteins to maintain the context of the ribosomal binding sites and translation initiation⁴³ (d) Fd_{f1}-FRB and FKBP-Fd_{f2} can function as a three-input AND gate that uses aTc, IPTG, and rapamycin to control electron flow. Each box shows cell density from cultures with the growth rate (h⁻¹) indicated by the color. The white lines and shaded region within each box represent the cell density mean and s.d., respectively (n=3 biologically independent samples).

Figure 3. Using domain insertion to create a Fd switch.

(a) The ER-LBD was inserted after residue 35 in *MI-Fd* to create *sFd-35-ER*. (b) Effect of 4-HT concentration on the complementation of *E. coli* EW11 by *sFd-35-ER* ($k_{1/2} = 2.11 \mu\text{M}$ 4-HT), *MI-Fd*, and *MI-Fd-C42A*. Growth of *sFd-35-ER* is enhanced significantly above the C42A mutant in the presence of 6.25 μM 4-HT (p-values [6.25 μM - 6×10^{-3} , 12.5 μM - 4×10^{-4} , 25 μM - 1×10^{-4} , 50 μM - 2×10^{-5}] using a two-tailed, independent t-test). Symbols and error bars represent the mean and s.d., respectively ($n=3$ biologically independent samples). (c) The effect of 4-HT on the relative fluorescence of cells expressing *sFd-35-ER*.

RFP. Upon addition of 4-HT in ethanol, there was no significant increase in fluorescence compared to ethanol alone (p-value > 0.01 using a two-tailed, independent t-test). Symbols and error bars represent the mean and s.d., respectively (n=3 biologically independent samples). All fluorescent reporters were amended to the C-terminus of proteins to maintain the context of the ribosomal binding sites and translation initiation⁴³. **(d)** Effect of estrogen receptor modulators (50 μ M) on complementation of *E. coli* EW11 by sFd-35-ER with 0 ng/mL aTc (white bars) or 200 ng/mL aTc (gray bars). The agonists (diethylstilbesterol, hexesterol, and 17 β -estradiol) do not significantly enhance growth of EW11 cells compared to DMSO alone (p-value > 0.01 using a two-tailed, independent t-test), while the antagonists (4-hydroxytamoxifen, raloxifene, and lasoxifene) as well as bisphenol-A significantly enhance growth compared to DMSO alone. * represents significant growth (p-values [4-hydroxytamoxifen- 1×10^{-4} , raloxifene- 5×10^{-6} , lasoxifene- 3×10^{-3} , bisphenol-A- 2×10^{-4}] using a two-tailed, independent t-test). Bars and error bars represent the average and s.d., respectively (n=3 biologically independent samples).

Figure 4. Using purified sFd-35-ER to control metabolite production in cell lysates.

(a) The absorbance and (b) circular dichroism spectra of purified sFd-35-ER and *Ml*-Fd (150 μ M each) in the presence (solid line) or absence (dashed line) of 4-HT display features characteristic of 2Fe-2S proteins³⁶. (c) The midpoint potentials of *Ml*-Fd and sFd-35-ER in and presence (solid line) or absence (dashed line) of 4-HT. Raw square wave voltammetry data are plotted in gray, baseline-subtracted data are plotted in black. Experiments depicted in a-c were performed twice using two independent protein preparations yielded the same results. (d) Lysates generated from *E. coli* EW11 constitutively expressing FNR and SIR

were mixed with purified sFd-35-ER (50 nM), NADPH (40 μ M), SO_3^{2-} (10 mM), and SF7 (2 μ M). Following mixing, 4-HT in ethanol (line) or an equivalent amount of ethanol (dashed line) were added to reactions, and fluorescence was monitored at 37°C. (e) The sFd-35-ER switch displayed a significant increase in sulfide production rate upon 4-HT addition ($p = 0.027$, using a two-tailed, independent t-test) relative to the ethanol carrier. The black lines and bars represent the mean while the gray shaded region and error bars represent the s.d. ($n = 3$ independent experiments).