Published in final edited form as:

Behav Brain Res. 2020 January 27; 378: 112156. doi:10.1016/j.bbr.2019.112156.

Impacts of sex and the estrous cycle on associations between post-fear conditioning sleep and fear memory recall

Ihori Kobayashi, Mark Hatcher, Camille Wilson, Linda Boadi, Milan Poindexter, Joanne S. Allard, Eva K. Polston*

Howard University College of Medicine, 520 W St., 20059 Washington, DC, United States

Abstract

Women are at greater risk than men for developing posttraumatic stress disorder (PTSD) after trauma exposure. Sleep, especially rapid-eye-movement sleep (REMS), has been considered a contributing factor to the development of PTSD symptoms through its effects on the processing of emotional memories. However, it remains unknown if sex and sex hormones play a role in the hypothesized impact of sleep on the development of PTSD. Animal models have methodological advantages over human studies in investigating this research question; however, animal models of sleep in PTSD have been tested only with males. C57BL/6 mice (7 males and 15 females) were exposed to 15 footshocks in a footshock chamber, and 5 min after the last footshock, were returned to their home cages for telemetric electroencephalographic sleep recording. Nine to thirteen days later, mice were returned to the footshock chamber for 10 min without footshocks. Fear recall rates were computed by comparing freezing behaviors in the footshock chamber immediately after the footshocks to those during fear context reexposure. Males had significantly lower recall rates compared to metestrous females (that received footshocks on metestrus). Overall, males slept more than both proestrous females (that received footshocks on proestrus) and metestrous females during the dark period. Regression analyses revealed that average REMS episode durations after footshocks were differentially associated with recall rates across groups, such that the association was positive in males, but negative in proestrous females. Results suggest that both sex and the estrous cycle modulate the associations between REMS continuity and fear memory consolidation.

Keywords

PTSD; Sleep; Fear conditioning; Sex difference; Estrous cycle; Mouse

1. Introduction

Posttraumatic stress disorder (PTSD) is a stress-related disorder some people develop after exposure to a traumatic event [1]. Sleep disturbance, one of the most commonly reported symptoms of PTSD, is also a risk factor for PTSD [2,3]. Sleep disturbance prior to and immediately after trauma exposure predicts the development of PTSD [2,4], and it has been suggested that sleep may contribute to the development of PTSD through its effects on

*Corresponding author.: epolston@howard.edu (E.K. Polston).

emotional memory learning. Women are more likely than men to develop PTSD after experiencing a traumatic event (11–20% vs. 4–8%) [5,6]. Sex differences in sleep and effects of circulating sex hormones on sleep have been reported [7–11]; however, the potential role of sleep in sex differences in PTSD risk has not been well explored. Fear conditioning and fear memory extinction have been used as models for the development of and recovery from PTSD in both human and animal studies. Given the methodological challenges in human studies related to recording sleep and controlling for the menstrual cycle immediately after trauma, animal models serve as a powerful tool for elucidating the mechanisms through which posttrauma sleep and circulating sex hormones interact to influence fear memory learning processes. However, to the best of our knowledge, only male animals have been used in studies that examined post-stress sleep in fear conditioning paradigms.

1.1. Sex differences in fear conditioning

Findings of studies examining sex differences in fear conditioning memories in animals and humans vary. In rodents, sex differences were most consistently found in fear responses during context reexposure following cued or contextual fear conditioning. Male rats and mice exhibited greater freezing in footshock chambers used for fear conditioning training than female animals in several studies [12–16], although a few studies did not find sex differences in rats [14,17,18]. Among the small number of studies that examined effects of the estrous cycle, one study found that females conditioned and tested during estrus, when estradiol and progesterone levels are low, had freezing responses equivalent to males and higher than females conditioned and tested during proestrus, when estradiol levels are high and progesterone is increasing [13]. A second study did not find any differences between females across different estrous cycle stages in response to the fear context reexposure [16].

In studies of healthy men and women, men had higher differential skin conductance response (i.e., differences between responses to conditioned stimuli (CSs) paired with an unconditioned stimulus and CSs not paired with an unconditioned stimulus) compared to naturally cycling women during fear conditioning and/or extinction learning [19–22]. These findings may reflect greater difficulty for women in differentiating between danger and safety cues [21], though there are also studies that did not find sex differences [19,22–24]. Inslicht and colleagues' [25] study of men and women with full or subsyndromal PTSD found sex differences in the opposite direction to those reported in studies of healthy participants. Women exhibited elevated skin conductance response to conditioned cues as well as higher differential responses compared to men. Overall, effects of the menstrual cycle stage and endogenous plasma estradiol levels were not observed [19,21–24].

To summarize, in rodent and healthy human studies that did find sex differences, greater reactivity to fear conditioning context or cues in males was indicated. Effects of the estrous cycle on fear conditioning memories in rodents are inconclusive due to the limited number of studies.

1.2. Sex differences in sleep

Women are more likely than men to report insomnia symptoms such as trouble falling and staying asleep, starting during late puberty and continuing throughout adulthood [26–29]. Effects of the menstrual cycle on some aspects of sleep have been suggested by a few polysomnographic studies with small samples of healthy naturally cycling women. These studies reported changes in the proportion of different sleep stages across the menstrual cycle [7,8]. For example, Parry and colleagues found that women in the follicular phase had less shallow sleep and more deep sleep and rapid-eye-movement (REM) sleep compared to women in the luteal phase [8].

In mice, intact females tend to have less overall and non-REM (NREM) sleep than intact males during dark periods [30–32]. Studies have shown that the estrous cycle also affects sleep/wakefulness during dark periods. In intact Sprague-Dawley rats and C57BL/6 J mice, less REM and more wakefulness were observed on proestrus, when estradiol levels are decreasing and progesterone levels are increasing, compared to dark periods during other estrous cycle stages [33–36]. These effects were not observed during the light periods.

1.3. Traumatic stress and sleep

Effects of exposure to traumatic stress or analogues of trauma on sleep have been shown in both rodents and humans. Rodent studies have shown that exposure to footshock stress during fear conditioning often decreased REM sleep [37]; however, these studies were conducted with only male animals. There have been a limited number of human studies that examined sleep soon after trauma exposure [38,39]. A single polysomnographic study that recorded sleep within a month of trauma exposure found that individuals who were exposed to trauma had shorter sleep, more wake after sleep onset, and shorter mean REM segment duration compared to non-trauma-exposed controls [38]. In addition, shorter mean REM segment duration was associated with the subsequent development of PTSD [38]. These findings suggest that altered REM sleep after trauma exposure may contribute to the development of PTSD.

In both humans and rodents, sleep, particularly REM sleep, has been implicated in consolidation of fear-conditioned memories [40,41]. However, limited investigation has been conducted to examine sex differences in associations of post-trauma sleep with the development of PTSD symptoms. Our group has demonstrated previously that sleep, as assessed a few weeks after trauma exposure, was differentially associated with subsequent PTSD symptom severity between men and women [4,42]. Impaired sleep initiation and maintenance predicted more severe PTSD symptoms in men, whereas increased REM sleep predicted more severe PTSD symptoms in women [4,42]. Effects of the menstrual cycle were not examined.

1.4. Experimental aims

The overarching goal of this study was to investigate whether or not sex and circulating gonadal steroids interact with post-stress sleep to influence the consolidation of a conditioned fear memory after a footshock stressor in mice. As mentioned above, studies examining post-footshock sleep in rodents have included only male animals, and only a

limited number of human studies have measured sleep within a few weeks of trauma exposure. Examining the effects of the menstrual cycle on sleep and PTSD symptom development immediately after trauma exposure has not been done in humans due to methodological challenges. We therefore used an animal model of sleep after trauma-exposure to examine if 1) differences in fear memory recall rate exist between males and females, including female groups exposed to stress at different stages of the estrous cycle; 2) measures of post-footshock sleep are different between males and females; 3) sex differences in post-footshock sleep can account for sex differences in fear memory recall rates; and 4) there are sex differences in the associations between post-footshock sleep and fear memory recall rates. Our studies provide important insight into the roles of sex and circulating ovarian hormones in PTSD and other stress-related disorders.

2. Materials and methods

2.1. Experimental overview

Male, proestrous female (P-female), and metestrous female (M-female) C57BL/6 mice were exposed to 15 footshocks in a footshock chamber and 5 min after the last footshock, were returned to their home cages for telemetric electroencephalographic (EEG) sleep recording. Nine to thirteen days later, mice were returned to the footshock chamber for a 10 min context reexposure test. Fear recall rates were computed by comparing freezing behaviors in the footshock chamber immediately after the footshocks to those during fear context reexposure, and the effects of sex and/or day of the estrous cycle and measures of sleep on subsequent conditioned freezing responses were assessed. An overall timeline for the experimental protocol is presented in Table 1.

2.2. Animals

Experimental animals were 11 male and 23 female adult C57BL/6 mice purchased in batches of 6 (2 males and 4 females) at 3–5 months of age from Harlan Laboratories, Inc. (Frederick, MD). Animals were included in the data analyses only if they maintained good health and, in the case of females, continuous estrous cycles throughout the entire experimental protocol. Inclusion also required that the interval between the two test sessions was between 9 and 13 days. The final animal numbers were 7 males and 15 females. Animals were housed individually within the Veterinary Services vivarium at the Howard University College of Medicine (HUCM) with *ad libitum* access to standard mouse chow and drinking water, and were maintained on a reversed light cycle (lights on 1300 h - 0100 h). Mice are a nocturnal species, and for the purposes of presentation in this paper, the light period refers to the time when lights were on in the housing room and the majority of sleep behavior occurred. Conversely, dark period refers to the time when lights were off in the housing room and the majority of wake time was observed. Upon arrival at the vivarium, animals were habituated to their individual home cages for at least 7 days, during which they were not handled or disturbed. Female mice were assigned to one of two groups, a proestrous group (P-female) that was exposed to both the footshock session and the fear memory recall test on the day of proestrus ($n = 8$) or a metestrous group (M-female), exposed to the footshock session and the fear memory recall test on the day of metestrus ($n = 7$). All experimental procedures were consistent with National Institute of Health

guidelines and approved by the Howard University Institutional Animal Care and Use Committee.

2.3. Preparation of telemetric EEG transmitters

Four implantable telemetric transmitters (Model ETA-F10; Data Sciences International [DSI], St. Paul, MN) were purchased to allow recording of sleep EEGs on one pair of mice while habituation and surgical implantation procedures were performed on a second pair. Pairs of experimental animals were alternated across the three experimental groups by pairing a male with a female (P-female and M-female alternately) and then pairing one female from each group every third pair. Prior to implantation, transmitters were cleaned with 0.9% saline and then submerged for 1–2 h at 25 °C in an aseptic cleaning solution (Cidex; Vitality Medical, Salt Lake City, UT), as recommended by the manufacturer. Following sterilization, transmitters were rinsed at least three times in sterile saline, re-rinsed in sterile distilled water and stored in a sterilized glass vial until surgical implantation. The following methods for transmitter implantation, sleep recording, and administration of the footshock stressor were previously published by Sanford and colleagues' team [43–45].

2.4. Surgical implantation of telemetric transmitters

Surgical implantation of the transmitters was performed in the surgical suite of the HUCM under isoflurane anesthesia. Surgical methods were in accordance with the Surgical Manual for Rodents published by DSI and were based on those reported previously by other groups [43–45]. Immediately after anesthetization, animals were treated prophylactically with antibiotics (penicillin, 100 IU/g s.c., gentamycin, 5 mg/kg s.c.). Animals were then placed in the prone position on a sterile surgical pad on top of a heating pad to prevent hypothermia from anesthesia, and an approximately 2 cm longitudinal incision was made along the dorsal midline of the skull. The skull incision was then covered in sterile gauze soaked in physiological saline to prevent drying and the animal was turned over into a supine position. A 1.5–2 cm incision was made through the skin and the abdominal wall along the ventral midline. The transmitter was inserted into the abdominal cavity and positioned parallel to the long axis of the body with the bipotential leads directed rostrally. To exteriorize the bipotential leads, the abdominal wall was perforated approximately 5 mm lateral to the margin of the abdominal incision using an 18-gauge needle and the leads were gently fed through the hole with a small blunt forceps. A subcutaneous tunnel from the ventral abdomen incision to the dorsal cranial incision was then made with a trocar and the bipotential leads were fed through the subcutaneous canal and externalized at the caudal end of the cranial incision. The abdominal muscle and skin incisions were then closed using 5–0 absorbable and silk suture, respectively. The animal was then returned to the prone position and the periosteal meninges of the skull were removed using a cotton-tipped applicator soaked in sterile physiological saline. Once the skull was completely clean and dry, a 0.9 mm microdrill bit was used to drill two holes for subcranial insertion of the bipotential EEG leads (Bregma: AP:1.0, ML:–1.0 for the negative lead and AP:3.0, L:3.0 for the positive lead). The leads were cut to the appropriate length allowing approximately 5 mm of additional length to ensure unrestricted head movement. EEG leads were stripped to expose the metal, and the exposed leads were inserted into the skull holes. A cranial set screw (Plastics One, Inc., Roanoke, VA) was inserted into the skull to stabilize a head cap made of

dental acrylic that was used to cover the EEG leads. Upon discontinuation of anesthetic, animals were maintained in a holding cage on the heating pad until they regained full mobility. An antibiotic ointment was applied to the cranial and abdominal incision sites to prevent infection. Post-operative anti-inflammatory medication (meloxicam 1 mg/kg s.c.) and analgesics (buprenorphine 1.5 mg/kg s.c.) were administered every 6–12 h for at least 24 h and continued as deemed necessary through close observation of the animals for signs of distress (hunched posture, loss of thirst or appetite, aggression, lethargy, etc.). In addition, postoperative nutritional supplement (Nutrical) was made available in the cage during the first postoperative week.

2.5. Postoperative monitoring of estrous cycles

Mice were allowed a 10-day recuperative period after the surgery, during which estrous cycles of the females were monitored through daily vaginal lavage at the beginning of each light period. As a control, males were handled similarly to the females; a drop of saline was deposited on the mid-scrotal skin and then removed using small lavaging pipettes. Vaginal cytology was examined with a light microscope, and used to characterize the day of the estrous cycle using previously published methods [46]. Vaginal smears for females and mock smears for males were continued on a daily basis throughout the remainder of the study

2.6. Sleep EEG recordings

Following the 10-day postoperative recovery period, pre-footshock baseline sleep recording in home cages was initiated. Pre-footshock sleep recordings were collected for 1–2 estrous cycles in females and for a comparable 5–10 day period in males. Human studies have demonstrated enhanced emotional memory consolidation not only during night sleep but also during short naps taken during the day [47], so EEG recordings were collected during both the dark and light periods each day throughout the remainder of the experiment.

2.7. Fear conditioning

Fear conditioning footshock sessions and fear recall test sessions were conducted in the early light period, when animals would normally be initiating sleep. This allowed for us to observe the immediate effects of footshock on same-day sleep parameters and also to perform the procedures before the late light period progesterone surge in P-females. On the proestrus (P-female) or metestrus (M-female) day of the second or third estrous cycle, or after 5–10 days in males, all animals were exposed to a single footshock session. Mice were placed in a footshock chamber (Coulbourn Instruments, Holliston, MA), and after a 10-min habituation period, 15 footshocks (0.5 mA, 0.5 s duration) were delivered at 1.0-min intervals over the course of 14 min [44,45]. Five minutes after the last shock, the mouse was returned to its home cage, and the footshock chamber was cleaned thoroughly with 95% EtOH. Sleep recording was reinitiated immediately upon return of the mice to their home cages and was continued for the remainder of the experiment. Fear recall tests were conducted after two complete post-footshock estrous cycles for females and after an equivalent time period in males. In three cases, females displayed one or two extended estrous cycles (6–7 days) following the footshock stressor. To ensure that this variability would be similar across all groups, three males were also allowed a 12–13 day interval

between the initial footshock session and the subsequent fear recall test. Thus, the inter-test interval spanned 9–13 days for all animals. After the post-footshock recording period was completed, subjects were reexposed to the footshock chamber for a 10-min fear recall test. Animals were videotaped throughout the initial footshock session and during the reexposure session for subsequent assessment of freezing behaviors.

2.8. Sleep assessment

Data from the implanted telemetric transmitters were collected through telemetric signal receiver plates (DSI, Model RPC-1) placed under the animals' home cages and stored on a PC. EEG and locomotor activity data were displayed on a computer screen using REMlogic (Embla, Ontario Canada) or NeuroScore (DSI) sleep scoring software and visually scored in 10-sec epochs following the scoring rules for mouse sleep used in prior studies [48]. Wakefulness, NREM, and REM sleep were determined based on EEG and locomotor activity data. The following sleep parameters were obtained for two light periods and two dark periods during the baseline sleep recording and again during the first two days immediately following the footshock session: percentage of sleep (total sleep duration divided by recording time), percentage of NREM sleep (total NREM sleep duration divided by recording time), percentage of REM sleep (total REM sleep duration divided by recording time), average number of REM sleep episodes per hour (the number of REM episodes divided by the number of hours of recording), and average REM sleep episode duration (total REM sleep duration divided by the number of REM episodes). Percentages rather than absolute amount of sleep or each sleep stage were computed to adjust for differences in sleep recording time during the light period immediately following the footshock session. For baseline measures, sleep parameter data obtained during two light periods and two dark periods were averaged to get one light period value and one dark period value. For females, a proestrous light and dark period and a metestrous light and dark period were averaged. For males, two light and two dark periods were selected randomly from the baseline recording period for averaging.

2.9. Fear memory assessment

Videotaped animal behaviors during the fear conditioning session and the fear context reexposure session were independently reviewed by two trained scorers who were blind to the animal's group assignment for scoring of freezing. Freezing was defined as the lack of movement except for respiration and heartbeat [49]. The scorers determined whether a freezing behavior was exhibited at least once in each 5-second epoch [50], and any epochs for which the scorers' decisions were discrepant were reviewed by the scorers in consultation with an additional investigator to achieve consensus. The percentages of epochs containing freezing during the 5-min period immediately following the last footshock delivery during the fear conditioning session and the first 5 min of the fear context reexposure session were computed. Fear memory recall rates were calculated by dividing the percentage of epochs with freezing during the context reexposure session by the percentage of epochs with freezing during the 5 min immediately after the last footshock delivery.

2.10. Statistical analyses

All statistical analyses were performed using IBM SPSS Statistics Version 25 (Armonk, NY) with an alpha level of .05. Initially, data distributions were examined using Shapiro-Wilk test, and square-root or natural log transformation was applied to some sleep variables including the percentage of REM sleep and the number of REM episodes during certain light/dark periods as their distributions were non-normal. Parametric tests were performed for normally distributed data, and non-parametric tests were applied to variables that could not be normalized by transformation.

To test group differences in fear memory recall rates (Aim 1), we performed Welch's one-way ANOVA followed by Games-Howell post hoc tests as variances were heterogeneous. To test effects of fear conditioning on sleep parameters (Aim 2), a 3 (group: Male, P-female, and M-female) x 3 (Time: baseline, Post-shock Day 1, and Post-shock Day 2) mixed ANOVA with time as a repeated measure factor was performed separately for each normally distributed sleep parameter. Paired comparisons using LSD were conducted following a significant main effect. For non-normally distributed data, Wilcoxon signed ranks test was performed to compare differences between time points (baseline vs. Post-shock Day 1; baseline vs. Post-shock Day 2; Post-shock Day 1 vs. Post-shock Day 2); and Kruskal Wallis test were used to examine group differences at each time point. For the percentage of REM sleep and the average number of REM episodes during dark period, data were not normal when divided into groups, but data distributed normally when groups were combined. Therefore, a repeated measures ANOVA was performed instead of Wilcoxon signed ranks test to examine differences across the time points combining all groups.

To examine whether post-footshock sleep differences account for group differences in fear memory recall rates (Aim 3) and whether the relationships between post-footshock sleep and fear memory recall differ depending on group (Aim 4), hierarchical regression analysis was performed separately for each sleep parameter. Specifically, at Step 1, group dummy codes (two dummy codes: P-females compared to males and M-females compared to males) were entered as predictors, at Step 2, a standardized sleep parameter (i.e. z-score) was entered, and at Step 3, interaction between a standardized sleep parameter and the P-female dummy code and interaction between a standardized sleep parameter and the M-female dummy code were entered. Regression analysis was not performed with Post-shock Day 1 NREM sleep during the light period due to non-normally distributed data.

3. Results

3.1. Group differences in fear memory recall rates

During the 5-min period immediately after the last footshock delivery, no significant group differences in percentage of epochs containing freezing were observed ($F = 0.28$, $p = .755$), indicating that all groups equally acquired fear memory. Therefore, the rest of the analyses focused on fear memory recall rates. Welch's one-way ANOVA revealed a significant group difference ($F = 5.62$, $p = .020$) in fear memory recall rates. Games-Howell post hoc analyses showed that males had significantly lower recall rates than M-females (mean = 42.41 vs.

91.25; $p = .044$) (Fig. 1). Fear memory recall rates of P-females did not differ significantly from either of the other two groups.

3.2. Effects of fear conditioning on sleep

Figs. 2 and 3 present means ($\pm$ s.e.m.) of each sleep variable before and after fear conditioning. As we did not find any significant main effects of group or group $\times$ time interaction effects, the three groups were combined for all sleep parameters for light periods and for percentage of REM sleep and the average number of REM episodes per hour for dark periods. Data for overall percentage of sleep and percentage of NREM sleep are depicted in Fig. 2. Regarding overall percentage of sleep during light periods (Fig. 2A), a 3 (Group) $\times$ 3 (Time: Baseline, Post-shock Day 1, and Post-shock Day 2) mixed ANOVA revealed a significant main effect of time ($F = 12.97$, $p < .001$). Post-hoc analyses indicated a significant increase in percentage of sleep from Baseline to Post-shock Day 1 ($p = .002$) and a significant decrease from Post-shock Day 1 to Post-shock Day 2 ($p < .001$). Neither the main effect of group ($F = 0.60$, $p = .561$) nor the interaction between group and time ($F = 0.78$, $p = .544$) was significant. For percentage of sleep during dark periods (Fig. 2B), significant main effects of time ($F = 3.95$, $p = .028$) and group ($F = 4.72$, $p = .022$) were found. Post-hoc analyses for the effect of time indicated a significant increase from Baseline to Post-shock Day 1 ($p = .017$). Post-hoc analyses unpacking the main effect of group showed that males had a higher percentage of sleep than both P-females ($p = .011$) and M-females ($p = .022$). The interaction between group and time was not significant ($F = .036$, $p = .839$).

Data for NREM sleep are presented in graphs 2C and 2D. As shown for percentage of NREM sleep during light periods (Fig. 2C), Wilcoxon signed ranks test revealed a significant increase from Baseline to Post-shock Day 1 ($z = -2.94$, $p = .003$) and a significant decrease from Post-shock Day 1 to Post-shock Day 2 ($z = -3.91$, $p < .001$). To examine if there were any group differences in percentage of NREM sleep at each time point, one-way ANOVAs were performed for Baseline and Post-shock Day 2, and Kruskal Wallis tests were performed for Post-shock Day 1. No significant group differences were indicated at any of the time points ($p > .142$).

Regarding the percentage of NREM sleep during dark periods (Fig. 2D), a 3 $\times$ 3 mixed ANOVA revealed a significant main effect of time ($F = 3.70$, $p = .034$) and a significant main effect of group ($F = 4.51$, $p = .025$). Post-hoc analyses indicated a significant increase in percentage of NREM sleep from Baseline to Post-shock Day 1 ($p = .020$). Post-hoc analyses of the main effect of group showed that males had higher percentage of NREM sleep than both P-females ($p = .011$) and M-females ($p = .030$). The interaction between group and time was not significant ($F = 0.34$, $p = .853$).

Data on REM sleep parameters are presented in Fig. 3. With regard to percentage of REM sleep during light period (Fig. 3A), a 3 $\times$ 3 mixed ANOVA indicated a significant main effect of time ($F = 4.53$, $p = .017$). Post-hoc analyses revealed a significant decrease in percentage of REM sleep from Post-shock Day 1 to Post-shock Day 2 ($p = .001$). Neither the main effect of group ($F = 0.21$, $p = .813$) nor a group $\times$ time interaction ($F = 0.19$, $p = .941$) was significant. During the dark periods (Fig. 3B), a repeated measure ANOVA revealed no

significant differences across the time points ($F = 3.08$, $p = .057$). To examine if there were any group differences at each time point, one-way ANOVAs were performed for Baseline and Post-shock Day 1, and Kruskal Wallis tests were performed for Post-shock Day 2. No significant group differences were indicated ($p = .119$).

As revealed by a 3×3 mixed ANOVA, there was a significant main effect of time on the average number of REM episodes during light periods (Fig. 3C; $F = 4.65$, $p = .016$). Post-hoc analyses showed that there were significantly fewer REM episodes during Post-shock Day 2 than both Baseline ($p = .039$) and Post-shock Day 1 ($p = .006$). Neither the main effect of group ($F = 0.63$, $p = .542$) nor the interaction between group and time ($F = 0.97$, $p = .396$) was significant. With regard to the average number of REM episodes during dark periods (Fig. 3D), a repeated measure ANOVA showed no significant differences across the time points ($F = 2.37$, $p = .106$). To further examine if there were any group differences at each time point, one-way ANOVAs were performed for Baseline and Post-shock Day 1, and Kruskal Wallis tests were performed for Post-shock Day 2. One-way ANOVA revealed a significant group difference at Baseline ($F = 3.68$, $p = .045$), and post-hoc analyses indicated that the average number of REM episode counts was lower in M-females than males ($p = .015$, not shown). Group differences were not found on Post-shock Day 1 or Post-shock Day 2 ($p = .098$). Regarding average REM episode durations during light and dark periods (Fig. 3E and F), 3×3 mixed ANOVAs showed no significant main effects or significant interactions ($p = .248$).

3.3. Hierarchical regression analyses examining main effects of group and sleep and interactions between group and sleep

Entering the two group dummy codes (P-female vs. male, and M-female vs. male) at Step 1 revealed that M-female significantly positively predicted fear memory recall rates ($\beta = .70$, $p = .004$), indicating that M-female was associated with higher recall rates compared to males. P-female did not significantly predict fear memory recall rates ($\beta = .38$, $p = .094$). M-female remained significant (p between .002 - .020) after entering each sleep parameter at Step 2, indicating that group differences in sleep did not account for the group difference in fear memory recall rates. Among all of the sleep parameters, only average REM episode duration during the dark period of Post-shock Day 1 ($R^2 = .23$, $\beta = .48$, $p = .005$) and percentage of NREM sleep during the dark period of Post-shock Day 2 ($R^2 = .20$, $\beta = -.44$, $p = .011$) significantly accounted for unique variance in fear memory recall rates at Step 2.

Table 2 presents results of the full model (Step 3) of the hierarchical regression analysis for each sleep parameter. The regression coefficients for the main effects of M-females were significant and positive in all of the analyses. A significant positive main effect of the average duration of REM episode was found in analyses for the Post-shock Day 1 light and dark periods, and the Post-shock Day 2 light period. We also found a significant negative main effect of the percentage of NREM sleep during the Post-shock Day 2 dark period. Scatter plots with fitting lines indicating the significant main effects of dark period REM episode duration and NREM percent sleep on fear recall are presented in Fig. 4 (A and C).

As indicated in Table 2, the P-female dummy code interacted with the average duration of REM episodes during the Post-shock Day 1 light period to predict recall rates. Both P-

female and M-female dummy codes interacted with average duration of REM episodes during the Post-shock Day 2 light period to predict fear recall rates. These findings indicate that associations between the average REM episode duration during post-shock light periods and fear memory recall rates were moderated by group, whereas average REM episode duration during the Post-shock Day 1 dark period and the percentage of NREM sleep during the Post-shock Day 2 dark period predicted fear memory recall rate regardless of group. Fig. 4B and D present scatter plots with fitting lines for each group. Different angles of fitting lines for each group reflect the interaction effects (See Table 2).

4. Discussion

The present study contributes to the field of animal models of sleep and fear conditioning, predominantly built on results obtained from male animals, by examining the roles of sex and stage of the estrous cycle on post-conditioning sleep and its associations with fear memory. As related to our experimental aims, we found that: 1) female mice fear conditioned and tested for recall on metestrus had higher fear memory recall rates than males during context re-exposure 9–13 days later; 2) overall, males slept more and had more NREM sleep during dark periods compared to females; 3) sex differences in post-shock sleep did not account for sex differences in fear memory recall; and 4) REM sleep continuity during post-shock light periods differentially predicted fear memory recall rates between sexes. Longer average REM sleep episode duration was associated with greater fear memory recall in males (i.e., positive coefficient for Sleep in the final regression model) but in females, average REM sleep episode duration negatively predicted fear memory recall rates (i.e., negative interaction coefficients).

Our finding of greater fear memory recall in females fear conditioned and tested on metestrus compared to males is consistent with clinical observations indicating that women are more likely to develop PTSD symptoms than men following trauma exposure. The group difference was significant only between males and metestrous females, and was not seen with proestrous females, which may suggest that low estradiol levels enhanced and high estradiol levels hindered fear memory learning and/or recall in females. This finding is consistent with those of Barha and colleagues [51]. In their study, injection of a low dose of 17 β -estradiol that is equivalent to circulating levels of estradiol during diestrus into ovariectomized rats 30 min before fear conditioning resulted in increased fear memory recall during contextual reexposure 24 h later. Conversely, administration of a high dose of 17 β -estradiol that is equivalent to circulating levels on the morning of proestrus resulted in decreased contextual fear memory compared to control rats. In the present experiment, estradiol levels would have differed between P-females and M-females both at the time of fear conditioning and during the fear recall test. Further studies are required to determine if differences in hormone levels at the time of the fear recall test may also have influenced freezing behavior.

Many prior rodent studies have reported that male animals showed greater freezing during exposure to the fear context than females [12–16]. The differences in the results between this study and prior studies may be due partially to methodological differences, such as the use of rats in other studies [12–14,16], administration of only 1–3 footshocks during

conditioning [12,14–16], and a 24 -h interval between conditioning and recall [12,14,13–16]. Differential variability across groups may also be a contributing factor. In the present study, the male group displayed greater variability in fear recall rates than the female groups. Recall rates of some of our males were as high as females whereas other males had zero or markedly low fear recall rates. It has been historically accepted in the fields of neuroscience and biomedical research that males exhibit lower variability than females due to cyclical hormonal fluctuations across the estrous cycle in females [52]. However, meta-analytic and systematic reviews examining sex differences in variability of neurobiological, behavioral, and other biomedical outcomes do not support this premise [52–54]. Our observation that some males may be resilient to fear conditioning is consistent with prior reports indicating individual differences in response to CSs or fear context in male rats [55,56]. Our finding suggests that sex differences in PTSD prevalence in humans may be due to greater resilience in a subpopulation of men rather than an overall heightened vulnerability in women. Factors that contribute to the attenuation of fear memory consolidation and/or recall observed in resilient males have yet to be determined.

Overall, male mice in our study slept more and had more NREM sleep during dark periods compared to females. The non-significant Time x Group interaction indicates that stress exposure did not differentially affect sleep between the sexes or across changing levels of circulating female hormones. In addition, results of our regression analyses showed that the sex difference in post-footshock sleep did not account for the group difference in fear memory recall rates. Prior studies have indicated that males had more overall sleep and NREM sleep during dark periods under non-stressed conditions [30–32], and the present study extends our knowledge about sex differences in sleep by demonstrating that sex differences during post-stress sleep were similar to those seen under non-stressed conditions.

Our results are not consistent with findings of other studies examining post-stress sleep in male rodents. Those studies have shown decreased REM sleep immediately following exposure to footshocks [37]. For example, male C57BL/6 mice exposed to 15 footshocks had decreased REM sleep during light and dark periods and increased or unchanged NREM sleep during light periods immediately following footshocks [44,45]. Changes in the amount of REM sleep after stress exposure in rodents can be influenced by experimental conditions, including types of stressor and daily handling [37,57]. Tang and colleagues [57] found that rats that went through 3 sessions of 5-minute manual restraint before receiving 20 footshocks did not exhibit post-footshock reduction of REM sleep. As our protocol required daily vaginal or mock swabs prior to and following the footshock session, it is possible that differences in daily handling methods between this study and prior studies using only male animals might have contributed to these differential results.

Importantly, the present study also revealed sex differences in associations between post-stress REM sleep continuity and fear memory recall. The average duration of REM episodes during post-shock light periods positively predicted fear memory recall rates in males, but in females, average REM episode duration negatively predicted recall rates. These findings suggest that greater REM continuity promotes fear memory consolidation in males, but not in females.

Mechanisms through which REM sleep influences consolidation of fear-conditioned memory have been actively investigated, but limited attention was paid to effects of sex or the menstrual/estrous cycle on those possible mechanisms. It has been hypothesized that coherent theta activity during REM sleep in the hippocampus and the amygdala enhances consolidation of emotional memory by promoting neural plasticity in these regions [58]. In male mice, optogenetically silencing theta rhythm oscillation of GABA neurons in the medial septum, which is a pacemaker for hippocampal theta activity, during REM sleep immediately after fear conditioning training impaired fear-conditioned contextual memory [59]. However, silencing theta rhythm oscillation during NREM sleep did not affect fear context memory [59]. REM sleep is the sleep stage when levels of acetylcholine in the cortex and the hippocampus are highest [60], and acetylcholine, along with GABA, contributes to generation of theta rhythm [61]. Sex differences in acetylcholine levels in the hippocampus have been reported; male rats had higher basal levels and greater stress-induced increase of acetylcholine than cycling females [62,63]. It is therefore possible that longer REM sleep episodes provide an acetylcholine-rich environment that promoted coherent theta activity in our male mice.

In addition to REM sleep effects, we found that the percentage of NREM sleep during the dark period on Post-shock Day 2 negatively predicted fear recall rates regardless of sex or the estrous cycle. Although the role of slow-wave sleep in consolidation of emotional memories has been less extensively studied than REM sleep, other studies have shown that slow-wave sleep may contribute to emotional memory consolidation. For example, in male C57BL/6 mice, exposure to an odor used as a CS during post-conditioning NREM sleep enhanced fear memory recall [64]. This finding indicates that circumstances during NREM sleep can influence consolidation of fear memories. The present finding of negative correlations between NREM sleep and fear recall rates may indicate that greater NREM sleep can promote resilience when the CS is absent during post-conditioning sleep. As shown in Fig. 4C, the resilient subset of males had markedly high percentages of NREM sleep during the dark period on Post-shock Day 2, suggesting that NREM sleep may have been protective against fear conditioning in those animals.

There were certain limitations to our experimental approach, and the present results should be interpreted in that context. We determined the estrous phase based on daily examination of vaginal cytology, but we did not measure hormone levels, and the intervals between fear conditioning and recall tests were not uniform among mice, but ranged between 9–13 days. The difference in recall rates could be influenced by the estrous phase at the time of recall session as well as the phase at the time of fear conditioning. Further, we did not examine effects of sex and the estrous cycle on fear extinction memory consolidation and post-extinction sleep, and the development of psychopathologies such as PTSD following stress exposure is known to be influenced by both fear extinction learning and fear conditioning. Future studies are therefore required to fully clarify the roles of sex and the estrous cycle in post-extinction sleep and its possible relationship with fear extinction memory processes.

To the best of our knowledge, the present study is the first to examine effects of sex and the estrous cycle in post-stress sleep and its associations with fear conditioned memory in rodents. Results of this study suggest that sex dependent mechanisms associated with REM

sleep continuity may contribute to sex differences in fear memory recall. Future studies exploring neurophysiological mechanisms contributing to sex dependent associations between REM sleep continuity and fear memory processes, including investigation into fear extinction, will elucidate sleep-related mechanisms that contribute to sex differences in stress-related psychopathology.

Acknowledgements

We would like to express our sincere gratitude to Dr. Kebreten Manaye for providing access to the fear conditioning footshock chambers and software used for administration of the footshock stimuli. We also wish to thank Dr. Larry Sanford and his collaborators, Dr. Laurie Wellman and Dr. Mayumi Machida for the time and effort they put forth in training us on the surgical procedures and sleep scoring required for this study, along with Dr. Thomas Mellman for providing ongoing consultation and advice about our experimental design. We also wish to thank the Veterinary Service staff at Howard University College of Medicine (HUCM) for their assistance with the care and housing of our animals. This research was supported by HUCM Bridge Funds and Pilot Study Award (U400045) and Faculty Seed Award (U400022) to EKP, NIMH Mentored Research Scientist Award (K01MH110647) to IK, NCATS grants (KL2TR001432-01 and UL1TR001409) to the Georgetown Howard Universities Center for Clinical Translational Science, and NIH RCMI grant (G12MD00759727) to HUCM.

References

- [1]. American Psychiatric Association Diagnostic and Statistical Manual of Mental Disorders, 5th ed., American Psychiatric Association, Washington, DC, 2013.
- [2]. Bryant RA, Creamer M, O'Donnell M, Silove D, McFarlane AC, Sleep disturbance immediately prior to trauma predicts subsequent psychiatric disorder, *Sleep* 33 (1) (2010) 69–74. [PubMed: 20120622]
- [3]. Galea S, Resnick H, Ahern J, Gold J, Bucuvalas M, Kilpatrick D, et al., Posttraumatic stress disorder in Manhattan, New York City, after the September 11th terrorist attacks, *J. Urban Health* 79 (3) (2002) 340–353. [PubMed: 12200503]
- [4]. Kobayashi I, Mellman TA, Gender differences in sleep during the aftermath of trauma and the development of posttraumatic stress disorder, *Behav. Sleep Med* 10 (3) (2012) 180–190. [PubMed: 22742436]
- [5]. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB, Posttraumatic stress disorder in the national comorbidity survey, *Arch. Gen. Psychiatry* 52 (1995) 1048–1060. [PubMed: 7492257]
- [6]. Breslau N, Peterson EL, Schultz LR, A second look at prior trauma and the posttraumatic stress disorder effects of subsequent trauma: a prospective epidemiological study, *Arch. Gen. Psychiatry* 65 (4) (2008) 431–437. [PubMed: 18391131]
- [7]. Driver HS, Dijk DJ, Werth E, Biedermann K, Borbely AA, Sleep and the sleep electroencephalogram across the menstrual cycle in young healthy women, *J. Clin. Endocrinol. Metab* 81 (2) (1996) 728–735. [PubMed: 8636295]
- [8]. Parry BL, Mostofi N, LeVeau B, Nahum HC, Golshan S, Laughlin GA, et al., Sleep EEG studies during early and late partial sleep deprivation in premenstrual dysphoric disorder and normal control subjects, *Psychiatry Res.* 85 (2) (1999) 127–143. [PubMed: 10220004]
- [9]. Goel N, Kim H, Lao RP, Gender differences in polysomnographic sleep in young healthy sleepers, *Chronobiol. Int* 22 (5) (2005) 905–915. [PubMed: 16298775]
- [10]. Roehrs T, Kapke A, Roth T, Breslau N, Sex differences in the polysomnographic sleep of young adults: a community-based study, *Sleep Med.* 7 (1) (2006) 49–53. [PubMed: 16194623]
- [11]. Redline S, Kirchner HL, Quan SF, Gottlieb DJ, Kapur V, Newman A, The effects of age, sex, ethnicity, and sleep-disordered breathing on sleep architecture, *Arch. Intern. Med* 164 (4) (2004) 406–418. [PubMed: 14980992]
- [12]. Maren S, De Oca B, Fanselow MS, Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP and contextual learning, *Brain Res.* 661 (1–2) (1994) 25–34. [PubMed: 7834376]

- [13]. Markus EJ, Zecevic M, Sex differences and estrous cycle changes in hippocampus-dependent fear conditioning, *Psychobiology* 25 (3) (1997) 246–252.
- [14]. Pryce CR, Lehmann J, Feldon J, Effect of sex on fear conditioning is similar for context and discrete CS in Wistar, Lewis and Fischer rat strains, *Pharmacol. Biochem. Behav* 64 (4) (1999) 753–759. [PubMed: 10593198]
- [15]. Wiltgen BJ, Sanders MJ, Behne MS, Fanselow MS, Sex differences, context preexposure, and the immediate shock deficit in Pavlovian context conditioning with mice, *Behav. Neurosci* 115 (1) (2001) 26–32. [PubMed: 11256449]
- [16]. Chang YJ, Yang CH, Liang YC, Yeh CM, Huang CC, Hsu KS, Estrogen modulates sexually dimorphic contextual fear extinction in rats through estrogen receptor beta, *Hippocampus* 19 (11) (2009) 1142–1150. [PubMed: 19338017]
- [17]. Gupta RR, Sen S, Diepenhorst LL, Rudick CN, Maren S, Estrogen modulates sexually dimorphic contextual fear conditioning and hippocampal long-term potentiation (LTP) in rats, *Brain Res.* 888 (2) (2001) 356–365. [PubMed: 11150498]
- [18]. Baran SE, Armstrong CE, Niren DC, Hanna JJ, Conrad CD, Chronic stress and sex differences on the recall of fear conditioning and extinction, *Neurobiol. Learn. Mem* 91 (3) (2009) 323–332 3 2009. [PubMed: 19073269]
- [19]. Milad MR, Goldstein JM, Orr SP, Wedig MM, Klibanski A, Pitman RK, et al., Fear conditioning and extinction: influence of sex and menstrual cycle in healthy humans, *Behav. Neurosci* 120 (6) (2006) 1196–1203. [PubMed: 17201462]
- [20]. Milad MR, Zeidan MA, Contero A, Pitman RK, Klibanski A, Rauch SL, et al., The influence of gonadal hormones on conditioned fear extinction in healthy humans, *Neuroscience* 168 (3) (2010) 652–658. [PubMed: 20412837]
- [21]. Lonsdorf TB, Haaker J, Schümann D, Sommer T, Bayer J, Brassen S, et al., Sex differences in conditioned stimulus discrimination during context-dependent fear learning and its retrieval in humans: the role of biological sex, contraceptives and menstrual cycle phases, *J. Psychiatry Neurosci* 40 (6) (2015) 368–375. [PubMed: 26107163]
- [22]. Antov MI, Stockhorst U, Stress exposure prior to fear acquisition interacts with estradiol status to alter recall of fear extinction in humans, *Psychoneuroendocrinology* 2014 (49) (2014) 106–118 11.
- [23]. Merz CJ, Tabbert K, Schweckendiek J, Klucken T, Vaitl D, Stark R, et al., Neuronal correlates of extinction learning are modulated by sex hormones, *SCAN* 7 (7) (2012) 819–830. [PubMed: 21990419]
- [24]. White EC, Graham BM, Estradiol levels in women predict skin conductance response but not valence and expectancy ratings in conditioned fear extinction, *Neurobiol. Learn. Mem* 134 (2016) 339–348. [PubMed: 27544848]
- [25]. Inslicht SS, Metzler TJ, Garcia NM, Pineles SL, Milad MR, Orr SP, et al., Sex differences in fear conditioning in posttraumatic stress disorder, *J. Psychiatr. Res* 47 (1) (2013) 64–71. [PubMed: 23107307]
- [26]. Zhang J, Chan NY, Lam SP, Li SX, Liu Y, Chan JW, et al., Emergence of sex differences in insomnia symptoms in adolescents: a large-scale school-based study, *Sleep*. 39 (8) (2016) 1563–1570. [PubMed: 27091537]
- [27]. Jaussent I, Dauvilliers Y, Ancelin M, Dartiques J, Tavernier B, Touchon J, et al., Insomnia symptoms in older adults: associated factors and gender differences, *Am. J. Geriatr. Psychiatry* 19 (1) (2011) 88–97. [PubMed: 20808113]
- [28]. Groeger JA, Zijlstra FRH, Dijk DJ, Sleep quantity, sleep difficulties and their perceived consequences in a representative sample of some 2000 British adults, *J. Sleep Res.* 13 (4) (2004) 359–371. [PubMed: 15560771]
- [29]. Lindberg E, Janson C, Gislason T, Bjornsson E, Hetta J, Boman G, Sleep disturbances in a young adult population: can gender differences be explained by differences in psychological status? *Sleep* 20 (6) (1997) 381–387. [PubMed: 9302720]
- [30]. Paul KN, Dugovic C, Turek FW, Laposky AD, Diurnal sex differences in the sleep-wake cycle of mice are dependent on gonadal function, *Sleep* 29 (9) (2006) 1211–1223. [PubMed: 17040009]

- [31]. Franken P, Dudley CA, Estill SJ, Barakat M, Thomason R, O'Hara BF, et al., NPAS2 as a transcriptional regulator of non-rapid eye movement sleep: genotype and sex interactions, *Proc. Natl. Acad. Sci. U. S. A* 103 (18) (2006) 7118–7123. [PubMed: 16636276]
- [32]. Ehlen JC, Hesse S, Pinckney L, Paul KN, Sex chromosomes regulate nighttime sleep propensity during recovery from sleep loss in mice, *PLoS One* 8 (5) (2013) e62205. [PubMed: 23658713]
- [33]. Koehl M, Battle SE, Turek FW, Sleep in female mice: a strain comparison across the estrous cycle, *Sleep* 26 (3) (2003) 267–272. [PubMed: 12749544]
- [34]. Schwierin B, Borbély AA, Tobler I, Sleep homeostasis in the female rat during the estrous cycle, *Brain Res.* 811 (1–2) (1998) 96–104. [PubMed: 9804908]
- [35]. Fang J, Fishbein W, Sex differences in paradoxical sleep: influences of estrus cycle and ovariectomy, *Brain Res.* 734 (1–2) (1996) 275–285. [PubMed: 8896835]
- [36]. Colvin GB, Whitmoyer DI, Lisk RD, Walter DO, Sawyer CH, Changes in sleep-wakefulness in female rats during circadian and estrous cycles, *Brain Res.* 7 (2) (1968) 173–181. [PubMed: 5689033]
- [37]. Pawlyk AC, Morrison AR, Ross RJ, Brennan FX, Stress-induced changes in sleep in rodents: models and mechanisms, *Neurosci. Biobehav. Rev* 32 (1) (2008) 99–117. [PubMed: 17764741]
- [38]. Mellman TA, Bustamante V, Fins AI, Pigeon WR, Nolan B, REM sleep and the early development of posttraumatic stress disorder, *Am. J. Psychiatry* 159 (10) (2002) 1696–1701. [PubMed: 12359675]
- [39]. Klein E, Koren D, Arnon I, Lavie P, Sleep complaints are not corroborated by objective sleep measures in post-traumatic stress disorder: a 1-year prospective study in survivors of motor vehicle crashes, *J. Sleep Res.* 12 (2003) 35–41. [PubMed: 12603785]
- [40]. Menz MM, Rihm JS, Salari N, Born J, Kalisch R, Pape HC, et al., The role of sleep and sleep deprivation in consolidating fear memories, *Neuroimage* 75 (2013) 87–96. [PubMed: 23501052]
- [41]. Graves LA, Heller EA, Pack AI, Abel T, Sleep deprivation selectively impairs memory consolidation for contextual fear conditioning, *Learn. Mem* 10 (3) (2003) 168–176. [PubMed: 12773581]
- [42]. Kobayashi I, Delahanty DL, Gender differences in subjective sleep after trauma and the development of posttraumatic stress disorder symptoms: a pilot study, *J. Trauma. Stress* 26 (4) (2013) 467–474. [PubMed: 23861181]
- [43]. Tang X, Orchard SM, Sanford LD, Home cage activity and behavioral performance in inbred and hybrid mice, *Behav. Brain Res.* 136 (2) (2002) 555–569. [PubMed: 12429418]
- [44]. Sanford LD, Yang L, Tang X, Influence of contextual fear on sleep in mice: a strain comparison, *Sleep* 26 (5) (2003) 527–540. [PubMed: 12938805]
- [45]. Sanford LD, Tang X, Ross RJ, Morrison AR, Influence of shock training and explicit fear-conditioned cues on sleep architecture in mice: strain comparison, *Behav. Genet* 33 (1) (2003) 43–58. [PubMed: 12645821]
- [46]. Byers SL, Wiles MV, Dunn SL, Taft RA, Mouse estrous cycle identification tool and images, *PLoS One* 7 (4) (2012) e35538. [PubMed: 22514749]
- [47]. Nishida M, Pearsall J, Buckner RL, Walker MP, REM Sleep, prefrontal theta, and the consolidation of human emotional memory, *Cereb. Cortex* 19 (5) (2009) 1158–1166. [PubMed: 18832332]
- [48]. Tang X, Sanford LD, Telemetric recording of sleep and home cage activity in mice, *Sleep.* 25 (6) (2002) 677–685. [PubMed: 12224847]
- [49]. Radulovic J, Kammermeier J, Spiess J, Relationship between Fos production and classical fear conditioning: effects of novelty, latent inhibition, and unconditioned stimulus preexposure, *J. Neurosci* 18 (18) (1998) 7452–7461. [PubMed: 9736664]
- [50]. Wellman LL, Yang L, Tang X, Sanford LD, Contextual fear extinction ameliorates sleep disturbances found following fear conditioning in rats, *Sleep* 31 (7) (2008) 1035–1042. [PubMed: 18652099]
- [51]. Barha CK, Dalton GL, Galea L, Low doses of 17 alpha-estradiol and 17 bold italic beta-estradiol facilitate, whereas higher doses of estrone and 17 alpha-and 17 bold italic beta-estradiol impair, contextual fear conditioning in adult female rats, *Neuropsychopharmacology* 35 (2010) 547–559. [PubMed: 19847162]

- [52]. Prendergast BJ, Onishi KG, Zucker I, Female mice liberated for inclusion in neuroscience and biomedical research, *Neurosci. Biobehav. Rev* 40 (2014) 1–5. [PubMed: 24456941]
- [53]. Becker JB, Prendergast BJ, Liang JW, Female rats are not more variable than male rats: a meta-analysis of neuroscience studies, *Biol. Sex Differ* 7 (2016) 34. [PubMed: 27468347]
- [54]. Itoh Y, Arnold AP, Are females more variable than males in gene expression? Meta-analysis of microarray datasets, *Biol. Sex Differ* 6 (1) (2015) 18. [PubMed: 26557976]
- [55]. Bush DEA, Sotres-Bayon F, LeDoux JE, Individual differences in fear: isolating fear reactivity and fear recovery phenotypes, *J Traum Stress* 20 (4) (2007) 413–422.
- [56]. Graham BM, Richardson R, Individual differences in the expression of conditioned fear are associated with endogenous fibroblast growth factor 2, *Learn. Mem* 23 (1) (2015) 42–45. [PubMed: 26670186]
- [57]. Tang X, Yang L, Sanford LD, Interactions between brief restraint, novelty and footshock stress on subsequent sleep and EEG power in rats, *Brain Res.* 1142 (2007) 110–118. [PubMed: 17300767]
- [58]. Hutchison IC, Rathore S, The role of REM sleep theta activity in emotional memory, *Front. Psychol* 6 (2015) 1439. [PubMed: 26483709]
- [59]. Boyce R, Glasgow SD, Williams S, Adamantidis A, Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation, *Science* 352 (6287) (2016) 812–816. [PubMed: 27174984]
- [60]. Marrosu F, Portas C, Mascia MS, Casu MA, Fà M, Giagheddu M, et al., Microdialysis measurement of cortical and hippocampal acetylcholine release during sleep-wake cycle in freely moving cats, *Brain Res.* 671 (2) (1995) 329–332. [PubMed: 7743225]
- [61]. Li S, Topchiy I, Kocsis B, The effect of atropine administered in the medial septum or hippocampus on high-and low-frequency theta rhythms in the hippocampus of urethane anesthetized rats, *Synapse* 61 (6) (2007) 412–419. [PubMed: 17372965]
- [62]. Masuda J, Mitsushima D, Funabashi T, Kimura F, Sex and housing conditions affect the 24-h acetylcholine release profile in the hippocampus in rats, *Neuroscience* 132 (2) (2005) 537–542. [PubMed: 15802204]
- [63]. Mitsushima D, Takase K, Takahashi T, Kimura F, Activational and organisational effects of gonadal steroids on sex-specific acetylcholine release in the dorsal hippocampus, *J. Neuroendocrinol* 21 (4) (2009) 400–405. [PubMed: 19356199]
- [64]. Rolls A, Makam M, Kroeger D, Colas Dd L Lecea De, Heller HC, Sleep to forget: interference of fear memories during sleep, *Mol. Psychiatry* 18 (11) (2013) 1166. [PubMed: 24081009]

Fig. 1. Fear memory recall rates for each group. Error bars indicate 95% CI. *Significant difference between groups; $p < 0.05$.

Fig. 2.

Parameters for overall sleep (A and B) and NREM sleep (C and D) across time points (mean $\pm$ s.e.m.). Significant differences between groups; *p 0.05, **p 0.01, ***p 0.001.

Fig. 3. REM sleep parameters across time points (mean $\pm$ s.e.m.). Significant differences between groups; *p 0.05, **p 0.01, ***p 0.001.

Fig. 4.

Associations between sleep and fear memory recall rate. The lines in graphs A and C, which represent associations between sleep and fear memory recall for all subjects during the dark period, indicate a main effect of sleep; interactions between sleep and group were not significant. The lines in graphs B and D indicate the associations for each group during the light period; both main effects of sleep and interactions between sleep and group were significant (see Table 2).

Table 1

Time course of the experimental procedures for all animal groups. P-females received the footshock and fear recall sessions on the day of proestrus; M-females received both procedures on the day of metestrus.

Group	n	Baseline Sleep Recording	Footshock Session	Post-shock Sleep Recording	Fear Recall Session
Male	7	5–10 days	Day 6–11	9–13 days	Post-shock Day 9–13
P-Female	8	1–2 cycles	Day 6–11 (Proestrus)	9–13 days	Post-shock Day 9–13 (Proestrus)
M-Female	7	1–2 cycles	Day 6–11 (Metestrus)	9–13 days	Post-shock Day 9–13 (Metestrus)

Table 2

Results of the final model of the hierarchical regression analyses predicting fear memory recall rate (%) from group, sleep, and group by sleep interactions. Significant differences: * $p < .05$, ** $p < .01$, *** $p < .001$. Note: ^aMale group is the reference group. +Analysis was not performed due to non-normal data.

Sleep parameter→ Predictor↓	Light period				Dark period			
	NREM %	REM %	# of REM episodes	Avg. REM duration	NREM %	REM %	# of REM episodes	Avg. REM duration
	<i>b</i>	<i>b</i>	<i>b</i>	<i>b</i>	<i>b</i>	<i>b</i>	<i>b</i>	<i>b</i>
Post-shock Day 1 Sleep								
P-females ^a	+	25.1	24.5	26.7	20.3	27.1	26.3	22.6
M-females ^a	+	49.6**	48.8**	51.2**	44.8*	57.0**	48.5*	42.5**
Sleep	+	12.9	<0.1	23.6*	-12.1	9.4	4.4	30.8**
P-females x Sleep	+	-9.5	10.4	-30.3*	26.3	4.1	4.3	-20.8
M-females x Sleep	+	-9.4	0.1	-11.4	24.5	-0.6	-8.6	-17.1
Adjusted R ²	+	.22	.20	.43*	.30	.26	.20	.55**
Post-shock Day 2 Sleep								
P-females ^a	19.9	24.6	23.9	15.1	4.8	21.2	17.5	33.0
M-females ^a	44.4*	48.4**	48.8**	37.2**	35.1*	44.6*	40.6*	57.3**
Sleep	-7.3	10.2	1.2	44.9***	-23.4**	-6.7	-10.7	-12.4
P-females x Sleep	0.3	-9.9	3.8	-52.8***	9.5	-0.3	2.3	8.9
M-females x Sleep	6.9	-8.8	-1.4	-38.8**	23.7	2.6	5.2	4.0
Adjusted R ²	.20	.20	.17	.63***	.49**	.21	.26	.24