Metagenome sequencing of fingermillet-associated microbial consortia provides insights into structural and functional diversity of endophytes

M. K. Prasannakumar¹ · H. B. Mahesh² · Radhika U. Desai¹ · Bharath Kunduru¹ · Karthik S. Narayan³ · Kalavati Teli¹ · M. E. Puneeth¹ · R. C. Rajadurai⁴ · Buella Parivallal¹ · Gopal Venkatesh Babu³

Received: 21 May 2019 / Accepted: 3 December 2019 / Published online: 10 December 2019
© King Abdulaziz City for Science and Technology 2019

Abstract

Endophytes confer unique ecological advantages to their host plants. In this study, we have characterized the diversity of endophytic consortia associated with the GPU-28 (GPU) and Udumallige (UM) finger millet varieties, which are resistant and susceptible to the blast disease, respectively. Whole genome metagenome sequencing of GPU and UM helped to identify 1029 species (includes obligate endophytes) of microbiota. Among them, 385 and 357 species were unique to GPU and UM, respectively. Remaining 287 species were common to both the varieties. Actinobacteria and other plant-growth promoting bacteria were abundant in GPU as compared to UM. Functional annotation of genes predicted from genomes of endophytes associated with GPU variety showed that many genes had functional role in stress response, secondary metabolism, aromatic compounds, glutathione, and cysteine synthesis pathways as compared to UM. Based on in vitro and in planta studies, *Bacillus cereus* and *Paenibacillus* spp. were found to be effective in suppressing the growth of blast disease pathogen *Magnaporthe grisea* (strain MG03). In the future, these strains could serve as potential biocontrol agents to reduce the incidence of blast disease in finger millet crop.

Keywords Endophytes · Metagenome sequencing · Blast disease · *Magnaporthe* · *Finger millet*

Introduction

Finger millet [*Eleusine coracana* (L.) Gaertn. subsp. *coracana*] is a highly nutritive cereal crop of the family Poaceae and is an allotetraploid (AABB) species with $2n = 4x = 36$ chromosomes. It is an important small millet of tropics and plays a prominent role in the subsistence agriculture of arid and semi-arid areas of Africa and South Asia. Finger millet is a hardy crop, which can be cultivated optimally under low-moisture regimes without affecting the net productivity (Bhatt et al. 2011). It is cultivated in more than 25 countries of Africa and Asia and occupies about 12% of the total global millet area (Vetriventhan et al. 2015). Finger millet is a rich source of essential amino acids like tryptophan and methionine and is also abundant in calcium, iron, and fiber. Its low glycemic index (GI) value and superior malting quality makes it a preferred choice of food for diabetics. Owing to its hardy nature and superior nutritional characteristics, finger millet assumes a significant role in addressing the nutritional infirmity in dryland areas of tropics.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s13205-019-2013-0>) contains supplementary material, which is available to authorized users.

✉ M. K. Prasannakumar
babu_prasanna@rediffmail.com

✉ H. B. Mahesh
maheshhbg@gmail.com

¹ Plant Pathogenomic Laboratory, Department of Plant Pathology, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India

² Department of Genetics and Plant Breeding, College of Agriculture, Mandya, Karnataka, India

³ Department of Centre for Advanced Study in Botany, University of Madras, Chennai, India

⁴ AgriGenome Laboratory, Cochin, Kerala, India

Blast disease is a major constraint in finger millet cultivation as this crop is highly susceptible to the disease. The blast disease in finger millet is caused by *Magnaporthe grisea* (Hebert) Barr. (Anamorph: *Pyricularia grisea* Sacc.), which is highly devastating and adversely affects the finger millet productivity. The pathogen infects all aerial parts of finger millet plant causing leaf, neck, and finger blast resulting in yield losses, which may often exceed 50% (Sastri 1950; Esele 2002). Although fungicides like carbendazim, mancozeb, and tricyclazole are effective in controlling blast disease (Netam et al. 2014), synthetic fungicides are hazardous to human health and also leaves a negative impact on the environment (Budnik and Baur 2009). Repeated application of chemicals may also induce resistance in the pathogen. Moreover, chemical control in blast disease management in finger millet is not viable for resource-poor farmers and in organic production systems. The pathogen is highly variable and breeding for resistant varieties could only control disease in the short run. In this perspective, it is desirable to identify more eco-friendly, economical, and farmer-friendly approaches to control the blast disease in finger millet, which could be equally applied to other crops to control various diseases.

The use of biological agents in plant disease management in agricultural crop production is already in practice. However, in recent decades, the novel research on biological control of plant diseases investigated the role of a particular class of microbes that colonize the internal tissues of the host plant, referred to as endophytes, and hypothesized that they confer host plant resistance against fungal pathogens (Arnold et al. 2003; Gunatilaka 2006; Mousa and Raizada 2013). These endophytes do not cause any apparent disease symptoms by themselves (Braun and Hirsch 1992). However, considerable evidence has been accumulated supporting the beneficial role of endophytes in host plants in promoting plant growth (Verma et al. 2001; Sziderics et al. 2007) improved nutrient acquisition from soil (Upson et al. 2009) abiotic stress tolerance (Rodriguez et al. 2008), and enhanced production of plant defensive compounds by up-regulating the gene expression of biochemical pathways involved in plant defense mechanisms (Gond et al. 2015). Various studies have reported several mechanisms by which the host inhabiting endophytes confer host plant resistance against multiple plant pathogens. They include competition for space and nutrition (Bolwerk et al. 2005), induction of host resistance genes (Waller et al. 2005), promotion of plant growth and physiology (Chaturvedi and Singh 2016), hyperparasitism, predation (Grosch et al. 2006; Dutta et al. 2014), and stimulation/production of natural biostatic and biocide compounds (Shankar et al. 1994).

In the present study, we investigated the diversity and abundance of different endophytic communities colonizing the host tissues of two different indigenous finger millet

genotypes viz. GPU-28 (GPU) and Udurumallige (UM). The GPU is highly resistant to the blast disease with < 2% incidence of both neck and finger blast (Nagaraja et al. 2008), whereas UM is highly susceptible (Shirke et al. 2016) to the disease. The whole genome metagenome (WMG) sequencing approach was used in this study to understand the diversity and function of microbial niches, which could not be addressed using conventional methods (Handelsman 2005; Schloss and Handelsman 2005; Ravin et al. 2015). This metagenomic study also helped to identify the putative endophytes contributing specific to blast disease resistant variety GPU. Endophytes isolated from these varieties were subsequently evaluated in vitro and in planta for their antagonistic effect on *M. grisea* strain MG03, which was whole genome characterized (Shirke et al. 2016).

Materials and methods

Leaf sample collection

Three healthy plants each of GPU and UM cultivars were collected randomly 21 days post sowing from the experimental plots in the Department of Plant Pathology, GKVK, UAS, Bengaluru, India (13° 05" N Latitude, 77° 34" E Longitude). The seedlings were raised under natural field conditions and the use of any plant protection chemicals was strictly avoided during the study. The collected samples were sealed in sterile bags maintained at 4 °C and processed immediately upon return to the laboratory.

Surface sterilization and genomic DNA extraction

The plant samples were surface sterilized and washed three times in sterile double-distilled water. An aliquot (100 µL) from final wash was plated onto the potato dextrose agar (PDA) and nutrient agar (NA) media to confirm the efficiency of surface sterilization (epiphytic check). The inoculated plates were maintained at 28 ± 2 °C for 3 days and only the tissue samples with no growth on epiphytic check plates were used for isolation of endophytes. The surface-sterilized leaf, stem and root fragments were processed individually as per Sessitch et al. (2012) for extraction of endophyte cells from plant tissues. Extracted endophyte cells were collected separately and utilized for DNA extraction using the DNASure® Plant Mini Kit (Genetix Biotech Asia Pvt. Ltd.) as per the manufacturer's instructions. The quality and quantity of DNA were assessed using both the Qubit™ dsDNA BR Assay Kit (Thermo Fisher Scientific, USA) and Nanodrop® (DS-11 FX+, DeNovix® Spectrophotometer/Fluorometer, DeNovix Inc., USA). The DNA was pooled and whole genome metagenome sequencing was performed.

Metagenome library preparation and sequencing

NEBNext® Ultra™ DNA Library Prep Kit for Illumina® (NewEngland BioLabs® Inc., USA) was used to prepare the whole genome libraries of GPU and UM from the DNA samples. The genomic DNA was fragmented using Covaris. The end-repaired samples were ligated with adapters and size selected. The library was further enriched using nine cycles of PCR. After purification, using Agencourt® AMPure® XP beads (Beckman Coulter), the library was quality checked using the Agilent 4200 TapeStation System (Agilent Technologies Inc., USA) and the average library size was confirmed to be about > 600 bp. Final libraries were sequenced in AgriGenome Labs (Cochin, Kerala, India) using HiSeq 2500 (Illumina Inc., USA).

Sequence data quality check, pre-processing and assembly

The raw fastq paired-end reads were quality checked using the FastQC program (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and the low-quality sequence bases (PHRED score $Q \leq 20$) were trimmed. Adapter dimers, GC-content, overlapping sequences, PCR duplicates, ambiguous bases, homopolymers, and chimeras were checked to avoid errors in the assembly. The Illumina adapters at 5' and 3' ends of sequence reads were trimmed using the Cutadapt tool version 1.8.1 (Martin 2011). The pre-processed reads were assembled using the Ray Meta metagenome assembler with default parameters (Boisvert et al. 2012). The contigs with a length < 150 bp were discarded to reduce false positives during downstream analyses like gene prediction and annotation.

Prediction of open reading frames and functional annotation

Assembled contigs were submitted to the MetaGeneAnnotator (Noguchi et al. 2008) to predict open reading frames (ORFs). The read depth of all predicted ORFs was quantified to reveal the taxonomic abundance. Functional annotation was performed by aligning ORF sequences against the NCBI-nr protein database using the DIAMOND tool version 0.7.9.58 (Buchfink et al. 2015). The DIAMOND output files were analyzed using MEGAN5 software (Huson et al. 2011). The gene ontology (cellular, biological and molecular functions) was assigned for annotated ORFs based on SEED (Overbeek et al. 2014) classification methods available in MEGAN5. The key genes involved particularly in hydrocarbon degradation were searched from the annotation results. All the functionally annotated ORFs were assigned the taxonomic hierarchy from phylum to species level (including Bacteria, Archaea, Eukaryota, Viruses, and unclassified

sequences). Rarefaction analysis was performed using the Lowest Common Ancestor (LCA) algorithm by binning reads to taxa based on their BLAST-hits and rooted taxon tree was constructed based on unique operational taxonomic units (OTUs) used in the analysis. Taxonomic profiling of endophytes was performed using the NCBI taxonomy data sets and phylogenetic trees at species level were generated based on the neighbor-joining method using MEGAN (Meta Genome ANalyzer) software. Figtree (version 1.4.3) software was used to construct the phylogenetic tree.

Comparative metagenome analysis

For functional profiling of genes associated with endophytes, SEED subsystems and KEGG databases were applied (Kanehisa et al. 2012). The clustered OTU BIOM table was normalized using the `normalize_by_copy_number.py` script and processed with the `predict_metagenomes.py` script. The predicted metagenome was exported to BIOM format for further statistical analysis. Statistical Analysis of Metagenomic Profiles (STAMP) software version 2.1.3 was used for studying the taxon difference. Potential gene comparison between samples was performed using two-sided Fisher's exact test and Storey's false discovery rate (FDR) method (Parks and Beiko 2010). *p* values were adjusted using Bonferroni correction. Comparative metagenomics such as taxonomic-to-phenotypic mapping and multivariate analysis including principal component analysis (PCA), clustering, and classification were performed using the METAGENassist web server by uploading BIOM files. R-package was used for generating the *p* value significant plots (Arndt et al. 2012).

Isolation and morphological characterization of endophytes

After epiphytic check, the surface-sterilized plant samples of GPU and UM were used to isolate endophytes using the serial dilution (Hung and Annapurna 2004) and tissue bits (Gayathri and Muralikrishnan 2013) methods. In serial dilution method, 100 µL aliquot from each dilution (10^{-3} and 10^{-4} dilutions for fungi; 10^{-5} to 10^{-7} dilutions for bacteria) of tissue extracts crushed in 0.9% NaCl solution were uniformly spread on potato dextrose agar (PDA) and nutrient agar (NA) media plates and incubated at 28 ± 2 °C to obtain colony growth. In case of the tissue bits method, the surface-sterilized leaf, stem, and root fragments were first imprinted on the control plates (epiphytic check) (Schulz et al. 1998) containing PDA and NA media for 5 s. Later, tissue fragments were removed and placed on fresh culture plates in identical positions to the control plates. Four bits were placed on medium in two replicates and incubated at 28 ± 2 °C until endophyte growth appeared on the periphery

of tissue fragments. The culture plates showing no growth on control plates were considered for further use.

Following overnight incubation, morphologically distinct bacterial colonies were transferred to fresh NA medium plates using the quadrant streak plate method. The pure cultures were stored in NA slants and glycerol stocks for further use. Similarly, pure cultures of distinct fungal endophytes were obtained using PDA medium plates and subsequently preserved in PDA slants and glycerol stocks.

16S rDNA and 18S rDNA analysis of bacterial and fungal endophytes

The genomic DNA of bacterial endophytes was extracted using AMPurE Bacterial Genomic DNA Mini Spin kit (Amnion Biosciences, India). The isolates were first inoculated on to a nutrient broth (NB) medium and incubated overnight in an orbital shaker at 150 rpm at 28 ± 2 °C. The bacterial cells were harvested from broth by centrifugation for 2 min at $14,000 \times g$ and the pelleted cells were processed for DNA isolation as per manufacturer's instructions. RNA contamination was removed by treating with 5 µL of RNase A (20 mg/mL) (HIMEDIA®, India) solution. The DNA quality was checked on both 0.8% agarose gel and using Nanodrop®. The isolated bacterial genomic DNA was used as template to amplify the 16S rDNA region using universal primers for bacterial endophytes' identification and characterization. Each 25-µL reaction mixture contained 2.5 µL (50 ng/µL) of template gDNA, 1.25 µL (10 pmol/µL) each of forward (5' GAGTTTGATCCTGGCTCAG 3') and reverse (5' AGAAAGGAGGTGATCCAGCC 3') primers, and 17.5 µL EmeraldAmp® GT PCR Master Mix (TaKaRa Bio Inc., Japan). Final volume was made up to 25 µL using nuclease-free water. The thermal cycling program on the Eppendorf Mastercycler® pro (Eppendorf AG, Hamburg, Germany) included an initial denaturation at 94 °C for 3 min followed by 35 cycles of 94 °C for 45 s, 58 °C for 1 min, and 72 °C for 90 s and a final extension at 72 °C for 10 min.

Genomic DNA from fungal isolates was extracted as per Moller Bahnweg et al. (1992). The quality of extracted DNA was assessed on a 0.8% agarose gel and quantified using a Nanodrop®. The 18S rDNA region in the fungal genome was amplified using ITS2 (5' GCTGCGTTCTTCATCGAT GC 3') and ITS4 (5' TCCTCCGCTTATTGATATGC 3') universal primers (White et al. 1990) for fungal endophytes' characterization. The composition of the PCR mixture was same as mentioned earlier and the amplification program included an initial denaturation at 94 °C for 4 min followed by 35 cycles of 94 °C for 30 s, 55 °C for 30 s, and 72 °C for 1 min, and a final extension at 72 °C for 5 min. The PCR amplicons of 16S rDNA (for bacterial endophytes) and 18S rDNA (for fungal endophytes) were sequenced by Sanger dideoxy sequencing technology. The 16S rDNA and 18S

rDNA sequences were searched with those on the NCBI database using the BLAST tool (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) for species identification.

In vitro and in planta evaluation of endophytes against rice blast pathogen *Magnaporthe grisea*

The in vitro interaction of different endophyte strains against MG03 (*Magnaporthe* strain, Shirke et al. 2016) was studied using dual-culture assay. The MG03 strain was inoculated on one side of Luria–Bertani (LB) agar medium and a loopful of bacterial endophyte culture was streaked/inoculated on the other end of culture plate (Singh and Varma 2015). The fungal endophytes were inoculated as per Benfradj et al. (2016). A culture plate with MG03 alone was maintained as a control. For the positive control, culture plate with actively growing MG03 was inoculated with *Bacillus velezensis* strains A6 and P42, a known strains antagonistic to *M. grisea* (Lim et al. 2017). Three replicates were maintained for each endophyte as well as for positive and negative controls, and the culture plates were incubated at 28 ± 2 °C for 5 days. The percentage inhibition of pathogen was calculated using the formula given by Vincent (1947):

$$I = \frac{(C - T) \times 100}{C},$$

where, I = percentage inhibition; C = radial growth in control plate (mm); T = radial growth in treatment (mm) plate.

The seeds of GPU and UM were bio-primed with endophytes to study their potential role in suppressing the blast disease in finger millet *in planta*. Endophyte-free seedlings of GPU and UM were obtained as per the previously established procedures (Pandey et al. 2016, 2018). The seeds were thoroughly washed in sterile ddH₂O and incubated for 24 h in 0.2% benomyl and 70 ppm each of streptomycin sulphate and tetracycline solutions to eliminate the seed-borne endophytes. The treated seeds were washed five times with sterile ddH₂O and an aliquot of seeds were homogenized using pestle and mortar. The homogenate was plated on PDA and NA agar media plates and incubated at 28 ± 2 °C for 7 days. When no microbial growth was observed on the incubated plates, the remaining treated seeds were considered endophyte-free and used for bio-priming. For bio-priming, all the bacterial and fungal endophytes were individually inoculated into sterile NB and potato dextrose broth (PDB) media and incubated for 24 h and 7 days, respectively. For each endophyte to be treated, 50 surface-sterilized treated seeds each of GPU and UM were soaked in these pre-inoculated NB and PDB media for 24 h. An equal number of chemical-treated seeds of GPU and UM soaked in sterile broth media was kept as a control (C_0). The soaked seeds were air-dried under aseptic conditions and later sown in

pots (15 cm diameter) filled with sterile growth medium and maintained in a greenhouse. Each treatment was replicated twice and the temperature was maintained at 28 ± 2 °C during day and 25 ± 2 °C at night. The plants were transferred to the inoculation chamber 1 day prior to pathogen inoculation for acclimatization (28 ± 5 °C day and 25 ± 5 °C night temperature, and relative humidity > 90%). The seeds obtained from C_0 were sown as a control.

Pathogenicity test and disease scoring

Sporulation was induced in MG03 culture plates and the spore suspension was sprayed onto 20-day-old GPU and UM seedlings maintained in growth chamber (Mahesh et al. 2016). The treated seedlings were further covered with polythene bags for 1 week to maintain high humidity (~94%) conducive for disease development. The polythene bags were removed on sixth day after inoculation (DAI) and seedlings were scored for disease severity as per the IRRI SES scale (Table S1) at 30 DAI. The percent disease severity (PDI) was calculated as per Chiang et al. (Chiang et al. 2017):

$$PDI = \frac{\sum \text{Sum of all numerical scores} \times 100}{\text{Total number of leaflets observed} \times \text{maximum score}}$$

After disease development, the infected leaves were collected, surface sterilized, and small infected leaf bits were placed on PDA medium to re-isolate the test pathogen. The culture plates were incubated at 28 ± 2 °C to promote the growth of test pathogen. Later, the pathogen was purified and its identity with original culture of MG03 was confirmed based on morphological characters. Thus, Koch postulates were established.

Statistical analysis

The data were analyzed using the appropriate statistical tools and percent values were arcsine transformed (Gomez et al. 1984) before subjecting the data to analysis of variance (ANOVA). The critical difference (CD) values were calculated at 5% probability level.

Results

Metagenome sequencing, assembly and identification of endophytes in finger millet cultivars

The WMG sequencing of the field samples of GPU and UM on Illumina HiSeq platform yielded 4,307,574 and 4,438,968 raw reads, respectively. We assembled the high-quality reads into 7,76,412 and 4,95,014 contigs (GPU and

UM, respectively) using Ray Meta de novo assembler (version 2.3.1) and only contigs of size > 150 bp were retained for downstream analysis (Table 1). Totally, 5,96,312 and 3,74,417 contigs having length > 150 bp among which only 5,62,769 (94.37%) and 3,50,015 (93.48%) contigs exclusively belonging to bacteria and fungi were retained for downstream analyses in GPU and UM, respectively. The contigs not classified within bacterial and fungal taxa were excluded from downstream analysis. The rarefaction curves (Fig. S1) indicated that GPU had high species richness as compared to UM.

Taxonomic profiling of the metagenomes of GPU and UM revealed that bacterial classes, Alphaproteobacter, showed eight genera of potential biocontrol agents that were seen in both GPU and UM; however, GPU showed more species richness (Fig. 1). Bacteria were the most abundant domain among the endophytic consortia colonizing both the genotypes followed by the domains of eukaryotes and archaea. Among the bacterial domain, the phylum Proteobacteria was the most abundant in both GPU and UM followed by Actinobacteria and Bacteroidetes. The remaining phyla viz., Cyanobacteria (blue green algae), Ascomycota and Basidiomycota (fungi) were less prominent in the metagenomes of GPU and UM. The domain Archaea in both the samples was represented by two phyla viz, Euryarchaeota and Thaumarchaeota (Fig. 2).

Out of the total 1029 species of microbiota collectively identified in GPU and UM, 287 species were identified as common, while 385 and 357 species were specific to GPU and UM, respectively. The plant growth-promoting bacteria like *Sorangium cellulosum*, *Lechevalieria aerocolonigenes*, *Stenotrophomonas maltophilia*, and *Streptomyces* spp. were widely distributed in both GPU and UM, but their relative abundance was much higher in GPU as compared to UM (Fig. 3). The genera *Streptomyces* in both UM and GPU was dominated by the species, *Streptomyces galbus*, which is a strong producer of antifungal compounds (Paul and Banerjee 1983) and this organism was more prevalent in GPU than UM. The other species in this group with documented antifungal activity and identified exclusively in GPU were *S. griseus* (Anitha and Rebeeth 2009; Rabeeth et al. 2011), *S. halstedii* (Frandsen et al. 2000), *S. anulatus* (Mander et al. 2016), and *S. cluvuligerus* (Doran et al. 1990). Bacilli were

Table 1 Statistics of metagenome assembly

Category	GPU	UM
Raw reads	4,307,574	4,438,968
Total Contigs	7,76,412	4,95,014
Contigs < 150 bp	1,80,100	1,20,597
Contigs 150–500 bp	5,95,995	3,71,101
Contigs > 500 bp	317	3316

Fig. 1 Phylogenetic diversity of metagenomes identified in the two finger millet genotypes. The circular phylogenetic trees (left—GPU and right—UM) representing various taxa identified from metage-

nome datasets of GPU and UM. The genus highlighted in colors is reported as biocontrol agents against plant pathogens. The classes are represented in the outer ring

another important group which are important in the context of this study. The species, *Bacillus cereus*, *Paenibacillus mucilaginosus*, and *Lactobacillus* sp. were identified exclusively in GPU and these species were reported to produce siderophores which promote plant growth and also possess biocontrol activity against phytopathogenic fungi (Tokpah et al. 2017; Amruta et al. 2018).

Functional annotation of assembled metagenome contigs

The SEED classification of annotated contigs affiliated to endophytes helped to identify 23 major Level-1 sub-systems in the metagenome samples of GPU and UM (Fig. 4). Carbohydrates, protein metabolism, DNA metabolism, and respiration were among the major Level-1 sub-systems identified in both the samples. The other important Level-1 sub-systems relevant to the present study were stress response, clustering-based sub-systems, sulfur metabolism, phosphorous metabolism, metabolism of aromatic compounds, and secondary metabolism. Comparative analysis of these functional sub-systems (Fig. S2) in GPU and UM showed that the reads encoding stress response (GPU = 76; UM = 63), clustering-based sub-systems (GPU = 81; UM = 70), sulfur metabolism (GPU = 29;

UM = 22), phosphorous metabolism (GPU = 22; UM = 12), metabolism of aromatic compounds (GPU = 63, UM = 48), and secondary metabolism (GPU = 11; UM = 12) were quantitatively higher in GPU as compared to UM (Fig. 4).

The metabolic functions of genes present in the endophytic microbiota were identified through KEGG analysis and revealed seven major metabolic and enzyme pathways at the metagenome level (Fig. S3). The contigs pertaining to metabolism (carbohydrate metabolism, energy metabolism, lipid metabolism, nucleotide metabolism, amino acid metabolism, and biosynthesis of secondary metabolites), genetic information processing, and environmental information processing were the major categories detected in both the cultivars. However, contigs belonging to many metabolism pathways hold a significant importance in key processes involved in disease resistance mechanism. Among the different metabolic pathways, reads belonging to cysteine synthase (GPU = 11; UM = 7) and glutathione synthase (GPU = 4 and UM = 1) bearing ECs '2.5.1.47' and '6.3.2.3,' respectively, were more in GPU as compared to UM. Glutathione synthase (Dubreuil-Maurizi and Poinssot 2012) and cysteine synthase (Álvarez et al. 2012) enzymes are reported to play an important role in plant defense mechanism.

Fig. 2 Relative abundance of different taxonomic groups identified in GPU and UM samples. Different phyla were mapped in the pie chart as per their relative abundance in anti-clock wise pattern

Morphological characterization of culturable endophytes

Using serial dilution and tissue bits isolation methods, a total of 75 bacterial and 5 fungal endophytes were isolated from the leaf, stem, and root tissues of GPU and UM. Out of 75 bacterial isolates, 52 and 23 strains were obtained from GPU and UM, respectively (Table 2). Maximum isolates were obtained from the leaf (22) tissues of GPU and the lowest from stem (6) tissues of UM (Fig. 5).

A rich microbial diversity was observed among the bacterial isolates with respect to their colony morphology, color, shape, and size (Table S2). The gram staining procedure differentiated the 75 bacteria into 45 Gram negative (G^{-ve}) and 30 Gram positive (G^{+ve}) isolates. Among them, 19 and 11 G^{+ve} ; 33 and 12 G^{-ve} isolates were obtained from GPU and UM, respectively. The oxidase test indicated that the majority of isolates were anaerobic (negative

reaction). In GPU, four and in UM, three endophytes were found to be aerobic (positive reaction).

Based on the morphological characteristics like mycelium type, spore characteristics and colony morphology, five fungal endophytes were found belonging to genera viz. *Lasioidiplodia*, *Talaromyces*, *Phoma*, and *Penicillium* (Table S3).

Molecular characterization of culturable endophytes

The amplified products of 16S rDNA (bacteria) and 18S rDNA (fungi) were sequenced and based on maximum query coverage, the species were identified and accession numbers were obtained from NCBI (Table S4). Based on molecular phylogeny, the 75 bacterial endophytes were assigned to 12 genera among which, six, four, and two genera were associated to Gammaproteobacteria, Bacilli, and Actinobacteria, respectively (Table 2).

Fig. 3 Comparative distribution of top 33 species identified in the metagenome of GPU and UM. The relative abundance is mapped based on the number of reads pertaining to each species identified in

the two samples. The two icons on each radius of the circular chart correspond to the respective genotypes indicated in the key

In GPU, the class Gammaproteobacteria (63.46%) was abundantly distributed followed by Bacilli (32.69%). The genera Cronobacter, Enterobacter, Escherichia, Pantoea, Acetobacter, and Pseudomonas were identified in Gammaproteobacteria, whereas Bacillus, Paenibacillus, and Lactobacillus were found among the Bacilli group. In case of UM, Bacilli (52.17%) was the predominant taxa followed by Gammaproteobacteria (47.83%). Three genera viz. Enterobacter, Pantoea, and Pseudomonas were identified in Gammaproteobacteria, while Bacillus and Lysinibacillus were detected in Bacilli group. Among the different culturable endophytes isolated in the present study, we found that the *Enterobacter* spp. and *Bacillus* spp. were the dominant bacterial taxa colonizing GPU and UM (Table 2).

Based on the maximum query coverage value and percentage identity, the fungal isolates were identified as *Penicillium polonicum*, *Lasiodiplodia theobromae*, *Talaromyces trachyspermus*, and *Phoma* sp. Among the five fungal genera identified, only *Phoma* was found common to both GPU and UM (Table S4).

In vitro study of interaction between endophytes and *M. grisea*

The dual plate assay (Fig. 6) revealed that, among 56 endophytes (52 bacteria and 4 fungi) isolated from GPU, 12 isolates inhibited the growth of *M. grisea* (strain MG03) by more than 50% (Table 3). The positive control, *B. velezensis* strains P42 and A6 inhibited the MG03 by 78.83% and 80%, respectively. The strains, *B. cereus* GPUL8 (71.93%), *Bacillus licheniformis* GPUL23 and *B. cereus* GPUR10 (66.67% each) inflicted maximum percentage inhibition (PI) on MG03. The least inhibitory effect on the pathogen was observed with the strain *Enterobacter aerogenes* GPUL5 (20.83%). Similarly, among the 24 endophytes (23 bacteria and one fungi) isolated from UM, seven isolates inhibited the growth of MG03 by more than 50%. The strain, *Bacillus mojaviensis* UMR9 imposed maximum inhibitory effect (65%) on MG03 followed by the strains *Bacillus* sp. UML10, *Bacillus* sp. UMS5, and *Pseudomonas aeruginosa* UML11 (61.67% each). The strains *Enterobacter mori* UML4 and *Enterobacter* sp.

Fig. 4 SEED classification-based functional annotation of contigs of GPU and UM endophytes. Relative abundance of contigs of GPU and UM endophytes affiliated to 23 functional categories identified through SEED classification. The cofactors like vitamins, prosthetic

groups, and pigments are represented by asterisk. The phages like prophages, transposable elements, and plasmids are represented by double asterisk

Table 2 Bacterial genera identified in GPU and UM cultivars

S. no.	Class	Genus	Genotypes	
			GPU	UM
1	Gammaproteobacteria (Phylum: Proteobacteria)	Acenitobacter	7 (13.46%)	–
2		Cronobacter	1 (1.92%)	–
3		Enterobacter	21 (40.38%)	7 (30.43%)
4		Escherichia	1 (1.92%)	–
5		Pantoea	2 (3.84%)	2 (8.70%)
6	Bacilli (Phylum: Firmicutes)	Pseudomonas	1 (1.92%)	2 (8.70%)
7		Bacillus	14 (26.92%)	10 (43.78%)
8		Paenibacillus	2 (3.84%)	–
9		Lysinibacillus	–	2 (8.70%)
10		Lactobacillus	1 (1.92%)	–
11	Actinobacteria (Phylum: Actinobacteria)	Microbacterium	1 (1.92%)	–
12		Micrococcus	1 (1.92%)	–
Total			52	23

The values in the parentheses indicate the percentage abundance of the genus in the genotype

UMR2 showed minimum growth inhibition (21.67% each) of MG03. Among the five fungal endophytes evaluated, *L. theobromae* had the highest inhibition effect on MG03

(44.44%). In general, the genera *Bacillus* and *Paenibacillus* were highly antagonistic and inhibited the growth of MG03 under in vitro condition (Table 3).

Fig. 5 Diversity and distribution of culturable endophytes isolated from tissues of GPU and UM genotypes. **a** Relative richness of different bacterial genera isolated from the leaf, stem, and root tissues. **b** Number of culturable fungal and bacterial endophytes isolated

from tissues. **c** Summary of richness of bacterial genera and number of bacterial species isolated from leaf, stem, and root tissues of GPU and UM

Fig. 6 *In-vitro* antimicrobial assay of endophytes against *M. grisea* (MG03). The positive control **a** is MG03 (*M. grisea*) and P42 (*Bacillus velezensis*) strains are co-cultured and similarly, **b** MG03 and A6 strains are co-cultured [P—MG03 pathogen, B1—*Bacillus velezensis* strain P42, B2—*Bacillus velezensis* strain A6, T Test endophyte strains viz., UMR1 and GPU13 isolated from UM and GPU, respectively]

Table 3 In vitro and in planta evaluation of bacterial and fungal endophytes isolated from GPU and UM against *M. grisea* pathogen

Sl no.	Strains	Bacterium	PI (%)	PDI (%)	
				UM	GPU
Bacteria					
1	UML2	<i>Lysinibacillus</i> sp.	48.33 (44.04)	41.67 (40.20)	36.11 (36.92)
2	UML4	<i>E. mori</i>	21.67 (27.72)	69.44 (56.65)	33.33 (35.19)
3	UML5	<i>Enterobacter</i> sp.	33.33 (35.26)	36.11 (36.92)	25.00 (29.97)
4	UML6	<i>E. cloacae</i> subsp. <i>dissolvens</i>	26.67 (31.09)	22.22 (27.95)	47.22 (43.41)
5	UML10	<i>Bacillus</i> sp.	61.67 (51.75)	52.78 (46.59)	27.78 (31.81)
6	UML11	<i>P. aeruginosa</i>	61.67 (51.75)	61.11 (51.42)	27.78 (31.69)
7	UML12	<i>Bacillus</i> sp.	48.33 (44.04)	30.56 (33.54)	25.00 (29.97)
8	UMS1	<i>B. tequilensis</i>	56.67 (48.83)	63.89 (53.08)	22.22 (28.13)
9	UMS2	<i>B. mojavensis</i>	48.33 (44.04)	50.00 (45.00)	36.11 (36.92)
10	UMS3	<i>P. agglomerans</i>	33.33 (35.26)	33.33 (35.26)	25.00 (29.97)
11	UMS5	<i>Bacillus</i> sp.	61.67 (51.75)	58.33 (49.80)	33.33 (35.19)
12	UMS6	<i>Pseudomonas</i> sp.	28.33 (32.15)	75.00 (60.03)	75.00 (60.03)
13	UMS7	<i>Bacillus</i> sp.	41.67 (40.13)	52.78 (46.59)	63.89 (53.08)
14	UMR2	<i>Enterobacter</i> sp.	21.67 (27.59)	41.67 (40.20)	8.33 (16.55)
15	UMR3	<i>E. cloacae</i>	36.67 (37.25)	16.67 (24.09)	44.44 (41.79)
16	UMR4	<i>Bacillus</i> sp.	60.00 (50.77)	61.11 (51.42)	8.33 (16.55)
17	UMR5	<i>E. cloacae</i>	31.67 (34.18)	33.33 (35.19)	13.89 (20.88)
18	UMR6	<i>P. agglomerans</i>	31.67 (34.24)	44.44 (41.79)	55.56 (48.21)
19	UMR9	<i>B. mojavensis</i>	65.00 (53.78)	36.11 (36.92)	47.22 (43.41)
20	UMR10	<i>Bacillus</i> sp.	55.00 (47.87)	52.78 (46.59)	55.56 (48.21)
21	UMR12	<i>Bacillus</i> sp.	42.11 (40.43)	44.40 (41.81)	69.44 (56.46)
22	UMR13	<i>Lysinibacillus</i> sp.	40.35 (39.42)	66.67 (54.81)	58.33 (49.80)
23	UMR15	<i>Enterobacter</i> sp.	43.86(41.39)	63.89 (53.08)	61.11 (51.42)
24	GPUL1	<i>E. cloacae</i>	28.50 (32.26)	38.89 (38.58)	44.44 (41.81)
25	GPUL2	<i>E. dissolvens</i>	28.50 (32.26)	16.67 (23.80)	13.89 (21.78)
26	GPUL3	<i>A. baumannii</i>	24.40 (29.60)	47.22 (43.41)	22.22 (28.13)
27	GPUL4	<i>Enterobacter</i> sp.	31.25 (33.88)	30.56 (33.54)	16.67 (23.80)
28	GPUL5	<i>E. cloacae</i>	27.85 (31.82)	36.11 (36.92)	36.11 (36.92)
29	GPUL6	<i>E. cloacae</i>	37.50 (37.76)	33.33 (35.19)	44.44 (41.81)
30	GPUL7	<i>Enterobacter</i> sp.	31.25 (33.99)	33.33 (35.26)	25.00(29.68)
31	GPUL8	<i>B. cereus</i>	71.93 (58.01)	41.67 (40.20)	47.22 (43.41)
32	GPUL9	<i>E. cloacae</i> subsp. <i>dissolvens</i>	21.88 (27.83)	58.33 (49.80)	25.00 (29.97)
33	GPUL11	<i>B. cereus</i>	28.33 (32.15)	41.67 (40.20)	33.33 (35.19)
34	GPUL12	<i>C. sakazakii</i>	51.11 (45.64)	52.78 (46.66)	27.78 (31.81)
35	GPUL13	<i>A. baumannii</i>	25.56 (30.32)	30.56 (33.54)	16.67(24.09)
36	GPUL14	<i>A. baumannii</i>	28.89 (32.50)	44.44 (41.81)	19.44 (26.11)
37	GPUL15	<i>Bacillus</i> sp.	55.56 (48.19)	33.33 (35.19)	27.78 (31.81)
38	GPUL16	<i>A. baumannii</i>	25.00 (29.97)	52.78 (46.59)	16.67 (23.80)
39	GPUL17	<i>B. cereus</i>	51.67 (45.96)	30.56 (33.54)	13.89 (21.78)
40	GPUL18	<i>A. baumannii</i>	27.85 (31.82)	44.44 (41.79)	36.11 (36.92))
41	GPUL19	<i>E. cloacae</i>	51.67 (45.96)	41.67 (40.13)	38.89 (38.54)
42	GPUL20	<i>M. lylae</i>	35.00 (36.27)	55.56 (48.21)	44.44 (41.81)
43	GPUL21	<i>B. cereus</i>	38.33 (38.25)	44.44 (41.79)	19.44 (26.11)
44	GPUL23	<i>B. licheniformis</i>	66.67 (54.74)	44.44 (41.79)	33.33 (35.19)
45	GPUL24	<i>P. stutzeri</i>	22.81 (28.46)	63.89 (53.08)	50.00 (45.00)
46	GPUS1	<i>E. cloacae</i>	20.00 (26.25)	38.89 (38.54)	19.44 (26.11)
47	GPUS2	<i>E. cloacae</i> subsp. <i>cloacae</i>	45.56 (42.45)	38.89 (38.54)	27.78 (31.81)

Table 3 (continued)

Sl no.	Strains	Bacterium	PI (%)	PDI (%)	
				UM	GPU
48	GPUS3	<i>Bacillus</i> sp.	41.11 (39.77)	38.89 (38.58)	13.89 (21.78)
49	GPUS4	<i>E. cloacae</i>	25.00 (29.68)	55.56 (48.27)	27.78 (31.69)
50	GPUS5	<i>A. calcoaceticus</i>	28.33 (32.15)	25.00 (29.68)	22.22 (28.13)
51	GPUS8	<i>Paenibacillus</i> sp.	41.67 (40.20)	33.33 (35.26)	16.67 (23.80)
52	GPUS9	<i>E. ludwigii</i>	25.00 (29.99)	38.89 (38.54)	16.67 (23.80)
53	GPUS10	<i>A. baumannii</i>	28.33 (32.15)	47.22 (43.41)	41.67 (40.20)
54	GPUS11	<i>Enterobacter aerogenes</i>	20.83 (27.15)	30.56 (33.54)	19.44 (26.11)
55	GPUS12	<i>A. baumannii</i>	28.88 (32.50)	27.78 (31.69)	22.22 (28.13)
56	GPUS13	<i>Paenibacillus</i> sp.	60.00 (50.77)	44.44 (41.81)	25.00 (29.97)
57	GPUS14	<i>Pantoea</i> sp.	21.67 (27.72)	58.33 (49.80)	13.89 (21.78)
58	GPUS16	<i>Bacillus</i> sp.	56.67 (48.83)	52.78 (43.38)	44.44 (41.81)
59	GPUS17	<i>Bacillus</i> sp.	45.00 (42.13)	58.33 (33.54)	30.56 (33.54)
60	GPUR1	<i>E. cloacae</i>	25.00 (29.89)	33.33 (35.26)	22.22 (27.95)
61	GPUR2	<i>E. coli</i>	25.00 (29.99)	41.67 (40.20)	22.22 (27.95)
62	GPUR3	<i>E. cloacae</i>	30.00 (33.18)	36.11 (36.92)	33.33 (35.26)
63	GPUR4	<i>Pantoea</i> sp.	28.33 (32.15)	55.56 (48.19)	58.33 (49.80)
64	GPUR5	<i>E. cloacae</i>	30.00 (33.21)	33.33 (35.19)	16.67 (24.09)
65	GPUR6	<i>E. cloacae</i>	21.67 (27.72)	41.67 (40.20)	27.78 (31.81)
66	GPUR7	<i>E. cloacae</i>	30.00 (33.21)	36.11 (36.92)	50.00 (45.00)
67	GPUR8	<i>E. cloacae</i>	26.67 (31.05)	47.22 (43.41)	27.78 (31.81)
68	GPUR9	<i>E. ludwigii</i>	30.00 (33.21)	38.89 (38.58)	52.78 (46.59)
69	GPUR10	<i>B. cereus</i>	66.67 (54.74)	47.22 (43.41)	22.22 (28.13)
70	GPUR11	<i>Bacillus</i> sp.	58.33 (49.80)	47.22 (43.41)	33.33 (35.26)
71	GPUR12	<i>Bacillus</i> sp.	60.00 (50.77)	38.89 (38.58)	16.67 (23.80)
72	GPUR13	<i>M. xylanilyticum</i>	33.30 (35.21)	52.78 (46.59)	47.22 (43.41)
73	GPUR14	<i>B. cereus</i>	41.67 (40.20)	47.22 (43.41)	38.89 (38.58)
74	GPUR15	<i>L. helveticus</i>	31.67 (34.24)	27.78 (31.69)	33.33 (35.19)
75	GPUR16	<i>B. cereus</i>	52.63(46.56)	44.40 (41.81)	30.56 (33.54)
Fungi					
1	YGL3	<i>L. theobromae</i>	45.00 (50.00)	37.78 (43.41)	27.22 (62.05)
2	PGL4	<i>P. polonicum</i>	34.75 (32.50)	52.78 (43.38)	47.22 (46.59)
3	PGR6	<i>T. trachyspermus</i>	35.45 (33.75)	38.89 (51.42)	31.11 (36.46)
4	PGL2	<i>Phoma</i> sp.	24.69 (17.50)	63.89 (48.21)	55.56 (53.08)
5	PUL1	<i>Phoma</i> sp.	18.05 (10.00)	58.33 (33.54)	30.56 (49.87)
	Benomyl treatment		—	47.22 (43.41)	30.56 (33.54)
	Streptomycin treatment		—	52.78 (46.59)	44.44 (41.79)
	Tetracycline treatment		—	50.00 (45.00)	55.56 (48.21)
	Negative control		0	55.56 (48.21)	52.56 (13.63)
	Positive control (a)		78.82 ()	—	—
	Positive control (b)		80.00 ()	—	—

The values in the parentheses are the arc-sine transformed values

The critical difference values are calculated at the 5% probability level

PI percentage inhibition, PDI percentage disease index

Pathogenicity test and blast disease scoring for blast in finger millet

Artificial inoculation of MG03 produced characteristic and typical spindle-shaped lesions on the leaves of GPU and UM seedlings at 6–7 DAI. The lesions initiated as tiny spots with yellowish margin and grayish-green center became whitish gray and disintegrated later. The percentage disease incidence (PDI) was recorded on the infected GPU and UM seedlings at 10 DAI and the disease severity was compared in both the finger millet genotypes. Among the 52 bacterial isolates of GPU, seed treatment with the strains viz. *Enterobacter dissolvens* GPUL2 (Disease severity 13.89% and 16.67%), *Enterobacter* sp. GPUL4 (16.67% and 30.56%), *B. cereus* GPUL17 (13.89% and 30.56%), *Acinetobacter baumannii* GPUL13 (16.67% and 30.56%), *Bacillus* sp. GPUR12 (16.67% and 38.89%), and *Paenibacillus* sp. GPUS13 (25% and 44.44%) showed less blast disease severity in GPU and UM cultivars (Table 3). The species belonging to *E. cloacae*, *A. baumannii*, *Lactobacillus helveticus*, *E. dissolvens*, and *Enterobacter* sp. were effective in suppressing the pathogen in planta in both GPU and UM; however, they exhibited a poor inhibition of the pathogen in in vitro assay. Among the different strains isolated from GPU, the species belonging to *B. cereus*, *A. baumannii*, *L. helveticus*, and *Paenibacillus* sp. were found unique to GPU.

Out of 23 bacterial isolates of UM, seed treatment with the strain *Bacillus* sp. UML12 resulted in lowest disease severity (25% and 30.56% in GPU and UM, respectively) in planta assay. However, seed treatment with the strains belonging to *E. cloacae* UMR5 (13.89% and 33.33%), *Enterobacter* sp. UML5 (25% and 36.11%), *Bacillus* sp. UML12 (25% and 30.56%) and *Pantoea agglomerans* UMS3 (25% and 33.33%) resulted in low PDI in in planta in both GPU and UM, respectively (Table 3). Among the five fungal endophytes identified, *L. theobromae* YGL3 and *T. trachyspermus* PGR6 showed low PDI in both GPU and UM. However, their antagonistic activity against the pathogen in in vitro was found to be less effective. In a control (C_0), the PDI was recorded as 52.56% and 55.56% in GPU and UM, respectively. An overview of the results from in vitro and in planta studies have revealed that the GPU strains, *B. cereus* GPUL8, *Bacillus* sp. GPUL15, *B. cereus* GPUL17, *Paenibacillus* sp. GPUS13, *Bacillus* sp. GPUR12, *B. cereus* GPUR10, *Bacillus* sp. GPUR11, *Bacillus* sp. GPUR12, and *B. cereus* GPUR16 were effectively suppressed the pathogen both in in vitro and in planta in GPU and UM cultivars.

Discussion

In the present study, metagenomics approach enabled to gain a deeper understanding of the diversity of endophytic consortia in the GPU and UM finger millet genotypes, which

are contrasting in their reaction towards blast disease, caused by Ascomycetes fungi *Magnaporthe grisea*. The WMG sequencing followed by metagenome assembly of GPU and UM resulted into 7,76,412 and 4,95,014 contigs in GPU and UM, respectively. The metagenome of both GPU and UM cultivars was dominated by the sequences belonging to the domain bacteria in which, the phyla Proteobacteria and Actinobacteria were abundant. The previous studies on the characterization of endophyte communities in rice (Sessitsch et al. 2012; Sengupta et al. 2017), which reported high abundance of Proteobacteria, Firmicutes, and Actinobacteria. Proteobacteria were abundant in UM, whereas the Actinobacteria were highly prominent in GPU. Actinomycetes are soil microbes that colonize the host plant tissues as endophytes (Madhurama et al. 2014; Jogaiah et al. 2016) and produce antibiotics, promote plant growth, and also protect against phytopathogenic fungi (Smith 1957; Fiedler et al. 2008; Ma et al. 2009; Vurukonda et al. 2018). Actinomycetes are the largest microbial source for different bioactive secondary metabolites, and among the genera of Actinomycetes group, the *Streptomyces* spp. alone contributes to more than 70% of the total antibiotic production (Ningthoujam et al. 2009). This suggests that the relative abundance of *Streptomyces* spp. in GPU as compared to UM may underpin the host-plant resistance in GPU against *M. grisea* pathogen. Similar antifungal activity of *Streptomyces* spp. against *M. grisea* was also reported in earlier studies (Hwang et al. 1996; Lee et al. 2002; Yoshii et al. 2012; Van Minh et al. 2015).

The species, *S. cellulorum*, identified in the metagenome samples of both GPU and UM, but more abundant in GPU, was known to produce many antifungal compounds like ambruticins, jerangolids (Gerth et al. 1995a; Knuat and Reichenbach 2000), soraphens (Gerth et al. 1994), ratjadon (Gerth et al. 1995b), and epothilones (Gerth et al. 1996), which inhibit several phytopathogenic fungi. Another important endophyte *S. maltophilia*, reported to hold a strong antagonistic activity against *M. grisea* (Jha and Subramanian 2013; Etesami and Alikhani 2016), was found in GPU. The production of alkaline serine proteolytic enzyme and host systemic acquired resistance induced by *S. maltophilia* might play a key role in its biocontrol activity against different phytopathogens (Selim et al. 2016). The *L. aerocolonigenes* was identified in GPU, and was reported to produce thiobutacin (Lee et al. 2004; Yeop et al. 2008) which is antagonistic against oomycetes and some ascomycetes fungi. The high abundance of these species in GPU might have a potential role in the resistant reaction of this cultivar against the blast pathogen. The functional annotation of genes of endophytes identified in GPU and UM through SEED classification revealed that the GPU associated endophytes harbored with genes for stress response, sulphur metabolism, phosphorus metabolism, secondary metabolites, and metabolism of

aromatic compounds. Earlier studies reported that phosphorous metabolism (Dordas 2008), secondary metabolites and metabolism of aromatic compounds (Aranega-Bou et al. 2014; Llorens et al. 2017), and sulphur metabolism (Romero et al. 2014) play an important role in inducing host-plant resistance against several phytopathogens. Our *in silico* analysis suggested that the microbial consortia of GPU may help to overcome the pathogen infection and impart disease resistance reaction.

Similarly, the KEGG pathway analysis revealed that the GPU accommodated more genes encoding glutathione synthase pathway as compared to UM. The glutathione is an important signaling compound involved in plant disease resistance (Ghanta and Chattopadhyay 2011; Gullner and Zechmann 2017) and is known to involve in antioxidative defense mechanism in plants. We also found that the genes involved in cysteine synthase pathway were abundant in GPU. For most sulphur-containing metabolites, a direct antifungal mode of action has been reported where it activates systemic acquired resistance thereby initiating hypersensitive reaction (HR). Cysteine is the main precursor for all sulfur-containing compounds and is directly linked to stress response through its function related to systemic acquired resistance (Luckner 2013). Cysteine displays a regulatory function in pathogen defense and it was reported that a specific cytosolic cysteine content is essential for the initiation of the plant immune response to pathogens and a link to the hypersensitive response (Álvarez et al. 2012).

To support the results of our metagenomics study, culturable endophytes were isolated from GPU and UM cultivars. 52 bacterial and four fungal endophytes from GPU; 23 bacterial endophytes and one fungal endophyte from UM were isolated. Molecular characterization of 75 bacterial (16S rDNA sequencing) and 5 fungal (18S rDNA sequencing) endophytes revealed that *Enterobacter* spp. and *Bacillus* spp. were the dominant bacterial phylotypes colonizing GPU and UM. Based on metagenome sequencing and wet lab experiments, we observed diversity with respect to the distribution of endophyte communities in these two host habitats, i.e. GPU and UM. *Enterobacter* spp. and *Bacillus* spp. were identified as more abundant in GPU as compared to UM. This indicated that the nature of cultivar may contribute to the structure of the endophytic communities associated with the finger millet.

Among all the culturable endophytes evaluated in both *in vitro* and *in planta*, the strains belonging to the GPU native endophytes like *B. cereus* and *Paenibacillus* sp. were found highly effective in suppressing the growth of blast pathogen. The fungistatic effect of *B. cereus* and *Paenibacillus* sp. on *M. grisea* identified in the present study was also reported in previous studies (Ladeuze et al. 2011; Xu et al. 2014; Huang et al. 2018). Liu et al. (2007) reported *A. baumannii* as a potential endophyte in *Cinnamomum camphora*

and described its antagonistic potential against several phytopathogenic fungi such as *Cryphonectria parasitica*, *Glomerella glycines*, *Phytophthora capsici*, *Fusarium graminearum*, *Botrytis cinerea*, and *Rhizoctonia solani*. Interestingly, metagenome data and isolation studies showed that the *A. baumannii* was present only in GPU cultivar. However, the *in vitro* and *in planta* studies showed variation in microbial composition with respect to genera and species in GPU and UM cultivars. The endophyte strains which effectively suppressed the pathogen *in vitro* (*Bacillus* sp. UML10, *Bacillus* sp. UMS5, *Bacillus* sp. UMR4, *Bacillus* sp. GPUS16, and *Cronobacter sakazakii* GPUL12) showed poor suppression of disease in *planta* and vice versa (*E. cloacae* GPUL5, *A. baumannii* GPUL13, *A. baumannii* GPUL14, *Bacillus* sp. GPUS3, *E. ludwigii* GPUS9, and *E. aerogenes* GPUS11, etc.). *In vitro*, the endophytes were directly exposed to pathogen on synthetic media resulting in direct interaction between the endophyte and the pathogen. But under *in planta*, each endophyte was evaluated under natural conditions in its original host where it becomes exposed to multitude of factors and this compound effect could possibly be responsible for the variance in results observed under the two experimental conditions. We also observed that the same endophyte strain produced contrasting results in GPU and UM against the blast pathogen *in planta*. Particularly, in case of the strains belonging to *Bacillus* spp. (*B. cereus*, *B. mojavensis*, *B. tequilensis*, *B. licheniformis*, and *Bacillus* sp.), we observed that the majority of the *Bacillus* sp. strains had high suppression of the blast disease in *planta* in GPU, while the same strains showed less blast incidence in UM. There could be a specific host-endophyte relationship between the *Bacillus* species and GPU, which could be derived through their evolutionary relationship and contributing GPU to take advantage of the colonization of the *Bacillus* species and use their antagonistic potential for the suppression of the blast pathogen during infection.

Conclusion

The information and knowledge of microbial community dynamics found in the two genotypes of finger millet, GPU-28 and Udurumallige, was limited. This metagenomic study had highlighted biocontrol agents, to provide a new genesis and inception of microbial diversity to protect the finger millet against the blast disease caused by *M. grisea*. The WMG sequencing of GPU and UM revealed high structural diversity and distribution of endophytic microbial communities in these two genotypes. This information was supported by the functional annotation of gene clusters of endophytic consortia which unveiled genes encoding stress response, secondary metabolites and metabolism of aromatic compounds, phosphorous metabolism, and sulphur metabolism to be in

greater fraction in GPU as compared to UM. This comparative study provides key insights into the structural and functional differences of endophytic communities associated with blast resistant and susceptible cultivars might have putative role in contrast disease reactions to blast pathogen. The unique endophytes of GPU like *B. cereus* and *Paenibacillus* sp. could be used as potential bio-control agents in reducing the incidence of blast disease in finger millet crop. The results reported in this study could enhance the current understanding of the application of endophytes in controlling diseases in agriculturally important crops.

Author contributions HBM and MKP conceived and conceptualized the study. HBM and RUD collected the plant samples and isolated genomic DNA. RUD and KT performed the in vitro experiments. RCP analyzed the metagenome data. HBM guided metagenome data analysis. MEP assisted in maintaining cultures of endophytes and sequencing. HBM, MKP and BK wrote the manuscript. HBM, KSN and GVB edited the manuscript. KSN, KB and BP contributed in preparing the figures. All authors read and approved the final manuscript for publication.

Availability of sequenced data The raw fastq files for the metagenome data were uploaded in NCBI SRA database with accession number SRP105033.

Compliance with ethical standards

Conflict of interest Author Rajadurai, R. C. was employed by company AgriGenome Laboratory. All other authors declare no competing interests.

References

- Álvarez C, Bermúdez MÁ, Romero LC et al (2012) Cysteine homeostasis plays an essential role in plant immunity. *New Phytol* 193:165–177. <https://doi.org/10.1111/j.1469-8137.2011.03889.x>
- Amruta N, Prasanna Kumar MK, Puneeth ME et al (2018) Exploring the potentiality of novel rhizospheric bacterial strains against the rice blast fungus *Magnaporthe oryzae*. *Plant Pathol J* 34:126–138. <https://doi.org/10.5423/PPJ.OA.11.2017.0242>
- Anitha A, Rebeeth M (2009) In vitro antifungal activity of *Streptomyces griseus* against phytopathogenic fungi of tomato field. *Acad J Plant Sci* 2:119–123
- Aranega-Bou P, de la Oleyva M, Finiti I et al (2014) Priming of plant resistance by natural compounds. Hexanoic acid as a model. *Front Plant Sci* 5:1–12. <https://doi.org/10.3389/fpls.2014.00488>
- Arndt D, Xia J, Liu Y et al (2012) METAGENassist: a comprehensive web server for comparative metagenomics. *Nucl Acids Res* 40:88–95. <https://doi.org/10.1093/nar/gks497>
- Arnold AE, Mejia LC, Kylo D et al (2003) Fungal endophytes limit pathogen damage in a tropical tree. *Proc Natl Acad Sci* 100:15649–15654. <https://doi.org/10.1073/pnas.2533483100>
- Benfradj N, Tounsi S, Boughalleb-MHamdi N (2016) In-vitro evaluation of Antagonists and Fungicides in Controlling Citrus Gummosis Caused by Phytophthora, Phytophthium and Pythium species in Tunisia. *Br Microbiol Res J* 16:1–14. <https://doi.org/10.9734/BMRJ/2016/27247>
- Bhatt D, Negi M, Sharma P et al (2011) Responses to drought induced oxidative stress in five finger millet varieties differing in their geographical distribution. *Physiol Mol Biol Plants* 17:347–353. <https://doi.org/10.1007/s12298-011-0084-4>
- Boisvert S, Raymond F, Godzaridis É et al (2012) Ray Meta: scalable de novo metagenome assembly and profiling. *Genome Biol*. <https://doi.org/10.1186/gb-2012-13-12-r122>
- Bolwerk A, Lagopodi AL, Lugtenberg BJJ, Bloemberg GV (2005) Visualization of interactions between a pathogenic and a beneficial *Fusarium* strain during biocontrol of tomato foot and root rot. *Mol Plant-Microbe Interact* 18:710–721. <https://doi.org/10.1094/MPMI-18-0710>
- Braun G, Hirsch U (1992) Communities of parasitic microfungi. In: Winterhoff W (ed) *Fungi in vegetation science*. Springer, Dordrecht, pp 225–250
- Buchfink B, Xie C, Huson DH (2015) Fast and sensitive protein alignment using DIAMOND. *Nat Methods* 12:59–60. <https://doi.org/10.1038/nmeth.3176>
- Budnik LT, Baur X (2009) The assessment of environmental and occupational exposure to hazardous substances by biomonitoring. *Dtsch Arztebl Int* 106:91–97. <https://doi.org/10.3238/arztebl.2009.0091>
- Chaturvedi H, Singh V (2016) Potential of bacterial endophytes as plant growth promoting factors. *J Plant Pathol Microbiol*. <https://doi.org/10.4172/2157-7471.1000376>
- Chiang KS, Liu HI, Bock CH (2017) A discussion on disease severity index values. Part I: warning on inherent errors and suggestions to maximise accuracy. *Ann Appl Biol* 171:139–154. <https://doi.org/10.1111/aab.12362>
- Doran JL, Leskiw BK, Aippersbach S, Jensen SE (1990) Isolation and characterization of a beta-lactamase-inhibitory protein from *Streptomyces clavuligerus* and cloning and analysis of the corresponding gene. *J Bacteriol* 172:4909–4918
- Dordas C (2008) Role of nutrients in controlling plant diseases in sustainable agriculture. A review. *Agron Sustain Dev* 28:33–46. <https://doi.org/10.1051/agro:2007051>
- Dubreuil-Maurizi C, Poinssot B (2012) Role of glutathione in plant signaling under biotic stress. *Plant Signal, Behav*
- Dutta D, Puzari KC, Gogoi R, Dutta P (2014) Endophytes: exploitation as a tool in plant protection. *Braz Arch Biol Technol* 57:621–629
- Esele JP (2002) Diseases of finger millet: a global overview. In: Murphy LJ, Montorsi F (eds) *Sorghum and finger millet diseases*. Iowa State Press, Iowa, pp 19–26
- Etesami H, Alikhani HA (2016) Suppression of the fungal pathogen *Magnaporthe grisea* by *Stenotrophomonas maltophilia*, a seed-borne rice (*Oryza sativa* L.) endophytic bacterium. *Arch Agron Soil Sci* 62:1271–1284. <https://doi.org/10.1080/03650340.2016.1139087>
- Fiedler H, Bruntner C, Riedlinger J et al (2008) ORIGINAL ARTICLE Proximicin A, B and C, Novel aminofuran antibiotic and anticancer compounds isolated from marine strains of the *Actinomycete Verrucosipora*. *J Antibiot* 61:158–163
- Frandsberg E, Petersson C, Lundgren LN, Schnurer J (2000) *Streptomyces halstedii* K122 produces the antifungal compounds bafilomycin B1 and C1. *Can J Microbiol* 46:753–758. <https://doi.org/10.1139/w00-050>
- Gayathri P, Muralikrishnan V (2013) Isolation and characterization of endophytic actinomycetes from mangrove plant for antimicrobial activity. *Int J Curr Microbiol Appl Sci* 2:78–89
- Gerth K, Bedorf N, Irschik H et al (1994) The soraphens: a family of novel antifungal compounds from *Sorangium cellulosum* (Myxobacteria). I. Soraphen A1 alpha: fermentation, isolation, biological properties. *J Antibiot (Tokyo)* 47:23–31. <https://doi.org/10.7164/antibiotics.47.23>

- Gerth K, Schummer D, Höfle G, Irschik H, Reichenbach H (1995a) Ratjadon: a new antifungal compound from *Sorangium cellulosum* (Myxobacteria) production, physico-chemical and biological properties. *J Antibiot* (Tokyo) 48:973–976. <https://doi.org/10.7164/antibiotics.48.973>
- Gerth K, Washausen P, Hfleff G et al (1995b) The jerangolids : a family of new antifungal compounds from *Sorangium cellulosum* (Myxobacteria) production, physico-chemical and biological properties of jerangolid A1. *Optim Antibiot Product* 49:71–75
- Gerth K, Bedorf N, Hflen G, Reichenbach H (1996) Epothilons A and B: antifungal and cytotoxic compounds from *Sorangium cellulosum* (Myxobacteria) production. *Physico-chem Biol Prop* 1:560–563
- Ghanta S, Chattopadhyay S (2011) Glutathione as a signaling molecule another challenge to pathogens. *Plant Signal Behav* 6:783–788. <https://doi.org/10.4161/psb.6.6.15147>
- Gomez KA, Gomez KA, Gomez AA (1984) Statistical procedures for agricultural research. Wiley, New York
- Gond SK, Bergen MS, Torres MS, White JF (2015) Endophytic *Bacillus* spp. produce antifungal lipopeptides and induce host defence gene expression in maize. *Microbiol Res* 172:79–87. <https://doi.org/10.1016/j.micres.2014.11.004>
- Grosch R, Scherwinski K, Lottmann J et al (2006) Fungal antagonists of the plant pathogen *Rhizoctonia solani*: selection, control efficacy and influence on the indigenous microbial community. *Mycol Res* 110:1464–1474. <https://doi.org/10.1016/j.mycres.2006.09.014>
- Gullner G, Zechmann B (2017) Glutathione in plant growth, development, and stress tolerance. Springer, Cham. <https://doi.org/10.1007/978-3-319-66682-2>
- Gunatilaka AAL (2006) Natural products from plant-associated microorganisms: distribution, structural diversity, bioactivity, and implications of their occurrence. *J Nat Prod* 69:509–526. <https://doi.org/10.1021/np058128n>
- Handelsman J (2005) Metagenomics: application of genomics to uncultured microorganisms. *Microbiol Mol Biol Rev* 69:195. <https://doi.org/10.1128/MMBR.69.1.195.2005>
- Huang W, Liu X, Zhou X, et al (2018) *Bacillus cereus* HS24 suppresses conidia germination of *Magnaporthe oryzae* by inhibiting the Ca²⁺ signaling pathway. <http://dx.doi.org/10.1101/310516>
- Hung PQ, Annapurna K (2004) Isolation and characterization of endophytic bacteria in soybean (*Glycine* Sp.). *Omonrice* 101:92–101
- Huson DH, Mitra S, Ruscheweyh H et al (2011) Integrative analysis of environmental sequences using MEGAN4. *Genome Res* 21:1552–1560. <https://doi.org/10.1101/gr.120618.111.Freely>
- Hwang BK, Lee JY, Kim BS, Moon SS (1996) Isolation, structure elucidation, and antifungal activity of a manumycin-type antibiotic from *Streptomyces flaveus*. *J Agric Food Chem* 44:3653–3657. <https://doi.org/10.1021/jf960084o>
- Jha Y, Subramanian RB (2013) Root associated bacteria from the local variety of rice GJ-17 antagonized the growth of *Magnaporthe grisea*. *Adv Biores* 4(1):70–77
- Jogaiah S, Kurjogi M, Govind SR et al (2016) Isolation and evaluation of proteolytic actinomycete isolates as novel inducers of pearl millet downy mildew disease protection. *Sci Rep* 6:1–7. <https://doi.org/10.1038/srep30789>
- Kanehisa M, Goto S, Sato Y et al (2012) KEGG for integration and interpretation of large-scale molecular data sets. *Nucl Acids Res* 40:109–114. <https://doi.org/10.1093/nar/gkr988>
- Knuat P, Reichenbach H (2000) On the mechanism of action of the myxobacterial fungicide ambruticin. *J Antibiot* (Tokyo) 53:1182–1190. <https://doi.org/10.7164/antibiotics.53.1182>
- Ladeuze S, Lentz N, Delbrassine L et al (2011) Antifungal activity displayed by cereulide, the emetic toxin produced by *Bacillus cereus*. *Appl Environ Microbiol* 77:2555–2558. <https://doi.org/10.1128/AEM.02519-10>
- Lee CH, Kim BJ, Choi GJ et al (2002) *Streptomyces* with antifungal activity against rice blast causing fungus, *Magnaporthe grisea*. *J Microbiol Biotechnol* 12:1026–1028
- Lee JY, Moon SS, Yun BS et al (2004) Thiobutacin, a novel antifungal and antioomycete antibiotic from *Lechevalieria aerocolonigenes*. *J Nat Prod* 67:2076–2078. <https://doi.org/10.1021/np049786v>
- Lim SM, Yoon M, Choi GJ et al (2017) Diffusible and volatile antifungal compounds produced by an antagonistic *Bacillus velezensis* G341 against various phytopathogenic fungi. *Plant Pathol J* 33:488–498. <https://doi.org/10.5423/PPJ.OA.04.2017.0073>
- Liu CH, Chen X, Liu TT et al (2007) Study of the antifungal activity of *Acinetobacter baumannii* LCH001 in vitro and identification of its antifungal components. *Appl Microbiol Biotechnol* 76:459–466. <https://doi.org/10.1007/s00253-007-1010-0>
- Llorens E, García-Agustín P, Lapeña L et al (2017) Advances in induced resistance by natural compounds: towards new options for woody crop protection. *Sci Agric* 74:90–100. <https://doi.org/10.1590/1678-992x-2016-0012>
- Luckner M (2013) Secondary metabolism in microorganisms, plants and animals. Springer Science & Business Media, Berlin
- Ma GBZ, Xia Z, Wu S (2009) Inhibiting effect of seven marine actinomycete strains against vegetable pathogenic microorganisms. *Crops* 5:3–9
- Madhurama G, Sonam D, Urmil PG, Ravindra NK (2014) Diversity and biopotential of endophytic actinomycetes from three medicinal plants in India. *Afric J Microbiol Res* 8:184–191. <https://doi.org/10.5897/AJMR2012.2452>
- Mahesh H, Meghana S, Shailaja H et al (2016) Acquisition of the grasshopper retro transposon by rice *Magnaporthe* isolates indicates a dynamic gene flow between rice and non-rice *Magnaporthe* population. *J Pathol Microbiol* 1(2):1011
- Mander P, Cho SS, Choi YH et al (2016) Purification and characterization of chitinase showing antifungal and biodegradation properties obtained from *Streptomyces anulatus* CS242. *Arch Pharm Res* 39:878–886. <https://doi.org/10.1007/s12272-016-0747-3>
- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10
- Moller Bahnweg G, Sander mann H, Geiger HHEM (1992) A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucl Acids Res* 20:6115–6116
- Mousa WK, Raizada MN (2013) The diversity of anti-microbial secondary metabolites produced by fungal endophytes: an interdisciplinary perspective. *Front Microbiol* 4:1–18. <https://doi.org/10.3389/fmicb.2013.00065>
- Nagaraja A, Gowda J, Krishnappa M, Gowda KTK (2008) GPU 28: a finger millet variety with durable blast resistance. *J Mycopathol Res* 46:109–111
- Netam RS, Tiwari RK, Bahadur ANDS (2014) In vitro and in vivo efficacy of fungicides against *Pyricularia grisea* causing finger millet blast disease. *Int J Plant Prot* 7:137–142
- Ningthoujam DS, Kshetri P, Sanasam S, Nimaichand S (2009) Screening, identification of best producers and optimization of extracellular proteases from moderately halophilic alkalithermotolerant indigenous actinomycetes. *World Appl Sci J* 7:907–916
- Noguchi H, Taniguchi T, Itoh T (2008) Meta gene annotator: detecting species-specific patterns of ribosomal binding site for precise gene prediction in anonymous prokaryotic and phage genomes. *DNA Res* 15:387–396. <https://doi.org/10.1093/dnares/dsn027>
- Overbeek R, Olson R, Pusch GD et al (2014) The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucl Acids* 42:206–214. <https://doi.org/10.1093/nar/gkt1226>
- Pandey SS, Singh S, Pandey H et al (2018) Endophytes of *Withania somnifera* modulate in planta content and the site of withanolide

- biosynthesis. *Sci Rep* 20:1–19. <https://doi.org/10.1038/s41598-018-23716-5>
- Parks DH, Beiko RG (2010) Identifying biologically relevant differences between metagenomic communities. *Bioinformatics* 26:715–721. <https://doi.org/10.1093/bioinformatics/btq041>
- Paul AK, Banerjee AK (1983) A new antifungal antibiotic produced by *Streptomyces galbus*. *Folia Microbiol (Praha)* 28:386–396. <https://doi.org/10.1007/BF02879488>
- Rabeeth M, Anitha A, Srikanth G (2011) Purification of an antifungal endochitinase from a potential biocontrol agent *Streptomyces griseus*. *Pak J Biol Sci* 14:788–797. <https://doi.org/10.3923/pjbs.2011.788.797>
- Ravin NV, Mardanov AV, Skryabin KG (2015) Metagenomics as a tool for the investigation of uncultured microorganisms. *Russ J Genet* 51:431–439. <https://doi.org/10.1134/S1022795415050063>
- Rodriguez RJ, Henson J, Van Volkenburgh E et al (2008) Stress tolerance in plants via habitat-adapted symbiosis. *ISME J* 2:404–416. <https://doi.org/10.1038/ismej.2007.106>
- Romero LC, Aroca MÁ, Laureano-Marín AM et al (2014) Cysteine and cysteine-related signaling pathways in *Arabidopsis thaliana*. *Mol Plant* 7:264–276. <https://doi.org/10.1093/mp/sst168>
- Sastri BN (1950) The Wealth of India. A dictionary of indian raw materials and industrial products. Raw materials. Wealth India A Dict Indian Raw Mater Ind Prod III:160–166
- Schloss PD, Handelsman J (2005) Metagenomics for studying unculturable microorganisms: cutting the Gordian knot. *Genome Biol* 6:6–9. <https://doi.org/10.1186/gb-2005-6-8-229>
- Schulz B, Guske S, Dammann U, Boyle C (1998) Endophyte-host interactions. II. Defining symbiosis of the endophyte-host interaction. *Symbiosis Phila Pa (USA)* 25:213–227
- Selim HMM, Gomaa NM, Essa AMM (2016) Antagonistic effect of endophytic bacteria against some phytopathogens. *Egypt J Bot* 56:613–626
- Sengupta S, Ganguli S, Singh PK (2017) Metagenome analysis of the root endophytic microbial community of Indian rice (*O. sativa* L.). *Genom Data* 12:41–43. <https://doi.org/10.1016/j.gdata.2017.02.010>
- Sessitsch A, Hardoim P, Döring J et al (2012) Functional characteristics of an endophyte community colonizing rice roots as revealed by metagenomic analysis. *Mol Plant-Microbe Interact* 25:28–36. <https://doi.org/10.1094/MPMI-08-11-0204>
- Shankar M, Kurtböke DI, Gillespie-Sasse LMJ et al (1994) Possible roles of competition for thiamine, production of inhibitory compounds, and hyphal interactions in suppression of the take-all fungus by a sterile red fungus. *Can J Microbiol* 40:478–483. <https://doi.org/10.1139/m94-077>
- Shirke MD, Mahesh HB, Gowda M (2016) Genome-wide comparison of *Magnaporthe* species reveals a host-specific pattern of secretory proteins and transposable elements. *PLoS One* 11:1–19. <https://doi.org/10.1371/journal.pone.0162458>
- Singh N, Varma A (2015) Antagonistic Activity of Siderophore Producing Rhizobacteria Isolated from the Semi-Arid Regions of Southern India. *Int J Curr Microbiol App Sci* 4(9):501–510
- Smith GE (1957) Inhibition of *Fusarium oxysporum* f. lycopersici by a species of Micromonospora isolated from tomato. *Phytopathology* 47:429–432
- Sziderics AH, Rasche F, Trognitz F et al (2007) Bacterial endophytes contribute to abiotic stress adaptation in pepper plants (*Capsicum annuum* L.). *Can J Microbiol* 53:1195–1202. <https://doi.org/10.1139/W07-082>
- Tokpah DP, Li H, Newmah JT et al (2017) Biological control of potential antagonistic bacteria isolates to restrict *Magnaporthe grisea* infection on rice. *Afr J Microbiol Res* 11:1108–1119. <https://doi.org/10.5897/AJMR2017.8562>
- Upton R, Read DJ, Newsham KK (2009) Nitrogen form influences the response of *Deschampsia antarctica* to dark septate root endophytes. *Mycorrhiza* 20:1–11. <https://doi.org/10.1007/s00572-009-0260-3>
- Van Minh N, Woo EE, Kim JY et al (2015) Antifungal substances from *Streptomyces* sp. A3265 antagonistic to plant pathogenic fungi. *Mycobiology* 43:333–338. <https://doi.org/10.5941/MYCO.2015.43.3.333>
- Verma SC, Ladha JK, Tripathi AK (2001) Evaluation of plant growth promoting and colonization ability of endophytic diazotrophs from deep water rice. *J Biotechnol* 91:127–141. [https://doi.org/10.1016/S0168-1656\(01\)00333-9](https://doi.org/10.1016/S0168-1656(01)00333-9)
- Vetriventhan M, Upadhyaya HD, Dwivedi SL et al (2015) Finger and foxtail millets. Elsevier Inc., Amsterdam
- Vincent JM (1947) Distortion of fungal hyphae in the presence of certain inhibitors. *Nature* 159:850
- Vurukonda SSKP, Giovanardi D, Stefani E (2018) Plant growth promoting and biocontrol activity of *Streptomyces* spp. As endophytes. *Int J Mol Sci*. <https://doi.org/10.3390/ijms19040952>
- Waller F, Achatz B, Baltruschat H et al (2005) The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and higher yield. *Proc Natl Acad Sci* 102:13386–13391. <https://doi.org/10.1073/pnas.0504423102>
- White TJ, Bruns T, Lee S, Taylor JL (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Academic Press, San Diego
- Xu SJ, Hong SJ, Choi W, Kim BS (2014) Antifungal activity of *Paenibacillus kribbensis* strain T-9 isolated from soils against several plant pathogenic fungi. *Plant Pathol J* 30:102–108. <https://doi.org/10.5423/PPJ.OA.05.2013.0052>
- Yeop LJ, Sherman DH, Hwang BK (2008) In vitro antimicrobial and in vivo antioomycete activities of the novel antibiotic thiobutacin. *Pest Management Science: formerly. Pestic Sci* 64(2):172–177
- Yoshii A, Moriyama H, Fukuhara T (2012) The novel kasugamycin 2'-N-acetyltransferase gene aac(2')-IIa, carried by the IncP island, confers kasugamycin resistance to rice-pathogenic bacteria. *Appl Env Microbiol* 78:5555–5564. <https://doi.org/10.1128/AEM.01155-12>