CORRESPONDENCE

The Prevalence, Comorbidity, Management, and Costs of Irritable Bowel Syndrome

An Observational Study Using Routine Health Insurance Data

by Prof. Dr. med. Winfried Häuser, Dr. med. Ursula Marschall, Prof. Dr. med. Peter Layer, and Dr. med. Thomas Grobe
in issue 27–28/2019

Anaerobic Cause

This article (1) on concomitant diagnoses, examination services, and drug prescriptions does not even mention one of the most common and very important causes of irritable bowel syndrome, namely colonization of the large intestine with anaerobes. A corresponding adequate therapy with a few doses of clindamycin can do wonders here, allowing the patients to heal and saving payers from paying inappropriate and meaningless psychotherapeutic measures.

DOI: 10.3238/arztebl.2019.0755a

Reference

1. Häuser W, Marschall U, Layer P, Grobe T:
The prevalence, comorbidity, management and costs of irritable bowel syndrome—an observational study using routine health insurance data. *Dtsch Arztebl Int* 2019; 116: 463–7.

Dr. med. Walter Neussel

Wittlich
Germany
walterneussel@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

“Psychogenic” is Not the Be-All and End-All

The topic of irritable bowel syndrome is repeatedly discussed in *Deutsches Ärzteblatt*, but unfortunately one-sided. I miss the groundbreaking study from 2014 by Prof. Annette Fritscher-Ravens, in which she showed that 22 of 36 patients with the diagnosis irritable bowel syndrome suffered from a specific food intolerance (1).

While the method used is certainly not suitable for routine examination, it should show us that we will have only limited success if we treat with a therapy that is based on a biopsychosocial disease model. Indeed, it underscores the need to develop methods that allow a simpler way to identify these food intolerances.

With the increasingly prevalent attitude that illnesses that cannot be explained using conventional methods should be classified as psychogenic, we are helping

neither our patients nor ourselves, but are rather causing psychological suffering and comorbidity—and also, of course, doctor hopping

DOI: 10.3238/arztebl.2019.0755b

References

1. Fritscher-Ravens A, Schuppan D, Ellrichmann M, et al.: Confocal endomicroscopy shows food-associated changes in the intestinal mucosa of patients with irritable bowel syndrome. *Gastroenterol* 2014; 147: 1012–20.
2. Häuser W, Marschall U, Layer P, Grobe T: The prevalence, comorbidity, management and costs of irritable bowel syndrome—an observational study using routine health insurance data. *Dtsch Arztebl Int* 2019; 116: 463–7.

Dr. med. Astrid Spangenberg

Zentrum für Seltene Erkrankungen
Medizinische Hochschule Hannover, Germany
spangenberg.astrid@mh-hannover.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We are grateful for the opportunity to comment on the two items of correspondence. Our article presents the administrative incidence, prevalence and comorbidities, and care and cost of irritable bowel syndrome (IBS) in the billing data from the German Barmer health insurance (*Barmer Ersatzkasse*, 1). We agree with Dr. Spangenberg that food intolerances in many patients contribute to symptoms of IBS. In the general population and in clinical trials, up to 50% of IBS patients report food intolerances (2). We discuss several possible reasons why the prevalence of cases coded for food intolerances (e.g. lactose intolerance in 8% of IBS

cases) was lower than that expected from epidemiological studies; these reasons include not having ICD-10 codes for some food intolerances as well as a lack of testing or coding for food intolerances by the practitioner (1).

Colonization of the colon with anaerobes is not an ICD-10 diagnosis and therefore could not be recorded in the billing data. A recent systematic review found a large heterogeneity of studies comparing the gut microbiota of IBS patients with healthy controls. A higher concentration of anaerobes in the stool of IBS patients was not found (3). As far as we know, no guideline for IBS recommends healing IBS with clindamycin, which

Dr. Neussel describes as being able to “do wonders.” A controlled trial of clindamycin for IBS was not found in PubMed.

Both correspondences comment critically on the possible role of psychological factors and psychotherapy in IBS. There is a high degree of interindividual variability in the importance of mental factors in the development and course of IBS (2)—as with food intolerances. Our observational study confirms the high rate of association of IBS with anxiety and depressive disorders (2).

In the four quarters prior to as well as the quarter in which IBS was diagnosed, 58% of persons with IBS had received psychotherapy services according to the current German guidelines. Additionally, 26% received outpatient psychiatry or specialist psychosomatic treatment. Billing data from health insurance carriers cannot make any statement as to whether the psychiatric–psychotherapeutic services were “inappropriate and meaningless”—as claimed by Dr. Neussel. In contrast to the therapy recommended by Dr. Neussel with clindamycin, the efficacy and safety of various psychotherapeutic procedures, such as gut-directed hypnosis and cognitive behavioral therapy, are well documented for IBS (4). The German S3 guideline on IBS recommends performing psychological diagnostics to detect psychosocial stress and/or mental comorbidity. Psychotherapeutic procedures should be based on the manifestation of the mental symptom burden (with psychosomatic primary care from a practitioner or gastroenterologist for mild cases, and specialist

psychotherapy in severe cases) and be integrated into a comprehensive medical approach that also includes nutritional advice and symptom-related drug therapy.

DOI: 10.3238/arztebl.2019.0755c

References

1. Häuser W, Marschall U, Layer P, Grobe T: The prevalence, comorbidity, management and costs of irritable bowel syndrome—an observational study using routine health insurance data. *Dtsch Arztebl Int* 2019; 116: 463–70.
2. Andresen V, Keller J, Pehl C, Schemann M, Preiss J, Layer P: Irritable bowel syndrome—the main recommendations. *Dtsch Arztebl Int* 2011;108:751–60.
3. Maharshak N, Ringel Y, Katibian D, et al.: Fecal and mucosa-associated intestinal microbiota in patients with diarrhea-predominant irritable bowel syndrome. *Dig Dis Sci* 2018; 63: 1890–9.
4. Ford AC, Lacy BE, Harris LA, Quigley EMM, Moayyedi P: Effect of antidepressants and psychological therapies in irritable bowel syndrome: an updated systematic review and meta-analysis. *Am J Gastroenterol* 2019; 114: 21–39.

On behalf of the authors

Prof. Dr. med. Winfried Häuser

Klinik für Innere Medizin I, Klinikum Saarbrücken, Klinik für Psychosomatische Medizin und Psychotherapie, Technical University of Munich, Germany
whaeuser@klinikum-saarbruecken.de

Prof. Dr. med. Peter Layer

Israelitisches Krankenhaus, Hamburg, Germany

Conflict of interest statement

Prof. Häuser has received author royalties for a CD with medical hypnosis for persons with IBS from Hypnos Verlag.

Prof. Layer has received consultant honoraria from Allergan.

CLINICAL SNAPSHOT

Acute Kidney Injury Following High-Dose Vitamin C Treatment

A 63-year-old man was admitted to our department with oliguric acute kidney injury 2 months after resection of ductal pancreatic cancer. The patient was a veterinarian and took an interest in alternative medicine. Serum creatinine, normal the week before, was measured at 8.2 mg/dL. Renal biopsy revealed osmotic tubulopathy (Figure, red asterisk) and extensive tubulointerstitial deposits of oxalate crystals (Figure, black asterisk). On questioning, the patient reported self-administration of intravenous high-dose ascorbic acid (veterinary preparation, dose unknown, contamination/damage not excluded). High-dose vitamin C is widely used as an adjuvant treatment for tumors in complementary medicine, despite the lack of reliable evidence for its efficacy. There are also indications that it may be effective in patients with sepsis or severe burns and in the context of coronary heart surgery. However, more research is required. Ascorbic acid dissolves in water and is therefore thought to be relatively safe. Acute kidney damage, especially with pre-existing kidney injury, is a rare but known adverse effect. Our patient's oliguria persisted, so hemodialysis was started. Dialysis was successfully discontinued 2 months later.

Dr. med. Annick Andres, Prof. Dr. med. Martin Zeier, Universitätsklinikum Heidelberg, Nephrologie, annick.andres@med.uni-heidelberg.de

Christoph Eckert, Universitätsklinikum Heidelberg, Pathologie

Conflict of interest statement: The authors declare that no conflict of interest exists.

Translated from the original German by David Roseveare.

Cite this as: Andres A, Zeier M, Eckert C: Acute kidney injury following high-dose vitamin C treatment. *Dtsch Arztebl Int* 2019; 116: 756. DOI: 10.3238/arztebl.2019.0756