Published in final edited form as:

J Intellect Disabil Res. 2016 May ; 60(5): 478–490. doi:10.1111/jir.12286.

Consonant Differentiation Mediates The Discrepancy Between Nonverbal And Verbal Abilities In Children With ASD

Alexandra P. Key^{1,2}, Paul J. Yoder^{1,2,3}, Wendy L. Stone⁴

¹Vanderbilt Kennedy Center

²Department of Hearing and Speech Sciences, Vanderbilt University School of Medicine

³Department of Special Education, Vanderbilt University

⁴Department of Psychology, University of Washington

Abstract

Background—Many children with autism spectrum disorders (ASD) demonstrate verbal communication disorders reflected in lower verbal than nonverbal abilities. The present study examined the extent to which this discrepancy is associated with atypical speech sound differentiation.

Methods—Differences in the amplitude of auditory event-related potentials elicited by contrasting consonant-vowel syllables during a passive listening paradigm were used to assess speech sound differentiation in 24 children with ASD and 18 chronological age-matched children with typical development (TD), *M* age 6.90 years (*SD*=1.39).

Results—Results revealed that compared to TD peers, children with ASD showed reduced consonant differentiation in the 84–308 ms period. Brain responses indexing consonant differentiation were negatively related to the degree of discrepancy in nonverbal-verbal ability and mediated the relationship between diagnostic group membership and the greater discrepancy.

Conclusions—We discuss the theoretical and clinical implications of the brain's response to speech sound contrasts possibly explaining the greater nonverbal vs. language ability in children with ASD compared to that in typically developing children.

Keywords

Autism; ERP; language; speech perception; consonant differentiation

Autism spectrum disorder (ASD) includes symptoms in the areas of social and communicative domains as well as repetitive behaviours (DSM 5; APA, 2013). Many children with ASD evidence receptive and expressive language abilities that are below their nonverbal cognitive abilities (Kwok et al., 2015; Loukas et al., 2008) and may demonstrate deficits similar to those of individuals with specific language impairment (SLI; Tager-Flusberg, 2006; Herbert & Kenet, 2007). This type of a verbal communication disorder is

particularly interesting because overall intellectual disability is unlikely to explain it (Stark & Tallal, 1981).

Although some form of language deficits has been reported in up to 60% of children with ASD (Kjelgaard & Tager-Flusberg, 2001), the estimates of the proportion of children with ASD who demonstrate higher nonverbal than verbal ability vary. Some have reported such discrepancy to be common but not universal (Lincoln et al., 2007). Others noted this profile to be present more often in younger (e.g., preschool) and lower-functioning (IQ < 80) children with ASD, with the extent of the discrepancy decreasing with age (Mayes & Calhoun, 2003; Joseph et al., 2002), mainly due to higher scores in the verbal domain for older children (Mayes & Calhoun, 2003). Tager-Flusberg and Joseph (2003) observed higher nonverbal than verbal abilities measured by Differential Ability Scales in 34% (14/47) of children with ASD (6–13 years), while 48% (35/73) of preschoolers with ASD showed a similar discrepancy (Joseph et al., 2002). Understanding the mechanisms underlying language deficits not due to intellectual disability will allow to develop novel treatment targets as well as diagnostic tools for better identification of early signs of communication delays.

Individual differences in the extent of language deficits relative to nonverbal ability are thought to have multiple origins. Difficulties in auditory information processing offer one possible explanation. Previous studies in children with SLI in the absence of ASD noted the importance of effective speech sound processing for proper language development (e.g., Benasich et al., 2002). They also reported deficits in basic auditory perception processes at the level of individual consonants or vowels (McArthur & Bishop, 2005; Shafer et al., 2005).

Event-related potentials (ERPs), which measure brain activity following presentation of a stimulus, have been widely used to examine auditory processing in children and adults. ERPs provide high temporal resolution that allows for the quantification of neural processing with great precision. Additionally, many ERP paradigms do not require active engagement from the participant, making this methodology especially valuable for use in populations who may be unable or unmotivated to provide reliable behavioural responses (Yoder et al., 2006) and suitable for use across multiple ages, including infants. Identifying the connection between neural evidence of atypical speech sound processing and language deficits could facilitate management of risk for poor developmental outcomes by providing objective evidence of altered responses to speech prior to the emergence of behavioral symptoms.

The majority of ERP studies examining auditory processing in individuals with ASD had used nonspeech tones (see Haesen et al. 2011, Hitoglou et al., 2010 for reviews). However, speech sound stimuli could be more relevant to understanding possible sources of language deficits, because speech sounds are more complex than tones, and their processing relies on brain mechanisms overlapping with, but not identical to, those engaged by nonspeech stimuli (Dehaene-Lambertz et al., 2005).

The existing auditory ERP studies in typical and atypical populations can be broadly categorised into those assessing speech sound processing and those measuring

differentiation of contrasting speech stimuli. Speech processing studies may include a single stimulus or multiple stimuli of interest and focus on ERP characteristics (e.g., amplitude or latency of a peak) associated with the individual sounds. In children with ASD, speech processing studies reported reduced amplitudes of P1/N1 responses following passive exposure to a single syllable (Russo et al., 2009) or showed atypical hemisphere lateralization of N1 latency and amplitude favouring the right hemisphere during an active syllable detection task (Dawson et al., 1989), suggesting altered sensory processes.

In contrast, speech sound differentiation studies use at least two stimuli that differ on one critical dimension (e.g. voice onset timing, place of articulation) and examine the difference in ERPs as the metric of stimulus differentiation. Prior speech differentiation studies in children with ASD included mainly vowels, and produced varied findings, suggesting intact or enhanced (Kemner et al., 1995; Lepisto et al., 2005, 2008; Whitehouse & Bishop, 2008), reduced (Ceponiene et al., 2003), or delayed (Oram Cardy et al., 2005) stimulus differentiation, as well as atypical right hemisphere lateralization of brain responses (Flagg et al., 2005; Lepisto et al., 2006). This variability in results could not be attributed to task demands, as the majority of the studies involved passive listening. Fewer studies in ASD have examined differentiation of consonant-vowel syllables, which are building blocks of English words. In two published reports, differences in ERPs elicited by the contrasting consonants were delayed (Jansson-Verkasalo et al., 2003) or reduced/absent (Kuhl et al., 2005).

Many of these studies employed an oddball stimulus presentation paradigm and its accompanying mismatch negativity (MMN) response, but this approach arguably has several limitations (see Kujala et al., 2007 for a review). In particular, the MMN response is not always detectable, even when behavioral responses evidence stimulus discrimination (e.g., Sharma et al., 2006; Wunderlich & Cone-Wesson, 2001). This could be due to masking by overlapping N1 or P3 responses (Duncan et al., 2009; Näätänen et al., 2005). Recently, Kujala et al. (2010) included consonant contrasts in a multi-feature MMN paradigm but observed no MMN responses in either the ASD or typical group (age 8–12 years); thus no conclusions about consonant differentiation could be made. The likelihood of an absent MMN responses increases with decreasing age (Cheour et al., 1998; Leppanen et al., 1997). Moreover, in young school-age children, the MMN response demonstrated poor test-retest reliability (Raikkonen et al., 2003; Uwer & von Suchodoletz, 2000), reducing its value for investigating neural mechanisms underlying verbal skills in young children with ASD.

The present study was designed to expand prior findings regarding speech sound processing in children with ASD in several ways: (1) by including a more diverse set of consonant-vowel syllables; (2) by employing equiprobable stimulus presentation that has been previously used to study speech sound differentiation in children with and without developmental disabilities (Molfese & Molfese, 1997; Espy et al., 2004; Key et al., 2009; Yoder et al., 2006; McArthur et al., 2010); and (3) by explicitly testing whether speech-sound differentiation ability (difference in ERPs to contrasting stimuli) could explain the difference in the extent of nonverbal vs. verbal ability discrepancy in children with ASD compared to chronological age-matched children with typical development.

We expected syllable differentiation to be reflected in amplitude differences of early auditory ERP responses (P1-N1-P2), known to be sensitive to stimulus physical characteristics and perceptual categorization. We hypothesised that children with ASD would process consonant-vowel syllables atypically, resulting in reduced differences in ERP responses to contrasting speech stimuli, which could in turn contribute to greater nonverbal-verbal discrepancies in children with ASD than typical peers. The comparison group matched on chronological age and nonverbal ability allowed us to focus on between-group differences in language deficits by controlling for factors related to maturation, perceptual experience, and intellectual ability, which may affect ERP responses to speech sounds.

Method

Participants

Forty-two children (35 males, 7 females) participated in the study. Eighteen children (4 females), M age 7.14 years ($SD=1.45$), were typically developing (TD) with no reported family history of ASD. Twenty-four children (3 females), M age 6.71 years ($SD=1.34$), had a diagnosis of ASD based on the DSM-IV definition (14 – autism, 3 – Asperger syndrome, 7 – pervasive developmental disorder—not otherwise specified). Diagnoses were made by licensed psychologists, and 16/24 (67%) children were diagnosed at a university-based autism clinic using the ADOS either alone or in combination with the ADI-R to support the clinical diagnosis. Of the remaining children, four (17%) were diagnosed by community professionals, and two (8%) by their school systems. Parents of two participants (8%) did not provide the information about the source of the diagnosis. The participants with ASD were required to have sufficient language skills and intellectual ability to understand and follow simple instructions associated with the study protocol.

The participant groups did not differ significantly in age ($p=.326$, $d=.308$) or nonverbal ability ($p=.178$, $d=.433$; see Table 1 for a summary of participant characteristics). However, the TD group had higher verbal abilities and higher overall cognitive abilities, as measured by Kaufman Brief Intelligence Test-2 (K-BIT2; Kaufman & Kaufman, 2004). Three children in each group were left-handed, the rest were right-handed, as determined by the Edinburgh Handedness Inventory (Oldfield, 1971), $M_{ASD} = .66$, $SD=.49$, $M_{TD} = .72$, $SD=.39$. All children were reported by parents to have normal hearing and vision.

Four additional children with ASD were excluded from the study due to the lack of cooperation with the electrode placement. Data from one TD child were excluded due to an insufficient number of artifact-free trials. The research was prospectively reviewed and approved by the university ethics committee. Parents provided written informed consent, and children provided their assent prior to completing the study procedures.

Procedure

All data were collected during a single visit. The behavioural testing preceded the ERP recording.

Behavioural testing—Kaufman Brief Intelligence Test-2 (K-BIT2) is a brief assessment of verbal and nonverbal abilities in persons 4–90 years of age. K-BIT2 provides Verbal and

Nonverbal standard scores and an overall Composite IQ score, each with a mean of 100 and a standard deviation of 15. The scores have high internal consistency (for children 4–18 years, mean=.88-.90). K-BIT2 is widely used in research settings and is suitable for typical individuals and those with intellectual and developmental disabilities, including ASD (Klinger et al., 2009). It was administered individually by a trained research assistant and lasted approximately 15 minutes.

ERP acquisition—ERP data were acquired individually in a sound-attenuated room using a 128-channel Geodesic Sensor Net with Ag/AgCl electrodes (EGI, Inc., Eugene, OR) connected to high-impedance amplifiers. Data were sampled at 250Hz with the filters set to . 1–30 Hz and electrode impedance levels below 40 kOhm. All electrodes were referred to vertex during recording and re-referenced offline to an average reference.

ERP stimuli—The stimuli included six consonant-vowel (CV) syllables /ba/, /da/, /ga/, /bu/, /du/, /gu/, originally employed by Stevens and Blumstein (1978) and accurately identified by the adult subjects in that study as members of their respective phonetic categories. A single five-formant token for each CV was synthesised on a Klatt (cascade) synthesiser, so that the amplitude of individual formants was modulated as a function of the respective formant frequencies, as in natural speech. The central frequencies of the steady-state portion of the formants were kept constant across the consonants and only varied as a function of the vowel sounds. Duration of F1 transition ranged between 15–45 ms across tokens depending on the initial consonant and the following vowel. Transition duration for all the other formants was 40 ms. Consonants were followed by a 250 ms steady state vowel. Rise and decay times were equivalent across sounds.

ERP paradigm—To reduce possible confounds related to motivation and attention, to facilitate comparisons with prior results obtained using passive listening paradigms, and based on recent findings suggesting that greater group differences in auditory processing between typical individuals and those with ASD may be observed in passive tasks (Dunn et al., 2007; Whitehouse & Bishop, 2008), the participants were instructed to sit quietly and listen to the speech sounds, but no active stimulus identification or behavioral responses were required. The syllables were presented at 75dB SPL(A) as measured at the ear level, through a speaker positioned approximately 100cm above the midline of the child's head. All speech sounds were presented in random order, 25 times each, for a total of 150 trials over the course of 10 minutes. Interstimulus intervals varied randomly between 1.6–2.6 seconds to prevent habituation to stimulus onset.

Recording of the brainwaves was controlled by Net Station software (v.4.3; EGI, Inc., Eugene, OR). E-Prime (v.1.2; PST, Inc., Pittsburgh, PA) controlled stimulus presentation. A researcher present in the testing room continuously monitored child's behaviour and electroencephalogram (EEG). If artifacts contaminated EEG for two consecutive trials, stimulus presentation could be manually suspended until the source of noise was addressed, and resumed when EEG signal was optimal.

Data Analysis

The EEG was segmented on stimulus onset to include a 100-ms prestimulus baseline and a 700-ms post-stimulus interval. Trials contaminated by ocular and movement artifacts were rejected from further analysis using an automated screening algorithm in NetStation followed by a manual review. Data from electrodes with poor signal quality were reconstructed using spherical spline interpolation (Srinivasan et al., 1998). If more than 15 electrodes (12% of the array) within a trial had poor signal, the entire trial was discarded. To be included in the statistical analyses, each participant had to have a minimum of 10 artifact-free trials per stimulus condition. Although on average slightly more trials were retained for children with ASD than with TD, $M_{ASD} = 15.72$, $SD = 3.72$; $M_{TD} = 13.73$, $SD = 2.33$, these group differences only approached significance ($p = .053$). There were no significant correlations between the number of trials retained and age, IQ, or nonverbal-verbal IQ discrepancy for the combined sample or for either of the participant groups.

Artifact-free ERPs were averaged and baseline-corrected by subtracting the average microvolt value across the 100-ms prestimulus interval from the post-stimulus segment. To reduce the number of potential significance tests (6 sounds x 128 electrodes x 175 time samples = 134,400 data points per participant), a spatial principal components analysis (sPCA) using a covariance matrix and Promax rotation was performed to group data from 128 electrodes into a small set of 'virtual electrodes', each representing a spatially contiguous group of electrodes with similar ERP waveforms (Spencer et al., 1999). Specific electrodes comprising each cluster were identified using the criterion of factor loadings of $|\cdot| \geq .6$ or greater. Data for individual electrodes within each cluster were averaged. Next, clustered data were submitted to a temporal PCA (tPCA) with Varimax rotation. This process reduced 175 time samples (700 ms) to a small set of non-correlated components corresponding to the temporal windows of maximal variability in the waveforms, such as slopes or peaks. The number of tPCA factors was determined using the Scree Test (Cattell, 1966), and temporal windows were labeled based on consecutive time samples with factor loadings of $|\cdot| \geq .6$ or greater.

To further limit the number of variables in the statistical analyses, only data for the anterior and posterior temporal scalp regions (overlapping T3/T4 and T5/T6 locations previously used in ERP studies of speech sound processing; Figure 1) were used as dependent variables. The repeated-measures ANOVA examined speech sound differentiation using a 4-factor design of Consonant (3: b, d, g) x Vowel (2: a, u) x Electrode (2: anterior temporal, posterior temporal) x Hemisphere (2: left, right), with Huynh-Feldt correction. Significant interactions involving Consonant or Vowel factors were further examined using pair-wise t -tests. The false discovery rate (FDR) approach (Benjamini & Hochberg, 1995) was used to control for multiple significance tests in the follow-up analyses. This approach was selected over the Bonferroni correction (Hochberg, 1988) because the former is more statistically powerful and fits the inter-correlational nature of ERP data better than the latter (Crowley et al., 2012).

To constrain the number of significance tests even further, ERPs of the TD sample were analyzed first to identify the particular temporal intervals and speech sound contrasts where typical children demonstrated statistically significant speech sound differentiation. We then

tested the hypotheses of atypical speech sound differentiation in children with ASD by examining the presence of stimulus-related differences only in that subset of ERPs presumably reflecting typical speech sound processing. Between-stimuli differences in factor scores were the dependent measures in one-way ANOVAs with Group (ASD vs. TD) as the between-subject factor.

To examine whether speech sound differentiation reflected in ERP response differences is related to nonverbal-verbal skill discrepancy, Pearson correlations were tested between ERP variables showing significant stimulus differentiation and the difference score between nonverbal and verbal standard scores on the K-BIT2. Finally, a mediation analysis (Tofighi & MacKinnon, 2011) was conducted to test the hypothesis that ERP measures showing significant group differences in speech sound differentiation mediate the relationship between the diagnostic group status (TD vs. ASD) and nonverbal-verbal skill discrepancy. Because typical children were expected to show greater ERP differentiation of speech sounds and reduced nonverbal-verbal ability discrepancies, a 1-tailed test was used to judge the significance of the indirect path in the mediation analysis.

Results

The temporal PCA applied to the entire data set identified five temporal windows accounting for 83.97% of the total variance in ERPs: 0–76 ms (Factor 5, 8.67%), 84–196 ms (Factor 4, 13.78%), 196–308 ms (Factor 2, 14.37%), 320–444 ms (Factor 3, 14.00%), 432–700 ms (Factor 1, 33.16%).

Typical speech sound processing

Within the TD group, the repeated-measures ANOVA identified consonant and/or vowel related differences in ERPs for four of the five temporal windows (excluding Factor 5 because it was too early for speech differentiation to affect). Of the remaining factors, the earliest interval, 84–196 ms, was characterised by a main effect of Consonant, $F(2,24)=5.135$, $p=.011$, partial $\eta^2=.232$, and interactions of Consonant x Vowel x Hemisphere, $F(2,34)=6.296$, $p=.005$, partial $\eta^2=.270$, and Consonant x Vowel x Electrode x Hemisphere, $F(2,34)=4.328$, $p=.021$, partial $\eta^2=.203$. Follow-up analyses revealed a number of significant speech sound contrasts, but only the consonant difference associated with the main effect remained significant following the FDR procedure: overall, /g/-initial syllables elicited more positive responses than /b/-initial syllables, $t(17)=3.238$, $p=.005$, $d=.763$ (Figure 2).

In the 196–308 ms interval, there was a Consonant x Electrode interaction, $F(2,34)=5.114$, $p=.013$, partial $\eta^2=.231$, where /b/-syllables elicited more positive amplitudes than /d/-syllables, $t(17)=3.650$, $p=.002$, $d=.860$, over the anterior temporal sites. There were no significant differences at the posterior electrodes, p 's=.072-.642.

The 320–444ms interval was characterised by a Vowel x Electrode interaction, $F(1,17)=7.472$, $p=.014$, partial $\eta^2=.305$. Follow-up analyses indicated that /a/-ending sounds elicited more positive amplitudes than /u/-ending stimuli at the anterior temporal sites, $t(17)=2.632$, $p=.017$, $d=.62$. As is common in ERP studies (Korzyukov et al., 2003; Woldorf

et al., 1998), the direction of these differences was reversed at the posterior temporal sites, $t(17)=2.193$, $p=.042$, $d=.52$.

In the final interval of 432–700 ms, there was a Consonant x Electrode interaction, $F(2,34)=4.439$, $p=.019$, partial $\eta^2=.207$, due to the /b/-syllables eliciting more positive waveforms than /g/-syllables at posterior temporal locations, $t(17)=3.074$, $p=.007$, $d=.725$. There was also a Consonant x Vowel x Hemisphere interactions, $F(2,34)=5.510$, $p=.010$, partial $\eta^2=.245$, but the follow-up analyses were not significant after the FDR correction.

In summary, stimuli differentiation was observed in the TD sample at 84–196 ms and 432–700 ms between /g/ and /b/, at 196–308 ms between /b/ and /d/, and at 320–444 ms between /a/ and /u/.

Between-Group Differences in Speech Sound Differentiation

Group differences between children with ASD and TD were observed for the /g/-/b/ contrast in 84–196 ms interval, $F(1,40)=6.994$, $p=.012$, $d=.848$, and for the /b/-/d/ contrast at the anterior temporal locations within 196–308 ms, $F(1,40)=3.976$, $p=.053$, $d=.623$ (Figure 3). In both cases, larger differences in ERPs elicited by the contrasting speech sounds were observed in the TD than ASD group. The /a/-/u/ difference in the 320–444 ms interval and the /g/-/b/ difference in 432–700 ms interval that were identified in the TD group did not yield significant between-group differences ($ps = .063-.094$).

Between-group differences in the discrepancy between nonverbal and verbal scores

The participant groups were statistically different in the extent of the nonverbal-verbal discrepancy, $F(1,40)=6.371$, $p=.016$, $d=.797$, with lower mean discrepancy values in children with TD than ASD. For 5- to 10- year-old children, a discrepancy of 15 points or greater between verbal and nonverbal standard scores on the K-BIT2 is considered to be a noteworthy difference (Kaufman & Kaufman, 2004). In the group of children with ASD, 29% (7/24) demonstrated such discrepancy favoring nonverbal performance, while 8% (2/24) had notably higher verbal than nonverbal scores. In the TD group, 11% (2/18) demonstrated the discrepancy toward higher nonverbal performance, and 28% (5/18) scored higher in the verbal than nonverbal domain.

Brain-behaviour correlations

For the combined sample (TD and ASD), the size of the /b/-/d/ difference at the anterior temporal locations within 196–308 ms was negatively correlated with the extent of discrepancy in K-BIT2 nonverbal vs. verbal standard scores, $r=-.399$, $p=.009$. This means that greater differences in ERP responses to speech sounds were associated with lower nonverbal-verbal difference score. The extent of discrepancy did not correlate with the overall IQ in the combined sample or in either of the groups ($ps=.383-.852$).

Mediation analysis

As demonstrated in Figure 4, difference in ERP responses for the /b/-/d/ contrast at the anterior temporal site during the 196–308 ms window mediated the relationship between diagnostic group membership and nonverbal-verbal score discrepancy. The 90% confidence

interval (the appropriate interval for a 1-tailed test) for the product of the “a” and “b” paths (i.e., the parameter for the indirect relation) did not include zero (.400 – 9.675). R-square change values for paths “a” and “b” were .15 and .08, respectively.

Discussion

The purpose of this study was to examine whether, as a group, children with ASD process consonant-vowel speech stimuli in an atypical way, and whether the brain’s ability to differentiate speech sounds relates to individual differences in language deficits, quantified as the degree of discrepancy in nonverbal-verbal abilities, and mediates the relationship between diagnostic group membership and such discrepancy. Our results indicated that children with ASD evidenced smaller than typical differences in ERP responses to consonant contrasts within 84–308 ms after stimulus onset, and the extent of speech sound differentiation is related to individual differences in nonverbal vs. verbal abilities.

We implemented a normative analytic approach, where ERPs of typical children were analysed first to establish the pattern of presumed typical speech sound processing in the brain, which then guided subsequent analyses. Consonant differences were detected within 84–196 ms (/b/-/g/), 196–308 ms (/b/-/d/), and 432–700 ms (/b/-/g/) periods, while vowel differentiation (/a/-/u/) was observed in 320–444 ms interval. Next, the differences in brain responses elicited by these sound contrasts were compared between the two diagnostic groups. Children with ASD evidenced reduced consonant differentiation in the 84–196 ms and 196–308 ms temporal windows, while no group differences reached significance for the remaining two intervals.

The pattern of diagnostic group differences in ERP responses is consistent with the previous studies reporting reduced consonant differentiation in children with ASD (Kuhl et al., 2005) but no group differences in vowel differentiation between children with ASD and TD (e.g., Kemner et al., 1995; Lepisto et al., 2005). This consistency in results obtained using different speech stimuli and ERP procedures is especially important because it demonstrates that reported findings are unlikely the result of multiple significance testing and that using consonant-vowel syllables in an equiprobable design is a construct valid way to evaluate speech sound processing in children with ASD. The passive nature of the equiprobable paradigm coupled with its relatively short duration makes it appealing for future use in younger children and in infants at risk for ASD or other communication disorders.

The presence of significant correlations between ERP differences for the contrasting speech sounds and behavioural evidence of nonverbal-verbal discrepancy supports the premise that language deficits that are not explained by cognitive abilities might be explained by individual differences in the brain’s ability to differentiate speech stimuli. Incidentally, the 29% rate of K-BIT2 score discrepancy toward higher nonverbal than verbal skills in 5- to 10- year-old children with ASD in the present study was very similar to the 34% rate previously reported for 6–13-year-olds with ASD using Differential Ability Scales, a more comprehensive measure of cognitive ability (Joseph et al., 2002; Tager-Flusberg & Joseph, 2003). This replication of the prevalence rates in a different sample with a different

assessment tool confirms that a noteworthy minority of children with ASD has verbal communication disorder in the absence of intellectual disability.

An exploratory mediation analysis further indicated that the brain's ability to respond differently to contrasting speech sounds mediates the relationship between the diagnostic group status and nonverbal-verbal discrepancy. This finding is new to the literature. While the design does not allow confident causal inferences, the significant indirect effect provides correlational support for the hypothesis that atypical speech sound differentiation in the brain explains, at least in part, greater instances of a substantial nonverbal-verbal ability discrepancy in children with ASD vs. TD. The reported association between the neural evidence of diminished consonant discrimination and the nonverbal-verbal skill discrepancy in children with ASD opens a new direction for the development of novel treatment targets as well as early screening tools that could be used in young children or even infants to manage risk for poor communicative outcomes.

While this study yielded novel findings, it has a number of limitations. Although not unusual in size for an ERP study, the typical participant sample was relatively small for the purpose of a true normative analysis, creating the possibility that brain responses of the TD group were not entirely representative of speech sound processing in the general population. Additionally, while the differences in nonverbal functioning between children with TD and ASD, were not statistically significant, some might consider them noteworthy. The TD sample could be representing a higher ability subgroup of the general population, as frequently happens in university-based samples of convenience. Yet, the average nonverbal K-BIT2 scores for both groups were very close to the mean, and the standardized between-group mean difference of .43 corresponds to a small-to-medium effect size, suggesting that the two groups were comparable.

Our sample of children with ASD was very heterogeneous, representing a spectrum of possible phenotypes. Compared to a more diagnostically or severity-level restricted sample, this high variability may have inflated the correlation between speech differentiation measured by ERPs and language deficits reflected by the nonverbal-verbal standard score discrepancy. Future studies with larger representative samples are needed to determine the external validity of the current findings. Related to the issue of the sample composition is the concern regarding diagnosis. While the majority of the participants in the present sample were diagnosed using ADOS/ADI-R, the instruments used to diagnose some children were unknown and thus may have not been assessed using these gold-standard tools, thereby contributing to increased variability within the ASD group. Finally, all intact group comparison and correlational results are subject to explanation by unmeasured third variables. The K-BIT2 provides a very basic assessment of children's verbal and nonverbal skills. Although it is considered to be acceptable for use with children with ASD 4 years of age or older (Klinger et al., 2009), future studies would need to include a wider range of standardised assessments in order to determine the degree to which discrepancies in verbal and nonverbal abilities are different between diagnostic groups and are related to brain measures of speech sound processing.

In summary, the current study is a step toward providing one explanation for language deficits unrelated to IQ (quantified as nonverbal-verbal skill discrepancy) in children with ASD. Improved understanding of the relationship between the brain's response to speech sound contrasts and nonverbal vs. language ability in ASD could lead in the future to the development of new approaches for identifying risks for language deficits at an earlier developmental stage.

Acknowledgments

The authors have no conflicts of interest to declare. This work was supported in part by NICHD Grant P30 HD15052 to Vanderbilt Kennedy Center and by a Vanderbilt Kennedy Center Hobbs Discovery Award to Dr. Alexandra P. Key. We are grateful to Susan M. Williams, Dorita Jones, and Amber Vinson for assistance with acquisition and processing of the ERP data.

Literature Cited

- American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders 5*. Washington, DC: Author.
- Benasich AA, Thomas J, Choudhury N, & Leppanen P (2002). The importance of rapid auditory processing abilities to early language development: Evidence from converging methodologies. *Developmental Psychobiology*, 40, 278–292. [PubMed: 11891639]
- Benjamini Y, & Hochberg Y (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, B*, 57, 289–300.
- Cattell RB (1966) The scree test for the number of factors. *Multivariate Behavioural Research* 1, 245–76.
- eponien R, Lepistö T, Shestakova A, Vanhala R, Alku P, Näätänen R, & Yaguchi K (2003). Speech-sound-selective auditory impairment in children with autism: They can perceive but do not attend. *Proceedings of the National Academy of Sciences of the United States of America*, 100(9), 5567–5575. [PubMed: 12702776]
- Cheour M, Ceponiele R, Lehtokoski A, Luuk A, Alli J, Alho K, & Näätänen R (1998). Development of language-specific phoneme representations in the infant brain. *Nature Neuroscience*, 1, 351–353. [PubMed: 10196522]
- Crowley MJ, Wu J, McCreary S, Miller KF, & Mayes LC (2012). Implementation of false discovery rate for exploring novel paradigms and trait dimensions with ERPs. *Developmental Neuropsychology*, 37(6), 559–577. doi:10.1080/87565641.2012.694513 [PubMed: 22889345]
- Dawson G, Finley C, Phillips S, & Lewy A (1989). A comparison of hemispheric asymmetries in speech-related brain potentials of autistic and dysphasic children. *Brain and Language*, 37(1), 26–41. [PubMed: 2473824]
- Dehaene-Lambertz G, Pallier C, Serniclaes W, Sprenger-Charolles L, Jobert A, & Dehaene S (2005). Neural correlates of switching from auditory to speech perception. *NeuroImage*, 24(1), 21–33. doi: 10.1016/j.neuroimage.2004.09.039 [PubMed: 15588593]
- Duncan CC, Barry RJ, Connolly JF, Fischer C, Michie PT, Näätänen R, et al. (2009). Event-related potentials in clinical research: Guidelines for eliciting, recording, and quantifying mismatch negativity, P300, and N400. *Clinical Neurophysiology*, 120(11), 1883–1908. [PubMed: 19796989]
- Dunn MA, Gomes H, & Gravel J (2007). Mismatch negativity in children with autism and typical development. *Journal of Autism and Developmental Disorders*, 38(1), 52–71. 10.1007/s10803-007-0359-3 [PubMed: 17624605]
- Espy K, Molfese D, Molfese V, & Modglin A (2004). Development of auditory event-related potentials in young children and relations to word-level reading abilities at age 8 years.. *Annals of Dyslexia*, 54(1), 9–38. [PubMed: 15765002]
- Flagg EJ, Cardy JEO, Roberts W, & Roberts TPL (2005). Language lateralization development in children with autism: Insights from the late field magnetoencephalogram. *Neuroscience Letters*, 386(2), 82–87. doi:10.1016/j.neulet.2005.05.037 [PubMed: 16046066]

- Haesen B, Boets B, & Wagemans J (2011). A review of behavioral and electrophysiological studies on auditory processing and speech perception in autism spectrum disorders. *Research in Autism Spectrum Disorders*, 5(2), 701–714. doi:10.1016/j.rasd.2010.11.006
- Herbert MR, & Kenet T (2007). Brain abnormalities in language disorders and in autism. *Pediatric Clinics of North America*, 54(3), 563–583. doi:10.1016/j.pcl.2007.02.007 [PubMed: 17543910]
- Hitoglou M, Ververi A, Antoniadis A, & Zafeiriou DI (2010). Childhood autism and auditory system abnormalities. *Pediatric Neurology*, 42(5), 309–314. doi:10.1016/j.pediatrneurol.2009.10.009 [PubMed: 20399382]
- Hochberg Y (1988). A sharper Bonferroni procedure for multiple significance testing. *Biometrika*, 75, 800–803.
- Jansson-Verkasalo E, eponien R, Kielinen M, Suominen K, Jäntti V, Linna S-L, Moilanen I, et al. (2003). Deficient auditory processing in children with Asperger Syndrome, as indexed by event-related potentials. *Neuroscience Letters*, 338(3), 197–200. doi:10.1016/S0304-3940(02)01405-2 [PubMed: 12581830]
- Joseph RM, Tager-Flusberg H, & Lord C (2002). Cognitive profiles and social-communicative functioning in children with autism spectrum disorder. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 43(6), 807–821.
- Kaufman A, & Kaufman N. (2004). *Kaufman Brief Intelligence Test*, Second Edition Circle Pines, MN: AGS Publishing.
- Kemner C, Verbaten MN, Cuperus JM, Camfferman G, & van Engeland H (1995). Auditory event-related brain potentials in autistic children and three different control groups. *Biological Psychiatry*, 38, 150–165. [PubMed: 7578658]
- Key A, Molfese D, O'Brien L, & Gozal D (2009). Sleep-disordered breathing affects auditory processing in 5–7 year-old children: evidence from brain recordings.. *Developmental Neuropsychology*, 34(5), 1–15. [PubMed: 19142764]
- Kjelgaard MM, & Tager-Flusberg H (2001). An investigation of language impairment in autism: Implications for genetic subgroups. *Language and Cognitive Processes*, 16(2–3), 287–308. doi: 10.1080/01690960042000058 [PubMed: 16703115]
- Klinger L, O'Kelley S, & Mussey J (2009). Assessment of intellectual functioning in autism spectrum disorders In Goldstein S, Naglieri J, & Ozonoff S (Eds.), *Assessment of autism spectrum disorders* (pp. 209–252), New York, NY: Guilford Press.
- Korzyukov OA, Winkler I, Gumenyuk VI, & Alho K (2003). Processing abstract auditory features in the human auditory cortex. *NeuroImage*, 20(4), 2245–2258. [PubMed: 14683726]
- Kujala T, Tervaniemi M, & Schröger E (2007). The mismatch negativity in cognitive and clinical neuroscience: Theoretical and methodological considerations. *Biological Psychology*, 74, 1–19. [PubMed: 16844278]
- Kuhl PK, Coffey Corina S, Padden D, & Dawson G (2005). Links between social and linguistic processing of speech in preschool children with autism: Behavioral and electrophysiological measures. *Developmental Science*, 8(1), F1–F12. [PubMed: 15647058]
- Kujala T, Kuuluvainen S, Saalasti S, Jansson-Verkasalo E, Wendt, von L, & Lepistö T (2010). Speech-feature discrimination in children with Asperger syndrome as determined with the multi-feature mismatch negativity paradigm. *Clinical Neurophysiology*, 121(9), 1410–1419. doi:10.1016/j.clinph.2010.03.017 [PubMed: 20382070]
- Kwok E, Brown HM, Smyth RE, & Cardy JO (2015). Meta-analysis of receptive and expressive language skills in autism spectrum disorder. *Research in Autism Spectrum Disorders*, 9 10.1016/j.rasd.2014.10.008
- Leppänen PHT, Eklund KM, & Lyytinen H (1997). Event-related brain potentials to change in rapidly presented acoustic stimuli in newborns. *Developmental Neuropsychology*, 13(2), 175–204. doi: 10.1080/87565649709540677
- Lepistö T, Silokallio S, Nieminen-von Wendt T, Alku P, Naatanen R, & Kujala T (2006). Auditory perception and attention as reflected by the brain event-related potentials in children with Asperger syndrome. *Clinical Neurophysiology*, 117, 2161–2171. [PubMed: 16890012]

- Lepistö T, Kajander M, Vanhala R, Alku P, Huotilainen M, Nääätänen R, & Kujala T (2008). The perception of invariant speech features in children with autism. *Biological Psychology*, 77(1), 25–31. doi:10.1016/j.biopsycho.2007.08.010 [PubMed: 17919805]
- Lepistö T, Kujala T, Vanhala R, Alku P, Huotilainen M, & Nääätänen R (2005). The discrimination of and orienting to speech and non-speech sounds in children with autism. *Brain Research*, 1066(1–2), 147–157. doi:10.1016/j.brainres.2005.10.052 [PubMed: 16325159]
- Lincoln A, Hansel E, & Quirmach L (2007). Assessing intellectual abilities of children and adolescents with autism and related disorders In Schopler E & Mesibov G (Eds.), *The clinical assessment of children and adolescents: A practitioner's handbook* (pp. 527–544). Mahwah, NJ: Erlbaum.
- Loucas T, Charman T, Pickles A, Simonoff E, Chandler S, Meldrum D, & Baird G (2008) Autistic symptomatology and language ability in autism spectrum disorder and specific language impairment. *Journal of Child Psychology and Psychiatry*, 49, 1184–1192 [PubMed: 19017030]
- Mayes S, & Calhoun S (2003). Analysis of WISC-III, Stanford-Binet:IV, and academic achievement test scores in children with autism. *Journal of Autism and Developmental Disorders*, 33, 329–341. [PubMed: 12908835]
- McArthur GM, & Bishop DVM (2005). Speech and non-speech processing in people with specific language impairment: A behavioral and electrophysiological study. *Brain and Language*, 94(3), 260–273. doi:10.1016/j.bandl.2005.01.002 [PubMed: 16098376]
- McArthur G, Atkinson C, & Ellis D (2010). Can training normalize atypical passive auditory ERPs in children with SRD or SLI?. *Developmental Neuropsychology*, 35(6), 656–678. [PubMed: 21038159]
- Molfese D, & Molfese V (1997). Discrimination of language skills at five years of age using event related potentials recorded at birth. *Developmental Neuropsychology*, 13, 135–156.
- Nääätänen R, Jacobsen T, & Winkler I (2005). Memory-based or afferent processes in mismatch negativity (MMN): A review of the evidence. *Psychophysiology*, 42(1), 25–32. [PubMed: 15720578]
- Oldfield RC (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia*, 9(1), 97–113. [PubMed: 5146491]
- Oram Cardy JE, Flagg EJ, Roberts W, & Roberts TPL (2005). Delayed mismatch field for speech and non-speech sounds in children with autism. *NeuroReport*, 16(5), 521–525. [PubMed: 15770164]
- Räikkönen K, Birkás E, Horváth J, Gervai J, & Winkler I (2003). Test-retest reliability of auditory ERP components in healthy 6-year-old children. *NeuroReport*, 14(16), 2121–2125. [PubMed: 14600509]
- Russo N, Zecker S, Trommer B, Chen J, & Kraus N (2009). Effects of background noise on cortical encoding of speech in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 39(8), 1185–1196. doi:10.1007/s10803-009-0737-0 [PubMed: 19353261]
- Shafer VL, Morr ML, Datta H, Kurtzberg D, & Schwartz RG (2005). Neurophysiological indexes of speech processing deficits in children with specific language impairment. *Journal of Cognitive Neuroscience*, 17(7), 1168–1180. [PubMed: 16138434]
- Sharma M, Purdy SC, Newall P, Wheldall K, Beaman R, & Dillon H (2006). Electrophysiological and behavioral evidence of auditory processing deficits in children with reading disorder. *Clinical Neurophysiology*, 117(5), 1130–1144. [PubMed: 16564738]
- Spencer KM, Dien J, & Donchin E (1999). A componential analysis of the ERP elicited by novel events using a dense electrode array. *Psychophysiology*, 36(3), 409–414. [PubMed: 10352565]
- Srinivasan R, Tucker DM, & Murias M (1998). Estimating the spatial Nyquist of the human EEG. *Behavior Research Methods, Instruments, and Computers*, 30, 8–19.
- Stark RE, & Tallal P (1981). Selection of children with specific language deficits. *The Journal of Speech and Hearing Disorders*, 46(2), 114–122. [PubMed: 7253588]
- Stevens KN, & Blumstein SE (1978). Invariant cues for place of articulation in stop consonants. *Journal of the Acoustical Society of America*, 64, 1358–1368. [PubMed: 744836]
- Tager-Flusberg H (2006). Defining language phenotypes in autism. *Clinical Neuroscience Research*, 6(3–4), 219–224. doi:10.1016/j.cnr.2006.06.007
- Tager-Flusberg H & Joseph RM (2003) Identifying neurocognitive phenotypes in autism. *Philosophical Transactions of the Royal Society, London. Ser. B* 358, 303–314.

- Tofighi D & MacKinnon DP (2011). RMediation: An R package for mediation analysis confidence intervals. *Behaviour Research Methods*, 43, 692–700.
- Uwer R, & von Suchodoletz W (2000). Stability of mismatch negativities in children. *Clinical Neurophysiology*, 111(1), 45–52. [PubMed: 10656510]
- Whitehouse AJO, & Bishop DVM (2008). Do children with autism “switch off” to speech sounds? An investigation using event-related potentials. *Developmental Science*, 11(4), 516–524. doi: 10.1111/j.1467-7687.2008.00697.x [PubMed: 18576959]
- Woldorff MG, Hillyard SA, Gallen CC, Hampson S, & Bloom FE (1998). Magnetoencephalographic recordings demonstrate attentional modulation of mismatch-related neural activity in human auditory cortex. *Psychophysiology*, 35, 283–292. [PubMed: 9564748]
- Wunderlich JL, & Cone-Wesson BK (2001). Effects of stimulus frequency and complexity on the mismatch negativity and other components of the cortical auditory-evoked potential. *Journal of the Acoustical Society of America*, 109(4), 1526–1537. [PubMed: 11325124]
- Yoder PJ, Camarata S, Camarata M, & Williams SM (2006). Association between differentiated processing of syllables and comprehension of grammatical morphology in children with Down syndrome. *American Journal of Mental Retardation*, 111(2), 138–152. [PubMed: 16466285]

Figure 1.
128-channel electrode layout and the electrode clusters used in the analyses.

Figure 2.

Average ERP waveforms in response to consonant contrasts at anterior and posterior temporal locations in typical children.

Figure 3.
Average ERP waveforms in response to consonant contrasts at anterior and posterior temporal locations in children with ASD.

Figure 4.

The putative mediated relation between diagnostic group membership and language impairment (nonverbal-verbal standard score discrepancy) through brain response to /b/-/d/ contrast. The standardised coefficient in parentheses is the effect of diagnostic group on K-BIT score discrepancy when brain measures are not included in the model.

* $p < .05$

$p < .06$

Table 1.

Participant sample characteristics: Means (SD) and ranges.

	ASD		TD		p-value
N (m/f)	24 (21/3)		18 (15/3)		
Age, yrs	6.71 (1.34)	5.08 – 9.92	7.14 (1.45)	5 – 10.17	0.326
Handedness	.66 (.49)	–.77 – 1	.72 (.39)	–.07 – 1	0.643
K-BIT2 IQ	93.88 (17.41)	60 – 135	110.44 (13.05)	88 – 136	0.001
Verbal	90.46 (17.48)	49 – 124	111.61 (12.29)	82 – 127	<.001
Nonverbal	98.54 (18.10)	61 – 138	105.78 (15.17)	80 – 138	0.178
Nonverbal-Verbal Discrepancy	8.08 (18.95)	–24 – 57	–5.83 (15.80)	–45 – 19	0.016
% Significant Nonverbal > Verbal *	29 (7/24)		11 (2/18)		

* Discrepancy of 15 points or more.