Total Mortality, Major Adverse Cardiac Events, and Echocardiographic-Derived Cardiac Parameters with Fragmented QRS Complex

Bojun Gong, M.D., and Zicheng Li, Ph.D.

From the Department of Cardiology, First Affiliated Hospital of Jinan University, Guangzhou, China

Background: Fragmented QRS complex (fQRS), an easily evaluated noninvasive electrocardiographic parameter, is associated with worse outcomes in patients with several cardiovascular conditions. The presence of fQRS on ECG may be an indicator of myocardial damage in patients with coronary artery disease (CAD). In this article, we performed a meta-analysis in order to characterize the presence of fQRS on ECG in patients with CAD.

Methods: We searched English-language randomized controlled trials involving fQRS on ECG in patients with CAD ($n = 3279$ patients, 12 trials). Two reviewers independently extracted data. Data on LVEF, LVESD, LVEDD, LVESV, LVEDV, total mortality, stroke, and MACE were collected. fQRS was performed a comparison with non-fQRS, calculating pooled relative risk (RRs) and weighted mean difference (WMD), and associated 95% confidence intervals (CIs).

Results: fQRS was associated with significant increased WMD of LVEDD (WMD, 2.26; 95%CI, 0.92 to 3.60, $P = 0.0009$), LVESD (WMD, 2.71; 95%CI, 1.23 to 4.19, $P = 0.0003$), LVEDV (WMD, 31.37; 95%CI, 24.82 to 37.92, $P < 0.00001$), and LVESV (WMD, 28.45; 95%CI, 22.92 to 33.98, $P < 0.00001$). As compared to non-fQRS, fQRS increased risk of total mortality (RR, 3.09; 95%CI, 1.76 to 5.44, $P < 0.0001$) and MACE (RR, 2.85; 95%CI, 1.98 to 4.09, $P < 0.00001$) in patients with CAD. However, a decreased trend was observed for LVEF (WMD, -3.59 ; 95%CI, -7.05 to -0.12 , $P = 0.04$). For the incidence of stroke, there was no difference between fQRS and non-fQRS group.

Conclusions: Our findings indicate that fQRS is a valuable factor to predict total mortality and MACE in patients with CAD.

Ann Noninvasive Electrocardiol 2016;21(4):404–412

fragmented QRS; coronary artery disease; mortality; meta-analysis

Fragmented QRS complexes (fQRS), novel electrocardiographic signals on routine 12-lead electrocardiography (ECG), which were defined by the presence of changes in QRS morphology including an additional R wave, a notching of the R wave or S wave, or the presence of >1 additional R wave in 2 contiguous leads.¹ fQRS is associated with myocardial fibrosis, adverse cardiac events, regional myocardial damage, and event-free survival. It has been related to a number of cardiac diseases including Brugada syndrome,² ventricular aneurysm, dilated cardiomyopathy, essential hypertension,³ coronary artery disease,⁴ tetralogy of Fallot, takotsubo cardiomyopathy,⁵

cardiac AL amyloidosis, heart failure, and acute aortic dissection. To investigate the relationship between fQRS and cardiac function as well as survival in patients with coronary artery disease, we hypothesized that fQRS on routine 12-lead electrocardiography would be associated with an increased risk of total mortality.

METHODS

Data Sources and Searches

We followed the quality of reporting of meta-analyses (QUOROM) and adhered to the

Address for correspondence: Bojun Gong, M.D., Department of Cardiology, First Affiliated Hospital of Jinan University, Guangzhou, 510632, China. Tel.: +86 20 38688665; Fax: +86 20 38688665; E-mail: bojungong@hotmail.com.

This research was supported by the Ministry of Science and Technology of Guangzhou (grant Nos. 2014Y2-00128).

© 2015 Wiley Periodicals, Inc.
DOI: 10.1111/anec.12525

Figure 1. Study flow diagram.

meta-analysis of observational studies in epidemiology (MOOSE) guidelines. A systemic literature search was conducted independently by two authors (B.J.G. and Z.C.L.). A search of the medical literature published in English and Chinese was searched from seven databases: (1) Web of Science (ISI) (1990 to 2015); (2) PubMed (1990 to 2015); (3) EMBASE (1980 to 2015); (4) China National Knowledge Infrastructure (CNKI) (1990 to 2015); (5) the Cochrane Library (2014); (6) China Biology Medical literature database (CBM) (1990 to 2015); (7) MEDLINE (1950 to 2015). To identify potential studies missed by our primary search strategy, a hand search of bibliographies of relevant publications was performed. Initial searches of database including terms such as: "fragmented QRS," "fQRS," "fragmented QRS complexes," "myocardial infarction," "electrocardiography," "electrocardiogram," "mortality," "primary percutaneous coronary intervention," "left ventricular systolic function," "major adverse cardiac events," "coronary artery disease," "ST-segment elevation myocardial infarction," and "acute coronary syndrome."

Inclusion and Exclusion Criteria

The articles were independently assessed by two investigators (B.J.G. and Z.C.L.). Disagreements were resolved by consensus with discussion.

The inclusion criterion for the analysis were: (1) randomized; (2) inclusion of patients with coronary artery disease; (3) comparison of fragmented QRS complexes with non-fQRS on routine 12-lead electrocardiography; and (4) outcomes.

fQRS was defined by the presence of an additional R wave (R') or notching in the nadir of the R wave or S wave, or the presence of >1 R' (fragmentation) in two contiguous leads, corresponding to a myocardial territory.

We excluded studies that (1) did not include the outcomes mentioned in the study; (2) were not published in English; (3) were published only in abstract form; (4) did not include a control group; and (5) were animal studies.

Outcomes

Study outcomes were total mortality, stroke, major adverse cardiac events, and echocardiographic-derived cardiac parameters. Major adverse cardiac events were cardiovascular mortality, reinfarction, advanced heart failure, and ventricular arrhythmias. Echocardiographic-derived cardiac parameters were left ventricular ejection fraction (LVEF), left ventricular end-systolic diameter (LVESD), left ventricular end-diastolic diameter (LVEDD), left ventricular end-systolic volume (LVESV), and left ventricular end-diastolic volume (LVEDV).

Methodological Quality

The bias assessments were shown in Figure 4 according to the risk of bias.

Data Synthesis

Meta-analysis models were created for the above outcomes: total mortality, stroke, major adverse cardiac events, and echocardiographic-derived cardiac parameters. Relative risks (RRs), weighted mean difference (WMD), and corresponding 95% confidence intervals (CIs) were calculated. Results for event rates were converted to relative risks. We use the Mantel-Haenszel method to calculate tests of heterogeneity. Heterogeneity was considered significant if $I^2 > 50\%$, when random-effect model (REM) was adopted as the pooling method; otherwise, fix-effect model (FEM) was used. A P-value of less than 0.05 was considered statistically significant. Statistical analysis was performed by using Review Manager 5.0 (The Cochrane Collaboration, Oxford, United Kingdom).

RESULTS

Study Selection and Characteristics

A total of 12 studies⁶⁻¹⁷ containing 3279 patients met the inclusion criteria (as shown in Fig. 1). The patient characteristics, outcomes, and survival findings for each study are summarized in Table 1. All studies were included patients with coronary artery disease. This meta-analysis included at least one of the following outcomes: LVEF, LVESD, LVEDD, LVESV, LVEDV, total mortality, stroke, major adverse cardiac events, cardiovascular mortality, reinfarction, advanced heart failure, and ventricular arrhythmias.

Primary Analyses

Echocardiographic-Derived Cardiac Parameters

The effect of fQRS versus non-fQRS on LVEF, LVEDD, LVESD, LVEDV, and LVESV in patients with coronary artery disease was shown in Figure 2. There were significant changes in LVEDD (WMD, 2.26; 95%CI, 0.92 to 3.60, $P = 0.0009$), LVESD (WMD, 2.71; 95%CI, 1.23 to 4.19, $P = 0.0003$), LVEDV (WMD, 31.37; 95%CI, 24.82 to 37.92, $P < 0.00001$), and LVESV (WMD, 28.45; 95%CI, 22.92 to 33.98, $P < 0.00001$). However,

a decreased trend was observed for LVEF (WMD, -3.59; 95%CI, -7.05 to -0.12, $P = 0.04$).

Safety Outcomes

Total Mortality

Pooled data for total mortality were presented in Figure 3. The pooled data revealed a significant difference in the risk of total mortality (RR, 3.09; 95%CI, 1.76 to 5.44, $P < 0.0001$).

Stroke

The incidence of stroke was available for 2 of 12 studies ($n = 956$). The pooled data revealed no significant difference in the risk of stroke (RR, 1.09; 95%CI 0.35 to 3.40, $P = 0.08$) (Fig. 3).

MACE

There was significant difference in the risk of MACE (RR, 2.85; 95%CI, 1.98 to 4.09, $P < 0.00001$), including cardiovascular mortality (RR, 4.29; 95%CI, 2.22 to 8.28, $P < 0.0001$), reinfarction (RR, 3.23; 95%CI, 2.25 to 4.65, $P < 0.00001$), advanced heart failure (RR, 2.71; 95%CI, 2.00 to 3.66, $P < 0.00001$), and ventricular arrhythmias (RR, 2.32; 95%CI, 1.08 to 4.99, $P < 0.003$), for patients with fQRS versus non-fQRS (Fig. 3).

DISCUSSION

This is, to the best of our knowledge, the second meta-analysis of clinical trials to assess the effects of fQRS in patients with coronary artery disease. This meta-analysis included 12 studies involving 3279 patients. When we compared fQRS with non-fQRS on routine 12-lead electrocardiography, fQRS was associated with an increased risk of total mortality. What's more, we also demonstrated an increased risk of MACE, and observed that fQRS was associated with significant positive effects on LVEDD, LVESD, LVEDV, and LVESV.

Although the mechanism of fQRS on routine 12-lead electrocardiography is still not clear, some studies have reported that fQRS may be associated with cardiac fibrosis¹⁸ and systemic inflammation. There is clinical plausibility to the hypothesis that fQRS on a surface 12-lead electrocardiography may correlate with survival among patients with coronary artery disease. A study published by Brenyo et al.¹⁹ in 2012 showed that fQRS in inferior

Table 1. Characteristics of Included Studies

Study	Year of Publication	Male, %	fQRS (+)/fQRS(-), n	Population	LVEF (fQRS (+)), %	LVEF (fQRS (-)), %	Follow-Up Duration	Outcomes
Lorgis et al. ⁶	2013	70	145/162	Acute myocardial infarction	NA	50 (45–60)	846±297 days	MACE, CV death, heart failure, reinfarction, ventricular arrhythmias, total mortality
Ahn et al. ⁷	2013	87	129/61	Acute myocardial infarction	53.3 ± 9.6	49.0 ± 8.1	Hosp	LVEF, LVESD, LVEDD
Cetin et al. ⁸	2012	64	116/237	Stable angina pectoris	51 ± 13	61 ± 10	Hosp	LVEF, LVESD, LVEDD, LVESV, LVEDV
Guo et al. ⁹	2012	64	110/73	Non-ST-elevation myocardial infarction	57.3 ± 10.3	57.1 ± 11.1	Hosp	LVEF
Erdem et al. ¹⁰	2013	83	45/55	Acute ST-elevated myocardial infarction	45 ± 8.4	49 ± 8.5	Hosp	LVEF
Stavileci ¹¹	2014	76	80/216	Acute ST-segment myocardial infarction	37 ± 11	44 ± 13	Hosp	LVEF, CV death, total mortality
Akgul et al. ¹²	2014	79	91/323	ST elevation myocardial infarction	49 ± 8	43.9 ± 10	1 year	LVEF, MACE, stroke, CV death, heart failure, reinfarction, ventricular arrhythmias, total mortality

(Continued)

Table 1. Continued

Study	Year of Publication	Male, %	fQRS (+)/fQRS(-), n	Population	LVEF (fQRS (+)), %	LVEF (fQRS (-)), %	Follow-Up Duration	Outcomes
Uslu et al. ¹³	2014	80	153/389	ST elevation myocardial infarction	45.20 ± 9.40	49.34 ± 10	563.7 ± 168.7 days	LVEF, LVESD, LVEDD, MACE, stroke, CV death, heart failure, reinfarction, ventricular arrhythmias, Total mortality
Ari et al. ¹⁴	2012	82	34/51	Acute ST-segment myocardial infarction	39.5 ± 5.4	43.01 ± 5.4	6.6 ± 2.3 months	LVEF, LVESD, LVEDD, MACE, CV death, heart failure, reinfarction, ventricular arrhythmias
Mahenthiran et al. ¹⁵	2007	52	155/254	Coronary artery disease	55 ± 14	64 ± 9	Hosp	LVEF, LVESV, LVEDV
Sheng ¹⁶	2014	68	169/131	Acute myocardial infarction	NA	NA	15.1 ± 0.8 days	total mortality
Akbarzadeh et al. ¹⁷	2013	84	30/70	Acute myocardial infarction	45.6 ± 12.9	54.3 ± 9.9	6 months	LVEF, CV death, total mortality

fQRS = fragmented QRS complexes; NA = not applicable; Hosp = during hospitalization; LVEF = left ventricular ejection fraction; LVESD = left ventricular end-systolic diameter; LVEDD = left ventricular end-diastolic diameter; LVESV = left ventricular end-systolic volume; LVEDV = left ventricular end-diastolic volume; MACE = major adverse cardiac events; CV = cardiovascular.

Figure 2. Meta-analysis for the comparison of echocardiographic-derived cardiac parameters in fQRS group versus non-fQRS group.

leads was a predictor of sudden cardiac death, appropriately implantable cardioverter defibrillators (ICD) shock, and total mortality in patients with ischemic cardiomyopathy, especially in patients with left bundle branch block. In trial published in 2015, Narayanan et al.²⁰ showed that fQRS in the lateral territory was associated with sudden cardiac death in obese/overweight subjects. Similarly, another study published in 2015 by Debonnaire et al.²¹ found that in patients with hypertrophic cardiomyopathy fQRS in three territories was also associated with sudden cardiac death. A study researched by Pietrasik et al.²² in 2007 reported that in patients with acute myocardial infarction an fQRS appeared on routine 12-lead electrocardiography during the first 48 hours or

later 2 months was associated with long-term mortality as a significant predictor. What's more, in patients with acute coronary syndrome, it is demonstrated that fQRS was a significant predictor of mortality.²³ Moreover, Das et al.²⁴ found that the rate of total mortality was significantly higher in fQRS group than that in non-fQRS group in patients with coronary artery disease. It is reported that chronic Chagas' cardiomyopathy (CChC) is associated with sudden cardiac death and implantable cardioverter defibrillators (ICD) therapy is associated with reducing sudden cardiac death. Compared with patients with CAD or dilated cardiac myopathy implanted with ICDs, the prevalence of fQRS was lower (23%) than that in CChC implanted patients with ICDs (almost 60%).

Figure 3. Meta-analysis for the comparison of safety outcomes in fQRS group versus non-fQRS group.

The FECHA study investigators also showed that in patients with CChC and ICD implantation fQRS is a poor predictor of appropriate therapies.²⁵ Our meta-analysis showed that the presence of fQRS on routine 12-lead electrocardiography is associated with total mortality in patients with coronary artery disease.

Previous studies reported that fQRS may be a predictor of MACE in patients with coronary artery disease. It is reported that the relationship between MACE and the presence of the fQRS on

routine 12-lead electrocardiography showed not only in patients with ischemic cardiomyopathy but also with acute myocardial infarction. Two studies published in 2013⁶ and 2014,¹² respectively reported that MACE was more frequent in fQRS group in patients with myocardial infarction. Ari et al.²⁶ showed that in patients underwent primary percutaneous coronary intervention MACE rate was higher in fQRS group. A recent meta-analysis showed MACE was significantly higher in fQRS group than that in non-fQRS group.²⁷ In our

Figure 4. Methodological quality of included studies according to the Cochrane Collaboration's tool for assessing risk of bias.

meta-analysis, it showed an increased risk of MACE in patients with coronary artery disease.

Previous studies reported that fQRS was associated with deterioration of systolic function.^{8,28} The first meta-analysis of 16 studies published in 2015 and including 3997 patients with coronary artery disease reported that LVEF was significantly lower in fQRS group than that in non-fQRS group on routine electrocardiogram.²⁴ In patients with ST elevation myocardial infarction, some studies found that fQRS was inversely associated with LVEF.^{11,29–31} Gardner et al.¹⁸ showed that in patients with coronary artery disease LVEF in fQRS (+) group was lower than that in fQRS (–) group. In 2012, Cetin et al.⁸ reported that in patients underwent coronary angiography and diagnosed with stable angina pectoris fQRS was negatively related to LVEF when they performed multiple logistic regression analysis. It is demonstrated that in patients with myocardial infarction who underwent cardiac magnetic resonance imaging and percutaneous coronary intervention fQRS was positively associated with left ventricular end-systolic dimension and negatively related to LVEF.⁷ Compared with non-fQRS group, LVEF was significantly lower in fQRS group in patients with takotsubo cardiomyopathy⁵ and non-ischemic dilated cardiomyopathy.³² Similarly, we demonstrated that LVEF was lower in fQRS group than that in non-fQRS group. However, LVEDD, LVESD, LVEDV, and LVESV were higher in fQRS group than that in non-fQRS group.

LIMITATIONS

Several potential limitations of present meta-analysis should be taken into account. First,

there is publication bias on considering only published studies. Second, outcome data including mortality, MACE, and echocardiographic-derived cardiac parameters were incomplete for most trials. Third, language was restricted to English and Chinese limiting the result. Fourth, the present meta-analysis has not reached out to all racial and ethnic groups. In addition, more large scale randomized controlled trials should be included.

CONCLUSION

This meta-analysis demonstrated that fQRS is consistently associated with total mortality, MACE (including cardiovascular mortality, reinfarction, advance heart failure, and ventricular arrhythmias).

REFERENCES

1. Das MK, Khan B, Jacob S, et al. Significance of a fragmented QRS complex versus a Q wave in patients with coronary artery disease. *Circulation* 2006;113(21):2495–2501.
2. Morita H, Kusano KF, Miura D, et al. Fragmented QRS as a marker of conduction abnormality and a predictor of prognosis of Brugada syndrome. *Circulation* 2008;118(17):1697–1704.
3. Zhang B, Zhen Y, Shen D, et al. Significance of fragmented QRS complexes for identifying left ventricular hypertrophy in patients with hypertension. *Ann Noninvasive Electrocardiol* 2015;20(2):175–180.
4. Yilmaz H, Gungor B, Kemaloglu T, et al. The presence of fragmented QRS on 12-lead ECG in patients with coronary slow flow. *Kardiol Pol* 2014;72(1):14–19.
5. Shimizu M, Nishizaki M, Yamawake N, et al. J wave and fragmented QRS formation during the hyperacute phase in Takotsubo cardiomyopathy. *Circ J* 2014;78(4):943–949.
6. Lorgis L, Jourda F, Hachet O, et al. Prognostic value of fragmented QRS on a 12-lead ECG in patients with acute myocardial infarction. *Heart Lung* 2013;42(5):326–331.
7. Ahn MS, Kim JB, Yoo BS, et al. Fragmented QRS complexes are not hallmarks of myocardial injury as detected by cardiac magnetic resonance imaging in patients with acute myocardial infarction. *Int J Cardiol* 2013;168:2008–2013.

8. Cetin M, Kocaman SA, Canga A, et al. The independent relationship between systemic inflammation and fragmented QRS complexes in patients with stable angina pectoris. *Kardiol Pol* 2012;70(7):668–676.
9. Guo R, Li Y, Xu Y, et al. Significance of fragmented QRS complexes for identifying culprit lesions in patients with non-ST-elevation myocardial infarction: A single-center, retrospective analysis of 183 cases. *BMC Cardiovasc Disord* 2012;12:44.
10. Erdem FH, Tavil Y, Yazici H, et al. Association of fragmented QRS complex with myocardial reperfusion in acute ST-elevated myocardial infarction. *Ann Noninvasive Electrocardiol* 2013;18(1):69–74.
11. Stavileci B, Cimci M, Ikitimur B, et al. Significance and usefulness of narrow fragmented QRS complex on 12-lead electrocardiogram in acute ST-segment elevation myocardial infarction for prediction of early mortality and morbidity. *Ann Noninvasive Electrocardiol* 2014;19(4):338–344.
12. Akgul O, Uyarel H, Pusuroglu H, et al. Predictive value of a fragmented QRS complex in patients undergoing primary angioplasty for ST elevation myocardial infarction. *Ann Noninvasive Electrocardiol* 2015;20(3):263–272.
13. Uslu N, Gul M, Cakmak HA, et al. The assessment of relationship between fragmented QRS complex and left ventricular wall motion score index in patients with ST elevation myocardial infarction who underwent primary percutaneous coronary intervention. *Ann Noninvasive Electrocardiol* 2015;20(2):148–157.
14. Ari H, Cetinkaya S, Ari S, et al. The prognostic significance of a fragmented QRS complex after primary percutaneous coronary intervention. *Heart Vessels* 2012;27(1):20–28.
15. Mahenthiran J, Khan BR, Sawada SG, et al. Fragmented QRS complexes not typical of a bundle branch block: A marker of greater myocardial perfusion tomography abnormalities in coronary artery disease. *J Nucl Cardiol* 2007;14:347–353.
16. Sheng QH, Hsu CC, Li JP, et al. Correlation between fragmented QRS and the short-term prognosis of patients with acute myocardial infarction. *J Zhejiang Univ Sci B* 2014;15(1):67–74.
17. Akbarzadeh F, Pourafkari L, Ghaffari S, et al. Predictive value of the fragmented QRS complex in 6-month mortality and morbidity following acute coronary syndrome. *Int J Gen Med* 2013;6:399–404.
18. Gardner PI, Ursell PC, Fenoglio JJ Jr, et al. Electrophysiologic and anatomic basis for fractionated electrograms recorded from healed myocardial infarcts. *Circulation* 1985;72(3):596–611.
19. Brenyo A, Pietrasik G, Barsheshet A, et al. QRS fragmentation and the risk of sudden cardiac death in MADIT II. *J Cardiovasc Electrophysiol* 2012;23(12):1343–1348.
20. Narayanan K, Zhang L, Kim C, et al. QRS fragmentation and sudden cardiac death in the obese and overweight. *J Am Heart Assoc* 2015;4(3):e001654.
21. Debonnaire P, Katsanos S, Joyce E, et al. QRS Fragmentation and QTc duration relate to malignant ventricular tachyarrhythmias and sudden cardiac death in patients with Hypertrophic Cardiomyopathy. *J Cardiovasc Electrophysiol* 2015;26(5):547–555.
22. Pietrasik G, Goldenberg I, Zdzienicka J, et al. Prognostic significance of fragmented QRS complex for predicting the risk of recurrent cardiac events in patients with Q-wave myocardial infarction. *Am J Cardiol* 2007;100(4):583–586.
23. Das MK, Michael MA, Suradi H, et al. Usefulness of fragmented QRS on a 12-lead electrocardiogram in acute coronary syndrome for predicting mortality. *Am J Cardiol* 2009;104(12):1631–1637.
24. Das MK, Saha C, EI Masry H, et al. Fragmented QRS on a 12-lead ECG: A predictor of mortality and cardiac events in patients with coronary artery disease. *Heart Rhythm* 2007;4(11):1385–1392.
25. FECHA Study Investigators. Fragmented surface ECG was a poor predictor of appropriate therapies in patients with Chagas's cardiomyopathy and ICD implantation (Fragmented ECG in CHAgas' Cardiomyopathy Study). *Ann Noninvasive Electrocardiol* 2014;19(1):43–49.
26. Ari H, Cetinkaya S, Ari S, et al. The prognostic significance of a fragmented QRS complex after primary percutaneous coronary intervention. *Heart Vessels* 2012;27(1):20–28.
27. Xu Y, Qiu Z, Xu Y, et al. The role of fQRS in coronary artery disease. A meta-analysis of observational studies. *Herz* 2015;40 Suppl 1:8–15.
28. Korhonen P, Husa T, Konttila T, et al. Fragmented QRS in prediction of cardiac deaths and heart failure hospitalizations after myocardial infarction. *Ann Noninvasive Electrocardiol* 2010;15(2):130–137.
29. Uslu N, Gul M, Cakmak HA, et al. The assessment of relationship between fragmented QRS complex and left ventricular wall motion score index in patients with ST elevation myocardial infarction who underwent primary percutaneous coronary intervention. *Ann Noninvasive Electrocardiol* 2015;20(2):148–157.
30. Stavileci B, Cimci M, Ikitimur B, et al. Significance and usefulness of narrow fragmented QRS complex on 12-lead electrocardiogram in acute ST-segment elevation myocardial infarction for prediction of early mortality and morbidity. *Ann Noninvasive Electrocardiol* 2014;19(4):338–344.
31. Erdem FH, Tavil Y, Yazici H, et al. Association of fragmented QRS complex with myocardial reperfusion in acute ST-elevated myocardial infarction. *Ann Noninvasive Electrocardiol* 2013;18(1):69–74.
32. Ahn MS, Kim JB, Joung B, et al. Prognostic implications of fragmented QRS and its relationship with delayed contrast-enhanced cardiovascular magnetic resonance imaging in patients with non-ischemic dilated cardiomyopathy. *Int J Cardiol* 2013;167(4):1417–1422.