EDITORIAL

It is Time to Establish a Subspecialty in Noninvasive Electrocardiology

Arthur J. Moss, M.D.

From the Heart Research Follow-up Program, University of Rochester Medical Center, Rochester, New York, USA

During the past half century, cardiology has incorporated a wide array of diagnostic techniques and therapeutic procedures into its specialty. The widespread clinical use of electrocardiography began in the 1940s. The work of Dr. Robert Grant helped integrate vectorcardiographic concepts into the rational interpretation of the scalar ECG in health and disease. The Coronary Care Unit with bedside ECG-rhythm monitoring, the external defibrillator, the ambulatory (Holter) ECG, and the exercise testing with ECG monitoring were introduced in the 1960s at about the time that implanted pacemakers were first used for the treatment of heart block. The pharmacological expansion of new therapeutic agents for the treatment of cardiovascular disease began in the 1960s with the introduction of oral diuretics (chlorthiazide), and then over the next 40 years with a remarkable number of effective cardiovascular agents. Cardiac catheterization was widely utilized in the 1960s for the hemodynamic study of valvular heart disease. With the introduction of coronary angiography and coronary bypass graft surgery in the late 1960s, the emphasis shifted from rheumatic valvular disease, which was declining as a result of the appropriate early treatment of group A streptococcal infections, to the diagnosis and treatment of coronary disease. Myocardial ischemia and ventricular arrhythmias dominated cardiology in the beginning in the 1970s, and this emphasis gave rise to increased utilization of exercise tolerance testing and Holter monitoring. Percutaneous coronary angioplasty and subsequently stent angioplasty became the domain of the cardiologist.

Invasive electrocardiology began with the implantable pacemaker in the early 1960s, and sub-

sequently incorporated electrophysiological testing and radiofrequency ablation into its diagnostic and therapeutic armamentarium. The implanted cardioverter defibrillator (ICD) was first implanted in 1980, and the technology has subsequently evolved so that the ICD unit now incorporates an implanted monitor that can be interrogated noninvasively to decode stored episodes of cardiac arrhythmias that have been terminated by antitachycardia pacing and/or defibrillatory shocks. The molecular genetics of a number of inherited arrhythmic disorders has been identified, and noninvasive electrocardiographic techniques have been utilized in genotype-phenotype studies of the long QT syndrome, hypertrophic cardiomyopathy, Brugada syndrome, and arrhythmogenic right ventricular dysplasia. More recently, we have seen the introduction of the automatic external defibrillator (AED) and the wearable defibrillator.

These advances in cardiology have contributed to the remarkable decline in the morbidity and mortality of cardiac disease in the last half century. Along with this progress, the discipline of cardiology has matured into a variety of subspecialties with separation into hemodynamic and electrophysiologic activities; invasive and noninvasive procedures; electrocardiographic, echocardiographic, and radionucleotide testing; and research involving laboratory investigations and patient-related population studies. The subspecialty of noninvasive electrocardiology has emerged from this smorgasbord of cardiological activities. The noninvasive electrocardiologist is primarily a bedside cardiologist involved in the management of patients with a variety of cardiac diseases, especially those with coronary heart disease. Patients with

myocardial ischemia, hypertension, and arrhythmias make up a majority of the practice of physicians who identify themselves as noninvasive electrocardiologists. These physicians have special expertise in electrocardiography, ambulatory (Holter) ECG monitoring, the signal-averaged ECG, exercise tolerance testing, T-wave alternans detection, and the monitoring of patients with implanted pacemakers and defibrillators. The noninvasive electrocardiologist uses various electrophysiologic testing procedures in the diagnosis of ischemic and rhythm-related cardiac disorders.

The practicing clinical cardiologist who identifies himself/herself as a noninvasive electrocardiologist has different expertise than the research electrophysiologist, usually with a Ph.D. degree, who is more involved in the cellular and molecular mechanisms of disordered electrophysiology or in the biomedical engineering of electrophysio-

logic testing. The electrocardiologist and the electrophysiologist are, on occasion, the same individual, but usually they are different individuals who are working together in an academic setting that spans the disciplines of electrocardiology and electrophysiology. This type of joint coinvestigational research forms the underpinning of translational research.

Noninvasive electrocardiology requires a broad array of diagnostic and therapeutic expertise. A subspecialty in noninvasive electrocardiology is needed, and a medical subspecialty board in noninvasive electrocardiology should be established in countries throughout the world. The International Society for Noninvasive Electrocardiology (ISHNE) is the proper organization to establish the guidelines and requirements for such a subspecialty, and I strongly recommend that ISHNE take on this responsibility.