Published in final edited form as:

Physiol Behav. 2020 January 01; 213: 112693. doi:10.1016/j.physbeh.2019.112693.

Divergent immune responses in behaviorally-inhibited vs. non-inhibited male rats

Kerry C. Michael^a, Robert H. Bonneau^b, Rebecca A. Bourne^c, LaDara Godbolt^d, Michael J. Caruso^{c,e}, Christine Hohmann^d, Sonia A. Cavigelli^{c,e,f,*}

^aDepartment of Psychology, University of Minnesota, Morris

^bDepartment of Microbiology and Immunology and Department of Pediatrics, The Pennsylvania State University Hershey Medical Center

^cDepartment of Biobehavioral Health, The Pennsylvania State University

^dDepartment of Biology, Morgan State University

^eCenter for Brain, Behavior, and Cognition, The Pennsylvania State University

^fHuck Institutes of the Life Sciences, The Pennsylvania State University

Abstract

Stable behavioral traits (temperament, personality) often predict health outcomes. Temperament-specific differences in immune function could explain temperament-specific health outcomes, however, we have limited information on whether immune function varies by personality. In the present study, we examined the relationship between a basic behavioral trait (behavioral-inhibition vs. non-inhibition) and two immune responses (innate inflammation and delayed-type hypersensitivity, DTH) in a rodent model. In humans, behavioral inhibition (fearful temperament) is associated with altered stress physiology and allergies. In laboratory rats, the trait is associated with elevated glucocorticoid production. We hypothesized that behavioral inhibition is associated with glucocorticoid resistance and dampened T-helper 1 cell responses often associated with chronic stress and allergies. Further, this immune profile would predict poorly-regulated innate inflammation and dampened DTH. In male Sprague-Dawley rats, we quantified consistent behavioral phenotypes by measuring latency to contact two kinds of novelty (object vs. social), then measured lipopolysaccharide(LPS)-induced innate inflammation or keyhole limpet hemocyanin(KLH)-induced DTH. Behaviorally-inhibited rats had heightened glucocorticoid and interleukin-6 responses to a low/moderate dose of LPS and reduced DTH swelling to KLH re-exposure compared to non-inhibited rats. These results suggest that behavioral inhibition is associated with a glucocorticoid resistant state with poorly regulated innate inflammation and dampened cell-mediated immune responses. This immune profile may be associated with exaggerated T-helper 2 responses, which could set the stage for an allergic/asthmatic/atopic

*Corresponding Author: Sonia A. Cavigelli, The Pennsylvania State University, 219 Biobehavioral Health Building, University Park, PA 16802, Phone: 814-863-0210, Fax: 814-863-7525, sac34@psu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

predisposition in inhibited individuals. Human and animal models of temperament-specific immune responses represent an area for further exploration of mechanisms involved in individual differences in health.

Keywords

innate immune response; delayed type hypersensitivity; temperament; personality; individual differences; inflammation

1. Introduction

Specific temperaments (behavioral phenotypes or personalities) have been associated with specific health outcomes. One mechanism by which temperament may moderate health outcomes is through differential temperament-specific immune regulation (Capitanio et al., 2008; Zozulya et al., 2008; Rangassamy et al., 2016). To date, we have relatively limited information on the relationship between temperament and immune function. Many studies have investigated the immune parameters of a variety of animal species and populations in response to environmental demands, such as parasite load (Koprivnikar et al., 2012; Barber & Dingemanse, 2010) and social competition/aggression (Azpiroz et al., 2008; Capitanio et al., 2008, 2011; Chun et al., 2017). This has been researched at the behavioral level (e.g. behaviors that affect exposure to parasites/antigens) and at the population level, but potential physiological mechanisms at the individual level are still unclear.

In the current study, we examined immune responses in laboratory rats that were characterized as behaviorally-inhibited, non-inhibited, or mixed. At the individual level, inhibition and exploration are behavioral traits that can be relatively stable over time and across conditions and may involve stable differences in underlying physiology (Cavigelli & McClintock, 2003; Roberts et al., 2006; Réale et al., 2007; Bell et al., 2009; Cavigelli et al., 2009). In humans, Behavioral Inhibition (BI) is defined as a fearful or avoidant response to a wide array of novel experiences and is evident in approximately 20% of children studied in the US (Garcia-Coll et al., 1984; Kagan et al., 1984). BI humans are prone to develop environmental allergies and asthma, conditions that are broadly considered to reflect a T-helper 2 cell (T_H2) immune polarization (Kagan et al., 1991; Bell, 1992; Kim et al., 2006; Gulec et al., 2010; Heffner et al., 2014). BI individuals also show elevated basal glucocorticoid production (Kagan et al., 1987; Schmidt et al., 1997; Smider et al., 2002; Baugh et al., 2017) associated with negative health and immune outcomes (summarized in Chrousos & Kino, 2009; Miller et al., 2002; Webster Marketon & Glaser, 2008; Picard et al., 2014). In animals, BI/neophobia/exploration (defined by approach latency, locomotion, and/or time interacting with novel objects) are relatively stable traits over time and across contexts (Verbeek et al., 1994; Wilson et al., 1994; Dingemanse et al., 2002; Cavigelli & McClintock, 2003; Mason et al., 2006; Cavigelli et al., 2007; Krajl-Fiser et al., 2007; Bell et al., 2009; Caruso et al., 2014). Like in humans, BI male rats, defined as having consistently slower-than-median approach latencies in two novel conditions, have low-grade elevation in glucocorticoid production which is associated with shortened life span (Cavigelli & McClintock, 2003; Cavigelli et al., 2007; Cavigelli et al., 2009). An animal model of

temperament-associated differences in immune responses provides a system to experimentally test origins and mechanisms that underlie the relationship between BI/fearful temperament and health risks.

Studies that have linked *individual* behavioral profiles to immune system competence do not converge on a clear pattern. In free-ranging house finches, low-exploratory individuals invested less in constitutive defenses in favor of inducible defenses (i.e. occasionally spending a lot of energy to fight infections rather than investing in a continuous low-level prevention; Zylberberg et al. 2014), but in fairy wrens, low-exploratory individuals had *higher* investment in constitutive defenses (Jacques-Hamilton et al., 2017). In greenfinches, reactive individuals mounted higher innate and adaptive immune responses compared to proactive birds (Sild et al., 2011). In farm pigs, low-active animals had decreased lymphocyte proliferation in response to KLH exposure and lower expression of transcripts associated with innate immunity at baseline, with no T_H1/T_H2 shift (Bolhuis et al., 2003; Oster et al., 2015; cf. Geverink et al. 2004). In lab-reared zebrafish, individuals neophobic to novel objects demonstrated a heightened basal pro-inflammatory profile (i.e. increased brain IL-1 β and reduced IL-10 expression), and fish that had less preference for social partners had reduced brain IFN- γ expression, indicating dampened cell-mediated immunity (Kirsten et al., 2018). Similarly, in wild male mice exposed to repeated social confrontations, anxious/neophobic mice showed a comparatively lower T helper/regulatory T cell ratio in the spleen indicating general immunosuppression compared to less fearful mice (Rangassamy et al., 2016). These results suggest that inhibition/exploration behavioral phenotypes are associated with different immune regulation, but that the specific direction of this regulation is unclear.

Population-level studies have shown that inhibited or low-exploration species tend to invest less in innate immune responses and more in adaptive immunity (Lee, 2006; Martin et al., 2006; 2007; c.f. Johnson et al., 2012). But on the individual level, the opposite has been proposed: less exploratory individuals invest more in innate immunity (Zuk & Stoehr 2002; Lee, 2006; Adelman & Martin, 2009), perhaps at the cost of immune tolerance (e.g. Medzhitov et al., 2012). Some of the complications in this area are that free-ranging individuals with different temperaments expose themselves to different kinds and amounts of antigens; and in more controlled settings, immune function is quantified in many different ways, including estimates of immune function in the absence of an immune challenge. In addition, there are a variety of ways to define and measure behavioral phenotypes. This is particularly true for exploration-related behaviors, which could be driven by two different underlying motivations - i.e. harm avoidance vs. novelty seeking (Budaev 1998; Ray & Hansen, 2004). To test whether immune responses are functionally different among behavioral phenotypes, we need well-characterized and conceptualized behavioral phenotypes and tests of immune function that use standardized antigens presented in relatively antigen-free environments (i.e. low bacterial, viral, and parasite loads). In the current study, we examined how a basic behavioral trait (Inhibition vs. Non-Inhibition) related to two basic functional immune responses (innate inflammation and delayed-type hypersensitivity, DTH) in a social mammal, the Sprague-Dawley rat (*Rattus norvegicus*).

Briefly, the immune system can be divided into two broad domains: innate and adaptive. Cells involved in the innate system are the first to respond at a site of potential infection and can mount an inflammatory response within hours (Copeland et al., 2005). They are non-specific but recognize a “library” of pathogen-associated molecular patterns (PAMPs), of which lipopolysaccharide (LPS) is a prominent example. LPS is an endotoxin found in the cell walls of Gram-negative bacteria and is detected by a variety of innate immune cells that secrete cytokines that in turn rapidly attract and activate other immune cells; notably macrophage-secreted interleukin-6 (IL-6) which triggers the sickness response (Papanicolaou et al., 1998; Bluthé et al., 2000; Visintin et al., 2001). Adaptive responses are slower than innate responses and are mediated by T-helper cells: intracellular pathogens stimulate dendritic cells to signal naive T-helper cells to mature into T-helper 1 cells (T_H1) that guide cell-mediated adaptive immunity, while extracellular pathogens stimulate T-helper cells to mature into T-helper 2 cells (T_H2) that guide humoral adaptive immunity. Typically, once T_H1/T_H2 differentiation occurs, resulting cytokines promote continued similar T-helper cell differentiation. Some have argued that individuals can develop a bias toward a greater T_H1 or T_H2-mediated adaptive response, e.g. an allergic phenotype is often referred to as T_H2 polarization. In terms of adaptive responses, delayed-type hypersensitivity (DTH) is an easily-observed T_H1 cell-mediated response that can be measured non-invasively. DTH involves a sensitization phase where naive T-helper cells develop into antigen-specific T_H1 cells, and an effector phase where antigen re-exposure causes antigen-specific T_H1 cells to secrete cytokines that recruit large numbers of non-specific inflammatory cells to the antigen which results in localized induration within 24-72 hours (Punt et al., 2018).

Experimentally-induced chronically-elevated glucocorticoid exposure can alter both innate and T_H1-mediated immune function. Chronic stress can induce glucocorticoid resistance in macrophages and contribute to an innate pro-inflammatory profile (Stark et al., 2001). Glucocorticoid drugs can inhibit macrophage-induced T_H1 response with little effect on macrophages, thus stimulating exaggerated and prolonged innate immune activity (Mosser & Edwards, 2008; van de Garde et al., 2014; Kelly et al., 2018). On the flip side, IL-6 stimulates HPA axis activity in the short term, though over time may blunt the ACTH response (Lyson & McCann, 1991; Mastorakos et al., 1993). The DTH immune response is also affected by environmental stressors and glucocorticoid activity. Chronic or inescapable stress in rodents leads to elevated glucocorticoid production and suppressed DTH response, that is partially glucocorticoid-mediated, with a resulting skew toward increased T_H2-associated responses and reduced glucocorticoid sensitivity (Dhabhar & McEwen, 1997; 1999; Jasnow et al., 2001; Bartolomucci et al., 2003; Fleshner et al., 1995; Gazda et al., 2003; Bailey et al., 2009; Li et al., 2013; reviewed in Haczku & Panettieri, 2010). Last, chronic corticosteroid overexposure (*in vitro* and *in vivo*) reduces IL-12 production thereby reducing T_H1 proliferation - which indirectly drives T_H2 polarization (DeKruyff et al., 1998). Failure to suppress T_H2 development may result in the heightened allergic responses seen in behaviorally inhibited people.

In the current study, we examined whether rat behavioral inhibition, which is associated with increased basal glucocorticoid production, is also associated with altered immune function. The working hypothesis is that behavioral inhibition is comparable to chronic stress and/or long-term overexposure to glucocorticoids, and thus is associated with immunological

profiles associated with chronic glucocorticoid elevations, even in the context of low-stress. We predicted that behaviorally-inhibited individuals will show signs of innate inflammatory immune cell glucocorticoid resistance and T_H2 polarization which will be displayed as elevated systemic innate pro-inflammatory response and attenuated T_H1 response like localized DTH (Stark et al., 2001; Elenkov, 2004). To test this hypothesis, we compared immune responses of behaviorally-inhibited (BI), non-inhibited (NI), and mixed behavior rats in two broad functional immune responses that are known to be affected by chronic stress and/or chronic glucocorticoid overexposure: lipopolysaccharide (LPS)-induced innate pro-inflammatory IL-6 production and keyhole limpet hemocyanin (KLH)-induced DTH swelling.

2. Methods

We conducted four studies to compare immune responses among behaviorally-inhibited, non-inhibited, and mixed behavior male rats. In the first two studies, we assessed the innate immune response by quantifying circulating IL-6 responses to LPS. In the third and fourth study we quantified T_H1 -mediated DTH by measuring localized swelling following sensitization with and reintroduction of KLH. The methods for housing rats and for classifying rat temperament were the same for all studies, as described below. For this initial study on temperament and immune function in a controlled laboratory condition, we focused on male rats because behavioral inhibition has been associated with elevated glucocorticoid production and a truncated life span in males (Cavigelli & McClintock, 2003; Cavigelli et al., 2007; 2009) and because sex hormones differentially affect behavior and T_H1 and T_H2 responses (Ray and Hansen, 2004; Roved et al., 2017).

2.1. Animals

Young male Sprague-Dawley rats (55-60 days of age) were purchased from Charles River Laboratories (Raleigh, NC), individually-housed in solid bottom plastic cages ($43.5 \times 23.5 \times 20.5$ cm), and maintained on a 12L:12D lighting schedule (lights off at 10:00 h and on at 22:00 h EST). Food and water were available *ad libitum* and cages were cleaned once a week by animal facility personnel trained in animal care and handling. The colony room was maintained at 22°C with ~50% humidity. Rats were allowed to acclimate to laboratory housing and daily handling for 1-2 weeks prior to testing. All methods in this study were approved by the Pennsylvania State University Institute for Animal Care and Use Committee and adhered to methods specified in the *Guide for the Care and Use of Laboratory Animals* (National Research Council, 2011). Across the 4 studies we used a total of 113 rats (specific numbers in each study given below).

2.2. Behavioral Response to Novelty

To determine generalized and consistent trait-like inhibited/non-inhibited behavior, we measured approach latency in two different low-threat novel environments: one with rat-sized objects ('novel object arena'; Figure 1a) and one with an unfamiliar rat ('novel social arena', Figure 1b) (Cavigelli et al. 2007; 2009). Behavior in one arena does not predict behavior in the other arena, but individuals that are consistently slow to approach novelty in both arenas maintain this behavioral tendency over time (Cavigelli et al. 2007; 2009). This

generalized measure of slow novelty engagement across different contexts is based on early studies of behavioral inhibition in children where this behavioral designation was used to identify those that had reliably slow approaches to a wide array of novel experiences, both social and non-social (Garcia-Coll et al., 1984; Kagan et al., 1984). In prior studies, we have found that inhibition in the social arena is more predictive of basal and reactive circulating glucocorticoid levels, thus, in the current study we also considered whether inhibition in the social vs. object arena was more closely related to immune function (Cavigelli et al., 2007). Behavioral tests were conducted at 70-75 days of age during the active period (13:00-15:00 h) in a room adjacent to the colony room, illuminated with two 25-watt red bulbs that provided ~6 lux of illumination at the center of the test arena. These two behavioral tests were conducted with all rats in all four studies, with 3-4 days between each test and the 'novel object arena' tested before the 'novel social arena'.

For both arenas, latency to approach novelty was coded as the time from placement in the arena to first contact with a novel object or social partner. In the novel social arena, 3 males never approached novelty and their latency was coded as 310 seconds (i.e. 10 seconds longer than test duration). Rat temperament was assigned according to latencies in the two novel arenas: *Behavioral-Inhibition* was defined as slower-than-median in both arenas, *Non-Inhibited* as faster-than-median on both arenas, and *Mixed-Behavior* as faster-than-median in one arena and slower-than-median in the other (Cavigelli et al., 2007; 2009). Using these criteria, 20-30% of rats tested were characterized as Inhibited or Non-Inhibited and the remaining 70-80% characterized as Mixed-Behavior. This method of characterizing individuals allows us to identify consistently inhibited vs non-inhibited rats across different contexts, as has been done with children.

2.2.1. Novel Object Arena.—The arena was designed to be minimally anxiety-provoking with familiar colony-room odors, low-light conditions, high walls, and a Plexiglas cover (Cavigelli & McClintock, 2003). To provide a complex rat odor, the floor was covered with clean corn cob bedding and sprinkled with bedding (feces removed) from all cages in the colony room. Beyond removal of new feces, the arena floor was not further altered between each rat. The square test arena (120 cm × 120 cm) had white polypropylene walls (46 cm high) and a clear plastic cover. Three of the four corners contained a novel rat-sized object (a plastic tube, an inverted bowl, or a wire tunnel; Figure 1a). During testing, rats were placed into a protected container (clean ceramic bowl with 5 cm sides or red acrylic tube) and lowered into the empty arena corner. Rats were videotaped for 5 min with a camera placed above the arena and returned to their home cage immediately after testing.

2.2.2. Novel Social Arena.—The novel social arena was the same size and height as the Novel Object Arena, but instead of rat-sized novel objects, the arena contained two cages – an empty cage and one with an unfamiliar same-sex rat of similar size and age as the test rat (Figure 1b) (Cavigelli et al., 2007; 2009). Rats were introduced to the arena, videotaped, and start bowl cleaned as described for the novel object arena.

2.3 Innate Immune Response – Rapid circulating IL-6 response to lipopolysaccharide (LPS).

To quantify the innate pro-inflammatory response, we injected rats with LPS and measured circulating IL-6 levels at regular intervals during the following 10 hrs. For this initial set of studies on rat temperament and immune function, we limited analyses to IL-6. The first of the two studies was used to identify an intermediate LPS dose to mimic a common immune challenge where individual variability in responses can be documented (as opposed to commonly-used high doses used to stimulate strong sickness behavior responses).

2.3.1 Innate Immune Response, Study 1: Identify Appropriate Dose.—We used 4 different doses: a commonly-used high dose (125 µg/kg), two intermediate doses (5 and 25 µg/kg), and a control dose (0 µg/kg) (Givalois et al., 1994; Salome et al., 2008). LPS, derived from *Escherichia coli*, serotype 0111:B4, with the same lot number within each study (cat. no. L3012; Sigma-Aldrich, St. Louis, MO) was diluted with sterile endotoxin-free (USP) saline (Cat. No. 2F7124; Baxter, Deerfield, IL) to arrive at an injection volume of 0.5 ml. At ~80 days of age, all rats ($N=26$) were injected intraperitoneally (IP) with one of the four LPS doses 1 hr before the dark phase of the light cycle. Six rats were assigned to each of the 0 and 25 µg/kg conditions and 7 rats to each of the 5 and 125 µg/kg conditions. To measure circulating IL-6, blood was collected by tail-tip amputation immediately before LPS injection and at 2, 4, 6, and 10 hours post-injection. Blood was collected into untreated tubes, stored on ice for approximately an hour, then centrifuged and serum aliquots stored at -80°C until assayed for IL-6.

Serum IL-6 concentrations were measured with a commercial ELISA kit (Rat IL-6 Quantikine, cat no. R6000B; R&D Systems, Minneapolis, MN) with a standard range of 62.5–4000 pg/ml and a sensitivity of 36 pg/ml. Samples were processed according to instructions supplied with the kit. Briefly, 75 µl of frozen rat serum was thawed and diluted 1:2 with diluent supplied with the kit and duplicate samples (50 µl) analyzed for each rat time point. Mean intra-assay and inter-assay coefficients of variation were 7.14 and 14.52.

2.3.2 Innate Immune Response, Study 2: Compare Innate Pro-inflammatory Responses among Behaviorally-Inhibited (BI), Non-Inhibited (NI), and Mixed-Behavior Rats.—We used 30 rats that had been characterized as either BI ($n=6$), NI ($n=7$), or Mixed ($n=17$) (according to methods in section 2.2). Methods were identical to LPS Study 1, except only the 25 µg/kg LPS dose was used, and an additional blood sample at 8 hrs post-LPS injection was collected based on results from the prior study. In addition, to test the glucocorticoid resistance hypothesis, we compared post-LPS IL-6 to glucocorticoid secretion; blood samples were analyzed for both IL-6 and corticosterone. Corticosterone concentration was analyzed with a commercial radioimmunoassay kit (cat. no. 07-120103; MP Biomedicals, Solon, OH) with a standard range of 25–1000 ng/ml and a sensitivity of 25 ng/ml. Samples were processed in duplicate according to instructions supplied with the kit, with some minor alterations. Intra- and inter-assay coefficients of variation were 3.44 and 15.79 for a low control sample and 0.78 and 4.65 for a high control sample across 3 assays.

2.4 Delayed-Type Hypersensitivity – Peripheral swelling response to keyhole limpet hemocyanin (KLH) re-exposure.

KLH is an extremely large heme protein found in the giant keyhole limpet *Megathura crenulata*. It is highly immunogenic, and because KLH is not homologous to any vertebrate proteins, it is unfamiliar to mammals commonly used in research and thus ideal for examining immune responses to a novel antigen. In addition, KLH causes dendritic cells to sensitize T_H1 cells but macrophages do not trigger a large inflammatory response. KLH is therefore an ideal antigen to experimentally-induce a delayed-type hypersensitivity reaction. KLH is introduced systemically, then reintroduced locally several weeks later using a subcutaneous injection, and DTH response estimated from relative swelling at the injection site. This provides a minimally-invasive measure of immune response to antigen re-exposure, which is ideal for longitudinal and lifespan studies.

2.4.1 Delayed-Type Hypersensitivity Response, Study 1: Identify Initial Dose/Injection Site and Re-Exposure Swelling Response Dynamics.—

In prior studies with KLH-induced DTH responses, several different injection doses and sites have been used for initial KLH exposure, and to estimate DTH swelling, induration measured at different times after re-exposure. To identify the best method to document individual variability in DTH responses, we compared swelling at 1, 2, and 3 days post-re-exposure after initial exposure at one of three locations/doses (Exon et al., 1990; Wood et al., 1993; Fleshner et al., 1995; Gazda et al., 2003). We used 27 male Sprague-Dawley rats classified as Behaviorally-Inhibited ($n=8$), Non-Inhibited ($n=8$), and Mixed-Behavior ($n=9$) with rats in each temperament group evenly distributed across three KLH initial injection sites/doses: (1) 500 µg/kg intraperitoneal (IP; injection volume 0.5 ml; $n=8$), (2) 3mg/kg subcutaneous in caudal tail fold (SC; injection volume 0.2 ml; $n=9$), or (3) both sites (IP+SC; $n=10$; KLH cat. no. 374825, Calbiochem, San Diego CA). Within each injection site, KLH dose/volume was the same for all animals. The initial KLH injection occurred at 95 days of age and rats were re-exposed to heat-aggregated KLH (100 µl injection at 20 mg/ml concentration) injected into the right hind footpad thirty days later (Exon et al., 1990). As a control, an equal volume of saline was injected into the left hind footpad. Right and left footpad swelling was measured 1, 2, or 3 days following re-exposure with a spring-loaded micrometer (Mitutoyo, catalog no. 7301). Analyses were conducted on the difference in swelling between right (KLH-treated) and left (saline-treated) footpads.

2.4.2 Delayed-Type Hypersensitivity Response, Study 2: Compare Delayed-Type Hypersensitivity Swelling among Behaviorally-Inhibited (BI), Non-Inhibited (NI), and Mixed-Behavior Rats.—

Based on the results of Study 1, for Study 2 we used only the IP method of KLH sensitization (using 500 µg/kg, IP, injection volume 0.5 ml) and measured foot swelling 1 day after KLH re-exposure ($N=30$ male rats). Sensitization occurred at 75 days of age and re-exposure 30 days later with non-heat-aggregated KLH using the same protocol as KLH Study 1. Behavioral and foot-swelling measurements were conducted as in DTH Study 1.

2.5 Statistical Analyses.

In the Innate Immune Response Study 1, a repeated measures ANOVA was used to compare IL-6 responses (2, 4, 6, 10 hr levels) between doses (low, mid, high) as a fixed factor, with IL-6 levels at the time of injection (0 hr) as a covariate. To compare IL-6 responses among temperaments, overall IL-6 production in response to low and mid LPS doses was estimated using area-under-the-curve with respect to ground (AUCg; Pruessner et al., 2003), then AUCg compared across temperaments using ANOVA with temperament (BI, NI, Mixed) and dose (low, mid) as fixed factors, IL-6 AUCg as dependent variable, and IL-6 concentration at the time of injection as a covariate. To further compare IL-6 responses across temperaments a regression analysis was used to compare mean approach latency across the two novel arena tests (or latency on either test) and LPS dose (predictor variables) to IL-6 AUCg (dependent variable). In the Innate Immune Response Study 2, similar analyses were conducted (excluding dose as a variable) to compare IL-6 responses to a mid-LPS dose among temperaments. In both studies, several LPS-injected rats had peak IL-6 levels that were very low and comparable to those in control (saline-injected) rats (< 200 pg/ml at any time point as compared to a mean peak of 2925 pg/ml in Study 2). Non-responders were defined as LPS-injected rats that had IL-6 levels less than 200 pg/ml at all time points, and these individuals were excluded from analyses (4 LPS-injected rats in Study 1 – all in the 5µg/kg and 25µg/kg groups – and 3 rats in Study 2, 2 BI, 1 NI). In Study 2, corticosterone responses to LPS injections were compared among temperaments using similar methods described above for IL-6; a one-factor ANOVA with temperament as a factor, corticosterone AUCg as the dependent variable, and concentration at the time of LPS injection as a covariate. We also used a repeated measures ANOVA with temperament as factor, corticosterone concentrations at each time point after LPS injection as the dependent variable and corticosterone levels at the time of injection as a covariate. Total glucocorticoid production was linearly compared to temperament using a correlation analysis with mean approach latency across novel arenas and corticosterone AUCg as variables. To compare the relative relationship between temperament and glucocorticoid and IL-6 production following LPS challenge, we used a multiple regression analysis with mean latency on both novel arenas and glucocorticoid concentration at the time of LPS injection and during the 10 hours after LPS (AUCg) as predictor variables, and IL-6 AUCg as the dependent variable. To determine if a specific form of inhibition (i.e. social vs. non-social) was most closely associated with immune function, these analyses were repeated with latency on each arena separately instead of mean latency across both arenas.

DTH responses were estimated from relative footpad swelling calculated as the difference in cross-section between the saline- and KLH-treated feet. In DTH Study 1, swelling response differences between the IP, SC, and IP+SC initial injection methods were assessed by ANOVA. To compare swelling responses among temperaments in DTH Study 2, an ANOVA was used with temperament as the fixed effect factor and swelling response as dependent variable. Two rats were identified as outliers with footpad swelling difference scores further than two standard deviations from the mean; these two males were excluded from analysis (2 NI), leaving 9 BI, 6 NI, and 13 Mixed rats. In addition, correlational analyses were conducted to compare swelling response to mean latency on the two arenas and to latency on each arena separately. For all analyses, dependent variables were inspected for normality and

for those variables that showed a right skew, natural-log-transformation values were used in statistical analyses to meet normal distribution requirements. For clarity, means and SEMs of raw values are presented in figures.

3. Results

3.1. Innate Immune Response Results

3.1.1. Study 1: Dose effects of LPS on IL-6 response.—As expected, increasing LPS doses led to increased circulating IL-6 concentrations (Dose: $F(3,22) = 8.39$, $p < 0.001$; Time: $F(4,19) = 305.48$, $p < 0.0001$; Dose $\times$ Time: $F(12,53) = 5.80$, $p < 0.0001$; Figure 2). In the control group (0 $\mu\text{g/kg}$), IL-6 levels did show a mild increase over time, likely a response to the injury of the tail clip required for repeat blood collection ($F(4,20) = 24.81$, $p < 0.0001$). There was no significant difference in IL-6 responses between the 5 and 25 $\mu\text{g/kg}$ doses ($F(1,7) = 1.35$, $p = 0.283$), and thus some analyses were conducted merging across doses. The 125 $\mu\text{g/kg}$ dose created the expected ceiling effect in all animals.

3.1.2. Study 1 & 2: Behaviorally-Inhibited (BI) rats have a greater interleukin-6 and glucocorticoid response to moderate LPS stimulation than Non-Inhibited (NI) rats.—In Study 1, at the moderate doses of 5 $\mu\text{g/kg}$ and 25 $\mu\text{g/kg}$, Behaviorally-Inhibited rats – those with the longest approach latencies in the novel arenas – had significantly higher circulating IL-6 AUCg values (mean latency across arenas: adjusted $R^2 = 0.85$, $n = 9$, $t = 3.81$, $p = 0.0089$). Latencies within each arena were less predictive of individual rat IL-6 AUCg after LPS, with object approach latencies more closely related to IL-6 responses than social approach latencies (object: adjusted $R^2 = 0.73$, $n = 9$, $t = 2.36$, $p = 0.056$; social: adjusted $R^2 = 0.58$, $n = 9$, $t = 1.21$, $p = 0.275$). At the higher LPS dose (125 $\mu\text{g/kg}$), there was no relationship between approach latencies and IL-6 responses to LPS ($R^2 = 0.09$, $n = 7$, $t = -0.71$, $p = 0.509$).

Study 2 confirmed this accentuated IL-6 response in Behaviorally-Inhibited rats. Basal circulating IL-6 concentrations at the time of LPS injection were no different among temperaments ($F(2,24) = 0.361$, $p = .701$, $\eta^2 = 0.029$; Figure 3a). However, in response to the LPS challenge, BI rats had peak and AUCg circulating IL-6 levels that were 2-3 times greater than NI and Mixed rats (IL-6 AUCg ANOVA: $F(2,23) = 5.044$, $p < 0.05$, $\eta^2 = 0.305$; repeated measures ANOVA: $F(2,23) = 3.802$, $p < .05$, $\eta^2 = 0.248$; Figure 3a). Tukey's HSD tests indicate that the BI rats were significantly different from both the NI and Mixed groups (p s = 0.012 and 0.020). There was a significant positive correlation between mean latency on the novelty arenas and IL-6 production (AUCg) in response to the moderate LPS dose ($r = 0.453$, $n = 27$, $p = 0.009$). Latency on the novel social arena was more closely related to IL-6 production than latency on the novel object arena ($r = 0.404$, $n = 27$, $p = 0.018$ vs. $r = 0.225$, $n = 27$, $p = 0.129$).

There was also a trend toward temperament-specific differences in glucocorticoid production in response to the moderate LPS challenge. At the time of LPS challenge, basal corticosterone levels did not differ among temperament groups ($F(2,23) = 0.868$, $p = 0.434$, $\eta^2 = 0.070$; Figure 3b), but there was a significant temperament difference in corticosterone secretion after LPS injection. BI rats had significantly greater corticosterone AUCg

compared to Mixed rats (corticosterone AUCg ANOVA: $F(2,22) = 4.755$, $p = 0.019$, $\eta^2 = 0.302$; repeated measures ANOVA: $F(2,22) = 4.201$, $p = 0.028$, $\eta^2 = 0.276$; Figure 3b). In the case of glucocorticoid production, there were no significant linear relationships between approach latency in the novel arenas and total corticosterone production (AUCg) after LPS stimulation (mean latency in both arenas: $r = 0.163$, $n = 27$, $p = 0.208$; novel social: $r = 0.049$, $n = 27$, $p = 0.403$; novel object: $r = 0.172$, $n = 27$, $p = 0.195$). Variance in IL-6 response to LPS was more closely related to behavioral measures of temperament (mean latency to approach) than to basal corticosterone level at the time of LPS challenge or to corticosterone responses (AUCg) to the challenge (regression statistics for mean approach latency, basal corticosterone, and corticosterone AUCg to LPS: $\beta = 0.436$, $t = 2.377$, $p = 0.026$; $\beta = -0.233$, $t = 1.052$, $p = 0.304$, $\beta = 0.202$, $t = 0.903$, $p = 0.376$).

3.2. Delayed-Type Hypersensitivity Results

3.2.1 Study 1: Temporal dynamic of footpad swelling after KLH re-injection did not differ depending on location of initial injection.—

In Study 1, on the day after KLH re-exposure, the KLH-treated footpad was thicker than the saline-treated footpad (paired $t(8) = 6.502$, $p < 0.0001$; KLH vs. saline mean $\pm$ SD: 5.032 ± 0.227 vs. 4.658 ± 0.166 mm). This difference in swelling between the right (KLH-treated) and left (saline-treated) footpads did not differ depending on the location of the initial KLH injection (intraperitoneal, caudal fold, or both; $F(2,21) = 0.513$, $p = 0.606$) but did differ according to days following re-exposure ($F(2, 22) = 7.453$; $p = 0.003$). Footpad swelling was greatest 1 day after re-exposure, intermediate 2 days after re-exposure, and resolved by day 3 (mean $\pm$ SD on Days 1, 2, 3: 0.375 ± 0.172 , 0.192 ± 0.221 , -0.008 ± 0.222 mm). Based on these results, analyses of temperament-related swelling were conducted with footpad swelling measured on Day 1 after KLH re-exposure.

3.2.2. Study 1 & 2: Behaviorally-Inhibited (BI) rats had a dampened (or slower) DTH response to KLH than Non-Inhibited (NI) rats.—

In Study 1, rats that were fast to approach novelty (NI) had more footpad swelling one day after KLH re-exposure than rats that were slower to approach novelty (BI) ($r = -0.65$, $n = 10$, $p = 0.041$). Two days after KLH re-exposure, there was a trend for greater KLH-induced footpad swelling in BI vs. NI males ($r = 0.67$, $n = 8$, $p = 0.069$), suggesting a possible delayed DTH response in BI. Study 2 outcomes were similar; NI males had a heightened swelling response one day after KLH footpad re-exposure compared to Mixed- and BI males ($F(2,25) = 3.53$, $p = 0.045$; Figure 4). There was a significant negative correlation between mean approach latencies on the two novelty arenas and footpad swelling responses ($r = -0.534$, $n = 28$, $p = 0.003$). Again, like for the IL-6 response to LPS, latency on the novel social arena was more closely related to footpad swelling than latency on the novel object arena ($r = -0.47$, $n = 28$, $p = 0.012$ vs. $r = -0.33$, $n = 28$, $p = 0.083$).

4. Discussion

In the current study, we found that behavioral inhibition, a trait associated with moderate, chronic glucocorticoid overproduction, was also associated with a specific immune response profile. Male rats that showed behavior inhibition had an elevated innate inflammatory

response with concurrent elevated circulating glucocorticoids, and dampened T_H1 -mediated delayed-type hypersensitivity response compared to non-inhibited rats. These results suggest that behavioral inhibition involves innate cell glucocorticoid resistance and a skew toward less T_H1 -mediated immune responses. Taken together, immune function was systematically related to a behavioral trait even in the absence of differential antigen exposure among behavioral types.

The results of the current study are similar to those documented in prior studies with mice, primates, and cattle (Azpiroz et al., 2008; Capitanio et al., 2011; Hulbert et al., 2011; Kirsten et al., 2018) and differ from other studies that have shown increased innate immunity in free-ranging avian risk-takers and heightened DTH responses in socially-inhibited human adults (e.g. Cole et al., 1999; Zylberberg et al., 2014). Different findings across studies may result from different behavioral categorization methods, species differences in evolutionary histories, sex differences, disease history, genetic predispositions, and/or different environmental conditions across studies. For example, laboratory-bred, farm-bred, and wild-caught animals all have vastly different breeding histories, life experiences, and pathogen exposures. Innate immune responses are associated with significant caloric cost, and for wild-living animals sickness behavior may lead to lost opportunities to forage, socialize, and mate (Rauw, 2012; McDade, 2005; Lochmiller & Deerenberg, 2000) and may cause tissue damage (i.e. immunopathology; Medzhitov et al., 2012; Peiris et al., 2010; La Gruta et al., 2007). Thus, innate immunity as it relates to temperament may differ between free-ranging and lab/farm-raised animals.

Additionally, variance among studies may be driven by differential behavioral characterization methods. Behavioral response to novelty is likely driven by two distinct motivations: harm avoidance vs. sensation-seeking; specific testing conditions may access one motivation more readily than the other (e.g. Ray & Hansen 2004). Tests to access these two underlying motivations are not strongly-established in rodents. In the current study, we used tests that mimic those used to measure behavioral inhibition (fearful temperament) in children: low-threat, moderately-complex testing environments that maximize movement in most animals. Sensation-seeking in humans has been defined as willing to take risks for the sake of varied, novel, and complex sensations and experiences (Zuckerman 1979). Thus, by testing in a low-threat environment (low light, protected start area, covered arena with high walls), we likely did not have a strong metric of sensation-seeking but rather of harm avoidance/fearful temperament/behavioral inhibition. Variance in immune profiles as they relate to exploration-related behaviors may result from a high degree of behavior test variance, both within and between species. Interestingly, studies that used similar behavioral characterization methods as those used in the current study, showed similar glucocorticoid and immune response biases (passive-reactive male mice, Azpiroz et al., 2008; nervous infant rhesus monkeys, Capitanio et al., 2011; temperamental Brahman bulls, Hulbert et al., 2011; high responder zebrafish, Kirsten et al., 2018).

Another source of variance among studies may result from sex of the study animals. Behavioral patterns and basal and stimulated glucocorticoid circulation and immune function is quite different between males and females and this difference must be taken into account when studying the relationship between behavioral and immune phenotypes

(Cavigelli et al. 2008; Immonen et al., 2018; Tarka et al., 2018). For example, in female humans social-inhibition was associated with elevated DTH responses (Cole et al., 1999) whereas the current study with male rats showed the opposite result. Further work is required to determine if immune function is systematically associated with behavioral phenotypes, and if so how these systematic immune differences affect health and fitness outcomes, in both free-ranging and domesticated animals, males and females, and harm avoiding vs. sensation-seeking individuals.

Prior work has demonstrated effects of chronic stress and glucocorticoid production on a variety of immune functions, including innate immunity and delayed-type hypersensitivity responses. The present study tested whether a simple behavioral trait associated with elevated glucocorticoid production was also associated with a biased immune response profile. Inhibited rats had heightened IL-6 and glucocorticoid responses to a low/moderate dose of lipopolysaccharide, and a dampened swelling response to keyhole limpet hemocyanin re-exposure compared to non-inhibited rats. Behavioral measures of temperament were linearly-related to rapid IL-6 responses to LPS, but glucocorticoid production was not, suggesting that behavior-associated immune alterations are not solely mediated by glucocorticoid profiles (e.g. Sloan et al., 2008; Azpiroz et al., 2008; Capitanio et al., 2011). On the other hand, a heightened IL-6 and corticosterone response to LPS suggests that behavioral inhibition may involve some degree of glucocorticoid resistance. Glucocorticoid administration can inhibit macrophage-triggered T_H1 inflammatory responses, although macrophages quickly develop glucocorticoid resistance, which leads to enhanced innate immune activity and dysregulation of immune balance (Mosser and Edwards, 2008; Stark et al., 2001; van de Garde et al., 2014). This dysregulation occurs in disorders like autoimmune diseases, chronic infections, major depression, and atherosclerosis – many of the health conditions that have been associated with behavioral inhibition (Calcagni & Elenkov, 2006). In addition, long-term exposure to elevated glucocorticoids can produce tolerance in specific immune cells, which trigger runaway inflammation and T_H2 responses (i.e. macrophages, via innate lymphoid cells) while other cell types remain inhibited (Schmidt et al., 2010; Banuelos & Lu, 2016). Chronically-elevated corticosterone levels in inhibited individuals may inhibit T_H1 adaptive immune responses but allow for elevated innate inflammatory and T_H2 responses.

T_H2 polarization may be adaptive for coping with acute stressors - a strong T_H2 response could help moderate an elevated T_H1 response initiated to cope with skin wounding and infection (Calcagni & Elenkov, 2006; Munck & Náray-Fejes-Tóth, 1994; Dhabhar & McEwen, 1997; Gause et al., 2013). However, in the face of chronic stress or chronically-elevated glucocorticoid production, and in the absence of wounds, this regulatory response may lead to long-term T_H2 activation – a risk-factor for allergic and atopic responses (Elenkov & Chrousos, 1999). Additionally, a major function of the immune system is also to defend against damaged self-cells via the DNA damage response and repair (DDR/R) pathway. Immune dysregulation toward T_H2 inhibits this DDR/R cascade, allowing cancerous cells to proliferate unchecked (Pateras et al., 2015). Although not specifically tested in the current study, a dampened T_H1 -mediated response in behaviorally-inhibited males may be linked to elevated T_H2 -mediated responses – a strong anti-parasite defense in

wild-living animals, but a hallmark of allergies/atopic disorders/asthma in clean, protected lab-living animals and humans (see Maggi, 1998).

Future mechanisms to explore include the function of regulatory T cells (T_{reg}). T_{reg} cell activity reduces allergic airway inflammation by suppressing T_H2 cell differentiation (Duan et al., 2011) or modulating the activity of T_H1 and T_H2 cells (Akdis et al., 2004). In humans with asthma and in allergic asthma mouse models, the ability of T_{reg} cells to influence monocyte activity and IL-6 production is altered (Lee et al., 2007; Matsumoto et al., 2009; Hamzaoui et al., 2010). Inhibited recruitment of T_{reg} cells in response to allergen can result in excessive T_H2 responses in sensitized asthmatics (Wang et al., 2009). In addition, T_{reg} cells are inhibited by T_H17 cells, which are resistant to glucocorticoid suppression (Schmidt et al., 2010; Banuelos & Lu, 2016). Thus, chronic glucocorticoid production may have minimal impact on these processes. Future work on temperament-associated immune biases might focus on T_{reg} cell function in fearful humans as a possible mechanism underlying risk for allergy-related health outcomes.

The results of the current study provide support that a rodent model of human behavioral inhibition is possible. This and prior studies indicate that male rats that are slow to explore a novel, complex, protected environment have a shortened lifespan, elevated basal and stress-reactive levels of glucocorticoids, a heightened pro-inflammatory innate immune response, and dampened delayed-type hypersensitivity compared to rats that are not inhibited in this setting (Cavigelli & McClintock, 2003; Cavigelli et al. 2007; 2009). Chronic glucocorticoid overexposure, such as that associated with behavioral inhibition, may lead to these specific immune alterations, although this requires further study. An animal model of fearful or harm avoidant or behavioral inhibition temperament with similar underlying physiological biases as those seen in similarly-behaving humans provides a system in which to experimentally-test how genetic, environmental, and developmental conditions affect the relationship between temperament and health.

An important aspect of this model is that latency to approach a novel conspecific (social partner) was more predictive of immune dysregulation than latency to approach a novel object. Social inhibition, rather than inhibition in non-social contexts, may be a more appropriate model of temperament-specific immune dysfunction. Social stress has been shown to enhance the T_H2 response to allergen and reduce glucocorticoid sensitivity in many studies (Bailey et al., 2009; Li et al., 2013; reviewed in Haczku & Panettieri, 2010). Zebrafish showing low social exploration had reduced expression of IFN- γ (Kirsten et al., 2018). Mice deficient in adaptive immunity (SCID mice) have been shown to exhibit defects in social behavior that can be reversed by IFN- γ , a cytokine critical for T_H1 development (Filiano et al., 2016). In humans, loneliness and social isolation are associated with negative health outcomes, including mortality (Steptoe et al., 2013; Ellwardt et al., 2015; reviewed in Holt-Lunstad et al., 2015), and social inhibition predicted DTH responses during social engagement (in adults with autoimmune disorders; Cole et al. 1999). Thus, an animal model of social inhibition may provide the best system in which to study fearful temperament-specific physiology and health.

Further work is required to identify mechanisms by which temperament-specific immune alterations occur. Future studies should include measures of central nervous system function, and experimental manipulation of glucocorticoid activity to determine the relative role of these mechanisms in temperament-related immune response biases. An important caveat for this future work is that experimentally-manipulated exogenous glucocorticoid exposure should mimic glucocorticoid profiles seen in distinct temperaments, recognizing that endogenous glucocorticoid profile mimicry can be difficult with exogenous glucocorticoids (Pruett et al. 1999; Fleshner et al., 2001). Specific aspects of central nervous system alterations include serotonergic function (Takano et al., 2007; Donner et al., 2016), hippocampal and prefrontal cortex glucocorticoid signaling (Cerqueira et al., 2005; Patel et al., 2008; Sorrells et al., 2014), the kynurenine pathway (Brooks et al., 2016), and pro-inflammatory cytokine regulation throughout the brain (Capitanio et al., 2008; reviewed in Ménard et al., 2016). Several cytokines, particularly IL-1 and TNF α , can affect central nervous system functioning (reviewed in Rochfort & Cummins, 2015). In turn, central nervous system processes can affect immune function by triggering secretion of immunomodulating hormones or autonomic stimulation of lymph nodes, all of which may be associated with an inhibited temperament (e.g. Sloan et al., 2008). Further research on mechanisms could provide insight into targeted interventions in order to improve both psychological and physical well-being in behaviorally-inhibited people.

Acknowledgments

We would like to acknowledge the many students that assisted with this project, in particular Wael Bahar, Nicole Chirichella, Brian Coleman, Mario DeNicola, Olivia Francois, Alisa Inthavongsa, Sammy Leathers, and Christina Ragan. Funding was provided by the National Institutes of Health (R03MH071406, R25GM058904), and The Pennsylvania State University (Pennsylvania State University Children, Youth and Families Consortium, and Pennsylvania State Institute for Neuroscience). We would like to dedicate this paper to Robert H. Bonneau; without his expert advice and guidance, this study was not possible. Rob was instrumental in the planning of this study and he provided unwavering support to SAC and KCM and to many other researchers both within and outside his laboratory. Rob was the ideal colleague – full of ideas and energy, supportive, and with high expectations. He is greatly missed.

References

- Adelman JS & Martin LB (2009). Vertebrate sickness behaviors: Adaptive and integrated neuroendocrine immune responses. *Integr Comp Biol.*, 49, 202–214. doi: 10.1093/icb/icp028 [PubMed: 21665814]
- Akdis M, Verhagen J, Taylor A, Karamloo F, Karagiannidis C, Crameri R, Thunberg S, Deniz G, Valenta R, Fiebig H, Kegel C, Disch R, Schmidt-Weber CB, Blaser K & Akdis CA (2004). Immune responses in healthy and allergic individuals are characterized by a fine balance between allergen-specific T regulatory 1 and T helper 2 cells. *J Exp Med.*, 199, 1567–1575. doi:10.1084/jem.20032058 [PubMed: 15173208]
- Azpiroz A, De Miguel Z, Fano E & Vegas O (2008). Relations between different coping strategies for social stress, tumor development and neuroendocrine and immune activity in male mice. *Brain Behav. Immun.*, 22, 690–698. doi:10.1016/j.bbi.2007.10.007 [PubMed: 18061400]
- Bailey MT, Kierstein S, Sharma S, Spaitis M, Kinsey SG, Tliba O, Amrani Y, Sheridan JF, Panettieri RA & Haczku A (2009). Social stress enhances allergen-induced airway inflammation in mice and inhibits corticosteroid responsiveness of cytokine production. *J Immunol.*, 182, 7888–7896. doi: 10.4049/jimmunol.0800891 [PubMed: 19494313]
- Banuelos J & Lu NZ (2016). A gradient of glucocorticoid sensitivity among helper T cell cytokines. *Cytokine Growth Factor Rev.*, 31, 27–35. doi:10.1016/j.cytogfr.2016.05.002 [PubMed: 27235091]

- Barber I & Dingemanse NJ (2010). Parasitism and the evolutionary ecology of animal personality. *Philos Trans R Soc Lond B Biol Sci.*, 365, 4077–4088. doi: 10.1098/rstb.2010.0182. [PubMed: 21078659]
- Bartolomucci A, Sacerdote P, Panerai AE, Pederzani T, Palanza P & Parmigiani S (2003). Chronic psychosocial stress-induced down-regulation of immunity depends upon individual factors. *J Neuroimmunol.*, 41, 58–64. doi:10.1016/S0165-5728(03)00220-0
- Baugh AT, Senft RA, Firke M, Lauder A, Schroeder J, Meddle SL, van Oers K & Hau M (2017). Risk-averse personalities have a systemically potentiated neuroendocrine stress axis: A multilevel experiment in *Parus major*. *Horm Behav.*, 93, 99–108. doi: 10.1016/j.yhbeh.2017.05.011 [PubMed: 28545898]
- Bell IR (1992). Allergens, physical irritants, depression, and shyness. *J. Appl. Dev. Psychol*, 13, 125–133. doi:10.1016/0193-3973(92)90022-A
- Bell AM, Hankison SJ & Laskowski KL (2009). The repeatability of behaviour: a meta-analysis. *Animal Behaviour*, 77, 771–783. doi:10.1016/j.anbehav.2008.12.022 [PubMed: 24707058]
- Bluthé RM, Michaud B, Poli V & Dantzer R (2000). Role of IL-6 in cytokine-induced sickness behavior: a study with IL-6 deficient mice. *Physiol Behav.*, 70, 367–373. doi: 10.1016/S0031-9384(00)00269-9 [PubMed: 11006436]
- Bolhuis JE, Parmentier HK, Schouten WG, Schrama JW & Wiegant VM (2003). Effects of housing and individual coping characteristics on immune responses of pigs. *Physiol Behav.*, 79, 289–296. doi:10.1016/S0031-9384(03)00090-8 [PubMed: 12834801]
- Brooks AK, Lawson MA, Smith RA, Janda TM, Kelley KW & McCusker RH. (2016). Interactions between inflammatory mediators and corticosteroids regulate transcription of genes within the Kynurenine Pathway in the mouse hippocampus. *J Neuroinflammation*, 13, 98. doi: 10.1186/s12974-016-0563-1 [PubMed: 27142940]
- Budaev SV (1998). How many dimensions are needed to describe temperament in animals: A factor reanalysis of two data sets. *Int J Comp Psychol.*, 11, 17–29.
- Calcagni E & Elenkov I (2006). Stress system activity, innate and T helper cytokines, and susceptibility to immune-related diseases. *Ann N Y Acad Sci.*, 1069, 62–76. doi: 10.1196/annals.1351.006 [PubMed: 16855135]
- Capitanio JP, Abel K, Mendoza SP, Blozis SA, McChesney MB, Cole SW & Mason WA (2008). Personality and serotonin transporter genotype interact with social context to affect immunity and viral set-point in simian immunodeficiency virus disease. *Brain Behav. Immun*, 22, 676–689. doi: 10.1016/j.bbi.2007.05.006 [PubMed: 17719201]
- Capitanio JP, Mendoza S & Cole SW (2011). Nervous temperament in infant monkeys is associated with reduced sensitivity of leukocytes to cortisol's influence on trafficking. *Brain Behav. Immun*, 25, 151–159. doi: 10.1111/j.1708-8208.2012.00444.x [PubMed: 20868738]
- Caruso MJ, McClintock MK & Cavigelli SA (2014). Temperament moderates the influence of periadolescent social experience on behavior and adrenocortical activity in adult male rats. *Horm. Behav*, 66, 517–524. doi:10.1016/j.yhbeh.2014.07.010 [PubMed: 25066485]
- Cavigelli SA & McClintock MK (2003). Fear of novelty in infant rats predicts adult corticosterone dynamics and an early death. *Proc. Natl. Acad. Sci. USA* 100, 16131–16136. doi: 10.1073/pnas.2535721100 [PubMed: 14673078]
- Cavigelli SA, Stine MM, Kovacsics CE, Jefferson A, Diep MN & Barrett CE (2007). Behavioral inhibition and glucocorticoid dynamics in a rodent model. *Physiol. Behav* 92, 897–905. doi: 10.1016/j.physbeh.2007.06.016 [PubMed: 17673266]
- Cavigelli SA, Bennett JM, Michael KC, Klein LC (2008). Female temperament, tumor development and life span: relation to glucocorticoid and tumor necrosis factor α levels in rats. *Brain Behav. Immun*, 22, 727–735. doi: 10.1016/j.bbi.2007.10.014 [PubMed: 18155400]
- Cavigelli SA, Ragan CM, Michael KC, Kovacsics CE & Bruscke AP (2009). Stable behavioral inhibition and glucocorticoid production as predictors of longevity. *Physiol. Behav* 98, 205–214. doi: 10.1016/j.physbeh.2009.05.012 [PubMed: 19477191]
- Cerqueira JJ, Pêgo JM, Taipa R, Bessa JM, Almeida OF & Sousa N (2005). Morphological correlates of corticosteroid-induced changes in prefrontal cortex-dependent behaviors. *J Neurosci.*, 25, 7792–7800. doi: 10.1523/JNEUROSCI.1598-05.2005 [PubMed: 16120780]

- Chrousos GP & Kino T (2009). Glucocorticoid signaling in the cell: Expanding clinical implications to complex human behavioral and somatic disorders. *Glucocorticoids and Mood: Ann. N.Y. Acad. Sci.*, 1179, 153–166. doi:10.1111/j.1749-6632.2009.04988.x [PubMed: 19906238]
- Chun K, Capitanio JP, Lamkin DM, Sloan EK, Arevalo JM & Cole SW (2017). Social regulation of the lymph node transcriptome in rhesus macaques (*Macaca mulatta*). *Psychoneuroendocrinology*, 76, 107–113. doi: 10.1016/j.psyneuen.2016.10.029 [PubMed: 27902946]
- Cole SW, Kemeny ME, Weitzman OB, Schoen M, Anton PA (1999). Socially inhibited individuals show heightened DTH response during intense social engagement. *Brain Behav. Immunity*, 13, 187–200. doi: 10.1006/brbi.1998.0543
- Copeland S, Shaw Warren H, Lowry SF, Calvano SE, Remick D & the Inflammation and the Host Response to Injury Investigators (2005). Acute inflammatory response to endotoxin in mice and humans. *Clin. Vaccine Immunol*, 12, 60–67. doi:10.1128/CDLI.12.1.60-67.2005
- DeKruyff RH, Fang Y & Umetsu DT (1998). Corticosteroids enhance the capacity of macrophages to induce Th2 cytokine synthesis in CD4⁺ lymphocytes by inhibiting IL-12 production. *J Immunol.*, 160, 2231–2237. doi: <https://www.jimmunol.org/content/160/5/2231> [PubMed: 9498762]
- Dhabhar FS & McEwen BS (1997). Acute stress enhances while chronic stress suppresses cell-mediated immunity in vivo: a potential role for leukocyte trafficking. *Brain Behav Immun.*, 11, 286–306. doi:10.1006/brbi.1997.0508 [PubMed: 9512816]
- Dhabhar FS & McEwen BS (1999). Enhancing versus suppressive effects of stress hormones on skin immune function. *Proc Natl Acad Sci USA*, 96, 1059–1064. doi:10.1073/pnas.96.3.1059 [PubMed: 9927693]
- Dingemanse NJ, Both C, Drent PJ, van Oers K & van Noordwijk AJ (2002). Repeatability and heritability of exploratory behaviour in great tits from the wild. *Animal Behaviour*, 64, 929–938. doi:10.1006/anbe.2002.2006
- Donner NC, Siebler PH, Johnson DT, Villarreal MD, Mani S, Matti AJ & Lowry CA (2016). Serotonergic systems in the balance: CRHR1 and CRHR2 differentially control stress-induced serotonin synthesis. *Psychoneuroendocrinology*, 63, 178–190. doi: 10.1016/j.psyneuen.2015.09.024. [PubMed: 26454419]
- Duan W, So T, Mehta AK, Choi H & Croft M (2011). Inducible CD4⁺LAP⁺Foxp3⁺ regulatory T cells suppress allergic inflammation. *J Immunol.*, 187, 6499–6507. doi:10.4049/jimmunol.1101398 [PubMed: 22079987]
- Elenkov IJ (2004). Glucocorticoids and the Th1/Th2 Balance. *Ann. NY Acad. Sci.*, 1024, 138–146. doi: 10.1196/annals.1321.010 [PubMed: 15265778]
- Elenkov IJ & Chrousos GP (1999). Stress hormones, Th1/Th2 patterns, pro/anti-inflammatory cytokines and susceptibility to disease. *Trends Endocrinol Metab.*, 10, 359–368. doi:10.1016/S1043-2760(99)00188-5 [PubMed: 10511695]
- Ellwardt L, van Tilburg T, Aartsen M, Wittek R & Steverink N (2015) Personal networks and mortality risk in older adults: a twenty-year longitudinal study. *PLoS One*, 10, e0116731. doi: 10.1371/journal.pone.0116731. doi:10.1371/journal.pone.0116731 [PubMed: 25734570]
- Exon JH, Bussierem JL & Mather GG (1990). Immunotoxicity testing in the rat: An improved multiple assay model. *Int. J. Immunopharmac*, 12, 699–701. doi:10.1016/0192-0561(90)90108-Y
- Filiano AJ, Xu Y, Tustison NJ, Marsh RL, Baker W, Smirnov I, Overall CC, Gadani SP, Turner SD, Weng Z, Peerzade SN, Chen H, Lee KS, Scott MM, Beenhakker MP, Litvak V & Kipnis J (2016). Unexpected role of interferon- γ in regulating neuronal connectivity and social behaviour. *Nature*, 535, 425–429. doi: 10.1038/nature18626 [PubMed: 27409813]
- Fleshner M, Hermann J, Lockwood LL, Laudenslager ML, Watkins LR & Maier SF (1995). Stressed rats fail to expand the CD45RC⁺CD4⁺ (Th1-like) T cell subset in response to KLH: Possible involvement of IFN- γ . *Brain Behav. Immun*, 9, 101–112. doi: 10.1006/brbi.1995.1011 [PubMed: 7549034]
- Fleshner M, Deak T, Nguyen KT, Watkins LR, Maier SF (2001). Endogenous glucocorticoids play a positive regulatory role in the anti-keyhole limpet hemocyanin *in vivo* antibody response. *J. Immunol*, 166, 3813–3819. doi: 10.4049/jimmunol.166.6.3813 [PubMed: 11238624]
- Garcia-Coll C, Kagan J & Reznick JS (1984). Behavioral inhibition in young children. *Child Dev.*, 55, 1005–1019. doi: 10.1001/archpsyc.1990.01810130023004

- Gause WC, Wynn TA & Allen JE (2013). Type 2 immunity and wound healing: evolutionary refinement of adaptive immunity by helminths. *Nat Rev Immunol*, 13, 607–614. doi: 10.1038/nri3476 [PubMed: 23827958]
- Gazda LS, Smith T, Watkins LR, Maier SF & Fleshner M (2003). Stressor exposure produces long-term reductions in antigen-specific T and B cell responses. *Stress*, 6, 259–267. doi: 10.1080/10253890310001594441 [PubMed: 14660058]
- Geverink NA, Parmentier HK, de Vries Reilingh G, Schouten WG, Gort G & Wiegant VM (2004). Effect of response to backtest and housing condition on cell-mediated and humoral immunity in adult pigs. *Physiol Behav.*, 80, 541–546. doi: <https://doi.org/10.1016/j.physbeh.2003.10.013> [PubMed: 14741239]
- Givalois L, Dornand J, Mekaouche M, Solier MD, Bristow AF, Ixart G, Siaud P, Assenmacher I & Barbanel G (1994). Temporal cascade of plasma level surges in ACTH, corticosterone, and cytokines in endotoxin-challenged rats. *Am. J. Physiol*, 266, R164–R170. doi: 10.1152/ajpregu.1994.267.1.R164 [PubMed: 7508208]
- Gulec MY, Gulec H, Oztuna F & Kose S (2010). Cloninger's temperament and character dimension of personality in patients with asthma. *Int. J. Psychiatry Med*, 40, 273–87. [PubMed: 21166338]
- Haczku A & Panettieri RA Jr. (2010). Social stress and asthma: the role of corticosteroid insensitivity. *J. Allergy Clin. Immunol*, 125, 550–558. doi:10.1016/j.jaci.2009.11.005 [PubMed: 20153032]
- Hamzaoui A, Ammar J & Hamzaoui K (2010). Regulatory T cells in induced sputum of asthmatic children: association with inflammatory cytokines. *Multidisp. Respir. Med*, 5, 22–30. doi: 10.1186/2049-6958-5-1-22
- Heffner KL, Kiecolt-Glaser JK, Glaser R, Malarkey WB & Marshall GD (2014). Stress and anxiety effects on positive skin test responses in young adults with allergic rhinitis. *Ann. Allergy Asthma Immunol*, 113, 13–18. doi:10.1016/j.anai.2014.03.008 [PubMed: 24726195]
- Holt-Lunstad J, Smith TB, Baker M, Harris T & Stephenson D (2015). Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspect Psychol Sci*, 10, 227–237. doi: 10.1177/1745691614568352 [PubMed: 25910392]
- Hulbert LE, Carroll JA, Burdick NC, Randel RD, Brown MS & Ballou MA (2011). Innate immune responses of temperamental and calm cattle after transportation. *Vet. Immunol. Immunopathol*, 143, 66–74. doi:10.1016/j.vetimm.2011.06.025 [PubMed: 21726904]
- Immonen E, Hämäläinen A, Schuett W & Tarka M (2018). Evolution of sex-specific pace-of-life syndromes: genetic architecture and physiological mechanisms. *Behav Ecol Sociobiol.*, 72, 60. doi: 10.1007/s00265-018-2462-1 [PubMed: 29576676]
- Jacques-Hamilton R, Hall ML, Buttemer WA, Matson KD, Gonçalves da Silva A, Mulder RA & Peters A (2017). Personality and innate immune defenses in a wild bird: Evidence for the pace-of-life hypothesis. *Horm Behav.*, 88, 31–40. doi: 10.1016/j.yhbeh.2016.09.005. [PubMed: 27633460]
- Jasnow AM, Drazen DL, Huhman KL, Nelson RJ & Demas GE (2001). Acute and chronic social defeat suppresses humoral immunity of male Syrian hamsters (*Mesocricetus auratus*). *Horm. Behav* 40, 428–433. doi:10.1006/hbeh.2001.1708 [PubMed: 11673916]
- Johnson PT, Rohr JR, Hoverman JT, Kellermanns E, Bowerman J, Lunde KB (2012). Living fast and dying of infection: host life history drives interspecific variation in infection and disease risk. *Ecol Lett.*, 15, 235–242. doi: 10.1111/j.1461-0248.2011.01730.x [PubMed: 22221837]
- Kagan J, Snidman N, Julia-Sellers M & Johnson MO (1991). Temperament and allergic symptoms. *Psychosom Med.*, 53, 332–340. [PubMed: 1882014]
- Kagan J, Reznick JS, Clarke C, Snidman N, & Garcia-Coll C (1984). Behavioral inhibition to the unfamiliar. *Child Dev.* 55, 2212–2225. doi: 10.2307/1129793
- Kagan J, Reznick JS & Snidman N (1987). The physiology and psychology of behavioral inhibition in children. *Child Dev.* 58, 1459–73. [PubMed: 3691195]
- Kelly KA, Michalovicz LT, Miller JV, Castranova V, Miller DB & O'Callaghan JP (2018). Prior exposure to corticosterone markedly enhances and prolongs the neuroinflammatory response to systemic challenge with LPS. *PLoS One*, 13, e0190546. doi: 10.1371/journal.pone.0190546. [PubMed: 29304053]

- Kim TS, Pae CU, Jeong JT, Kim SD, Chung KI, & Lee C (2006). Temperament and character dimensions in patients with atopic dermatitis. *J. Dermatol* 33, 10–15. doi: 10.1111/j.1346-8138.2006.00002.x [PubMed: 16469078]
- Kirsten K, Fior D, Kreutz LC & Barcellos LJG (2018). First description of behavior and immune system relationship in fish. *Sci. Rep.*, 8, 846. doi: 10.1038/s41598-018-19276-3 [PubMed: 29339805]
- Koprivnikar J, Gibson CH & Redfern JC (2012). Infectious personalities: behavioural syndromes and disease risk in larval amphibians. *Proc Biol Sci.*, 279, 1544–1550. doi: 10.1098/rspb.2011.2156. [PubMed: 22090390]
- Krajčl-Fiser S, Scheiber IB, Blejec A, Mostl E & Kotrschal K (2007). Individualities in a flock of free-roaming greylag geese: behavioral and physiological consistency over time and across situations. *Horm. Behav.*, 51, 239–48. doi: 10.1016/j.yhbeh.2006.10.006 [PubMed: 17196200]
- La Gruta NL, Kedzierska K, Stambas J & Doherty PC (2007). A question of self-preservation: immunopathology in influenza virus infection. *Immunol Cell Biol.*, 85, 85–92. doi:10.1038/sj.icb.7100026 [PubMed: 17213831]
- Lee KA (2006). Linking immune defenses and life history at the levels of the individual and the species. *Integr Comp Biol.*, 46, 1000–1015. doi: 10.1093/icb/icl049 [PubMed: 21672803]
- Lee JH, Yu HH, Wang LC, Yang YH, Lin YT & Chiang BL (2007). The levels of CD4+CD25+ regulatory T cells in paediatric patients with allergic rhinitis and bronchial asthma. *Clin. Exp. Immunol.*, 148, 53–63. doi: 10.1111/j.1365-2249.2007.03329.x [PubMed: 17349011]
- Li B, Duan XH, Wu JF, Liu BJ, Luo QL, Jin HL, DU YJ, Zhang HY, Cao YX & Dong JC (2013). Impact of psychosocial stress on airway inflammation and its mechanism in a murine model of allergic asthma. *Chin. Med. J. (Engl.)*, 126, 325–334. doi: 10.3760/cma.j.issn.0366-6999.20120685 [PubMed: 23324285]
- Lochmiller RL & Deerenberg C (2000). Trade-offs in evolutionary immunology: just what is the cost of immunity? *OIKOS*, 88, 87–98. doi: 10.1034/j.1600-0706.2000.880110.x
- Lyson K & McCann SM (1991). The effect of interleukin-6 on pituitary hormone release in vivo and in vitro. *Neuroendocrinology*, 54, 262–266. doi: 10.1159/000125884 [PubMed: 1658674]
- Maggi E (1998). The TH1/TH2 paradigm in allergy. *Immunotechnology*, 3, 233–244. doi: 10.1016/S1380-2933(97)10005-7 [PubMed: 9530557]
- Martin LB 2nd, Hasselquist D & Wikelski M (2006). Investment in immune defense is linked to pace of life in house sparrows. *Oecologia*, 147, 565–575. doi: 10.1007/s00442-005-0314-y [PubMed: 16450181]
- Martin LB 2nd, Weil ZM & Nelson RJ (2007). Immune defense and reproductive pace of life in *Peromyscus* mice. *Ecology*, 88, 2516–2518. doi: <https://doi.org/10.1890/07-0060.1> [PubMed: 18027755]
- Mason WA, Capitanio JP, Machado CJ, Mendoza SP & Amaral DG (2006). Amygdalectomy and responsiveness to novelty in rhesus monkeys (*Macaca mulatta*): generality and individual consistency of effects. *Emotion*, 6, 73–81. doi: 10.1037/1528-3542.6.1.73 [PubMed: 16637751]
- Mastorakos G, Chrousos GP & Weber JS (1993). Recombinant interleukin-6 activates the hypothalamic-pituitary-adrenal axis in humans. *J Clin Endocrinol Metab.*, 77, 1690–1694. doi: 10.1210/jcem.77.6.8263159 [PubMed: 8263159]
- Matsumoto K, Inoue H, Fukuyama S, Kan-O K, Eguchi-Tsuda M, Matsumoto T, Moriwaki A, Nakano T & Nakanishi Y (2009). Frequency of Foxp3+CD4CD25+ T cells is associated with the phenotypes of allergic asthma. *Respirology*, 14, 187–194. doi:10.1111/j.1440-1843.2008.01472.x [PubMed: 19192224]
- McDade TW (2005). Life history, maintenance, and the early origins of immune function. *Am J Hum Biol.*, 17, 81–94. doi: 10.1002/ajhb.20095 [PubMed: 15612049]
- Medzhitov R, Schneider DD & Soares MP (2012). Disease tolerance as a defense strategy. *Science*, 335, 936–941. doi: 10.1126/science.1214935 [PubMed: 22363001]
- Ménard C, Pfau ML, Hodes GE & Russo SJ (2016). Immune and neuroendocrine mechanisms of stress vulnerability and resilience. *Neuropsychopharmacology*, 42, 62–80. doi: 10.1038/npp.2016.90 [PubMed: 27291462]

- Miller GE, Cohen S & Ritchey AK (2002). Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health Psychol.*, 21, 531–541. doi: 10.1037//0278-6133.21.6.531 [PubMed: 12433005]
- Mosser DM & Edwards JP (2008). Exploring the full spectrum of macrophage activation. *Nat Rev Immunol.*, 8, 958–969. doi:10.1038/nri2448. [PubMed: 19029990]
- Munck A & Náray-Fejes-Tóth A (1994). Glucocorticoids and stress: permissive and suppressive actions. *Ann N Y Acad Sci.*, 746, 115–130. doi:10.1111/j.1749-6632.1994.tb39221.x [PubMed: 7825870]
- National Research Council (2011). *Guide for the Care and Use of Laboratory Animals*. Washington, DC: The National Academies Press. doi: 10.17226/12910
- Oster M, Scheel M, Muráni E, Ponsuksili S, Zebunke M, Puppe B & Wimmers K (2015). The fight-or-flight response is associated with PBMC expression profiles related to immune defence and recovery in swine. *PLoS One*, 10, e0120153. doi: 10.1371/journal.pone.0120153 [PubMed: 25793368]
- Papanicolaou DA, Wilder RL, Manolagas SC & Chrousos GP (1998). The pathophysiologic roles of interleukin-6 in human disease. *Ann. Intern. Med.*, 128, 127–137. doi: 10.7326/0003-4819-128-2-199801150-00009 [PubMed: 9441573]
- Patel PD, Katz M, Karssen AM & Lyons DM (2008). Stress-induced changes in corticosteroid receptor expression in primate hippocampus and prefrontal cortex. *Psychoneuroendocrinology*, 33, 360–367. doi: 10.1016/j.psyneuen.2007.12.003 [PubMed: 18222612]
- Pateras IS, Havaki S, Nikitopoulou X, Vougas K, Townsend PA, Panayiotidis MI, Georgakilas AG & Gorgoulis VG (2015). The DNA damage response and immune signaling alliance: Is it good or bad? Nature decides when and where. *Pharmacol. Ther.*, 154, 36–56. doi: 10.1016/j.pharmthera.2015.06.011 [PubMed: 26145166]
- Peiris JSM, Hui KPY, & Yen H-L (2010). Host response to Influenza virus: protection versus immunopathology. *Curr Opin Immunol.*, 22, 475–481. doi: 10.1016/j.coi.2010.06.003 [PubMed: 20594815]
- Picard M, Juster R-P & McEwen BS (2014). Mitochondrial allostatic load puts the ‘glue’ back in glucocorticoids. *Nat. Rev. Endocrinology*, 10, 303–310. doi:10.1038/nrendo.2014.22 [PubMed: 24663223]
- Pruessner JC, Kirschbaum C, Meinlschmid G & Hellhammer DH (2003). Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*, 28, 916–931. doi: 10.1016/j.bbi.2014.05.016 [PubMed: 12892658]
- Pruett SB, Collier S, Wu W-J, Fan R (1999). Quantitative relationships between the suppression of selected immunological parameters and the area under the corticosterone concentration vs. time curve in B6C3F1 mice subjected to exogenous corticosterone or to restraint stress. *Toxicol Sci.*, 49, 272–280. doi: 10.1093/toxsci/49.2.272 [PubMed: 10416272]
- Punt J, Stranford S, Jones P & Owen J (2018). *Kuby Immunology 8th Edition*: New York, NY W.H. Freeman and Company.
- Rangassamy M, Khaleghparast Athari S, Monclús R, Boissier MC, Bessis N & Rödel HG (2016). Personality modulates proportions of CD4⁺ regulatory and effector T cells in response to socially induced stress in a rodent of wild origin. *Physiol Behav.*, 167, 255–264. doi: 10.1016/j.physbeh.2016.09.016 [PubMed: 27666163]
- Rauw WM (2012). Immune response from a resource allocation perspective. *Front Genet.*, 3, 267. doi: 10.3389/fgene.2012.00267 [PubMed: 23413205]
- Ray J & Hansen S (2004). Temperament in the rat: sex differences and hormonal influences on harm avoidance and novelty seeking. *Behavioral Neuroscience*, 118, 488–497. doi: 10.1037/0735-7044.118.3.488 [PubMed: 15174926]
- Reale D, Reader SM, Sol D, McDougall PT & Dingemanse NJ (2007). Integrating animal temperament within ecology and evolution. *Biol Rev Camb Philos Soc.*, 82, 291–318. doi: 10.1111/j.1469-185X.2007.00010.x [PubMed: 17437562]

- Roberts BW, Walton KE & Viechtbauer W (2006). Patterns of mean-level change in personality traits across the life course: a meta-analysis of longitudinal studies. *Psychol. Bull*, 132, 1–25. doi: 10.1037/0033-2909.132.1.1 [PubMed: 16435954]
- Rochford KD & Cummins PM (2015). The blood-brain barrier endothelium: a target for pro-inflammatory cytokines. *Biochem Soc Trans.*, 43, 702–706. doi: 10.1042/BST20140319 [PubMed: 26551716]
- Roved J, Westerdahl H & Hasselquist D (2017). Sex differences in immune responses: Hormonal effects, antagonistic selection, and evolutionary consequences. *Horm Behav.*, 88, 95–105. doi: 10.1016/j.yhbeh.2016.11.017 [PubMed: 27956226]
- Salome N, Tasiemski A, Dutriez I, Wigger A, Landgraf R & Viltart O (2008). Immune challenge induces differential corticosterone and interleukin-6 responsiveness in rats bred for extremes in anxiety-related behavior. *Neuroscience*, 151, 1112–1118. doi:10.1016/j.neuroscience.2007.12.010 [PubMed: 18207648]
- Schmidt LA, Fox NA, Rubin KH, Sternberg EM, Gold PW, Smith CC & Schulkin J (1997). Behavioral and neuroendocrine responses in shy children. *Dev. Psychobiol* 30, 127–140. doi:10.1002/(SICI)1098-2302(199703)30:2<127::AID-DEV4>3.0.CO;2-S [PubMed: 9068967]
- Schmidt D, Reber SO, Botteron C, Barth T, Peterlik D, Uschold N, Männel DN & Lechner A (2010). Chronic psychosocial stress promotes systemic immune activation and the development of inflammatory Th cell responses. *Brain Behav. Immun.*, 24, 1097–1104. doi: 10.1016/j.bbi.2010.04.014. [PubMed: 20451603]
- Sild E, Sepp T & Hõrak P (2011). Behavioural trait covaries with immune responsiveness in a wild passerine. *Brain Behav Immun.*, 25, 1349–1354. doi: 10.1016/j.bbi.2011.03.020 [PubMed: 21473910]
- Sloan EK, Capitanio JP, Tarara RP & Cole SW (2008). Social temperament and lymph node innervation. *Brain Behav. Immun* 22, 717–726. doi:10.1016/j.bbi.2007.10.010 [PubMed: 18068331]
- Smider NA, Essex MJ, Kalin NH, Buss KA, Klein MH, Davidson RJ & Goldsmith HH (2002). Salivary cortisol as a predictor of socioemotional adjustment during kindergarten: a prospective study. *Child Dev.* 73, 75–92. doi:10.1111/1467-8624.00393 [PubMed: 14717245]
- Sorrells SF, Munhoz CD, Manley NC, Yen S & Sapolsky RM (2014). Glucocorticoids increase excitotoxic injury and inflammation in the hippocampus of adult male rats. *Neuroendocrinology*, 100, 129–140. doi: 10.1159/000367849 [PubMed: 25228100]
- Stark JL, Avitsur R, Padgett DA, Campbell KA, Beck FM & Sheridan JF (2001). Social stress induces glucocorticoid resistance in macrophages. *Am J Physiol Regulatory Integr. Comp. Physiol* 280, R1799–R1805. doi: 10.1152/ajpregu.2001.280.6.R1799
- Steptoe A, Shankar A, Demakakos P & Wardle J (2013). Social isolation, loneliness, and all-cause mortality in older men and women. *Proc Natl Acad Sci U S A.*, 110, 5797–5801. doi:10.1073/pnas.1219686110 [PubMed: 23530191]
- Takano A, Arakawa R, Hayashi M, Takahashi H, Ito H & Suhara T (2007). Relationship between neuroticism personality trait and serotonin transporter binding. *Biol Psychiatry*, 62, 588–592. doi: 10.1016/j.biopsych.2006.11.007 [PubMed: 17336939]
- Tarka M, Guenther A, Niemelä PT, Nakagawa S, & Noble DWA (2018). Sex differences in life history, behavior, and physiology along a slow-fast continuum: a meta-analysis. *Behav Ecol Sociobiol.*, 72, 132. doi: 10.1007/s00265-018-2534-2 [PubMed: 30100667]
- van de Garde MDB, Martinez FO, Melgert BN, Hylkema MN, Jonkers RE & Hamann J (2014). Chronic exposure to glucocorticoids shapes gene expression and modulates innate and adaptive activation pathways in macrophages with distinct changes in leukocyte attraction. *J. Immunol* 192, 1196–1208. doi:10.4049/jimmunol.1302138 [PubMed: 24395918]
- Verbeek MEM, Drent PJ & Wiepkema PR (1994). Consistent individual differences in early exploratory behaviour of male great tits. *Anim. Behav.*, 48, 1113–1121. doi:10.1006/anbe.1994.1344
- Visintin A, Mazzoni A, Spitzer JH, Wyllie DH, Dower SK & Segal DM (2001). Regulation of Toll-like receptors in human monocytes and dendritic cells. *J. Immunol* 166, 249–255. doi: 10.4049/jimmunol.166.1.249 [PubMed: 11123299]

- Wang LH, Lin YH, Yang J & Guo W (2009). Insufficient increment of CD4+CD25+ regulatory T cells after stimulation in vitro with allergen in allergic asthma. *Int. Arch. Allergy Immunol*, 148, 199–210. doi:10.1159/000161580 [PubMed: 18849611]
- Webster Marketon JI & Glaser R (2008). Stress hormones and immune function. *Cell. Immunol*, 252, 16–26. doi:10.1016/j.cellimm.2007.09.006 [PubMed: 18279846]
- Wilson DS, Clark AB, Coleman K & Dearstyne T (1994). Shyness and boldness in humans and other animals. *Trends in Ecology & Evolution*, 9, 442–446. doi:10.1016/0169-5347(94)90134-1
- Wood PG, Karol MH, Kusnecov AW & Rabin BS (1993). Enhancement of antigen-specific humoral and cell-mediated immunity by electric footshock stress in rats. *Brain Behav. Immun*, 7, 121–134. doi:10.1006/brbi.1993.1014 [PubMed: 8347894]
- Zozulya AA, Gabaeva MV, Sokolovm OY, Surkina ID & Kost NV (2008). Personality, coping style, and constitutional neuroimmunology. *J Immunotoxicol.*, 5, 221–225. doi: 10.1080/15476910802131444. [PubMed: 18569393]
- Zuckerman M (1979). Sensation seeking and risk-taking In: Izard CE, ed. *Emotions in Personality and Psychopathology*. Plenum Press, New York.
- Zuk M & Stoehr AM (2002). Immune defense and host life history. *Am Nat.*, 160 Suppl 4, S9–S22. doi: 10.1086/342131. [PubMed: 18707455]
- Zylberberg M, Klasing KC & Hahn TP (2014). In house finches, *Haemorrhous mexicanus*, risk takers invest more in innate immune function. *Anim. Behav*, 89, 115–122. doi:10.1016/j.anbehav.2013.12.021

HIGHLIGHTS:

- Different immune responses may explain temperament-specific health outcomes.
- Rats were tested for stable behavioral phenotypes by testing on two different arenas.
- Behaviorally-inhibited (BI) rats had a greater innate response than non-inhibited (NI).
- BI rats had a dampened delayed-type hypersensitivity response compared to NI rats.

a.

b.

Figure 1.

Overhead view of the novel object (a) and novel social (b) arenas. Test rat is placed in a clean ceramic bowl and lowered into the corner of the arena and allowed to explore freely for 5 minutes.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2. Male Sprague-Dawley rat circulating interleukin-6 (IL-6) concentration at baseline and in response (at 2, 4, 6, 10 hr) to 4 different doses of intraperitoneal lipopolysaccharide (LPS; 0, 5, 25, 125 µg/kg; n = 6-7 rats in each dosage group respectively). Circles indicate means, error bars indicate S.E.M.

Figure 3a,b.

(a) Mean circulating interleukin-6 (IL-6) concentration at baseline and after intraperitoneal lipopolysaccharide (LPS) injection for Behaviorally-Inhibited (BI), Non-Inhibited (NI), and Mixed-Behavior male rats ($n = 4, 6, 17$ respectively). Left-hand figure depicts mean IL-6 concentration at each sampling time, right-hand figure depicts the total IL-6 response to LPS represented as area-under-the-curve relative to ground (AUCg) during the 10 hrs following LPS injection. Error bars indicate S.E.M. (b) Mean circulating corticosterone concentration at baseline and after intraperitoneal lipopolysaccharide (LPS) injection for BI, NI, and Mixed male rats ($n = 4, 6, 17$ respectively; same rats as those whose data are depicted in Figure 3a). Left-hand figure depicts mean corticosterone concentration at each sampling time, right-hand figure depicts the total corticosterone response to LPS represented as area-under-the-curve relative to ground (AUCg) during the 10 hrs following LPS injection. Error

bars indicate S.E.M. For both AUCg figures, asterisk indicates significant differences ($p < .05$) between groups in post-hoc comparisons.

Figure 4.

Delayed-type hypersensitivity response, as measured by relative footpad swelling in response to keyhole limpet hemocyanin (KLH) re-exposure in Behaviorally-Inhibited (BI), Non-Inhibited (NI), and Mixed-Behavior male rats ($n = 9, 6, 13$ respectively). Relative swelling represents the difference in footpad thickness between the saline-exposed footpad and the KLH re-exposed footpad. Error bars indicate S.E.M. Asterisk indicates significant difference ($p < .05$) between groups in post-hoc comparisons.