Published in final edited form as:

Clin Rev Bone Miner Metab. 2019 June ; 17(2): 65–76. doi:10.1007/s12018-019-09260-4.

Low bone mineral density in anorexia nervosa: Treatments and challenges

Pouneh K. Fazeli, MD, MPH

Neuroendocrine Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA

Abstract

Anorexia nervosa, a psychiatric disease predominantly affecting women, is characterized by self-induced starvation and a resultant low-weight state. During starvation, a number of hormonal adaptations – including hypothalamic amenorrhea and growth hormone resistance – allow for decreased energy expenditure during periods of decreased nutrient intake, but these very same adaptations also contribute to the medical complications associated with chronic starvation, including low bone mass. Almost 90% of women with anorexia nervosa have bone mineral density (BMD) values more than one-standard deviation below the mean of healthy women at peak bone mineral density and this disease is associated with a significantly increased risk of fracture. Although multiple therapies have been studied for the treatment of low bone mass in anorexia nervosa, there are currently no approved therapies and few promising long-term therapeutic options. This review will outline the mediators of low bone mass in anorexia nervosa, discuss therapies that have been studied for the treatment of low BMD in this disorder, and highlight the important challenges that remain, including the differences in bone modeling in adolescents with anorexia nervosa as compared to adults, necessitating that potential therapies be tested in these two populations separately, and the paucity of long-term therapeutic strategies for treating bone loss in this disorder.

Keywords

anorexia nervosa; bone mineral density; estrogen

Introduction

Anorexia nervosa, a psychiatric disorder predominantly affecting women, is characterized by low body-weight due to self-induced starvation. The disease, which typically presents during

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <http://www.springer.com/gb/open-access/authors-rights/aam-terms-v1>

Corresponding Author: Pouneh K. Fazeli, MD, Neuroendocrine Unit, Massachusetts General Hospital, 55 Fruit Street, Bulfinch 457, Boston, MA 02114, Tel: 617-726-3870, Fax: 617-726-5072, pkfazeli@partners.org.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Conflict of Interest: The author reports no disclosures or conflicts of interest.

Ethical approval: Not applicable

Informed consent: Not applicable

adolescence [1], has a lifetime prevalence of approximately 2.2% [2]. Importantly, this is a chronic disorder in that only 50–60% of women with anorexia nervosa recover even twenty-years after diagnosis [3, 4]; therefore this disease is also prevalent in older women.

The most common medical complication associated with anorexia nervosa is low bone mass [5]. More than 50% of women with anorexia nervosa meet the definition of osteopenia, defined as a T-score between -1 and -2.5 compared to healthy women at peak bone mineral density (BMD), and an additional 34–38% meet the definition of osteoporosis (T-score -2.5) [5, 6]. This loss of bone mass is predominantly due to hormonal adaptations critical for reducing energy expenditure during periods of low caloric intake, including functional hypothalamic amenorrhea and growth hormone resistance. Although these adaptations advantageously divert energy away from reproduction and growth during periods of low nutrient intake, the long-term consequences of these adaptations include loss of bone mass and importantly, a significantly increased risk of fracture [7–10]. Approximately 30% of adolescent girls and women report a prior history of fracture [5, 9] and prospective data similarly demonstrate a seven-fold increased risk of fracture in young women with anorexia nervosa [7].

Given the chronic nature of this disorder and the fact that anorexia nervosa typically develops during adolescence, which is a critical time for bone accrual, the increased risk of fracture may persist even many years after initial diagnosis; a population-based study demonstrated a 57% cumulative incidence of fracture forty-years after the diagnosis of anorexia nervosa [11]. Therefore, finding effective, long-term treatment strategies for the prevention of bone loss in anorexia nervosa is critical. In this review, we will discuss possible determinants of the low bone mass and increased fracture risk in anorexia nervosa and we will review the efficacy of treatments that have been studied in this population. We will also discuss challenges that remain in this field, including the current lack of long-term therapeutic options for the treatment of bone loss in adult women with anorexia nervosa and the persistently heightened fracture risk in individuals who have recovered from this disorder.

Body composition and possible nutritional mediators of low bone mass

Lean Body Mass and Peripheral Fat Stores

Individuals with anorexia nervosa have less lean body mass [12] and lower subcutaneous and visceral fat stores [13] as compared to normal weight controls. In most studies, lean body mass has been found to be a strong positive predictor of BMD in both girls and women with anorexia nervosa [14–16] and lean mass, but not fat mass, has been shown to be positively associated with parameters of bone geometry, as assessed by hip structural analysis [17]. Similarly, in longitudinal studies, change in lean body mass, but not change in fat mass, is positively associated with change in BMD [18, 19], likely due to the increased mechanical loading on bone with increased muscle mass.

Given the positive association between lean body mass and BMD in anorexia nervosa, an important question is whether exercise should be encouraged as a possible treatment for low bone mass. Whereas some studies in adolescents and adults demonstrate positive

associations between physical activity and BMD [20–24], others have shown no association [14, 25–27]. Importantly, one study showed that women who exercised more than 6 hours/week had lower BMD compared to women who exercised between 1–6 hours/week, suggesting a possible negative effect of increased physical activity [28]. Physical activity has also been associated with amenorrhea [29] and will necessitate increasing caloric intake. Therefore, while moderate physical activity does not necessarily need to be discouraged, data to date do not support encouraging exercise for the treatment of low bone mass in anorexia nervosa.

Leptin

Although fat mass does not appear to be a significant determinant of BMD in anorexia nervosa, leptin --an anorexigenic hormone secreted by adipose tissue -- may be a determinant, in part because it is likely a key hormonal mediator in the development of hypogonadotropic hypogonadism in states of undernutrition. Compared to normal-weight controls, levels of leptin are low in girls and women with anorexia nervosa [30, 31]. Treatment with recombinant human leptin in women with functional hypothalamic amenorrhea has been shown to lead to recovery of menses when compared to placebo-treated individuals [32] and an increase in markers of bone formation [33]. Although leptin levels are positively associated with BMD in anorexia nervosa [34, 35], suggesting that low levels may be a contributor to low bone mass in this disorder, weight loss is a side effect of recombinant human leptin treatment, possibly due to its anorexigenic effects [32, 33]. Therefore, given the possible risk of further weight loss, leptin is not currently a potential treatment option for bone loss in individuals with anorexia nervosa.

Marrow adipose tissue

In contrast to peripheral adipose tissue stores, marrow adipose tissue may be a determinant of BMD in anorexia nervosa. Marrow adipose tissue is a component of the bone marrow microenvironment and in cross-sectional studies, levels of marrow adipose tissue are higher in anorexia nervosa as compared to controls [36, 37] and inversely associated with BMD [36]. In women who have recovered from anorexia nervosa, marrow adipose tissue levels appear to normalize and are similar to those of normal-weight controls [38]. Why this fat depot increases in size when peripheral fat stores are actively being used during chronic starvation is not known but in a recent longitudinal study, we demonstrate that levels of marrow adipose tissue increase with short-term weight gain and are positively associated with changes in BMD [39], suggesting that this fat depot may serve a different function with dynamic changes in weight as compared to the steady-state. Marrow adipose tissue has been associated with impaired bone integrity [40] and in anorexia nervosa, marrow adipose tissue is negatively associated with estimates of bone strength in adolescent girls [41], suggesting that it could be a determinant of increased fracture risk in this population.

Calcium, Vitamin D, and Protein intake

Although insufficient caloric intake is a defining characteristic of anorexia nervosa, inadequate intake of calcium, vitamin D, and/or protein do not appear to be important factors contributing to loss of bone mass in this population. Compared to controls, adolescents with anorexia nervosa are less frequently vitamin D deficient and consume more calcium and

vitamin D, primarily through supplement intake [42, 43]. Similarly, although isolated protein deficiency, independent of caloric intake, may be associated with decreased cortical bone thickness [44] and low BMD [45], adolescents with anorexia nervosa have similar protein intake compared to normal-weight controls [42, 46, 47]. Therefore, insufficient calcium, vitamin D or protein intake are unlikely to be significant factors contributing to low bone mass in anorexia nervosa but given the known positive association between low 25-OH vitamin D levels and low BMD [48], individuals with anorexia nervosa should be screened and treated for low 25-OH vitamin D levels.

Hyponatremia

Seven to 17% of women with anorexia nervosa have low sodium levels [5, 49]. This is likely due to a decreased ability to excrete free water in the setting of low solute intake. In rodent studies, hyponatremia is associated with reductions in trabecular and cortical bone [50] and in humans, hyponatremia is independently associated with an increased risk of fracture [51, 52]. In anorexia nervosa, women with a plasma sodium level of <135 mmol/L have lower BMD compared to those with a sodium level ≥135 mmol/L [53], suggesting that hyponatremia may be a determinant of low BMD in anorexia nervosa.

Hormonal adaptations to undernutrition (Table 1)

Functional hypothalamic amenorrhea

Hypoestrogenemia—Although it is no longer a component of the diagnostic criteria in the most recent edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5), amenorrhea, is a common finding in the majority of women with anorexia nervosa [54]. Amenorrhea, or the absence of at least three consecutive menstrual periods, is due to disruption of gonadotropin-releasing hormone (GnRH) secretion [55–59]. This shunting of energy away from the costly process of reproduction during periods of undernutrition is an adaptive, energy-preserving response, but the resultant hypoestrogenemia contributes to the loss of bone mass in states of chronic undernutrition such as anorexia nervosa. Low estrogen states are characterized by increased bone remodeling and subsequent bone loss [60–63] and in individuals with anorexia nervosa, longer duration of amenorrhea is associated with lower BMD [21, 25].

Despite the fact that duration of estrogen deficiency is significantly correlated with low BMD in anorexia nervosa, oral estrogen therapy has not been shown to improve BMD in this population. Whereas retrospective studies suggested beneficial effects of oral estrogen on BMD in anorexia nervosa [21], subsequent randomized and/or prospective studies demonstrated no overall benefit in BMD in individuals receiving oral estrogen versus placebo [64–66]. In contrast, physiologic doses of predominantly transdermal estrogen have been shown to be effective in increasing BMD in adolescent girls with anorexia nervosa [67]. BMD in the spine increased by 2.6% after 18 months of transdermal estrogen compared to < 0.5% in those not treated [67]. The reason for this difference in efficacy with low-dose transdermal estrogen as compared to oral estrogen may be due to suppression of insulin-like growth factor (IGF)-1 – a bone-trophic hormone that is already low in individuals with anorexia nervosa [68]. In contrast to transdermal estrogen, oral estrogen has

been shown to have greater IGF-1 suppressive effects in a population of postmenopausal women [69, 70]. Similarly, higher doses of oral estrogen, such as the doses found in oral contraceptive pills, have greater IGF-1 suppressing effects compared to lower, more physiologic doses of oral estrogen [70]. Importantly, IGF-1 has been significantly correlated with BMD in anorexia nervosa [15] and therefore further suppression of this bone anabolic hormone may be the critical reason for the difference in efficacy between oral and transdermal estrogen in anorexia nervosa.

Although transdermal estrogen is a promising treatment for low bone mass in anorexia nervosa, it has only been studied in adolescents with anorexia nervosa, and there are important differences in bone modeling/remodeling in adolescents as compared to adults with this disease that may influence efficacy of treatment. Compared to normal-weight controls, adolescents with anorexia nervosa have lower levels of bone formation markers and similar levels of bone resorption markers [27, 71, 72], suggesting an overall low bone remodeling state. In contrast, adult women with anorexia nervosa have similarly low levels of bone formation markers but *elevated* levels of bone resorption markers compared to controls [73–76]. This difference in bone modeling/remodeling in adolescents versus adults may be due in part to the fact that adult women with anorexia nervosa are estrogen deficient in comparison to healthy adult women and as detailed above, estrogen deficiency is associated with increased bone remodeling [61–63, 77]. Therefore, bone loss in adult women with anorexia nervosa may be due to decreased bone formation in the setting of accelerated bone remodeling, whereas in adolescents with anorexia nervosa, there is an overall state of decreased remodeling. These differences may help explain differences observed in adolescents versus adults with anorexia nervosa in response to treatments for low bone mass. For example, 12-months of treatment with a bisphosphonate significantly improved BMD in adult women with anorexia nervosa compared to patients treated with placebo [78], whereas 12-months of treatment with a bisphosphonate in adolescent girls did not result in a significant difference in change in BMD compared to placebo-treated patients [79]. Given differences in bone modeling/remodeling in adolescents versus adults, bisphosphonates, which decrease rates of bone remodeling [80], may be more effective in adults with anorexia nervosa with their accelerated rate of bone remodeling as compared to adolescents with their overall low remodeling state. Therefore, potential therapies for bone loss must be studied separately in adults and adolescents with anorexia nervosa and it is not yet known if transdermal estrogen is an effective treatment for bone loss in adult women with anorexia nervosa.

Low testosterone levels—Low testosterone levels, an androgen produced in part by the ovary, are also a characteristic finding in women with anorexia nervosa [81, 82]. Low levels of testosterone are associated with decreased BMD and impaired bone microarchitecture in this population [82, 83]. Despite these associations, transdermal testosterone treatment was not effective in increasing BMD in a randomized, placebo-controlled study in women with anorexia nervosa [78]. Therefore, decreased levels of testosterone may not be an important determinant of bone mass in anorexia nervosa.

Growth hormone resistance

Growth hormone (GH) is secreted by the pituitary gland and stimulates the liver to produce IGF-1 – a hormone which mediates the majority of the growth-promoting effects of GH. GH has additional functions, including fat mobilization [84], and the counter-regulatory effects of GH may be critical for sustaining euglycemia during periods of starvation [85]. Therefore, acquired GH resistance, which is signified by normal or elevated GH levels in the setting of low IGF-1 levels, allows for the counter-regulatory effects of GH to be maintained while energy is diverted away from the costly process of growth in states of starvation such as anorexia nervosa [86–89]. Yet similar to functional hypothalamic amenorrhea, this energy-preserving adaptation is also a likely contributor to the loss of bone mass.

Low IGF-1 levels—IGF-1 – a bone anabolic hormone -- is produced and secreted by the liver in response to GH, but this hormone is extremely responsive to nutritional cues as well. During an acute fast, IGF-1 levels decrease significantly after four days and this drop in IGF-1 levels is coupled with a decrease in bone formation markers [90]. In a more chronic state of starvation, such as anorexia nervosa, levels of IGF-1 are also significantly lower as compared to controls [68]. Importantly, in anorexia nervosa, levels of GH are normal or elevated in the setting of low IGF-1 levels [86–89], demonstrating the primary importance of nutritional status in the regulation of IGF-1. In fact, even supraphysiologic doses of recombinant human GH do not raise IGF-1 levels in women with anorexia nervosa [91].

These decreased IGF-1 levels are likely a significant contributor to the loss of bone mass in anorexia nervosa. IGF-1 has been shown to be positively associated with BMD in women with anorexia nervosa [15] and in prospective studies, treatment with recombinant human IGF-1 increases bone formation markers in both girls and women with anorexia nervosa [92, 93]. Treatment with recombinant human IGF-1 has also been shown to increase lumbar spine BMD in women with anorexia nervosa when combined with oral estrogen [66]. Importantly, given the differences in bone remodeling in adolescents versus adults with anorexia nervosa, the effects of IGF-1 on BMD in adolescent girls with anorexia nervosa are not yet known but this is an active area of investigation.

Fibroblast growth factor (FGF)21—Fibroblast growth factor (FGF)21 is a hormone induced during fasting in both mice [94, 95] and humans [96]. Although the mechanism underlying GH resistance in humans is not known, in murine models, FGF21 has been shown to be a mediator of GH resistance through a reduction in STAT5 – a transcription factor critical for GH signaling and induction of IGF-1 [97]. In a cross-sectional study, FGF21 was inversely associated with IGF-1 and positively associated with GH- area-under-the-curve in adolescents with anorexia nervosa, suggesting that FGF21 may also be a mediator of GH resistance in humans as well [98]. In murine models, FGF21 transgenic mice weigh less than wild-type littermates, have decreased bone mass, and an increase in the number and area of bone marrow adipocytes -- a potential determinant of BMD in anorexia nervosa [99, 100], and therefore FGF21 transgenic mice are phenotypically similar to individuals with anorexia nervosa. In a cross-sectional study in women with anorexia nervosa, FGF21 levels were inversely associated with bone microarchitecture parameters

and estimates of bone strength [101], suggesting a possible role for FGF21 in the impaired bone phenotype of women with anorexia nervosa.

Adrenal steroids

Hypercortisolemia—Elevated cortisol levels may also be an important mediator of low bone mass in anorexia nervosa. The negative effects of cortisol on bone are the result of a number of mechanisms including decreased calcium availability via a decrease in calcium absorption in the intestine and an increase in urinary calcium excretion [102–104], an impairment of bone formation via reduced osteoblast proliferation [105], and an increase in bone resorption due in part to inhibition of gonadotropin secretion [106]. Levels of cortisol are higher in individuals with anorexia nervosa, a physiologically stressful state, as compared to normal-weight controls [107–110] and cortisol is inversely correlated with bone formation markers and BMD in adolescent girls and in women with anorexia nervosa [25, 109, 110]. Therefore, cortisol may be an important mediator of bone loss in anorexia nervosa.

Dehydroepiandrosterone—Levels of the adrenal androgen precursor dehydroepiandrosterone (DHEA) and its sulfated form, DHEA-S [111, 112] are lower in individuals with anorexia nervosa compared to controls and these low levels are associated with both increased levels of bone resorption markers and decreased BMD [22, 82]. Studies investigating the effects of DHEA replacement in individuals with anorexia nervosa demonstrate that treatment with DHEA in combination with oral estrogen for 18 months leads to improvement in parameters of bone geometry, as assessed by hip structural analysis, and stabilization of BMD as compared to placebo [113, 114]. In contrast, treatment with DHEA for six months or one year without oral estrogen does not lead to improvements in BMD when compared to placebo-treated individuals or those treated with oral estrogen [115, 116]. Therefore, low DHEA levels may also contribute to the low bone mass in individuals with anorexia nervosa.

Other potential hormonal determinants

PYY

Similar to leptin, peptide YY (PYY), a hormone secreted by the intestine, is also an anorexigenic hormone. In contrast to leptin, PYY levels are elevated in girls and women with anorexia nervosa [47, 117], although whether these increased levels contribute to the decreased hunger reported in this disease is not known [118]. In murine models, compared to wildtype littermates, overexpression of PYY in female mice results in lower BMD and trabecular bone volume [119] and in anorexia nervosa, elevated PYY levels are associated with low BMD [117]. Therefore, elevated PYY levels may also be a mediator of low bone mass in anorexia nervosa.

Ghrelin

Ghrelin is an orexigenic hormone secreted by the stomach's fundal cells. Compared to healthy controls, individuals with anorexia nervosa have higher levels of ghrelin [120] but appear resistant to its appetite-stimulating effects, as women with anorexia nervosa report

decreased hunger compared to controls [118]. Ghrelin administration has been shown to increase BMD in a rodent model [121] and in normal-weight adolescent girls, ghrelin is positively correlated with BMD [122], suggesting that ghrelin has beneficial effects on bone. Yet in a longitudinal study in girls with anorexia nervosa, there was a negative association between baseline ghrelin levels and changes in BMD [123], suggesting that individuals with anorexia nervosa may also be resistance to ghrelin's positive bone effects. It is possible that this resistance to the effects of ghrelin are due to differences in the ratio of acylated ghrelin -- the active form of ghrelin which binds to the growth hormone secretagogue receptor 1a (GHSR1a) -- to desacylated ghrelin, as levels of desacylated ghrelin are significantly higher in individuals with anorexia nervosa as compared to controls [124]. In a double-blind, randomized, placebo-controlled study, we have recently shown that treatment with a GHSR1a-agonist was able to overcome resistance to the gastric motility effects of ghrelin in women with anorexia nervosa [125] and we also found a trend towards a decrease in marrow adipose tissue -- elevated levels of which are a potential determinant of low BMD in anorexia nervosa -- with treatment [126]. Further studies will be necessary to determine whether treatment with a GHSR1a-agonist can also overcome the resistance to the bone anabolic effects of ghrelin in individuals with anorexia nervosa.

Oxytocin

Oxytocin, a hormone secreted by the posterior pituitary known primarily for promoting uterine contractions during childbirth and milk ejection during lactation, has also been shown to have bone anabolic effects [127]. In an ovariectomized-osteoporotic mouse model, treatment with oxytocin led to improved trabecular bone volume fraction and a decrease in marrow adipocyte number [128]. Compared to controls, levels of oxytocin are lower in women with anorexia nervosa and positively associated with BMD [129]; therefore, low oxytocin levels may also contribute to the low bone mass phenotype in this population.

Preadipocyte Factor (Pref)-1

Preadipocyte factor (Pref)-1 is a member of the epidermal growth factor family of proteins and an inhibitor of osteoblast and adipocyte differentiation [130]. In anorexia nervosa, circulating levels of Pref-1 are elevated compared to controls and Pref-1 is inversely associated with BMD and positively associated with marrow adipose tissue [34]. Further studies are necessary to better understand the role of circulating Pref-1 and its association with bone mass and marrow adipose tissue in anorexia nervosa.

Sclerostin

Sclerostin is secreted by osteocytes and negatively affects bone formation via inhibition of Wnt signaling. In postmenopausal women, treatment with a sclerostin monoclonal antibody increases BMD [131] and reduces risk of fracture [132]. Sclerostin levels are higher in anorexia nervosa compared to controls [133], suggesting that it may be a determinant of low bone mass in this population, but elevated levels are not associated with low spine or hip BMD in anorexia nervosa [133] and changes in circulating sclerostin levels do not appear to mediate increases in BMD in response to six months of teriparatide in women with anorexia nervosa [134] or in response to 12 months of transdermal estrogen in adolescents [135].

Therefore, further study is necessary to better understand the potential role of sclerostin in the low bone mass phenotype of anorexia nervosa.

Challenges in the treatment of low BMD in anorexia nervosa

The most significant challenges in the treatment of loss of bone mass in anorexia nervosa are 1) the chronicity of the disease, 2) a lack of long-term therapeutic options, and 3) the persistence of low bone mass and increased fracture risk even after recovery [11, 136]. Because only 50–60% of women recover from anorexia nervosa decades after their initial diagnosis [3, 4], **long-term** therapeutic strategies for low bone mass are critical but currently lacking. We will review various pharmacologic therapies that have been studied for the treatment of bone loss in anorexia nervosa below (Table 2), but to date, the most effective long-term treatment for bone loss in anorexia nervosa is recovery from the disorder, inclusive of both weight recovery and a resumption of menses. Weight recovery accompanied by a resumption of menses results in an annual rate of increase in BMD of 3.1% in the spine and 1.8% in the hip [19], which is in sharp contrast to an annual rate of bone loss of 2.6% in the spine and 2.4% in the hip in women who remain amenorrheic and low-weight [19]. Yet even in patients who recover from anorexia nervosa, there may be a persistence of low BMD and increased fracture risk [11, 136] and there are currently no prospective studies investigating how to best treat this post-recovery population.

Given the association between hormonal adaptations to starvation, including functional hypothalamic amenorrhea and growth hormone resistance, and low bone mass, a number of hormonal treatment strategies have been studied in girls and women with persistent anorexia nervosa; these hormonal treatments include estrogen, DHEA, IGF-1, and testosterone. Physiologic doses of predominantly transdermal estrogen increase spine BMD by 2.6% in adolescent girls after 18-months of treatment [67] and therefore transdermal estrogen is a promising long-term treatment option for low bone mass in adolescents with anorexia nervosa. In contrast, oral estrogen, predominantly in the form of oral contraceptives, when used alone, does not improve BMD in women or adolescents with anorexia nervosa [64–66]. Yet when added to other hormonal treatments, including IGF-1 or DHEA, oral estrogen shows modest efficacy; treatment with recombinant human IGF-1 combined with oral contraceptives increases spine BMD by 1.8% after 9 months in adult women with anorexia nervosa [66] and the combination of oral estrogen and DHEA is effective in preserving BMD in a mixed population of adolescent girls and adult women with anorexia nervosa [113]. Importantly, given the differences in bone remodeling in adults versus adolescents with anorexia nervosa, the effects on BMD of IGF-1 combined with estrogen in adolescents and the effects of transdermal estrogen in adults are not known but are currently being investigated. The effects of 12 months of transdermal testosterone have also been studied in women with anorexia nervosa, but testosterone did not improve BMD as compared to placebo [78].

Additional treatments, including bisphosphonates and teriparatide, have demonstrated benefit in short-term studies in women with anorexia nervosa, but long-term use of these medications is limited. Six months of teriparatide increases spine BMD by 6–10% in women with anorexia nervosa [134], but teriparatide is only approved for 24 months of use and also

carries an FDA warning because of an increased risk of osteosarcoma observed in rat studies [137]. Therefore, teriparatide cannot be used long-term and should not be prescribed to individuals at increased risk of osteosarcoma, including individuals with open epiphyses. Similarly, compared with placebo, bisphosphonates increase spine and hip BMD by 2–4% after 12 months in adults [78], but in adolescents treated for 12 months, bisphosphonates do not increase BMD any more than placebo [79], emphasizing the need to separately test potential therapies for bone loss in both adolescent and adult populations. Importantly, long-term use of bisphosphonates has been associated with an increased risk of atypical femoral fractures in population-based studies [138], and in rodent studies, bisphosphonates cross the placenta and are associated with fetal abnormalities [139]. Given the long half-life of bisphosphonates in bone [140], caution must be taken when considering bisphosphonate therapy in premenopausal women, as the risks of bisphosphonates to the human fetus have not been well-established. Therefore, there are currently no promising BMD-improving therapies that can be used long-term in adults with anorexia nervosa.

Conclusions

Anorexia nervosa is a chronic psychiatric illness, predominantly affecting women and characterized by inadequate caloric intake [3, 4]. Low bone mass, due primarily to hormonal adaptations to the state of chronic undernutrition, is the most common medical complication associated with anorexia nervosa and approximately 90% of women have BMD values more than one standard deviation below the mean of comparably-aged women [5, 6]. Importantly, this low bone mass is associated with a significantly increased risk of fracture [5, 7] and therefore therapeutic strategies for preventing this morbidity are critical. Although there are currently no approved treatments for low bone mass in anorexia nervosa, physiologic, transdermal estrogen is a promising therapy for adolescents. The treatments which have demonstrated the greatest benefit in adult women with anorexia nervosa, notably bisphosphonates and teriparatide, are not potential long-term treatment options and therefore there remains a significant need for therapies for the treatment of low BMD in the adult population. It is also important to note that we do not know whether any of the treatments that improve BMD in anorexia nervosa also reduce the fracture rate in this population. Therefore, future studies are critically needed to evaluate the *long-term* efficacy and safety of potential treatment strategies for bone loss in anorexia nervosa.

Acknowledgements:

This project was supported by NIH grant R03 DK106410. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Funding: This project was supported by NIH grant R03 DK106410

References:

1. Lucas AR, Crowson CS, O'Fallon WM, Melton LJ 3rd. The ups and downs of anorexia nervosa. *The International Journal of Eating Disorders*. 1999;26(4):397–405. [PubMed: 10550780]
2. Keski-Rahkonen A, Hoek HW, Susser ES, Linna MS, Sihvola E, Raevuori A et al. Epidemiology and course of anorexia nervosa in the community. *The American Journal of Psychiatry*. 2007;164(8):1259–65. doi:10.1176/appi.ajp.2007.06081388. [PubMed: 17671290]

3. Lowe B, Zipfel S, Buchholz C, Dupont Y, Reas DL, Herzog W. Long-term outcome of anorexia nervosa in a prospective 21-year follow-up study. *Psychological medicine*. 2001;31(5):881–90. [PubMed: 11459385]
4. Eddy KT, Tabri N, Thomas JJ, Murray HB, Keshaviah A, Hastings E et al. Recovery From Anorexia Nervosa and Bulimia Nervosa at 22-Year Follow-Up. *The Journal of clinical psychiatry*. 2017;78(2): 184–9. doi:10.4088/JCP.15m10393. [PubMed: 28002660]
5. Miller KK, Grinspoon SK, Ciampa J, Hier J, Herzog D, Klibanski A. Medical findings in outpatients with anorexia nervosa. *Archives of internal medicine*. 2005;165(5):561–6. doi:10.1001/archinte.165.5.561. [PubMed: 15767533]
6. Grinspoon S, Thomas E, Pitts S, Gross E, Mickley D, Miller K et al. Prevalence and predictive factors for regional osteopenia in women with anorexia nervosa. *Annals of internal medicine*. 2000;133(10):790–4. [PubMed: 11085841]
7. Rigotti NA, Neer RM, Skates SJ, Herzog DB, Nussbaum SR. The clinical course of osteoporosis in anorexia nervosa. A longitudinal study of cortical bone mass. *JAMA : the journal of the American Medical Association*. 1991;265(9):1133–8. [PubMed: 1995999]
8. Vestergaard P, Emborg C, Stoving RK, Hagen C, Mosekilde L, Brixen K. Fractures in patients with anorexia nervosa, bulimia nervosa, and other eating disorders--a nationwide register study. *The International journal of eating disorders*. 2002;32(3):301–8. doi:10.1002/eat.10101. [PubMed: 12210644]
9. Faje AT, Fazeli PK, Miller KK, Katzman DK, Ebrahimi S, Lee H et al. Fracture risk and areal bone mineral density in adolescent females with anorexia nervosa. *The International journal of eating disorders*. 2014;47(5):458–66. doi:10.1002/eat.22248. [PubMed: 24430890]
10. Nagata JM, Golden NH, Leonard MB, Copelovitch L, Denburg MR. Assessment of Sex Differences in Fracture Risk Among Patients With Anorexia Nervosa: A Population-Based Cohort Study Using The Health Improvement Network. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2017;32(5):1082–9. doi:10.1002/jbmr.3068.
11. Lucas AR, Melton LJ 3rd, Crowson CS, O'Fallon WM. Long-term fracture risk among women with anorexia nervosa: a population-based cohort study. *Mayo Clinic proceedings Mayo Clinic*. 1999;74(10):972–7. doi:10.4065/74.10.972.
12. Grinspoon S, Thomas L, Miller K, Pitts S, Herzog D, Klibanski A. Changes in regional fat redistribution and the effects of estrogen during spontaneous weight gain in women with anorexia nervosa. *The American journal of clinical nutrition*. 2001;73(5):865–9. doi:10.1093/ajcn/73.5.865. [PubMed: 11333838]
13. Mayer L, Walsh BT, Pierson RN Jr, Heymsfield SB, Gallagher D, Wang J et al. Body fat redistribution after weight gain in women with anorexia nervosa. *The American journal of clinical nutrition*. 2005;81(6):1286–91. doi:10.1093/ajcn/81.6.1286. [PubMed: 15941877]
14. Kooh SW, Noriega E, Leslie K, Muller C, Harrison JE. Bone mass and soft tissue composition in adolescents with anorexia nervosa. *Bone*. 1996;19(2):181–8. [PubMed: 8853863]
15. Grinspoon S, Miller K, Coyle C, Krempin J, Armstrong C, Pitts S et al. Severity of osteopenia in estrogen-deficient women with anorexia nervosa and hypothalamic amenorrhea. *The Journal of clinical endocrinology and metabolism*. 1999;84(6):2049–55. doi:10.1210/jcem.84.6.5792. [PubMed: 10372709]
16. Turner JM, Bulsara MK, McDermott BM, Byrne GC, Prince RL, Forbes DA. Predictors of low bone density in young adolescent females with anorexia nervosa and other dieting disorders. *The International journal of eating disorders*. 2001;30(3):245–51. [PubMed: 11746283]
17. DiVasta AD, Beck TJ, Petit MA, Feldman HA, LeBoff MS, Gordon CM. Bone cross-sectional geometry in adolescents and young women with anorexia nervosa: a hip structural analysis study. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2007;18(6): 797–804. doi:10.1007/s00198-006-0308-6.
18. Soyka LA, Misra M, Frenchman A, Miller KK, Grinspoon S, Schoenfeld DA et al. Abnormal bone mineral accrual in adolescent girls with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2002;87(9):4177–85. doi:10.1210/jc.2001-011889. [PubMed: 12213868]

19. Miller KK, Lee EE, Lawson EA, Misra M, Minihan J, Grinspoon SK et al. Determinants of skeletal loss and recovery in anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2006;91(8):2931–7. doi:10.1210/jc.2005-2818. [PubMed: 16735492]
20. Rigotti NA, Nussbaum SR, Herzog DB, Neer RM. Osteoporosis in women with anorexia nervosa. *The New England journal of medicine*. 1984;311(25):1601–6. doi:10.1056/NEJM198412203112503. [PubMed: 6504095]
21. Seeman E, Szmukler GI, Formica C, Tsalamandris C, Mestrovic R. Osteoporosis in anorexia nervosa: the influence of peak bone density, bone loss, oral contraceptive use, and exercise. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 1992;7(12):1467–74. doi:10.1002/jbmr.5650071215.
22. Gordon CM, Goodman E, Emans SJ, Grace E, Becker KA, Rosen CJ et al. Physiologic regulators of bone turnover in young women with anorexia nervosa. *The Journal of pediatrics*. 2002;141(1):64–70. doi:10.1067/mpd.2002.125003. [PubMed: 12091853]
23. DiVasta AD, Feldman HA, O'Donnell JM, Long J, Leonard MB, Gordon CM. Effect of Exercise and Antidepressants on Skeletal Outcomes in Adolescent Girls With Anorexia Nervosa. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 2017;60(2):229–32. doi:10.1016/j.jadohealth.2016.10.003. [PubMed: 27939877]
24. Nagata JM, Carlson JL, Golden NH, Murray SB, Long J, Leonard MB et al. Associations between exercise, bone mineral density, and body composition in adolescents with anorexia nervosa. *Eat Weight Disord*. 2018. doi:10.1007/s40519-018-0521-2.
25. Biller BM, Saxe V, Herzog DB, Rosenthal DI, Holzman S, Klibanski A. Mechanisms of osteoporosis in adult and adolescent women with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1989;68(3):548–54. doi:10.1210/jcem-68-3-548. [PubMed: 2493036]
26. Bachrach LK, Guido D, Katzman D, Litt IF, Marcus R. Decreased bone density in adolescent girls with anorexia nervosa. *Pediatrics*. 1990;86(3):440–7. [PubMed: 2388792]
27. Soyka LA, Grinspoon S, Levitsky LL, Herzog DB, Klibanski A. The effects of anorexia nervosa on bone metabolism in female adolescents. *The Journal of clinical endocrinology and metabolism*. 1999;84(12):4489–96. doi:10.1210/jcem.84.12.6207. [PubMed: 10599707]
28. Joyce JM, Warren DL, Humphries LL, Smith AJ, Coon JS. Osteoporosis in women with eating disorders: comparison of physical parameters, exercise, and menstrual status with SPA and DPA evaluation. *J Nucl Med*. 1990;31(3):325–31. [PubMed: 2308003]
29. Frisch RE, Gotz-Welbergen AV, McArthur JW, Albright T, Witschi J, Bullen B et al. Delayed menarche and amenorrhea of college athletes in relation to age of onset of training. *JAMA : the journal of the American Medical Association*. 1981;246(14):1559–63. [PubMed: 7277629]
30. Grinspoon S, Gulick T, Askari H, Landt M, Lee K, Anderson E et al. Serum leptin levels in women with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1996;81(11):3861–3. [PubMed: 8923829]
31. Misra M, Miller KK, Kuo K, Griffin K, Stewart V, Hunter E et al. Secretory dynamics of leptin in adolescent girls with anorexia nervosa and healthy adolescents. *American journal of physiology Endocrinology and metabolism*. 2005;289(3):E373–81. doi:10.1152/ajpendo.00041.2005. [PubMed: 15811876]
32. Chou SH, Chamberland JP, Liu X, Matarese G, Gao C, Stefanakis R et al. Leptin is an effective treatment for hypothalamic amenorrhea. *Proceedings of the National Academy of Sciences of the United States of America*. 2011;108(16):6585–90. doi:10.1073/pnas.1015674108. [PubMed: 21464293]
33. Welt CK, Chan JL, Bullen J, Murphy R, Smith P, DePaoli AM et al. Recombinant human leptin in women with hypothalamic amenorrhea. *The New England journal of medicine*. 2004;351(10):987–97. doi:10.1056/NEJMoa040388. [PubMed: 15342807]
34. Fazeli PK, Bredella MA, Misra M, Meenaghan E, Rosen CJ, Clemmons DR et al. Preadipocyte factor-1 is associated with marrow adiposity and bone mineral density in women with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2010;95(1):407–13. doi:10.1210/jc.2009-1152. [PubMed: 19850693]

35. Legroux-Gerot I, Vignau J, Biver E, Pigny P, Collier F, Marchandise X et al. Anorexia nervosa, osteoporosis and circulating leptin: the missing link. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2010;21(10):1715–22. doi:10.1007/s00198-009-1120-x.
36. Bredella MA, Fazeli PK, Miller KK, Misra M, Torriani M, Thomas BJ et al. Increased bone marrow fat in anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2009;94(6):2129–36. doi:10.1210/jc.2008-2532. [PubMed: 19318450]
37. Ecklund K, Vajapeyam S, Feldman HA, Buzney CD, Mulkern RV, Kleinman PK et al. Bone marrow changes in adolescent girls with anorexia nervosa. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2010;25(2):298–304. doi:10.1359/jbmr.090805.
38. Fazeli PK, Bredella MA, Freedman L, Thomas BJ, Breggia A, Meenaghan E et al. Marrow fat and preadipocyte factor-1 levels decrease with recovery in women with anorexia nervosa. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2012;27(9):1864–71. doi:10.1002/jbmr.1640.
39. Fazeli PK, Faje A, Bredella MA, Polineni S, Russell S, Resulaj M et al. Changes in marrow adipose tissue with short-term changes in weight in premenopausal women with anorexia nervosa. *European journal of endocrinology / European Federation of Endocrine Societies*. 2018. doi:10.1530/EJE-18-0824.
40. Schellinger D, Lin CS, Hatipoglu HG, Fertikh D. Potential value of vertebral proton MR spectroscopy in determining bone weakness. *AJNR American journal of neuroradiology*. 2001;22(8):1620–7. [PubMed: 11559519]
41. Singhal V, Tulsiani S, Campoverde KJ, Mitchell DM, Slattery M, Schorr M et al. Impaired bone strength estimates at the distal tibia and its determinants in adolescents with anorexia nervosa. *Bone*. 2018;106:61–8. doi:10.1016/j.bone.2017.07.009. [PubMed: 28694162]
42. Misra M, Tsai P, Anderson EJ, Hubbard JL, Gallagher K, Soyka LA et al. Nutrient intake in community-dwelling adolescent girls with anorexia nervosa and in healthy adolescents. *The American journal of clinical nutrition*. 2006;84(4):698–706. [PubMed: 17023694]
43. Haagenen AL, Feldman HA, Ringelheim J, Gordon CM. Low prevalence of vitamin D deficiency among adolescents with anorexia nervosa. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2008;19(3):289–94. doi:10.1007/s00198-007-0476-z.
44. Garn SM, Rohmann CG, Behar M, Viteri F, Guzman MA. Compact Bone Deficiency in Protein-Calorie Malnutrition. *Science*. 1964;145(3639):1444–5. [PubMed: 14172641]
45. Fontana L, Shew JL, Holloszy JO, Villareal DT. Low bone mass in subjects on a long-term raw vegetarian diet. *Archives of internal medicine*. 2005;165(6):684–9. doi:10.1001/archinte.165.6.684. [PubMed: 15795346]
46. Fernstrom MH, Weltzin TE, Neuberger S, Srinivasagam N, Kaye WH. Twenty-four-hour food intake in patients with anorexia nervosa and in healthy control subjects. *Biol Psychiatry*. 1994;36(10):696–702. [PubMed: 7880939]
47. Misra M, Miller KK, Tsai P, Gallagher K, Lin A, Lee N et al. Elevated peptide YY levels in adolescent girls with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2006;91(3):1027–33. doi:10.1210/jc.2005-1878. [PubMed: 16278259]
48. Gatti D, El Ghoch M, Viapiana O, Ruocco A, Chignola E, Rossini M et al. Strong relationship between vitamin D status and bone mineral density in anorexia nervosa. *Bone*. 2015;78:212–5. doi:10.1016/j.bone.2015.05.014. [PubMed: 25980743]
49. Mehler PS, Blalock DV, Walden K, Kaur S, McBride J, Walsh K et al. Medical findings in 1,026 consecutive adult inpatient-residential eating disordered patients. *The International journal of eating disorders*. 2018;51(4):305–13. doi:10.1002/eat.22830. [PubMed: 29417593]
50. Verbalis JG, Barsony J, Sugimura Y, Tian Y, Adams DJ, Carter EA et al. Hyponatremia-induced osteoporosis. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2010;25(3):554–63. doi:10.1359/jbmr.090827.

51. Kinsella S, Moran S, Sullivan MO, Molloy MG, Eustace JA. Hyponatremia independent of osteoporosis is associated with fracture occurrence. *Clin J Am Soc Nephrol*. 2010;5(2):275–80. doi:10.2215/CJN.06120809. [PubMed: 20056759]
52. Jamal SA, Arampatzis S, Harrison SL, Bucur RC, Ensrud K, Orwoll ES et al. Hyponatremia and Fractures: Findings From the MrOS Study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2015;30(6):970–5. doi:10.1002/jbmr.2383.
53. Lawson EA, Fazeli PK, Calder G, Putnam H, Misra M, Meenaghan E et al. Plasma sodium level is associated with bone loss severity in women with anorexia nervosa: a cross-sectional study. *The Journal of clinical psychiatry*. 2012;73(11):e1379–83. doi:10.4088/JCP.12m07919. [PubMed: 23218167]
54. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-5). 2013;5th edition:Washington, DC.
55. Wiegmann W, Solbach HG. Effects of LH-RH on plasma levels of LH and FSH in anorexia nervosa. *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et métabolisme*. 1972;4(5):404. doi:10.1055/s-0028-1097104. [PubMed: 4566864]
56. Mecklenburg RS, Loriaux DL, Thompson RH, Andersen AE, Lipsett MB. Hypothalamic dysfunction in patients with anorexia nervosa. *Medicine*. 1974;53(2):147–59. [PubMed: 4593749]
57. Travaglini P, Beck-Peccoz P, Ferrari C, Ambrosi B, Paracchi A, Severgnini A et al. Some aspects of hypothalamic-pituitary function in patients with anorexia nervosa. *Acta endocrinologica*. 1976;81(2):252–62. [PubMed: 813472]
58. Nillius SJ, Fries H, Wide L. Successful induction of follicular maturation and ovulation by prolonged treatment with LH-releasing hormone in women with anorexia nervosa. *American journal of obstetrics and gynecology*. 1975;122(8):921–8. [PubMed: 1098466]
59. Boyar RM, Katz J, Finkelstein JW, Kapen S, Weiner H, Weitzman ED et al. Anorexia nervosa. Immaturity of the 24-hour luteinizing hormone secretory pattern. *The New England journal of medicine*. 1974;291(17):861–5. doi:10.1056/NEJM197410242911701. [PubMed: 4412035]
60. Riis BJ, Rodbro P, Christiansen C. The role of serum concentrations of sex steroids and bone turnover in the development and occurrence of postmenopausal osteoporosis. *Calcified tissue international*. 1986;38(6):318–22. [PubMed: 3089552]
61. Seeman E. Estrogen, androgen, and the pathogenesis of bone fragility in women and men. *Curr Osteoporos Rep*. 2004;2(3):90–6. [PubMed: 16036088]
62. Falahati-Nini A, Riggs BL, Atkinson EJ, O'Fallon WM, Eastell R, Khosla S. Relative contributions of testosterone and estrogen in regulating bone resorption and formation in normal elderly men. *The Journal of clinical investigation*. 2000;106(12):1553–60. doi:10.1172/JCI10942. [PubMed: 11120762]
63. Finkelstein JS, Lee H, Leder BZ, Burnett-Bowie SA, Goldstein DW, Hahn CW et al. Gonadal steroid-dependent effects on bone turnover and bone mineral density in men. *The Journal of clinical investigation*. 2016;126(3):1114–25. doi:10.1172/JCI84137. [PubMed: 26901812]
64. Klibanski A, Biller BM, Schoenfeld DA, Herzog DB, Saxe VC. The effects of estrogen administration on trabecular bone loss in young women with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1995;80(3):898–904. [PubMed: 7883849]
65. Golden NH, Lanzkowsky L, Schebendach J, Palestro CJ, Jacobson MS, Shenker IR. The effect of estrogen-progestin treatment on bone mineral density in anorexia nervosa. *J Pediatr Adolesc Gynecol*. 2002;15(3):135–43. [PubMed: 12106749]
66. Grinspoon S, Thomas L, Miller K, Herzog D, Klibanski A. Effects of recombinant human IGF-I and oral contraceptive administration on bone density in anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2002;87(6):2883–91. [PubMed: 12050268]
67. Misra M, Katzman D, Miller KK, Mendes N, Snelgrove D, Russell M et al. Physiologic estrogen replacement increases bone density in adolescent girls with anorexia nervosa. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2011;26(10):2430–8. doi:10.1002/jbmr.447.
68. Counts DR, Gwirtsman H, Carlsson LM, Lesem M, Cutler GB Jr The effect of anorexia nervosa and refeeding on growth hormone-binding protein, the insulin-like growth factors (IGFs), and the

- IGF-binding proteins. *The Journal of clinical endocrinology and metabolism*. 1992;75(3):762–7. [PubMed: 1381372]
69. Weissberger AJ, Ho KK, Lazarus L. Contrasting effects of oral and transdermal routes of estrogen replacement therapy on 24-hour growth hormone (GH) secretion, insulin-like growth factor I, and GH-binding protein in postmenopausal women. *The Journal of clinical endocrinology and metabolism*. 1991;72(2):374–81. doi:10.1210/jcem-72-2-374. [PubMed: 1991807]
 70. Kam GY, Leung KC, Baxter RC, Ho KK. Estrogens exert route- and dose-dependent effects on insulin-like growth factor (IGF)-binding protein-3 and the acid-labile subunit of the IGF ternary complex. *The Journal of clinical endocrinology and metabolism*. 2000;85(5):1918–22. doi: 10.1210/jcem.85.5.6527. [PubMed: 10843175]
 71. Saggese G, Bertelloni S, Baroncelli GI, Di Nero G. Serum levels of carboxyterminal propeptide of type I procollagen in healthy children from 1st year of life to adulthood and in metabolic bone diseases. *Eur J Pediatr*. 1992;151(10):764–8. [PubMed: 1425800]
 72. Heer M, Mika C, Grzella I, Heussen N, Herpertz-Dahlmann B. Bone turnover during inpatient nutritional therapy and outpatient follow-up in patients with anorexia nervosa compared with that in healthy control subjects. *The American journal of clinical nutrition*. 2004;80(3):774–81. doi: 10.1093/ajcn/80.3.774. [PubMed: 15321821]
 73. Hotta M, Fukuda I, Sato K, Hizuka N, Shibasaki T, Takano K. The relationship between bone turnover and body weight, serum insulin-like growth factor (IGF) I, and serum IGF-binding protein levels in patients with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2000;85(1):200–6. [PubMed: 10634387]
 74. Weinbrenner T, Zittermann A, Gouni-Berthold I, Stehle P, Berthold HK. Body mass index and disease duration are predictors of disturbed bone turnover in anorexia nervosa. A case- control study. *Eur J Clin Nutr*. 2003;57(10):1262–7. doi:10.1038/sj.ejcn.1601683. [PubMed: 14506487]
 75. Bolton JG, Patel S, Lacey JH, White S. A prospective study of changes in bone turnover and bone density associated with regaining weight in women with anorexia nervosa. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2005;16(12):1955–62. doi: 10.1007/s00198-005-1972-7.
 76. Viapiana O, Gatti D, Dalle Grave R, Todesco T, Rossini M, Braga V et al. Marked increases in bone mineral density and biochemical markers of bone turnover in patients with anorexia nervosa gaining weight. *Bone*. 2007;40(4):1073–7. doi:10.1016/j.bone.2006.11.015. [PubMed: 17240212]
 77. Hughes DE, Dai A, Tiffée JC, Li HH, Mundy GR, Boyce BF. Estrogen promotes apoptosis of murine osteoclasts mediated by TGF-beta. *Nat Med*. 1996;2(10):1132–6. [PubMed: 8837613]
 78. Miller KK, Meenaghan E, Lawson EA, Misra M, Gleysteen S, Schoenfeld D et al. Effects of risedronate and low-dose transdermal testosterone on bone mineral density in women with anorexia nervosa: a randomized, placebo-controlled study. *The Journal of clinical endocrinology and metabolism*. 2011;96(7):2081–8. doi:10.1210/jc.2011-0380. [PubMed: 21525157]
 79. Golden NH, Iglesias EA, Jacobson MS, Carey D, Meyer W, Schebendach J et al. Alendronate for the treatment of osteopenia in anorexia nervosa: a randomized, double-blind, placebo-controlled trial. *The Journal of clinical endocrinology and metabolism*. 2005;90(6):3179–85. doi:10.1210/jc.2004-1659. [PubMed: 15784715]
 80. Seeman E Reduced bone formation and increased bone resorption: rational targets for the treatment of osteoporosis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2003;14 Suppl 3:S2–8. doi:10.1007/s00198-002-1340-9.
 81. van Binsbergen CJ, Coelingh Bennink HJ, Odink J, Haspels AA, Koppeschaar HP. A comparative and longitudinal study on endocrine changes related to ovarian function in patients with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1990;71(3):705–11. [PubMed: 2203799]
 82. Miller KK, Lawson EA, Mathur V, Wexler TL, Meenaghan E, Misra M et al. Androgens in women with anorexia nervosa and normal-weight women with hypothalamic amenorrhea. *The Journal of clinical endocrinology and metabolism*. 2007;92(4):1334–9. doi:10.1210/jc.2006-2501. [PubMed: 17284620]

83. Lawson EA, Miller KK, Bredella MA, Phan C, Misra M, Meenaghan E et al. Hormone predictors of abnormal bone microarchitecture in women with anorexia nervosa. *Bone*. 2010;46(2):458–63. doi:10.1016/j.bone.2009.09.005. [PubMed: 19747572]
84. Gahete MD, Cordoba-Chacon J, Luque RM, Kineman RD. The rise in growth hormone during starvation does not serve to maintain glucose levels or lean mass but is required for appropriate adipose tissue response in female mice. *Endocrinology*. 2013;154(1):263–9. doi:10.1210/en.2012-1849. [PubMed: 23150490]
85. Zhao TJ, Liang G, Li RL, Xie X, Sleeman MW, Murphy AJ et al. Ghrelin O-acyltransferase (GOAT) is essential for growth hormone-mediated survival of calorie-restricted mice. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;107(16):7467–72. doi:10.1073/pnas.1002271107. [PubMed: 20231469]
86. Garfinkel PE, Brown GM, Stancer HC, Moldofsky H. Hypothalamic-pituitary function in anorexia nervosa. *Archives of general psychiatry*. 1975;32(6):739–44. [PubMed: 1130937]
87. Scacchi M, Pincelli AI, Caumo A, Tomasi P, Delitala G, Baldi G et al. Spontaneous nocturnal growth hormone secretion in anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1997;82(10):3225–9. [PubMed: 9329343]
88. Stoving RK, Veldhuis JD, Flyvbjerg A, Vinten J, Hangaard J, Koldkjaer OG et al. Jointly amplified basal and pulsatile growth hormone (GH) secretion and increased process irregularity in women with anorexia nervosa: indirect evidence for disruption of feedback regulation within the GH-insulin-like growth factor I axis. *The Journal of clinical endocrinology and metabolism*. 1999;84(6):2056–63. [PubMed: 10372710]
89. Misra M, Miller KK, Bjornson J, Hackman A, Aggarwal A, Chung J et al. Alterations in growth hormone secretory dynamics in adolescent girls with anorexia nervosa and effects on bone metabolism. *The Journal of clinical endocrinology and metabolism*. 2003;88(12):5615–23. [PubMed: 14671143]
90. Grinspoon SK, Baum HB, Peterson S, Klibanski A. Effects of rhIGF-I administration on bone turnover during short-term fasting. *The Journal of clinical investigation*. 1995;96(2):900–6. doi:10.1172/JCI118137. [PubMed: 7543494]
91. Fazeli PK, Lawson EA, Prabhakaran R, Miller KK, Donoho DA, Clemmons DR et al. Effects of recombinant human growth hormone in anorexia nervosa: a randomized, placebo-controlled study. *The Journal of clinical endocrinology and metabolism*. 2010;95(11):4889–97. doi:10.1210/jc.2010-0493. [PubMed: 20668047]
92. Grinspoon S, Baum H, Lee K, Anderson E, Herzog D, Klibanski A. Effects of short-term recombinant human insulin-like growth factor I administration on bone turnover in osteopenic women with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1996;81(11):3864–70. doi:10.1210/jcem.81.11.8923830. [PubMed: 8923830]
93. Misra M, McGrane J, Miller KK, Goldstein MA, Ebrahimi S, Weigel T et al. Effects of rhIGF-I administration on surrogate markers of bone turnover in adolescents with anorexia nervosa. *Bone*. 2009;45(3):493–8. doi:10.1016/j.bone.2009.06.002. [PubMed: 19523548]
94. Badman MK, Pissios P, Kennedy AR, Koukos G, Flier JS, Maratos-Flier E. Hepatic fibroblast growth factor 21 is regulated by PPARalpha and is a key mediator of hepatic lipid metabolism in ketotic states. *Cell metabolism*. 2007;5(6):426–37. doi:10.1016/j.cmet.2007.05.002. [PubMed: 17550778]
95. Inagaki T, Dutchak P, Zhao G, Ding X, Gautron L, Parameswara V et al. Endocrine regulation of the fasting response by PPARalpha-mediated induction of fibroblast growth factor 21. *Cell metabolism*. 2007;5(6):415–25. doi:10.1016/j.cmet.2007.05.003. [PubMed: 17550777]
96. Fazeli PK, Lun M, Kim SM, Bredella MA, Wright S, Zhang Y et al. FGF21 and the late adaptive response to starvation in humans. *The Journal of clinical investigation*. 2015;125(12):4601–11. doi:10.1172/JCI83349. [PubMed: 26529252]
97. Inagaki T, Lin VY, Goetz R, Mohammadi M, Mangelsdorf DJ, Kliewer SA. Inhibition of growth hormone signaling by the fasting-induced hormone FGF21. *Cell metabolism*. 2008;8(1):77–83. doi:10.1016/j.cmet.2008.05.006. [PubMed: 18585098]
98. Fazeli PK, Misra M, Goldstein M, Miller KK, Klibanski A. Fibroblast growth factor-21 may mediate growth hormone resistance in anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2010;95(1):369–74. doi:10.1210/jc.2009-1730. [PubMed: 19926712]

99. Kharitonov A, Shiyanova TL, Koester A, Ford AM, Micanovic R, Galbreath EJ et al. FGF-21 as a novel metabolic regulator. *The Journal of clinical investigation*. 2005;115(6):1627–35. doi: 10.1172/JCI23606. [PubMed: 15902306]
100. Wei W, Dutchak PA, Wang X, Ding X, Wang X, Bookout AL et al. Fibroblast growth factor 21 promotes bone loss by potentiating the effects of peroxisome proliferator-activated receptor gamma. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;109(8):3143–8. doi:10.1073/pnas.1200797109. [PubMed: 22315431]
101. Fazeli PK, Faje AT, Cross EJ, Lee H, Rosen CJ, Bouxsein ML et al. Serum FGF-21 levels are associated with worsened radial trabecular bone microarchitecture and decreased radial bone strength in women with anorexia nervosa. *Bone*. 2015;77:6–11. doi:10.1016/j.bone.2015.04.001. [PubMed: 25868802]
102. Morris HA, Need AG, O'Loughlin PD, Horowitz M, Bridges A, Nordin BE. Malabsorption of calcium in corticosteroid-induced osteoporosis. *Calcified tissue international*. 1990;46(5):305–8. [PubMed: 2110853]
103. Hansen JW, Gordan GS, Prussin SG. Direct measurement of osteolysis in man. *The Journal of clinical investigation*. 1973;52(2):304–15. doi:10.1172/JCI107186. [PubMed: 4346006]
104. Canalis E Clinical review 83: Mechanisms of glucocorticoid action in bone: implications to glucocorticoid-induced osteoporosis. *The Journal of clinical endocrinology and metabolism*. 1996;81(10):3441–7. [PubMed: 8855781]
105. Rauch A, Seitz S, Baschant U, Schilling AF, Illing A, Stride B et al. Glucocorticoids suppress bone formation by attenuating osteoblast differentiation via the monomeric glucocorticoid receptor. *Cell metabolism*. 2010;11(6):517–31. doi:10.1016/j.cmet.2010.05.005. [PubMed: 20519123]
106. Padmanabhan V, Keech C, Convey EM. Cortisol inhibits and adrenocorticotropin has no effect on luteinizing hormone-releasing hormone-induced release of luteinizing hormone from bovine pituitary cells in vitro. *Endocrinology*. 1983;112(5):1782–7. [PubMed: 6299709]
107. Walsh BT, Katz JL, Levin J, Kream J, Fukushima DK, Hellman LD et al. Adrenal activity in anorexia nervosa. *Psychosom Med*. 1978;40(6):499–506. [PubMed: 734026]
108. Boyar RM, Hellman LD, Roffwarg H, Katz J, Zumoff B, O'Connor J et al. Cortisol secretion and metabolism in anorexia nervosa. *The New England journal of medicine*. 1977;296(4):190–3. doi: 10.1056/NEJM197701272960403. [PubMed: 831089]
109. Misra M, Miller KK, Almazan C, Ramaswamy K, Lapcharoensap W, Worley M et al. Alterations in cortisol secretory dynamics in adolescent girls with anorexia nervosa and effects on bone metabolism. *The Journal of clinical endocrinology and metabolism*. 2004;89(10):4972–80. doi: 10.1210/jc.2004-0723. [PubMed: 15472193]
110. Lawson EA, Donoho D, Miller KK, Misra M, Meenaghan E, Lydecker J et al. Hypercortisolemia is associated with severity of bone loss and depression in hypothalamic amenorrhea and anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2009;94(12):4710–6. doi: 10.1210/jc.2009-1046. [PubMed: 19837921]
111. Zumoff B, Walsh BT, Katz JL, Levin J, Rosenfeld RS, Kream J et al. Subnormal plasma dehydroisoandrosterone to cortisol ratio in anorexia nervosa: a second hormonal parameter of ontogenic regression. *The Journal of clinical endocrinology and metabolism*. 1983;56(4):668–72. [PubMed: 6220026]
112. Winterer J, Gwirtsman HE, George DT, Kaye WH, Loriaux DL, Cutler GB Jr. Adrenocorticotropin-stimulated adrenal androgen secretion in anorexia nervosa: impaired secretion at low weight with normalization after long-term weight recovery. *The Journal of clinical endocrinology and metabolism*. 1985;61(4):693–7. doi:10.1210/jcem-61-4-693. [PubMed: 2993344]
113. Divasta AD, Feldman HA, Giancaterino C, Rosen CJ, Leboff MS, Gordon CM. The effect of gonadal and adrenal steroid therapy on skeletal health in adolescents and young women with anorexia nervosa. *Metabolism: clinical and experimental*. 2012;61(7):1010–20. doi:10.1016/j.metabol.2011.11.016. [PubMed: 22257645]
114. DiVasta AD, Feldman HA, Beck TJ, LeBoff MS, Gordon CM. Does hormone replacement normalize bone geometry in adolescents with anorexia nervosa? *Journal of bone and mineral*

research : the official journal of the American Society for Bone and Mineral Research. 2014;29(1):151–7. doi:10.1002/jbmr.2005.

115. Gordon CM, Grace E, Emans SJ, Feldman HA, Goodman E, Becker KA et al. Effects of oral dehydroepiandrosterone on bone density in young women with anorexia nervosa: a randomized trial. *The Journal of clinical endocrinology and metabolism*. 2002;87(11):4935–41. [PubMed: 12414853]
116. Bloch M, Ish-Shalom S, Greenman Y, Klein E, Latzer Y. Dehydroepiandrosterone treatment effects on weight, bone density, bone metabolism and mood in women suffering from anorexia nervosa-a pilot study. *Psychiatry Res*. 2012;200(2–3):544–9. doi:10.1016/j.psychres.2012.07.012. [PubMed: 22858403]
117. Utz AL, Lawson EA, Misra M, Mickley D, Gleysteen S, Herzog DB et al. Peptide YY (PYY) levels and bone mineral density (BMD) in women with anorexia nervosa. *Bone*. 2008;43(1):135–9. doi:10.1016/j.bone.2008.03.007. [PubMed: 18486583]
118. Halmi KA, Sunday S, Puglisi A, Marchi P. Hunger and satiety in anorexia and bulimia nervosa. *Ann N Y Acad Sci*. 1989;575:431–44; discussion 44–5. [PubMed: 2633673]
119. Wong IP, Driessler F, Khor EC, Shi YC, Horner B, Nguyen AD et al. Peptide YY regulates bone remodeling in mice: a link between gut and skeletal biology. *PLoS One*. 2012;7(7):e40038. doi:10.1371/journal.pone.0040038. [PubMed: 22792209]
120. Otto B, Cuntz U, Fruehauf E, Wawarta R, Folwaczny C, Riepl RL et al. Weight gain decreases elevated plasma ghrelin concentrations of patients with anorexia nervosa. *European journal of endocrinology / European Federation of Endocrine Societies*. 2001;145(5):669–73.
121. Fukushima N, Hanada R, Teranishi H, Fukue Y, Tachibana T, Ishikawa H et al. Ghrelin directly regulates bone formation. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 2005;20(5):790–8. doi:10.1359/JBMR.041237.
122. Misra M, Miller KK, Stewart V, Hunter E, Kuo K, Herzog DB et al. Ghrelin and bone metabolism in adolescent girls with anorexia nervosa and healthy adolescents. *The Journal of clinical endocrinology and metabolism*. 2005;90(9):5082–7. doi:10.1210/jc.2005-0512. [PubMed: 15998770]
123. Misra M, Prabhakaran R, Miller KK, Goldstein MA, Mickley D, Clauss L et al. Prognostic indicators of changes in bone density measures in adolescent girls with anorexia nervosa-II. *The Journal of clinical endocrinology and metabolism*. 2008;93(4):1292–7. doi:10.1210/jc.2007-2419. [PubMed: 18089697]
124. Hotta M, Ohwada R, Katakami H, Shibasaki T, Hizuka N, Takano K. Plasma levels of intact and degraded ghrelin and their responses to glucose infusion in anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2004;89(11):5707–12. doi:10.1210/jc.2004-0353. [PubMed: 15531532]
125. Fazeli PK, Lawson EA, Faje AT, Eddy KT, Lee H, Fiedorek FT, Breggia A, Gaal IM, DeSanti R, Klibanski A. Treatment with a Ghrelin Agonist in Outpatient Women with Anorexia Nervosa: A Randomized Clinical Trial. *Journal of Clinical Psychiatry*. 2018;in press.
126. Fazeli PK, Klibanski A. The paradox of marrow adipose tissue in anorexia nervosa. *Bone*. 2019;118:47–52. doi:10.1016/j.bone.2018.02.013. [PubMed: 29458121]
127. Tamma R, Colaianni G, Zhu LL, DiBenedetto A, Greco G, Montemurro G et al. Oxytocin is an anabolic bone hormone. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;106(17):7149–54. doi:10.1073/pnas.0901890106. [PubMed: 19369205]
128. Elabd C, Basillais A, Beaupied H, Breuil V, Wagner N, Scheideler M et al. Oxytocin controls differentiation of human mesenchymal stem cells and reverses osteoporosis. *Stem cells*. 2008;26(9):2399–407. doi:10.1634/stemcells.2008-0127. [PubMed: 18583541]
129. Lawson EA, Donoho DA, Blum JJ, Meenaghan EM, Misra M, Herzog DB et al. Decreased nocturnal oxytocin levels in anorexia nervosa are associated with low bone mineral density and fat mass. *The Journal of clinical psychiatry*. 2011;72(11):1546–51. doi:10.4088/JCP.10m06617. [PubMed: 21903023]
130. Abdallah BM, Jensen CH, Gutierrez G, Leslie RG, Jensen TG, Kassem M. Regulation of human skeletal stem cells differentiation by Dkk1/Pref-1. *Journal of bone and mineral research : the*

- official journal of the American Society for Bone and Mineral Research. 2004;19(5):841–52. doi: 10.1359/JBMR.040118.
131. McClung MR, Grauer A, Boonen S, Bolognese MA, Brown JP, Diez-Perez A et al. Romosozumab in postmenopausal women with low bone mineral density. *The New England journal of medicine*. 2014;370(5):412–20. doi:10.1056/NEJMoa1305224. [PubMed: 24382002]
 132. Cosman F, Crittenden DB, Adachi JD, Binkley N, Czerwinski E, Ferrari S et al. Romosozumab Treatment in Postmenopausal Women with Osteoporosis. *The New England journal of medicine*. 2016;375(16):1532–43. doi:10.1056/NEJMoa1607948. [PubMed: 27641143]
 133. Maimoun L, Guillaume S, Lefebvre P, Philibert P, Bertet H, Picot MC et al. Role of Sclerostin and Dickkopf-1 in the Dramatic Alteration in Bone Mass Acquisition in Adolescents and Young Women With Recent Anorexia Nervosa. *The Journal of clinical endocrinology and metabolism*. 2014;jc20132565. doi:10.1210/jc.2013-2565.
 134. Fazeli PK, Wang IS, Miller KK, Herzog DB, Misra M, Lee H et al. Teriparatide increases bone formation and bone mineral density in adult women with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 2014;jc20134105. doi:10.1210/jc.2013-4105.
 135. Faje AT, Fazeli PK, Katzman DK, Miller KK, Breggia A, Rosen CJ et al. Sclerostin levels and bone turnover markers in adolescents with anorexia nervosa and healthy adolescent girls. *Bone*. 2012;51(3):474–9. doi:10.1016/j.bone.2012.06.006. [PubMed: 22728230]
 136. Bachrach LK, Katzman DK, Litt IF, Guido D, Marcus R. Recovery from osteopenia in adolescent girls with anorexia nervosa. *The Journal of clinical endocrinology and metabolism*. 1991;72(3):602–6. doi:10.1210/jcem-72-3-602. [PubMed: 1997514]
 137. Vahle JL, Sato M, Long GG, Young JK, Francis PC, Engelhardt JA et al. Skeletal changes in rats given daily subcutaneous injections of recombinant human parathyroid hormone (1–34) for 2 years and relevance to human safety. *Toxicologic pathology*. 2002;30(3):312–21. doi: 10.1080/01926230252929882. [PubMed: 12051548]
 138. Schilcher J, Koeppen V, Aspenberg P, Michaelsson K. Risk of atypical femoral fracture during and after bisphosphonate use. *The New England journal of medicine*. 2014;371(10):974–6. doi: 10.1056/NEJMc1403799. [PubMed: 25184886]
 139. Patlas N, Golomb G, Yaffe P, Pinto T, Breuer E, Ornoy A. Transplacental effects of bisphosphonates on fetal skeletal ossification and mineralization in rats. *Teratology*. 1999;60(2):68–73. doi:10.1002/(SICI)1096-9926(199908)60:2<68::AID-TERA10>3.0.CO;2-H. [PubMed: 10440778]
 140. Lin JH. Bisphosphonates: a review of their pharmacokinetic properties. *Bone*. 1996;18(2):75–85. [PubMed: 8833200]

Table 1:

Potential mediators of low bone mass in anorexia nervosa.

	Association with bone in anorexia nervosa	References
Low IGF-1	Low IGF-I levels are associated with low BMD	[15]
Hypoestrogenemia	Duration of amenorrhea is associated with low BMD	[25]
Low testosterone	Low testosterone levels are associated with low BMD and impaired bone microarchitecture	[82, 83]
Elevated cortisol	High cortisol levels are associated with low BMD	[110]
Low DHEAS	Low DHEAS levels are associated with low BMD	[82]
FGF21	FGF-21 levels are associated with impaired bone microarchitecture	[101]
Low leptin	Low leptin levels are associated with low BMD and impaired bone microarchitecture	[34, 83, 35]
Elevated Ghrelin	Baseline ghrelin levels are inversely associated with change in BMD in adolescent girls	[123]
Elevated PYY	Elevated PYY levels are associated with low BMD	[117]
Low Oxytocin	Low oxytocin levels are associated with low BMD	[129]
Elevated Preadipocyte Factor 1 (Pref-1)	Elevated Pref-1 levels are associated with low BMD	[34]
Elevated levels of marrow adipose tissue	Elevated marrow adipose tissue stores are associated with low BMD	[36]

Table 2:

Therapies that been investigated for the treatment of low bone mass in adults and/or adolescents with anorexia nervosa and have shown efficacy in at least one of these populations

Therapy	Effect on BMD in adults	Duration of Treatment (adults)	Effect on BMD in adolescents	Duration of Treatment (adolescents)
Physiologic estrogen replacement (transdermal)	Not known	_____	2.6% increase in spine BMD [67]	18 months
IGF-1 + oral contraceptives	1.8% increase in spine BMD [66]	9 months	Not known	_____
DHEA + oral contraceptives	Maintenance of BMD (mixed population of adolescents and adults) [113]	18 months	Maintenance of BMD (mixed population of adolescents and adults) [113]	18 months
Bisphosphonates	2–4% increase in spine/hip BMD compared to placebo [78]	12 months	No significant difference compared to placebo [79]	12 months
Teriparatide	6–10% increase in spine BMD [134]	6 months	Not known	_____