cross-section in subjects with CTD-ILD and U-ILD. This supports the findings of Maher and colleagues, who showed that an increase in CA-125 over time was a strong predictor of progressive IPF (2).

This study had several limitations. First, despite our multicenter approach, the sample sizes were relatively small, especially for CTD subtypes, which undoubtedly left some biomarker analyses underpowered. Next, batch effects and protein degradation may have influenced our results, as plasma storage time varied substantially between centers. Finally, biomarker threshold precision could not be ascertained given the multiplex platform used for this analysis. Prospective studies using validated ELISA platforms are needed.

Conclusions

Circulating plasma biomarkers are among the most readily measurable biomarkers in the human body. A wealth of biomarker data have emerged in IPF, and our findings suggest that molecular markers of progressive IPF may in fact be markers of progressive PF, irrespective of ILD subtype. Additional research is needed to validate these findings and determine how such biomarkers can inform clinical decision-making across diverse forms of ILD. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Shehabaldin Alqalyoobi, M.D.*
University of California, Davis
Sacramento, California

Ayodeji Adegunsoye, M.D., M.S.*
University of Chicago
Chicago, Illinois

Angela Linderholm, Ph.D.
University of California, Davis
Sacramento, California

Cara Hrusch, Ph.D.
University of Chicago
Chicago, Illinois

Claire Cutting, M.D.
University of California, Davis
Sacramento, California

Shwu-Fan Ma, Ph.D.
University of Virginia
Charlottesville, Virginia

Anne Sperling, Ph.D.
University of Chicago
Chicago, Illinois

Imre Noth, M.D.
University of Virginia
Charlottesville, Virginia

Mary E. Strek, M.D.
University of Chicago
Chicago, Illinois

Justin M. Oldham, M.D., M.S.†
University of California, Davis
Sacramento, California
and

Northern California VA Medical Center
Sacramento, California

ORCID ID: 0000-0001-5602-2725 (S.A.).

*These authors contributed equally to this work.

†Corresponding author (e-mail: joldham@ucdavis.edu).

References

1. Vuga LJ, Tedrow JR, Pandit KV, Tan J, Kass DJ, Xue J, *et al*. C-X-C motif chemokine 13 (CXCL13) is a prognostic biomarker of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2014;189:966–974.
2. Maher TM, Oballa E, Simpson JK, Porte J, Habgood A, Fahy WA, *et al*. An epithelial biomarker signature for idiopathic pulmonary fibrosis: an analysis from the multicentre PROFILE cohort study. *Lancet Respir Med* 2017;5:946–955.
3. Richards TJ, Kaminski N, Baribaud F, Flavin S, Brodmerkel C, Horowitz D, *et al*. Peripheral blood proteins predict mortality in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2012;185:67–76.
4. Korthagen NM, van Moorsel CH, Barlo NP, Ruven HJ, Kruit A, Heron M, *et al*. Serum and BALF YKL-40 levels are predictors of survival in idiopathic pulmonary fibrosis. *Respir Med* 2011;105:106–113.
5. Ryerson CJ, Vittinghoff E, Ley B, Lee JS, Mooney JJ, Jones KD, *et al*. Predicting survival across chronic interstitial lung disease: the ILD-GAP model. *Chest* 2014;145:723–728.
6. Ansel KM, Ngo VN, Hyman PL, Luther SA, Förster R, Sedgwick JD, *et al*. A chemokine-driven positive feedback loop organizes lymphoid follicles. *Nature* 2000;406:309–314.
7. Agostini C, Calabrese F, Poletti V, Marcer G, Facco M, Miorin M, *et al*. CXCR3/CXCL10 interactions in the development of hypersensitivity pneumonitis. *Respir Res* 2005;6:20.
8. Klimatcheva E, Pandina T, Reilly C, Torno S, Bussler H, Scrivens M, *et al*. CXCL13 antibody for the treatment of autoimmune disorders. *BMC Immunol* 2015;16:6.
9. Long X, He X, Ohshimo S, Griese M, Sarria R, Guzman J, *et al*. Serum YKL-40 as predictor of outcome in hypersensitivity pneumonitis. *Eur Respir J* 2017;49:1501924.
10. Hozumi H, Fujisawa T, Enomoto N, Nakashima R, Enomoto Y, Suzuki Y, *et al*. Clinical utility of YKL-40 in polymyositis/dermatomyositis-associated interstitial lung disease. *J Rheumatol* 2017;44:1394–1401.

Copyright © 2020 by the American Thoracic Society

Impact of Past Research Experience on Subsequent Trials: A Caution

To the Editor:

One of the primary goals for recruitment into a clinical trial is to maximize enrollment while minimizing expenses (1). For inner-city minority populations, recruitment of participants can be especially challenging, and one of the strategies used by the National Institute of Allergy and Infectious Diseases (NIAID) Inner-City Asthma

Supported in whole or in part with funds from the National Institute of Allergy and Infectious Diseases, NIH, U.S. Department of Health and Human Services, under contracts U01 AI30751, AI30752, AI30756, AI30772, AI30773-01, AI30777, AI30779, AI30780, and N01 AI15105.

Originally Published in Press as DOI: 10.1164/rccm.201905-0926LE on September 16, 2019

Consortium research program has been to recruit from a pool of current participants in natural history or epidemiologic protocols. These pools of participants typically provide researchers with a well-characterized population within which eligible study participants can be more easily identified and recruited at a considerably lower cost.

This report assesses the impact that previous participation in a research study has on the response to an intervention. The NIAID Inner-City Asthma Consortium portfolio provides a unique opportunity to assess this question. The initial National Cooperative Inner-City Asthma Study (NCICAS) was a large-scale, descriptive epidemiologic study followed by an asthma intervention study (NCICAS II). Because nearly one-third of the approximately 1,500 original participants proceeded to enroll in the subsequent intervention, this question of prior research experience can be directly tested.

NCICAS phase II, a 2-year randomized controlled trial of a tailored intervention based on the family's asthma risk (2), enrolled 1,033 participants between 5 and 11 years of age who had a physician diagnosis of asthma and persistent asthma symptoms and/or evidence of emergency department visits or hospitalizations for asthma in the prior 6 months. The study was designed as a follow-up to NCICAS I, a 12-month epidemiologic observational study that enrolled 1,529 children, ages 4 through 9 years, with the goal of identifying risk factors for asthma morbidity (3); 457 (44%) of the children enrolled in NCICAS II were recruited from NCICAS I.

The primary clinical outcome measure in NCICAS II was self-reported maximum symptom days (MSDs), a validated asthma outcome (4) that has been used in numerous NIAID Inner-City Asthma Consortium studies (5–7). MSDs were defined as the maximum number of days in the prior 2 weeks during which participants experienced any of the following: wheezing, loss of sleep due to asthma, or reduction in play activity caused by asthma (2). Secondary clinical outcome measures included a 2-month recall of asthma-related hospitalizations and unscheduled visits to the doctor or emergency department. These three measurements were collected every 2 months over the course of 2 years. The MSDs were averaged over each of the 2 years of follow-up, and hospitalization data were defined as “ever” over the course of each year separately, and number of unscheduled visits was summed within each year.

We reanalyzed the NCICAS II data, using analysis of variance as reported previously (2), but stratified on prior research involvement (research naive vs. research experienced). Analyses of the primary and secondary endpoints were adjusted for site, MSDs (MSDs <4 vs. MSDs ≥4), and baseline variables, including those that differed significantly between the research-naive and research-experienced groups. Baseline data were compared with either analysis of variance or the Mantel-Haenszel chi-square test. Statistical analyses were performed with SAS version 9.4 (SAS Institute) and R version 3.2.2 software.

All participants in NCICAS phase II, regardless of recruitment source, met the same protocol entry criteria. Research-naive participants—those who have not been involved in prior NIAID-funded inner-city asthma trials—were significantly older, were less likely to be black, and had more MSDs than research-experienced participants (Table 1).

Table 1. Baseline Characteristics of Study Populations for NCICAS II

	Enrolled in NCICAS I		P Value
	Yes (n = 457)	No. (n = 576)	
Sex, M	63% (287)	65% (347)	0.48
Age, yr, mean ± SD	7.5 ± 1.7	7.8 ± 1.9	0.003
Race			
Hispanic	17% (76)	18% (102)	0.02
Black	78% (353)	72% (415)	
Other	6% (26)	10% (59)	
Income <\$15,000/yr	71% (320)	64% (362)	0.02
Maximum symptom days (randomization)	4.6 ± 4.2	5.5 ± 4.8	0.001
Any hospitalizations?*	3% (15)	5% (31)	0.10
Number of asthma medications, mean ± SD	2.3 ± 1.4	2.9 ± 1.5	<0.001

Definition of abbreviation: NCICAS = National Cooperative Inner-City Asthma Study.

*Number represents any hospitalizations within the prior 2 months.

Figure 1 illustrates the significant -0.9 MSD difference between intervention and control groups for the research-naive participants ($P < 0.001$) in the first year of follow-up. In contrast, the difference between treatment groups for the research-experienced participants was only -0.1 MSD ($P = 0.64$). The difference in the reduction of MSDs between the two subpopulations (i.e., the research experience-by-intervention interaction term of the model) was significant ($P = 0.037$). MSDs in the second year of follow-up also showed a strong intervention effect for the research-naive population (difference, -0.8 ; $P = 0.001$) versus the research-experienced participants (difference, -0.2 ; $P = 0.50$); the experience-by-intervention interaction trended toward significance ($P = 0.10$). Similar trends toward larger treatment effects in research-naive participants were observed for the secondary outcomes of hospitalization and unscheduled visits in both Years 1 and 2, with only unscheduled visits for asthma during the second year of follow-up approaching an interaction P value of 0.11 (data not shown).

During the recruitment phase of a trial, researchers often target potential participants who have been enrolled in a previous protocol at their institution or clinic. Benefits of enrolling these participants include the following: Their contact information is already known; the staff is familiar with the participants and their families; participants are familiar with study procedures; less staff time is used for screening; and clinical measurements needed to determine study eligibility may already have been collected in the recent past, thus reducing cost and burden. Previous research experience is generally considered exclusionary only if a previous drug or intervention is likely to have a direct effect on the intervention under investigation. However, it appears that patients with asthma who have participated in an NIAID Inner-City Asthma Consortium research protocol, even a noninterventional epidemiologic protocol, respond differently to a subsequent intervention than newly recruited participants.

The NCICAS II asthma counselor intervention was consistently less effective for the study-experienced participants across a range of

Figure 1. Postbaseline maximum symptom days, by research experience. Bar height represents the adjusted estimated mean (SE) postbaseline maximum symptom days for both control (black bars) and intervention participants (gray bars). The analysis is stratified by prior research experience (naive vs. experienced) and endpoint collection time (Years 1 and 2). The *n* value represents the number analyzed within each subgroup. The intervention effect (with 95% confidence interval) is shown at the top of each panel; significant within-group effects are shown in bold. The interaction *P* value is given in the column header. NCICAS = National Cooperative Inner-City Asthma Study.

outcomes, although the experience-by-intervention interaction effect was not always statistically significant. However, all analyses were *post hoc* and exploratory, and therefore they were not powered *a priori*.

We have observed similar trends in another set of clinical trials conducted by the NIAID Inner-City Asthma Consortium, but the sample size of participants who overlapped between the two trials was limited. ICATA (Inner-City Anti-IgE Therapy for Asthma), a 60-week, double-blind, placebo-controlled, randomized clinical trial conducted to determine if guideline-based asthma care would be further improved by the addition of anti-IgE therapy (omalizumab) (6) recruited 39 (9%) of its eligible participants from the ACE (Asthma Control Evaluation) trial, a 46-week, double-blind, randomized clinical trial that determined if exhaled nitric oxide-guided therapy improved asthma control when used as an adjunct to the National Asthma Education and Prevention Program–recommended guideline-based asthma care (8). Although 9% of the ICATA participants were recruited from ACE, we observed a trend consistent with what we report in the present letter from NCICAS using a similar stratified analysis (data not shown).

These findings suggest that research-experienced participants in a clinical trial tend to have fewer asthma symptoms. Increasing the proportion of the sample with reduced morbidity will reduce the

power of the study and may bias results toward the null hypothesis, thus warranting consideration when designing protocol entry criteria for intervention trials. Investigators may wish to consider limiting the recruitment of research-experienced participants in their intervention trials or at the very least carefully examining the impact of “research experience” on their outcome measures. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Miguel A. Villarreal, M.S.*
Jeremy J. Wildfire, M.S.
Rho, Inc.
Chapel Hill, North Carolina

Christine A. Sorkness, Pharm.D.
University of Wisconsin School of Medicine and Public Health
Madison, Wisconsin

Peter J. Gergen, M.D., M.P.H.
National Institutes of Health
Rockville, Maryland

Cynthia M. Visness, Ph.D., M.P.H.
Herman E. Mitchell, Ph.D.
Rho, Inc.
Chapel Hill, North Carolina

*Corresponding author (e-mail: miguel_villarreal@rhoworld.com).

References

1. Nicholson LM, Schwirian PM, Klein EG, Skybo T, Murray-Johnson L, Eneli I, *et al.* Recruitment and retention strategies in longitudinal clinical studies with low-income populations. *Contemp Clin Trials* 2011;32:353–362.
2. Evans R III, Gergen PJ, Mitchell H, Kattan M, Kercsmar C, Crain E, *et al.* A randomized clinical trial to reduce asthma morbidity among inner-city children: results of the National Cooperative Inner-City Asthma Study. *J Pediatr* 1999;135:332–338.
3. Mitchell H, Senturia Y, Gergen P, Baker D, Joseph C, McNiff-Mortimer K, *et al.* Design and methods of the national cooperative inner-city asthma study. *Pediatr Pulmonol* 1997;24:237–252.
4. Wu TD, Perzanowski M, Peng RD, Wise RA, Balcer-Whaley S, Newman M, *et al.* Validation of the maximum symptom day among children with asthma. *J Allergy Clin Immunol* 2019;143:803–805.e10.
5. Teach SJ, Gill MA, Togias A, Sorkness CA, Arbes SJ Jr, Calatroni A, *et al.* Preseasonal treatment with either omalizumab or an inhaled corticosteroid boost to prevent fall asthma exacerbations. *J Allergy Clin Immunol* 2015;136:1476–1485.
6. Busse WW, Morgan WJ, Gergen PJ, Mitchell HE, Gern JE, Liu AH, *et al.* Randomized trial of omalizumab (anti-IgE) for asthma in inner-city children. *N Engl J Med* 2011;364:1005–1015.
7. Zoratti EM, Krouse RZ, Babineau DC, Pongracic JA, O'Connor GT, Wood RA, *et al.* Asthma phenotypes in inner-city children. *J Allergy Clin Immunol* 2016;138:1016–1029.
8. Szeffler SJ, Mitchell H, Sorkness CA, Gergen PJ, O'Connor GT, Morgan WJ, *et al.* Management of asthma based on exhaled nitric oxide in addition to guideline-based treatment for inner-city adolescents and young adults: a randomised controlled trial. *Lancet* 2008;372:1065–1072.

Copyright © 2020 by the American Thoracic Society

Identifying “Real” Patients in the Real World

To the Editor:

I applaud Dr. Dempsey and colleagues on their recent article, “Clinical Effectiveness of Antifibrotic Medications for Idiopathic Pulmonary Fibrosis,” investigating the effects of nintedanib and pirfenidone in idiopathic pulmonary fibrosis (IPF) (1). This study provides the first description of these medications in real-world clinical practice and compares patients with IPF who received antifibrotic therapy with matched subjects who received no antifibrotic therapy. The article’s illustration of patients and their demographics was heartening, especially given pulmonologists’ experience in caring for patients significantly older than those included in the INPULSIS (Efficacy and Safety of Nintedanib in Idiopathic Pulmonary Fibrosis) and ASCEND (Efficacy and Safety of Pirfenidone in Patients with Idiopathic Pulmonary Fibrosis)

§This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgerm@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.201907-1458LE on August 16, 2019

trials (2, 3). Their findings support the use of antifibrotic therapies perhaps in a larger cohort than is typically perceived to derive benefit from them.

Despite the retrospective nature of this study, the authors were able to apply an extremely rigorous methodology to ascertain the effects of these medications by using propensity score matching and local International Classification of Diseases (ICD)-9 and ICD-10 validation strategies. One additional step that might shed further light on antifibrotic effects would be to control the analysis for treatment origin site (i.e., whether the prescribing physician originated from an academic center or a community practice). As interstitial lung disease physicians, we perceive high levels of discordance among interstitial lung disease diagnoses between academic and community physicians (4). Numerous drivers explain these discrepancies, including variability in regular multidisciplinary conference participation and the infrequency with which providers may see these complex patients (5, 6). Regular practices such as multidisciplinary conferences are associated with higher diagnostic accuracy rates (7).

One could speculate about what the actual rate of “true” IPF diagnoses is in the insurance database used in this article, given the retrospective nature of the study and the use of ICD codes to identify patients. The authors astutely recognize these limitations in their discussion, although there appear to be opportunities to further enrich this analysis. Accounting for the origination site of antifibrotic therapy would provide a greater degree of confidence that they are identifying patients with IPF, as those patients would run the gauntlet of the multidisciplinary conference and would be more likely to be cared for by practitioners experienced with IPF. One representative example of this point is the large number of patients with rheumatoid arthritis, which would suggest an alternative diagnosis in >8% of the dataset’s population. Although it is not a certainty, one could envision that such misinterpretations would not occur in a setting that has more experience with this kind of patient. It is unclear what effect this modification would have on the study results, but the degree of benefits may even be intensified given that academic center-affiliated clinicians would likely have more experience in prescribing these medications, as well as more expertise in managing their adverse effects.

I would again like to commend the authors for this tremendous article demonstrating the practical effects of antifibrotics over a half decade after their approval by the U.S. Food and Drug Administration. I hope the article spurs further research in the field of IPF and leads clinicians to offer more patients the option of antifibrotic therapy. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Sean J. Callahan, M.D.*
University of Utah Health
Salt Lake City, Utah
and

George E. Wahlen Department of Veterans Affairs Medical Center
Salt Lake City, Utah

*Corresponding author (e-mail: sean.callahan@hsc.utah.edu).