Published in final edited form as:

J Behav Med. 2016 October ; 39(5): 866–875. doi:10.1007/s10865-016-9756-4.

Racial and Socioeconomic Disparities in Body Mass Index among College Students: Understanding the Role of Early Life Adversity

David S. Curtis, M.S.^a, Thomas E. Fuller-Rowell, Ph.D.^a, Stacey N. Doan, Ph.D.^b, Aleksandra E. Zgierska, M.D., Ph.D.^c, Carol D. Ryff, Ph.D.^d

^aDepartment of Human Development and Family Studies, Auburn University, AL

^bDepartment of Psychology, Claremont McKenna College, Claremont, CA

^cDepartment of Family Medicine, University of Wisconsin—Madison, WI

^dDepartment of Psychology, University of Wisconsin—Madison, WI

Abstract

The role of early life adversity (ELA) in the development of health disparities has not received adequate attention. The current study examined differential exposure and differential vulnerability to ELA as explanations for socioeconomic and racial disparities in body mass index (BMI). Data were derived from a sample of 150 college students ($M_{age}=18.8$, $SD=1.0$; 45% African American; 55% European American) who reported on parents' education and income as well as on exposure to 21 early adverse experiences. Body measurements were directly assessed to determine BMI. In adjusted models, African American students had higher BMI than European Americans. Similarly, background socioeconomic status was inversely associated with BMI. Significant mediation of group disparities through the pathway of ELA was detected, attenuating disparities by approximately 40%. Furthermore, ELA was more strongly associated with BMI for African Americans than for European Americans. Efforts to achieve health equity may need to more fully consider early adversity.

Keywords

Health disparities; early life adversity; body mass index; differential exposure; differential vulnerability; college students

Racial and socioeconomic disparities in adolescent obesity are pronounced and appear to have long-term implications for health and well-being. In 2010, approximately 12% of White/European American (EA) adolescents and 21% of Black/African American (AA)

Address correspondence to: David S. Curtis, M.S., Department of Human Development and Family Studies, 203 Spidle Hall, Auburn University, Auburn, AL 36849; Tel: +1-801-803-3846, Fax: +1-334-844-4515, dsc0019@auburn.edu.

Conflict of Interest: The authors declare that they have no conflict of interest.

Compliance with Ethical Standards

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

adolescents were obese (Ogden CL, Carroll MD, Kit BK, & Flegal KM, 2012). Similarly, 8% of adolescents whose parents had at least a bachelor's degree were obese compared to nearly 25% of adolescents whose parents had a high school diploma or less, a disparity which has grown in recent years (Frederick, Snellman, & Putnam, 2014).

With established associations between adolescent BMI and subsequent morbidity and mortality (Juonala et al., 2011; Tirosh et al., 2011; Wong, Shapiro, Boscardin, & Ettner, 2002), disparities in BMI during adolescence are a likely contributing factor to racial and socioeconomic health disparities across the life span. Late adolescence/young adulthood is also a key period for understanding the emergence of weight problems and obesity because body weight is known to increase during this period, and health behavior patterns, which are likely to persist across the life span, are being formed (Laska, Pelletier, Larson, & Story, 2012; M. C. Nelson, Story, Larson, Neumark-Sztainer, & Lytle, 2008). Therefore, understanding early life risk factors for obesity may offer critically important insights into addressing health disparities

Adverse experiences during childhood and adolescence represent one such risk factor as early adversity has been consistently associated with obesity risk and morbidity outcomes across the life span (Felitti et al., 1998; Shonkoff et al., 2012). However, few studies have examined the degree to which early life adversity (ELA) accounts for racial or socioeconomic disparities in obesity and other physical health outcomes (differential exposure hypothesis) (Fuller-Rowell, Curtis, Doan, & Coe, 2015; Lee & Hicken, 2013; Wells, Evans, Beavis, & Ong, 2010). Similarly, limited research has examined the extent to which individuals from marginalized racial/ethnic groups, or disadvantaged socioeconomic backgrounds, are more susceptible to the adverse effects of ELA (differential vulnerability hypothesis). Focusing on the association between ELA and body mass index (BMI) in a young adult sample of Black/African American (AA) and White/European American (EA) college students, the current study sought to address these gaps in current knowledge.

The mediating role of ELA

Social conditions in the US are structured such that adverse experiences are unequally distributed among youth, with racial/ethnic minority and socioeconomically disadvantaged children being disproportionately exposed (Thoits, 2010; Wells et al., 2010), and thus at increased risk for the many adverse consequences (Felitti et al., 1998; Shonkoff et al., 2012). For example, AAs and individuals of low SES consistently score higher on measures of exposure to early life adverse events or high-risk contexts, such as living in a poor neighborhood, being exposed to family or neighborhood violence, having a parent with mental health problems, and experiencing childhood maltreatment (Chang, 2006; Felitti et al., 1998; Slopen et al., 2010; Thoits, 2010; Wells et al., 2010). In turn, such experiences and environments have been associated with obesity risk (Carroll-Scott et al., 2013; Danese & Tan, 2014; Garasky, Stewart, Gundersen, Lohman, & Eisenmann, 2009; Lohman, Stewart, Gundersen, Garasky, & Eisenmann, 2009; Nicholson & Browning, 2012). Despite existing evidence linking racial minority status and low SES to ELA, and ELA to poorer physical health, few studies have tested the mediating role of ELA in the development of group health disparities. Such tests are needed to demonstrate the degree to which early experiences may

lead to health inequities. Notably, ELA has, in fact, been shown to mediate the effects of poverty on BMI trajectories (Wells et al., 2010) and Black-White differences in obesity among females (Lee & Hicken, 2013) in adolescent samples.

Higher education is often seen as a way to escape disadvantaged social circumstances and their negative sequelae. Students from less advantaged backgrounds demonstrate resiliency by seeking higher education degrees (Schoon, Parsons, & Sacker, 2004). Interestingly, however, resiliency as indicated by academic success, may not extend to health status (T. F. Nelson, Gortmaker, Subramanian, Cheung, & Wechsler, 2007). In fact, evidence suggests that achievement in the face of adversity may even take a biological toll, contributing to racial and socioeconomic disparities (Chen, Miller, Brody, & Lei, 2014). Thus, the presence of socioeconomic and racial disparities in obesity among college students, and the role of ELA in accounting for these disparities are understudied topics relevant to forming policy and intervention.

Differential vulnerability

In addition to differences in exposure, vulnerability to the sequelae of ELA may also vary according to one's racial/ethnic group membership or socioeconomic position. In particular, early adverse experiences and disadvantaged social status may have multiplicative health-related effects (Slopen et al., 2010). Consistent with this idea, and especially relevant to high-achieving students, the "weathering hypothesis" postulates that when faced with chronic stressors and social exclusion, high-effort, active coping may be detrimental to physical health (Geronimus, 2001; Levine & Crimmins, 2014). Accordingly, emerging evidence suggests that AA youth who attain high academic competency or matriculate into college are more vulnerable to the effects of early adversity on physiologic dysregulation than their lower achieving AA peers (Brody et al., 2013; Chen et al., 2014). These findings suggest that the stress associated with making it to college for those from high adversity circumstances may carry physiological costs despite other benefits. Whether youth from disadvantaged socioeconomic backgrounds are also more vulnerable to ELA has not previously been considered; however, as is the case with AAs, youth of low SES are at elevated risk of being exposed to ELA (Thoits, 2010) and of being socially excluded (Fuller-Rowell, Evans, & Ong, 2012). Empirical investigation is therefore needed to consider whether background SES moderates the association between ELA and health. Additionally, although some evidence supports AAs' increased vulnerability to ELA (Slopen et al., 2010), to our knowledge, no studies have considered this hypothesis for BMI. Because of established group disparities in obesity as well as known associations between ELA and obesity, BMI is an important extension to prior research.

Hypotheses

The current study adds to existing literature by testing ELA as a mediator of racial and socioeconomic disparities in BMI, and by examining the moderating role of race and background SES on the association between ELA and BMI. We hypothesize that expected racial and socioeconomic differences in BMI among first and second year college students

will be mediated by their past exposure to ELA. Further, we predict that ELA will be more strongly associated with BMI for AAs and for those from less advantaged SES backgrounds.

Method

Data and sample

The data were derived from the College Student Health Study, an ongoing study of college students at a predominantly white university (3% African American, 4% Asian, 4% Hispanic/Latino, 1% American Indian) in the Midwestern United States. In 2012, 235 AA students and an equally sized stratified random sample of EA students were invited to participate. First-generation EA college students (neither parent graduated from college) were over-sampled to ensure that race and parent education would not be confounded. A total of 150 first and second year college students (mean age=18.8, SD=1.0; 45% AA, 57% female) participated in the study. The protocol was approved by the University's institutional review board and informed consent was obtained from all participants.

Measures

Demographic and control variables—Race was collected from university records and confirmed via self-report. Students also reported their sex, age, whether they currently smoke cigarettes (1 = yes, 0 = no), and the number of alcoholic drinks they consume “on a typical day when they are drinking”. A dichotomous variable indicating heavy alcohol use was included in analyses (≥ 5 alcoholic drinks = 1, <5 drinks = 0). Participants reported each parent's level of education on a seven-point scale ranging from 8th grade or less (coded as 1) to graduate degree (coded as 7), and parent education was coded as the mean education level for students from two parent families or the education level of a single parent. Income-to-needs ratio was calculated from youth reports of their parents' combined household income from the year prior to the baseline assessment, measured on a scale with 28 possible categories (1 = <\$5,000 to 28 = >\$200,000), and adjusted for household size, using 2011 US Federal poverty guidelines. A composite representing background socioeconomic status was scored as the mean of standardized parent education and household income-to-needs variables.

Early life adversity—The number of early life adversities experienced was measured using a 21-item inventory that was adapted for the current study from previous research (Felitti et al., 1998; Finkelhor D, Shattuck A, Turner H, & Hamby S, 2013). Twelve of the items were based on the Adverse Childhood Experiences Questionnaire (Felitti et al., 1998). Additional childhood risk factors with known influences on health were measured, including crime and violence exposure, family environment dynamics, and illness in the family (Finkelhor D et al., 2013; Garasky et al., 2009). The ELA inventory is not an exhaustive list of potential types of adversity, rather it is meant to index broadly one's exposure to early stressful events. Items measuring parental educational attainment and employment status were not included to avoid conceptual overlap with background socioeconomic status. Items were coded to indicate the occurrence of each risk factor (1=yes, 0=no) and a total scale score, which could range from 0–21, was calculated by summing all items ($\alpha=.81$).

Body size—Anthropometric measurements of weight, height, and hip circumference were collected by trained nursing staff. Body mass index (BMI) was calculated as weight (kg) divided by height (m)². Racial differences in BMI are limited because AAs, on average, have less abdominal visceral fat than EAs, which is not captured by the BMI (Carroll et al., 2008). To enhance our confidence in the findings, the body adiposity index (BAI), a direct estimate of percentage of body fat and an alternative method for between-race comparisons (Barreira, Harrington, Staiano, Heymsfield, & Katzmarzyk, 2011; Bergman et al., 2011), was also considered as a secondary outcome. BAI was calculated as hip circumference (cm) / height (m)^{1.5} – 18.

Statistical analyses

Univariate descriptive statistics and bivariate correlations were examined to consider unadjusted racial and socioeconomic disparities in key variables. A series of regression analyses were estimated using Mplus 7.11 (Muthén & Muthén, 2013) to test mediation and moderation hypotheses. Missing data were handled using Full Information Maximum Likelihood (FIML) estimation, which utilizes all available data to estimate model parameters and ensures a consistent sample size across models. Of the 150 individuals, 6 had missing data on background SES, 2 had missing data on ELA, 14 had missing data on heavy alcohol use, and all other variables had no missing data.

Initially, racial and socioeconomic differences in ELA and BMI were estimated in a multivariate regression model (Model 1). In addition to race and SES, BMI was adjusted for sex, age, current smoker status, and heavy alcohol use, and ELA was adjusted for sex. The full mediation model (Model 2) was then tested using path analysis, in which ELA was added as a predictor of BMI. Mediation was tested using the product of coefficients methods, and bias-corrected confidence intervals were estimated using a bootstrapping procedure (Hayes & Scharkow, 2013; MacKinnon, Fairchild, & Fritz, 2007). Lastly, to examine whether the association between ELA and BMI is conditioned by sociodemographic factors, ELA was interacted with race and SES and interaction terms were tested in a multiple regression model.

Results

Descriptive statistics for AAs and EAs are shown in Table 1. AA students had lower background SES ($M_{\text{difference}} = 0.32$, 95% CI [0.05, 0.60], $p = .023$) and reported having experienced greater ELA ($M_{\text{difference}} = 2.41$ events, 95% CI [1.35, 3.46], $p < .001$) than EAs. AAs also had higher BMI than did EAs ($M_{\text{difference}} = 1.77$ units, 95% CI [0.32, 3.21], $p = .017$). Bivariate correlations are presented in Table 2 and were in line with study hypotheses. In particular, background SES was inversely correlated with ELA ($p < .001$) and with BMI ($p = .029$). ELA was also correlated with BMI ($p < .001$). In addition, body adiposity index was strongly correlated with BMI and showed comparable associations to BMI with race, background SES, and ELA. Racial and socioeconomic differences in the likelihood of experiencing each adverse event were also tested. Odds ratios from logistic regression models and bivariate correlations between adverse events and BMI are presented in Table 3. Large racial differences in adverse events were detected for 8 of the 21 items with

AAs being more than 3 times as likely to experience each event, after controlling for SES. Moreover, differences by SES were detected for 6 of the 21 items, after controlling for race.

Results for racial and socioeconomic disparities in health and mediation hypotheses

First, we considered group differences in ELA and BMI controlling for sex, age, heavy alcohol use, and smoking status. AAs reported experiencing .60 *SD* units greater ELA ($B = 2.01, p < .001, 95\% \text{ CI } [1.02, 2.99]$) than did EAs. After adjustment for background SES and other covariates, AAs also had .41 *SD* units higher BMI ($B = 1.76, p = .014, 95\% \text{ CI } [0.36, 3.16]$) than EAs. Due to the racial disparity in BMI often being found only between females, a race by sex interaction term was also considered but it was not significant ($p = .58$) and was dropped from subsequent models. Adjusting for race and other covariates, background SES was significantly associated with ELA ($B = -1.15, p < .001, 95\% \text{ CI } [-1.74, -.55]$) and BMI ($B = -1.08, p = .018, 95\% \text{ CI } [-1.98, -.018]$). The magnitude of this effect suggests that one *SD* unit increase in background SES was associated with a .29 *SD* unit decrease in ELA and a .21 *SD* unit decrease in BMI. Because SES is often differentially associated with health outcomes between AAs and EAs (Fuller-Rowell et al., 2015), a race by background SES interaction was considered but was not found to be significant ($p = .79$).

Next, we tested the extent to which ELA explained (mediated) group differences in BMI, adjusting for all covariates in previous model (see Figure 1 for fitted path analysis). ELA was significantly associated with BMI ($B = 0.38, p < .001, 95\% \text{ CI } [.16, .61]$), accounting for 7% of the variance in BMI above all other covariates. Each one *SD* unit increase in ELA (3.3 events) was associated with a .30 *SD* unit increase in BMI. Estimated racial and socioeconomic differences in BMI were attenuated by 42% and 43% when accounting for ELA, respectively, and neither race ($p = .16$) nor income-to-needs ($p = .18$) remained significantly associated with BMI. Furthermore, significant indirect effects of race on BMI were detected via the pathway of ELA ($B = .77, 95\% \text{ CI } [.20, 1.97]$). Background SES was also associated with BMI through the pathway of ELA ($B = -.44, 95\% \text{ CI } [-1.08, -.010]$). In sum, results of the mediational analyses suggest that differential exposure to ELA is one pathway through which group disparities in BMI among college students develop.

Results for moderation hypotheses

We also tested whether the strength of the associations between ELA and BMI varied as a function of race or background SES. Results indicated that race significantly moderated the relation between ELA and BMI ($B = 0.45, p = .033, 95\% \text{ CI } [.04, .87]$), accounting for an additional 3% of the variance in BMI. Figure 2 demonstrates that ELA was more strongly associated with BMI for AAs compared to EAs. Specifically, no race difference was found for those who experienced low levels of ELA, yet a large racial gap existed for those coming from backgrounds high in adversity. Background SES was not a statistically significant moderator of the association between ELA and BMI ($p = .71$).

Supplementary analyses

Additional path models were fit using BAI (body adiposity index) rather than BMI to enhance our confidence in the results. Findings from these analyses were generally consistent with those presented for BMI. Significant indirect effects were detected for both

race ($B = 1.09$, 95% CI [.35, 2.52]) and background SES ($B = -.61$, 95% CI [-1.45, -.22]) as predictors of BAI through the pathway of ELA. Moderation findings were also consistent, such that the association between ELA and BAI was larger for AAs as compared to EAs ($p = .029$), and SES was not a significant moderator ($p = .37$).

Discussion

There is growing evidence for the long-term health consequences of ELA, with recent reports recommending action by Congress, pediatricians, and other health professionals aimed at reducing exposure to or the consequences of such adversity (Block et al., 2013; Garner et al., 2012; Shonkoff et al., 2012). Accordingly, the social patterning of ELA is a public health concern as well as a social justice issue. We demonstrated that even among college students, there are large racial and socioeconomic differences in exposure to early adverse events. Moreover, differences in early adversity account for a substantial portion of racial and socioeconomic disparities in BMI (and BAI). This finding is consistent with existing literature linking various early life adverse contexts to elevated BMI (Carroll-Scott et al., 2013; Suglia, Duarte, Chambers, & Boynton-Jarrett, 2012) and studies that have documented ELA as an explanation for group disparities in BMI (Lee & Hicken, 2013; Wells et al., 2010). These results extend this work to a college student population, showing that the sequelae of ELA are present among students despite the demonstrated resiliency of college enrollment.

Because weight gain is common during college, students are an important target demographic for obesity research (Vella-Zarb & Elgar, 2009, p. 5). Furthermore, university policies and regulations have influence on—and are able to change—the availability of healthy food choices and exercise opportunities as well as other health promoting resources (Horacek et al., 2013; Levitsky, Halbmaier, & Mrdjenovic, 2004). College populations are also relatively accessible for psychosocial and behavioral interventions due to shared spaces and course requirements (LaRose, Gorin, Clarke, & Wing, 2011). Educational contexts, and colleges in particular, are therefore critical settings in which to consider weight-related interventions (M. C. Nelson et al., 2008; Office of the Surgeon General, 2001).

Additionally, because many resources (e.g., campus recreation programs) are widely available to the college students of a particular campus, colleges have the potential to play an important role in achieving health equity by increasing access to health promoting resources. However, psychological stress—often a continuation or even proliferation of ELA (Nurius, Green, Logan-Greene, & Borja, 2015)—has also been shown to have an important influence on weight change and weight-related behaviors during this time (Freeman & Gil, 2004; Serlachius, Hamer, & Wardle, 2007), such that access to healthy environmental influences may not confer the same benefit to members of social groups experiencing disproportionate psychological stress. Accordingly, in order to reduce group disparities in obesity, effective weight-related interventions and policies, such as university courses on obesity and related health behaviors (Laska et al., 2012), may also benefit from incorporating or being supplemented by psychosocial or economic supports for disadvantaged students (e.g., mentoring programs or mindfulness-based interventions; O'Reilly, Cook, Spruijt-Metz, &

Black, 2014; Quintiliani & Whiteley, 2016). Research addressing the effectiveness of such multidimensional weight-related interventions is needed.

Our findings also support previous recommendations regarding effective screening of early adversity within the health care system in order to ameliorate its long-term consequences (Brcic, Eberdt, & Kaczorowski, 2011; Garner et al., 2012). For example, for patients who have experienced significant adversity, pediatricians and general practitioners may spend additional time probing about support networks and can make recommendations for counseling or other types of supportive interventions when warranted. Physicians at University medical clinics are especially well-positioned to be aware of university resources and programs.

The specific mechanisms linking ELA to obesity are beyond the scope of this paper but are an important topic of research. Current evidence suggests that the pathways of influence appear to be varied, including physiologic and behavioral processes. For example, ELA may alter epigenetic processes that govern inflammatory and stress physiology, leading to further physiologic dysregulation (Miller et al., 2009). In addition, the development of higher level cognitive structures and processes (i.e., executive functioning and self-regulatory skills) may be affected by ELA, which may influence health cumulatively across the life span through changes in health behaviors (Evans, Fuller-Rowell, & Doan, 2012). Moreover, in circumstances of high ELA, opportunities to develop and engage in healthy behaviors may be constrained. For example, high-crime, low-resource neighborhoods appear to be obesogenic due to poor access to healthy food, and fewer opportunities for physical recreation (Carroll-Scott et al., 2013). Indeed, differences in physical activity and television viewing have been shown to account for a portion of estimated black-white disparities in obesity (T. F. Nelson et al., 2007). Further research is needed to consider the primary mechanisms through which ELA affects obesity risk among college students, and in turn, whether intervening on these pathways leads to greater health equity.

This is one of the first studies to report on differential vulnerability to early life adversity as a function of race and SES. In particular, we show that the number of reported adverse events were more strongly related to the BMI of AA college students, as compared to EAs. Our results are congruent with past research showing increased stress vulnerability and accelerated aging among socially disadvantaged groups (Geronimus, 2001; Slopen et al., 2010; Tomiyama, Puterman, Epel, Rehkopf, & Laraia, 2013). For example, ELA has been shown to be associated with the proinflammatory physiology of AA adults, but not for EAs (Slopen et al., 2010). Differential vulnerability to the effects of *psychological stress* on BMI trajectories among AA and EA female adolescents has also been documented (Tomiyama et al., 2013). One potential explanation for AA adolescents' greater vulnerability to ELA may be that fewer resources exist during their youth, including social support systems and positive role models, likely due in part to racial biases as well as higher residence in at-risk communities (Noguera, 2003). As a result, achieving upward mobility in such contexts often requires overcoming several barriers (Hardaway & McLoyd, 2009). AA youth from high adversity backgrounds may therefore be required to engage in particularly high-effort, active coping during their youth in order to meet the demands associated with gaining entrance to a four-year university. Active coping (e.g., John Henryism) appears to be well suited for

attaining educational success, but comes with biological costs when also confronting early adversity (Brody et al., 2013; Chen et al., 2014). This research suggests that, in order to be optimally beneficial to physical health, academic achievement and upward mobility among AAs will require additional support during childhood and adolescence to offset exposure to ELA.

Contrary to our hypothesis, background SES did not moderate the association between ELA and BMI. One explanation for this null finding may be that the sample came from a “highly competitive plus” public university (~50% acceptance rate), according to Barron’s Profiles of American Colleges (2009), with relatively few students from families below the federal poverty line (7%). The stigma associated with disadvantaged social status may primarily be experienced by those from very poor households rather than on a SES gradient (Fuller-Rowell et al., 2012), thereby reducing our ability to detect the multiplicative effects of SES and ELA. Alternatively, the lack of moderation findings may be due to unmeasured adversity that is patterned by race to a greater degree than SES. For example, AA’s experience discrimination and live in areas of concentrated poverty at particularly high rates as compared to socioeconomically disadvantaged EAs (Chang, 2006; Fuller-Rowell et al., 2011; Kessler, Mickelson, & Williams, 1999; Lichter, Parisi, & Taquino, 2012). Additional research is needed to consider reasons for AA’s increased vulnerability to ELA, and whether very poor EA youth also show greater susceptibility.

Several limitations of the current study should be noted. First, although college students are an important and understudied demographic in disparities research, the findings on this particular population are limited by their generalizability to other groups. One important question for future research is whether non-college-bound AA youth are also vulnerable to the weight-related sequelae of ELA. Second, although the sample size was adequate based on the expected magnitude of associations, the number of participants and their recruitment from a single college campus are additional limitations. Larger and more representative samples of college students will increase the generalizability of findings and add additional possibilities to consider important moderators of risk (e.g., resilience factors). Third, because the study utilizes cross-sectional data, causality cannot be inferred and future research should utilize longitudinal data. However, while youth who are obese may experience greater adversity as a result of their weight, the majority of early adverse experiences assessed would likely not be affected by children’s body size, including each of the items for which a bivariate association with BMI was detected (see Table 3). A fourth limitation is that retrospective reports of ELA were used, leading to potential recall bias. However, adverse experiences measured are primarily objective events or major stressors that are less susceptible to recall bias (Hardt & Rutter, 2004), and research has shown retrospective and prospective reports of ELA to be comparably predictive of health outcomes (Patten et al., 2014).

In sum, the findings of this study demonstrate that racial and socioeconomic disparities in body size among college students are partly accounted for by differential exposure to ELA, and that, as compared to EAs, AAs are more vulnerable to ELA. These results support recent recommendations to incorporate assessment of early adverse experiences into pediatrician visits and public health initiatives. Both reducing exposure to ELA as well as

providing additional social and community supports to mitigate the deleterious consequences of ELA will likely be important steps to achieving health equity.

Acknowledgments

Funding: Support was provided by the Robert Wood Johnson Health and Society Scholars Program at the University of Wisconsin—Madison, and by the Clinical and Translational Science Award (CTSA) program, through the NIH National Center for Advancing Translational Sciences (NCATS), grant UL1TR000427.

The authors declare that they have no conflict of interest. Support for this project was provided by the Robert Wood Johnson Health and Society Scholars Program at the University of Wisconsin—Madison, and by the Clinical and Translational Science Award (CTSA) program, through the NIH National Center for Advancing Translational Sciences (NCATS), grant UL1TR000427. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

References

- Barreira TV, Harrington DM, Staiano AE, Heymsfield SB, & Katzmarzyk PT (2011). Body Adiposity Index, Body Mass Index and body fat in White and Black adults. *Journal of the American Medical Association*, 306(8), 828–830. 10.1001/jama.2011.1189 [PubMed: 21862743]
- Bergman RN, Stefanovski D, Buchanan TA, Sumner AE, Reynolds JC, Sebring NG, ... Watanabe RM (2011). A better index of body adiposity. *Obesity*, 19(5), 1083–1089. 10.1038/oby.2011.38 [PubMed: 21372804]
- Block RW, Dreyer BP, Cohen AR, Stapleton FB, Furth SL, & Bucciarelli RL (2013). An agenda for children for the 113th Congress: Recommendations from the Pediatric Academic Societies. *Pediatrics*, 131(1), 109–119. 10.1542/peds.2012-2661 [PubMed: 23277316]
- Brcic V, Eberdt C, & Kaczorowski J (2011). Development of a tool to identify poverty in a family practice setting: a pilot study. *International Journal of Family Medicine*, 2011. Retrieved from <http://www.hindawi.com/journals/ijfm/2011/812182/abs/>
- Brody GH, Yu T, Chen E, Miller GE, Kogan SM, & Beach SRH (2013). Is resilience only skin deep? Rural African Americans' socioeconomic status–related risk and competence in preadolescence and psychological adjustment and allostatic load at age 19. *Psychological Science*, 24, 1285–1293. 10.1177/0956797612471954 [PubMed: 23722980]
- Carroll JF, Chiapa AL, Rodriquez M, Phelps DR, Cardarelli KM, Vishwanatha JK, ... Cardarelli R (2008). Visceral fat, waist circumference, and BMI: Impact of race/ethnicity. *Obesity*, 16(3), 600–607. [PubMed: 18239557]
- Carroll-Scott A, Gilstad-Hayden K, Rosenthal L, Peters SM, McCaslin C, Joyce R, & Ickovics JR (2013). Disentangling neighborhood contextual associations with child body mass index, diet, and physical activity: The role of built, socioeconomic, and social environments. *Social Science & Medicine*, 95, 106–114. 10.1016/j.socscimed.2013.04.003 [PubMed: 23642646]
- Chang VW (2006). Racial residential segregation and weight status among US adults. *Social Science & Medicine*, 63(5), 1289–1303. 10.1016/j.socscimed.2006.03.049 [PubMed: 16707199]
- Chen E, Miller GE, Brody GH, & Lei M (2014). Neighborhood poverty, college attendance, and diverging profiles of substance use and allostatic load in rural African American youth. *Clinical Psychological Science*, 2167702614546639 10.1177/2167702614546639
- Danese A, & Tan M (2014). Childhood maltreatment and obesity: Systematic review and meta-analysis. *Molecular Psychiatry*, 19(5), 544–554. 10.1038/mp.2013.54 [PubMed: 23689533]
- Evans GW, Fuller-Rowell TE, & Doan SN (2012). Childhood cumulative risk and obesity: The mediating role of self-regulatory ability. *Pediatrics*, 129(1), e68–e73. 10.1542/peds.2010-3647 [PubMed: 22144695]
- Felitti MD, Vincent J, Anda MD, Robert F, Nordenberg MD, Williamson MS, ... others. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14(4), 245–258. [PubMed: 9635069]

- Finkelhor D, Shattuck A, Turner H, & Hamby S. (2013). Improving the Adverse Childhood Experiences study scale. *JAMA Pediatrics*, 167(1), 70–75. 10.1001/jamapediatrics.2013.420 [PubMed: 23403625]
- Frederick CB, Snellman K, & Putnam RD (2014). Increasing socioeconomic disparities in adolescent obesity. *Proceedings of the National Academy of Sciences*, 111(4), 1338–1342. 10.1073/pnas.1321355110
- Freeman LMY, & Gil KM (2004). Daily stress, coping, and dietary restraint in binge eating. *International Journal of Eating Disorders*, 36(2), 204–212. 10.1002/eat.20012 [PubMed: 15282690]
- Fuller-Rowell TE, Cogburn CD, Brodish AB, Peck SC, Malanchuk O, & Eccles JS (2011). Racial discrimination and substance use: Longitudinal associations and identity moderators. *Journal of Behavioral Medicine*, 35, 581–590. 10.1007/s10865-011-9388-7 [PubMed: 22113318]
- Fuller-Rowell TE, Curtis DS, Doan SN, & Coe CL (2015). Racial disparities in the health benefits of educational attainment: A study of inflammatory trajectories among African American and white adults. *Psychosomatic Medicine*, 77(1), 33–40. 10.1097/PSY.0000000000000128 [PubMed: 25490696]
- Fuller-Rowell TE, Evans GW, & Ong AD (2012). Poverty and health: The mediating role of perceived discrimination. *Psychological Science*, 23(7), 734–739. 10.1177/0956797612439720 [PubMed: 22700331]
- Garasky S, Stewart SD, Gundersen C, Lohman BJ, & Eisenmann JC (2009). Family stressors and child obesity. *Social Science Research*, 38(4), 755–766. [PubMed: 20645439]
- Garner AS, Shonkoff JP, Siegel BS, Dobbins MI, Earls MF, McGuinn L, ... Wood DL (2012). Early childhood adversity, toxic stress, and the role of the pediatrician: Translating developmental science into lifelong health. *Pediatrics*, 129(1), e224–e231. [PubMed: 22201148]
- Geronimus AT (2001). Understanding and eliminating racial inequalities in women's health in the United States: The role of the weathering conceptual framework. *Journal of the American Medical Women's Association*, 56(4), 133–136, 149–150.
- Hardaway CR, & McLoyd VC (2009). Escaping poverty and securing middle class status: How race and socioeconomic status shape mobility prospects for African Americans during the transition to adulthood. *Journal of Youth and Adolescence*, 38(2), 242–256. [PubMed: 19636721]
- Hardt J, & Rutter M (2004). Validity of adult retrospective reports of adverse childhood experiences: Review of the evidence. *Journal of Child Psychology and Psychiatry*, 45(2), 260–273. 10.1111/j.1469-7610.2004.00218.x [PubMed: 14982240]
- Hayes AF, & Scharkow M (2013). The relative trustworthiness of inferential tests of the indirect effect in statistical mediation analysis: Does method really matter? *Psychological Science*, 24(10), 1918–1927. 10.1177/0956797613480187 [PubMed: 23955356]
- Horacek TM, Erdman MB, Byrd-Bredbenner C, Carey G, Colby SM, Greene GW, ... others. (2013). Assessment of the dining environment on and near the campuses of fifteen post-secondary institutions. *Public Health Nutrition*, 16(7), 1186–1196. [PubMed: 23174458]
- Juonala M, Magnussen CG, Berenson GS, Venn A, Burns TL, Sabin MA, ... others. (2011). Childhood adiposity, adult adiposity, and cardiovascular risk factors. *New England Journal of Medicine*, 365(20), 1876–1885. [PubMed: 22087679]
- Kessler RC, Mickelson KD, & Williams DR (1999). The prevalence, distribution, and mental health correlates of perceived discrimination in the United States. *Journal of Health and Social Behavior*, 40(3), 208–230. [PubMed: 10513145]
- LaRose JG, Gorin AA, Clarke MM, & Wing RR (2011). Beliefs about weight gain among young adults: Potential challenges to prevention. *Obesity*, 19(9), 1901–1904. 10.1038/oby.2011.203 [PubMed: 21738239]
- Laska MN, Pelletier JE, Larson NI, & Story M (2012). Interventions for weight gain prevention during the transition to young adulthood: A review of the literature. *Journal of Adolescent Health*, 50(4), 324–333. 10.1016/j.jadohealth.2012.01.016 [PubMed: 22443834]
- Lee H, & Hicken MT (2013). Cumulative social risk and racial/ethnic disparities in obesity during the transition to adulthood. *Journal of Health Care for the Poor and Underserved*, 24(2), 907–927. [PubMed: 23728056]

- Levine ME, & Crimmins EM (2014). Evidence of accelerated aging among African Americans and its implications for mortality. *Social Science & Medicine*, 118, 27–32. 10.1016/j.socscimed.2014.07.022 [PubMed: 25086423]
- Levitsky DA, Halbmaier CA, & Mrdjenovic G (2004). The freshman weight gain: A model for the study of the epidemic of obesity. *International Journal of Obesity*, 28, 1435–1442. [PubMed: 15365585]
- Lichter DT, Parisi D, & Taquino MC (2012). The geography of exclusion: Race, segregation, and concentrated poverty. *Social Problems*, 59(3), 364–388.
- Lohman BJ, Stewart S, Gundersen C, Garasky S, & Eisenmann JC (2009). Adolescent overweight and obesity: Links to food insecurity and individual, maternal, and family stressors. *Journal of Adolescent Health*, 45(3), 230–237. 10.1016/j.jadohealth.2009.01.003 [PubMed: 19699418]
- MacKinnon DP, Fairchild AJ, & Fritz MS (2007). Mediation analysis. *Annual Review of Psychology*, 58, 593 10.1146/annurev.psych.58.110405.085542
- Miller GE, Chen E, Fok AK, Walker H, Lim A, Nicholls EF, ... Kobor MS (2009). Low early-life social class leaves a biological residue manifested by decreased glucocorticoid and increased proinflammatory signaling. *Proceedings of the National Academy of Sciences*, 106(34), 14716–14721. 10.1073/pnas.0902971106
- Muthén LK, & Muthén BO (2013). *Mplus: Statistical analysis with latent variables User's guide* (Version 7.11). Los Angeles, CA: Muthen and Muthen.
- Nelson MC, Story M, Larson NI, Neumark-Sztainer D, & Lytle LA (2008). Emerging adulthood and college-aged youth: An overlooked age for weight-related behavior change. *Obesity*, 16(10), 2205–2211. 10.1038/oby.2008.365 [PubMed: 18719665]
- Nelson TF, Gortmaker SL, Subramanian SV, Cheung L, & Wechsler H (2007). Disparities in overweight and obesity among US college students. *American Journal of Health Behavior*, 31(4), 363–373. 10.5993/AJHB.31.4.3 [PubMed: 17511571]
- Nicholson LM, & Browning CR (2012). Racial and ethnic disparities in obesity during the transition to adulthood: The contingent and nonlinear impact of neighborhood disadvantage. *Journal of Youth and Adolescence*, 41(1), 53–66. [PubMed: 21706260]
- Noguera PA (2003). The trouble with Black boys: The role and influence of environmental and cultural factors on the academic performance of African American males. *Urban Education*, 38(4), 431–459. 10.1177/0042085903038004005
- Nurius PS, Green S, Logan-Greene P, & Borja S (2015). Life course pathways of adverse childhood experiences toward adult psychological well-being: A stress process analysis. *Child Abuse & Neglect*, 45, 143–153. 10.1016/j.chiabu.2015.03.008 [PubMed: 25846195]
- Office of the Surgeon General. (2001). *The Surgeon General's call to action to prevent and decrease overweight and obesity*. Rockville, MD: US Department of Health and Human Services.
- Ogden CL, Carroll MD, Kit BK, & Flegal KM. (2012). Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *Journal of the American Medical Association*, 307(5), 483–490. 10.1001/jama.2012.40 [PubMed: 22253364]
- O'Reilly GA, Cook L, Spruijt-Metz D, & Black DS (2014). Mindfulness-based interventions for obesity-related eating behaviors: A literature review. *Obesity Reviews*, 15(6), 453–461. 10.1111/obr.12156 [PubMed: 24636206]
- Patten SB, Wilkes TCR, Williams JVA, Lavorato DH, El-Guebaly N, Schopflocher D, ... Bulloch AGM (2014). Retrospective and prospectively assessed childhood adversity in association with major depression, alcohol consumption and painful conditions. *Epidemiology and Psychiatric Sciences*, 1–8.
- Quintiliani LM, & Whiteley JA (2016). Results of a nutrition and physical activity peer counseling intervention among nontraditional college students. *Journal of Cancer Education*, 31(2), 366–374. [PubMed: 25994357]
- Schoon I, Parsons S, & Sacker A (2004). Socioeconomic adversity, educational resilience, and subsequent levels of adult adaptation. *Journal of Adolescent Research*, 19(4), 383–404. 10.1177/0743558403258856

- Serlachius A, Hamer M, & Wardle J (2007). Stress and weight change in university students in the United Kingdom. *Physiology & Behavior*, 92(4), 548–553. 10.1016/j.physbeh.2007.04.032 [PubMed: 17537466]
- Shonkoff JP, Garner AS, Siegel BS, Dobbins MI, Earls MF, Garner AS, ... Wood DL (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, peds.2011–2663. 10.1542/peds.2011-2663
- Slopen N, Lewis TT, Gruenewald TL, Mujahid MS, Ryff CD, Albert MA, & Williams DR (2010). Early life adversity and inflammation in African Americans and Whites in the Midlife in the United States Survey. *Psychosomatic Medicine*, 72(7), 694–701. 10.1097/PSY.0b013e3181e9c16f [PubMed: 20595419]
- Suglia SF, Duarte CS, Chambers EC, & Boynton-Jarrett R (2012). Cumulative social risk and obesity in early childhood. *Pediatrics*, 129(5), e1173–e1179. [PubMed: 22508921]
- Thoits PA (2010). Stress and health: Major findings and policy implications. *Journal of Health and Social Behavior*, 51(1 suppl), S41–S53. 10.1177/0022146510383499 [PubMed: 20943582]
- Tirosh A, Shai I, Afek A, Dubnov-Raz G, Ayalon N, Gordon B, ... Rudich A (2011). Adolescent BMI trajectory and risk of diabetes versus coronary disease. *New England Journal of Medicine*, 364(14), 1315–1325. 10.1056/NEJMoa1006992 [PubMed: 21470009]
- Tomiyama AJ, Puterman E, Epel ES, Rehkopf DH, & Laraia BA (2013). Chronic psychological stress and racial disparities in body mass index: Change between Black and White girls aged 10–19. *Annals of Behavioral Medicine*, 45(1), 3–12. 10.1007/s12160-012-9398-x [PubMed: 22993022]
- Vella-Zarb RA, & Elgar FJ (2009). The “freshman 5”: A meta-analysis of weight gain in the freshman year of college. *Journal of American College Health*, 58(2), 161–166. [PubMed: 19892653]
- Wells NM, Evans GW, Beavis A, & Ong AD (2010). Early childhood poverty, cumulative risk exposure, and body mass index trajectories through young adulthood. *American Journal of Public Health*, 100(12), 2507–2512. 10.2105/AJPH [PubMed: 20966374]
- Wong MD, Shapiro MF, Boscardin WJ, & Ettner SL (2002). Contribution of major diseases to disparities in mortality. *New England Journal of Medicine*, 347(20), 1585–1592. [PubMed: 12432046]

Figure 1.

Fitted mediation model ($N = 150$).

Note. Solid lines indicate significant paths. ELA and BMI are abbreviations for early life adversity and body mass index. Race is coded as European American=0, African American=1. Standardized coefficients and residual variances are presented. In order to demonstrate mediation, parameter estimates without ELA as a covariate are included below arrows. Controls include sex, age, heavy alcohol use, and current cigarette smoker. * $p < .05$. *** $p < .001$.

Figure 2. Estimated body mass index as a function of early life adversity for African American ($n = 68$) and European American ($n = 82$) college students.

Table 1.

Descriptives for African American (AA; $n = 68$) and European American (EA; $n = 82$) first and second year college students.

Variables	AA	EA	Race Difference
			<i>p</i>
Sex (female)	41 (60.3%)	43 (52.4%)	.34
Age (yrs.)	18.90 $\pm$ 1.01	18.74 $\pm$ 0.91	.33
Current cigarette smoker	5 (7.4%)	5 (6.1%)	.76
Heavy drinker (≥ 5 drinks)	17 (25.0%)	41 (50.0%)	.002
Socioeconomic status	-0.18 $\pm$ 0.86	0.14 $\pm$ 0.81	.022
Early life adversity	4.49 $\pm$ 3.66	2.09 $\pm$ 2.62	<.001
Body mass index (kg/m ²)	25.31 $\pm$ 5.36	23.54 $\pm$ 3.00	.017
Body adiposity index ^a	23.88 $\pm$ 7.19	21.44 $\pm$ 4.37	.016

Note. Values are expressed as mean $\pm$ SD or number of participants (%).

^aBody adiposity index is an estimate of percent body fat, calculated as hip circumference (cm)/height (m)^{1.5} - 18.

Table 2.

Bivariate correlations.

	1.	2.	3.	4.	5.	6.	7.	8.
1. Race	-							
2. Sex	.08	-						
3. Age	.08	-.22**	-					
4. Cigarette smoker	.03	-.03	.22**	-				
5. Heavy drinker	-.22**	-.22*	.10	.23**	-			
6. Socioeconomic status	-.19*	.01	-.20*	.04	.16*	-		
7. Early life adversity	.36***	.08	.16	.21**	-.10	-.32***	-	
8. Body mass index	.21*	.06	.02	-.05	-.06	-.18*	.34***	-
9. Body adiposity index	.21*	.43***	-.01	-.07	-.19*	-.18*	.36***	.62***

Note. Race is coded as European American=0, African American=1. Sex is coded as Male=0, Female=1.

* $p < .05$.

** $p < .01$.

*** $p < .001$.

Table 3.

Frequency of event occurrence for Early Life Adversity items for African American (AA) and European American (EA) college students, as well as racial and socioeconomic (SES) differences in likelihood of experiencing and predictive validity of adverse events.

Items	Race estimate ^a		SES estimate ^b		Predictive validity ^c	
	AA (%)	EA (%)	OR	OR	r with BMI	
Lived with someone who was alcoholic	14.9	9.9	0.85	0.32**		.15
Lived with someone who used street drugs	20.9	8.6	2.54	0.88		-.03
Lived with someone who was depressed	19.4	18.5	0.92	0.80		.16
Lived with someone who attempted suicide	1.5	6.2	0.25	0.88		-.10
Parents argued a lot	34.3	22.2	1.45	0.50**		.20*
Physical altercations between parents	10.4	4.9	1.90	0.67		.02
Reason to worry about parent's safety	25.4	6.2	3.63*	0.29**		.18*
You were sexually abused	9.0	3.7	1.40	0.39~		.31***
Parent verbally abused you	26.9	29.6	0.68	0.62*		.10
Parent physically threatened you	16.4	7.4	2.88	1.08		.08
Parent physically abused you	31.3	12.3	3.09*	0.80		.11
Parent injured you	10.4	6.3	1.77	1.22		.10
Parent spanked you with object	62.7	21.0	6.04***	0.44**		.16
Required to take care of younger family member	22.4	16.0	0.99	0.57		.24**
Immediate family member had illness	37.3	13.6	3.93**	0.98		.28**
Immediate family member died	23.9	3.7	7.09**	0.66		.21**
Immediate family member was shot, stabbed, or raped	6.0	0.0	N/A	0.62		.19*
Immediate family member was attacked or robbed	14.9	3.7	4.65*	0.67		.20*
Immediate family member went to prison	22.7	3.7	5.06*	0.54		.25**
Lived in a high crime neighborhood	23.9	4.9	4.75*	0.37**		.22**
House was broken into	14.9	6.2	1.98	0.57		.27**

Note.

^{a,b} Odds ratios are derived from logistic regression models using individual adverse events as outcomes and both race and SES as predictors.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

c Predictive validity of early life adversity items demonstrated using bivariate correlations with body mass index (BMI).

* $p < .05$.
** $p < .01$.
*** $p < .001$.