Weight stigma and physical health: an unconsidered ‘obesity’ cost

Letter to the Editor: Response to Singh et al. (2018). Is there more to the equation? Weight bias and the costs of obesity. *Canadian Journal of Public Health*. <https://doi.org/10.17269/s41997-018-0146-2>. Online first: 26 October 2018

Angela Meadows¹

Received: 7 November 2018 / Accepted: 8 May 2019 / Published online: 28 May 2019
© The Canadian Public Health Association 2019

Singh et al. (2018) outline the role that weight-related stigma plays in inflating the supposed economic costs of obesity. However, despite their thorough and helpful synthesis of the literature, one cost that was not well elucidated in their model was the direct impact of weight stigma on physical morbidity. In addition to indirect effects via delayed or inappropriate medical treatment, exposure to a chronic stigmatizing environment, even in the absence of personal discriminatory experiences, has a direct impact on physical health via the body’s dynamic neuroendocrine response to stress—allostasis. Allostasis maintains homeostasis in the face of physical, environmental or psychosocial stressors and involves multiple systems within the body, including the cardiovascular and immune systems. While allostasis represents an adaptive response to acute stress, chronic or repeated stress over time can result in multi-system biological dysregulation—a concept known as allostatic load, which is linked with a wide range of disease conditions, including cardiovascular disease, diabetes, hypertension and cancer (Beckie 2012). This situation is not specific to high-weight individuals. The wider stigma literature provides a wealth of evidence suggesting that exposure to personal and systemic prejudice and ill treatment places members of disadvantaged and marginalized groups at particular risk of developing these conditions (Hatzenbuehler et al. 2013; Pascoe and Smart Richman 2009). Hostile environments may also drive adverse physical health outcomes, for example, chronic pain conditions and poorer birth outcomes, via psychological distress (Brown et al. 2018; Lauderdale 2006).

In the case of high-weight individuals, experienced weight stigma has been linked with higher levels of cortisol and inflammatory markers (Jackson et al. 2016; Sutin et al. 2014) and impaired glucose metabolism (Tsenkova et al. 2011), with a recent longitudinal study supporting a causal relationship: in a cohort of 986 adults age 25–75 years in the Midlife Development in the US Biomarker Sub-study, perceived weight discrimination was associated with a twofold increase in the 10-year risk of high allostatic load (Vadiveloo and Mattei 2017). Experienced weight stigma and weight-related bullying were also linked with overt disease prevalence, including arteriosclerosis, diabetes and minor cardiac conditions, in a sample of over 21,000 high-weight adults in the National Survey of Alcohol and Related Conditions study (Udo et al. 2016). As many of these health problems are generally attributed to carrying excess weight, any calculation of the economic burden of ‘obesity’ must consider the direct causal role played by weight stigma in healthcare costs associated with physical as well as psychological morbidity.

Acknowledgements The support of the Economic and Social Research Council (UK) is gratefully acknowledged.

References

- Beckie, T. M. (2012). A systematic review of allostatic load, health, and health disparities. *Biological Research for Nursing*, 14(4), 311–346. <https://doi.org/10.1177/1099800412455688>.
- Brown, T. T., Partanen, J., Chuong, L., Villaverde, V., Chantal Griffin, A., & Mendelson, A. (2018). Discrimination hurts: the effect of discrimination on the development of chronic pain. *Social Science & Medicine*, 204, 1–8. <https://doi.org/10.1016/j.socscimed.2018.03.015>.
- Hatzenbuehler, M. L., Phelan, J. C., & Link, B. G. (2013). Stigma as a fundamental cause of population health inequalities. *American Journal of Public Health*, 103(5), 813–821. <https://doi.org/10.2105/AJPH.2012.301069>.
- Jackson, S. E., Kirschbaum, C., & Steptoe, A. (2016). Perceived weight discrimination and chronic biochemical stress: a population-based

✉ Angela Meadows
drameadows@gmail.com

¹ School of Psychology, University of Exeter, Exeter EX4 4QG, UK

- study using cortisol in scalp hair. *Obesity*, 24(12), 2515–2521. <https://doi.org/10.1002/oby.21657>.
- Lauderdale, D. S. (2006). Birth outcomes for Arabic-named women in California before and after September 11. *Demography*, 43(1), 185–201.
- Pascoe, E. A., & Smart Richman, L. (2009). Perceived discrimination and health: a meta-analytic review. *Psychological Bulletin*, 135(4), 531–554. <https://doi.org/10.1037/a0016059>.
- Singh, K., Russell-Mayhew, S., von Ranson, K., & McLaren, L. (2018). Is there more to the equation? Weight bias and the costs of obesity. *Canadian Journal of Public Health*. <https://doi.org/10.17269/s41997-018-0146-2>.
- Sutin, A. R., Stephan, Y., Luchetti, M., & Terracciano, A. (2014). Perceived weight discrimination and C-reactive protein. *Obesity*, 22(9), 1959–1961. <https://doi.org/10.1002/oby.20789>.
- Tsenkova, V. K., Carr, D., Schoeller, D. A., & Ryff, C. D. (2011). Perceived weight discrimination amplifies the link between central adiposity and nondiabetic glycemic control (HbA1c). *Annals of Behavioral Medicine*, 41(2), 243–251. <https://doi.org/10.1007/s12160-010-9238-9>.
- Udo, T., Purcell, K., & Grilo, C. M. (2016). Perceived weight discrimination and chronic medical conditions in adults with overweight and obesity. *International Journal of Clinical Practice*, 70(12), 1003–1011. <https://doi.org/10.1111/ijcp.12902>.
- Vadiveloo, M., & Mattei, J. (2017). Perceived weight discrimination and 10-year risk of allostatic load among US adults. *Annals of Behavioral Medicine*, 51(1), 94–104. <https://doi.org/10.1007/s12160-016-9831-7>.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.