Food insecurity and nutritional biomarkers in relation to stature in Inuit children from Nunavik

Catherine M. Pirkle, PhD,¹ Michel Lucas, PhD,^{1,2} Renée Dallaire, PhD,¹ Pierre Ayotte, PhD,^{1,2}
Joseph L. Jacobson, PhD,³ Sandra W. Jacobson, PhD,³ Eric Dewailly*, PhD,¹ Gina Muckle, PhD^{1,4}

ABSTRACT

OBJECTIVES: Inuit in Canada experience alarming levels of food insecurity, but nutritional and physiological consequences are poorly documented, especially in school-age children. The objective of this study was to assess the relation of food insecurity to iron deficiency and stature in school-aged Inuit children from Nunavik (Northern Quebec).

METHODS: Food insecurity, iron deficiency, and stature were assessed in a cohort of children. Food insecurity was determined by interviewing the children's mothers. Multiple logistic regression was used to evaluate the association of food insecurity to iron deficiency and short stature. We defined short stature as a height in the lowest tertile for age and sex, based on Canadian growth charts. The relation of food insecurity to height (cm) was analyzed with a general linear model. Statistical models controlled for age, sex, normal/overweight/obese status, prenatal lead exposure and postnatal polychlorinated biphenyls exposure.

RESULTS: Half of the children (49.7%, $n=145$) were food insecure, while one third were iron depleted, 12.6% had anaemia, and 8.7% had iron-deficiency anaemia. The multivariate odds ratio of anaemia was 1.82 (95% CI: 0.97, 3.42, $p=0.06$) for food-insecure children. Prevalence of short stature was 18.7%. Food-insecure children were an average of 2 cm shorter (95% CI: -0.48, -3.17) than food-secure children ($p<0.01$).

CONCLUSION: In this population, food-insecure children have greater burdens of nutritional deficiencies and slower linear growth. Considering the high prevalence of food insecurity among Inuit children in Nunavik, nutritional deficiencies and adverse effects on development should be carefully monitored.

KEY TERMS: Food insecurity; nutritional deficiencies; iron; growth; children; indigenous health

La traduction du résumé se trouve à la fin de l'article.

Can J Public Health 2014;105(4):e233-e238.

Food insecurity attains alarming levels in Inuit populations of northern Canada.¹ Measures of food insecurity traditionally encompass concepts of availability and access to sufficient food, as well as food variety and the ability to eat in a socially acceptable manner.² Among Inuit from Nunavut, Nanatsiavut, and the Inuvialuit settlement, 63% of households were food insecure and 27% were considered severely insecure in the 12 months preceding a survey of the region conducted in 2007-8.¹ The 2004 Nunavik Inuit Health Survey documented that a quarter of Nunavik households lacked food in the month preceding the survey.³ In contrast, approximately 12% of Canadian households experienced some food insecurity during the year of 2011. Throughout Canada, food insecurity is highest in households with children under the age of 18; on average, 17% of households with children are food insecure.⁴

The nutritional and physiological consequences of food insecurity are poorly documented in the Canadian Arctic, especially in children. Elsewhere, studies observed significant associations between food insecurity and nutritional deficiencies, particularly iron deficiency and anaemia.⁵⁻¹⁰ Such a relation is intuitive, because as defined, food insecurity entails insufficient and lower quality foods. According to one study of the non-Indigenous Canadian population, food insecurity in adults and adolescents correlated with inadequate nutrient intake. However, these authors failed to observe a similar relation in children.¹¹ In the United States (US), food insecurity was associated with three

times the odds of iron deficiency in children aged 3 to 19.⁶ Other studies from the US have found associations between food insecurity and iron-deficiency anaemia in toddlers.^{8,10} Among Canadian Inuit men, food insecurity was associated with 2.5 fold greater odds of low or depleted iron stores.⁷ With scant exception,¹² few studies have investigated the nutritional consequences of food insecurity among Inuit children.

Inadequate nutrition during childhood can lead to growth faltering and in the extreme, to stunting. In a recent meta-analysis of children five years and younger, micronutrient supplementation – with vitamin A, iron and zinc – improved linear growth,¹³ which suggests that certain micronutrients affect

Author Affiliations

1. Population Health & Optimal Health Practices Research Unit, CHU de Québec Research Centre, Québec City, QC
2. Department of Social & Preventive Medicine, Université Laval, Québec City, QC
3. Department of Psychiatry and Behavioral Neurosciences, Wayne State University School of Medicine, Detroit, MI
4. School of Psychology, Université Laval, Québec City, QC

Correspondence: Catherine McLean Pirkle, Axe Santé publique et pratiques optimales en santé, Centre de recherche du CHU de Québec, 2875, boulevard Laurier, Édifice Delta II, Bureau 600, 6e étage, Québec, QC G1V 2M2, Tel: 418-525-4444, E-mail: catherine-mclean.pirkle.1@ulaval.ca

Financial Support: This study was supported by grants from the National Institute of Environmental Health Sciences/NIH (R01-ES007902 to JLJ); the Northern Contaminants Program, Indian and Northern Affairs Canada (to GM); and the Joseph Young, Sr., Fund from the State of Michigan (to SWJ).

Conflict Of Interest: None to declare.

* Dr. Éric Dewailly passed away while this manuscript was being revised. He was an authority on environmental and human health in the circumpolar world, an exceptional mentor, and a brilliant mind.

Table 1. Number and (%) affirmative (or yes) responses of caregiver participants to the questions about household food security in preceding month (N=292)

Food security questions	Affirmative responses
Not enough food eaten by family	114 (39.0%)
Cut down the size of adult meals	116 (39.7%)
Cut down the size of children's meals	78 (26.7%)
Number of affirmative responses to above questions	
3	62 (21.2%)
2	101 (34.6%)
1	145 (49.7%)

growth trajectories. Because of the increased demand for iron during periods of rapid growth, iron deficiency may slow growth in children;¹⁴ although, the results are mixed for the dozens of randomized controlled trials that have assessed the effects of iron supplementation on child height.¹³ Household food insecurity has been shown to substantially retard growth in young children from Pakistan, Tanzania, Brazil, South Africa, Peru, Bangladesh, Nepal and India.^{15,16} In another study of children aged 24 months and less in Bangladesh, food insecurity was strongly associated with lower length for age scores.¹⁷ There was a distinct gradient from extremely to never food-insecure groups, in which the smallest children came from the extremely food-insecure households.¹⁷ Here, we assess the relation of food insecurity to iron status and stature in school-age children from Nunavik.

METHODS

Study population

Participants were Inuit children from Nunavik, a region located north of the 55th parallel in Quebec and about 1500 km from Montreal. Children were recruited before birth when their mothers participated in one of two cohort studies: the Cord Blood Monitoring Program (1993-1998) or the Environmental Contaminants and Infant Development Study (1996-2000). Both of these studies collected biological samples of contaminants and nutrients from umbilical cord blood samples (30 mL). Recruitment details for these studies have been provided elsewhere.¹⁸⁻²⁰ The current study, called the Nunavik Child Development Study (NCDS), is a 10-year follow-up of school-age children from the previous two cohorts. Eligibility criteria for the NCDS included: between 8 and 15 years of age, birth weight of ≥ 2.5 kg, gestational duration of ≥ 35 weeks, and no major birth defects or neurological or pervasive health disorders. Two hundred and ninety-four children participated in the study.

Study procedures

Between September 2005 and February 2010, the mothers of the children completed a detailed socio-demographic questionnaire. Each child provided a 20 mL blood sample for biological analyses. Blood samples were frozen in Nunavik at -80°C, transported by plane to the *Centre de toxicologie du Québec*, and sent on dry ice to other laboratories when necessary. In addition, our trained research nurses measured the child's height and weight. The research nurses conducted assessments in the three largest Nunavik villages. Participants who resided in other communities were transported by plane to one of the larger villages for data collection. Written informed consent was

Table 2. Characteristics of the study population according to food security status (N=292)

	Food		p-value†
	Secure (n=147)	Insecure (n=145)	
Age, mean (SD), yr	11.3 (0.8)	11.3 (0.8)	0.90
Girls, n (%)	78 (53.1)	69 (47.6)	0.35
Weight, mean (SD), kg	41.1 (10.4)	39.1 (9.4)	0.09
BMI status, n (%)			0.53
Normal	106 (72.1)	109 (75.2)	
Overweight	31 (21.1)	28 (19.3)	
Obese	10 (6.8)	8 (5.5)	
Height, mean (SD), cm	142.6 (7.7)	140.2 (7.0)	0.01
Short stature*, n (%)	22 (15.2)	32 (22.5)	0.11
Cord blood lead‡, mean (SD), µg/dL	2.4 (2.1)	3.0 (2.1)	0.03
Postnatal PCB 153‡, mean (SD), µg/kg	3.93 (0.89)	3.77 (0.93)	0.13
Serum ferritin, mean (SD), µg/L	25.6 (27.1)	22.5 (19.5)	0.26
<15 µg/L, n (%)§	42 (29.4)	55 (38.2)	0.11
Haemoglobin, mean (SD), g/L	131.4 (9.3)	127.6 (10.6)	<0.01
<120 µg/L, n (%)	13 (9.0)	24 (16.6)	0.06
Iron-deficiency anaemia, n (%)§	9 (6.3)	16 (11.1)	0.15

* Lowest height tertile for age and sex based on WHO growth charts for Canada.

† p-values were obtained by Student's t test for continuous variables and by chi-square statistics for categorical variables.

‡ Log transformed.

§ 5 missing values.

|| 7 missing values.

obtained from each participating mother, and oral consent was obtained from the child. This project was approved by the Laval University and Wayne State University ethics committees.

Assessment of food insecurity

Our exposure measure was moderate to severe food insecurity in the month preceding data collection. Moderate to severe food insecurity refers to a reduction in the quantity of food in a household due to lack of money. In contrast, marginal food insecurity, which we did not measure, encompasses anxieties about running out of food and/or limited food selection because of insufficient money.⁴ Four questions from the USDA (United States Department of Agriculture) Household Food Security Survey Module²¹ were included in the socio-demographic, interviewer-administered questionnaire. These questions, adapted to the Nunavik context, were: 1) Could you please describe the amount eaten by your family (enough/not enough)? 2) How many days in the previous month did your family not have food or money to buy food (0, ≥ 1)? 3) Do adults in your family have to cut down on the size of meals because there is not enough money (yes/no)? 4) Do children in your family have to cut down on the size of meals because there is not enough money (yes/no)? Questions one and two were highly collinear ($\kappa = 0.82$, $p=0.04$); none of the respondents who reported that they had enough to eat also stated that they went one or more days without food. Twenty-three participants (20%) reported that they did not have enough to eat but never went one or more days without food in the preceding month. Given the very high agreement for the responses to these questions, we excluded question two from subsequent analyses. Children whose mothers responded affirmatively to any of the three remaining food security questions were categorized as food insecure.

Assessment of nutritional deficiencies and short stature

Our outcome measures were: iron depletion, anaemia, iron-deficiency anaemia, and height (cm). Haematological measures

Table 3. Association of food insecurity to nutritional deficiencies and stature in 292 school-aged Inuit children from Nunavik, adjusting for age and sex (model 1); age, sex and BMI status (model 2); and age, sex, BMI status and environmental contaminants (model 3)*

Models/Outcomes	Food insecure							
	Iron depletion (n=97)		Anaemia (n=37)		Iron-deficiency anaemia (n=25)		Short stature (n=54)	
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
Multivariate model 1†	1.50 (0.91-2.45)	0.11	1.98 (0.97-4.07)	0.06	1.81 (0.77-4.28)	0.18	1.69 (0.92-3.11)	0.09
Multivariate model 2‡	1.45 (0.87-2.41)	0.15	1.97 (0.96-4.05)	0.07	1.76 (0.74-4.17)	0.20	1.66 (0.90-3.08)	0.11
Multivariate model 3§	1.28 (0.75-2.18)	0.36	1.75 (0.81-3.77)	0.16	1.59 (0.62-4.06)	0.34	1.49 (0.77-2.88)	0.23

* Odds ratios (OR) and *p*-values were obtained using logistic regression models. ORs represent the relation of food insecurity to nutritional deficiencies and short stature.

† Adjusted for age in years and sex.

‡ Further adjusted for BMI status (normal/overweight/obese).

§ Further adjusted for cord blood lead concentration (µg/L, continuous log transformed) and PCB 153 (µg/kg, continuous log transformed) measured at 10-year follow-up.

were analyzed by the laboratory of the *Centre hospitalier de l'Université Laval* (CHUL). Haemoglobin concentrations were determined by spectrophotometry using a Sysmex XT-2000i Automated Hematology Analyzer, whereas ferritin levels were quantified with a Modular Analytics E170 immunoassay analyzer (Roche Diagnostics). A child was categorized as iron depleted when serum ferritin was <15 µg/L²² and anaemic when haemoglobin level was below 120 g/L.²³ A child was considered to have iron-deficiency anaemia when haemoglobin was below 120 and serum ferritin was less than 15 or when haemoglobin was less than 120 and the transferrin saturation coefficient was less than 0.14.²⁴ For height, we used both the continuous measure and the WHO growth charts for Canada.²⁵ Based on the Canadian growth charts, a child was considered small of stature when his/her height was in the lowest growth tertile, based on age and sex.

Covariates assessment

We considered the following covariables: current age, sex, and normal/overweight/obesity status. Children were categorized as normal, overweight or obese according to their BMI, age and sex, using the International Obesity Task Force cut-off values.²⁶ We also considered cord blood lead and child plasma PCB 153 concentrations, as they were related to child growth in the same sample (unpublished data) and blood lead concentration may interfere with iron synthesis.²⁷ Total lead concentrations in children's blood samples were determined by inductively coupled plasma mass spectrometry (ICP-MS) using a PerkinElmer Sciex Elan 6000 and PE DRC II instrument. PCB congener 153 was measured in purified cord extracts using gas chromatography/mass spectrometry. Detection limits in cord samples were 0.2 µg/L for lead, and 0.02 µg/L for PCB 153. Upstream variables including household income, education and crowding were not considered because food insecurity is a consequence of these conditions and inclusion of these variables in statistical models would lead to over-adjustment. Finally, we did not adjust for inflammation (C-reactive protein ≥10.0 mg/L), which can affect the interpretation of serum ferritin levels,²⁸ because only 15 children (6.1%) had inflammation and there was no evidence that the distribution of these children differed by food security status.

Statistical analyses

Bivariate associations of food insecurity with nutritional deficiencies and stature were tested with the chi-square statistic or

Student's *t*-test. Logistic regression models were used to estimate odds ratio (OR) and 95% confidence intervals (CIs) of nutritional deficiencies (iron depletion, anaemia, iron-deficiency anaemia) and short stature in relation to food insecurity. Height (cm) was also examined using a general linear model. Multivariate Model 1 included food insecurity, age and sex. Model 2 was further adjusted for BMI status (normal/overweight/obesity). Because children were categorized as short for stature based on their age, we ran Model 2 with and without age for this outcome; results were identical for both models. Finally, we adjusted for exposure to environmental contaminants (Model 3). Cord blood lead and current blood PCB 153 concentrations were log transformed to reduce skewness. Because iron deficiency is associated with impaired growth,²⁹ our sensitivity analyses for height further adjusted for iron deficiency and anaemia. Statistical analyses were performed in SPSS version 21 (IBM Corp., Armonk, NY).

RESULTS

Participants ranged between 8.5 and 14.3 years of age. Age did not differ by sex. Half of the children lived in a household categorized as food insecure, and about one in five lived in households in which the caregiver reported not enough to eat and that both adults and children had cut their portion sizes (Table 1). Information on food security was missing for two children.

One third of participants had iron depletion, 12.6% were anaemic and 8.7% had iron deficiency anaemia (Table 2). The prevalence of iron depletion and anaemia was not statistically different between boys and girls. However, boys were significantly more likely to have iron-deficiency anaemia compared to girls (12.4% versus 4.9%, *p*=0.04). The mean height of the population was 141.4 cm (SD=7.4, range=118.9–167.8 cm). According to WHO growth charts for Canada, 19% of the children were in the lowest tertile for height. There were no significant differences in the mean heights of boys and girls. However, girls were significantly more likely to be short for their age (24% versus 13%, *p*=0.02) compared to boys.

Food security status was not related to child age, sex or BMI status (Table 2). Compared to food-secure children, food-insecure children were significantly shorter and had higher cord blood lead levels, whereas current blood PCB 153 concentrations were unrelated to food security status. Food-insecure children had significantly lower haemoglobin levels, but the other haematological measures did not differ by food security status.

Table 4. Adjusted associations between food insecurity and height in 276 school-aged Inuit children from Nunavik*

Models/Outcomes	Food insecure		R ²
	Height (cm)		
Multivariate model 1 [†]	$\beta^{\dagger}$ (95% CI)	P-value	
Multivariate model 2 [§]	-2.41 (-0.97 to -3.95)	<0.001	0.21
Multivariate model 3	-2.19 (-0.86 to -3.51)	0.001	0.41
Multivariate model 3	-1.82 (-0.48 to -3.15)	0.008	0.47
Sensitivity analysis			
Multivariate model 4 [¶]	-1.82 (-0.48 to -3.17)	0.008	0.47

* Estimates and *p*-values were obtained using univariate general linear models.
[†] β estimates represent the relation of food insecurity to height (cm).
 β estimates with negative values represent the mean number of cm smaller a food-insecure child would be after adjustment for the covariables in the multivariate models.
[§] Adjusted for age in years and sex.
^{||} Further adjusted for BMI status (normal, overweight or obese).
[¶] Further adjusted for cord blood lead concentration ($\mu\text{g/L}$, continuous log transformed) and PCB 153 measured at 10-year follow-up ($\mu\text{g/kg}$, continuous log transformed).
[¶] Further adjusted for iron depletion (y/n) and anaemia (y/n).

In the adjusted models, food insecurity was not associated with iron depletion or iron-deficiency anaemia (Table 3). In Model 1, food insecurity was marginally associated with anaemia and short stature. For both of these outcomes, there was a reduction in the strength of the association once contaminants were adjusted statistically (Model 3). In the fully-adjusted multivariate analysis of the continuous height measure, food-insecure children were on average about 2 cm ($\beta = -1.82$; 95% CI: -0.48 to -3.15) shorter than children who were food secure (Table 4). Iron depletion ($\beta = 1.06$; $p = 0.19$) and anaemia ($\beta = -0.96$; $p = 0.39$) were not independently associated with height and did not alter the coefficient for food security, or other variables in the model.

DISCUSSION

This is the first study to look at the effects of food insecurity on school-aged Inuit children and the only study of this population to examine the association between food insecurity and stature. Nearly half of the children in this study lived in households that were moderately to severely food insecure; that is, caregivers reported not having enough to eat and/or reducing meal portion sizes. Eighty percent of the households reporting not enough to eat went 1 day/month or more without food, and 27% of caregivers reported that they had cut the size of their children's meals because there was not enough money for food. By comparison, data from a recent representative study of Canadian households reported that less than 1% of caregivers cut children's portion sizes.⁴

This study suggests that food insecurity has implications for Inuit children's health and development. Children from food-insecure households in Nunavik were marginally more likely to be anaemic and had significantly lower mean haemoglobin levels. In the adjusted models, food insecurity was not statistically significantly associated with iron depletion or iron-deficiency anaemia, although the coefficient for iron-deficiency anaemia was similar to that of anaemia in Model 1. The small number of cases of iron-deficiency anaemia ($n = 25$) may have prevented us from observing a statistically significant association. When we adjusted for environmental contaminants (Model 3),

the odds ratio for the relation of food insecurity to anaemia was notably reduced (from 1.98 to 1.75). Exposure to environmental contaminants in this population is largely attributable to the consumption of traditional, or "country foods", which are also important sources of iron and other nutrients.³ Those with greater contaminant levels may have an overall more nutritious diet and thus, less nutritional deficiencies.

While not all iron-related nutritional deficiencies were significantly associated with food insecurity, they were nonetheless notably more prevalent in food-insecure children compared to food-secure children. Food insecurity may be a proxy or a consequence of other determinants – such as, household income, crowding and social support – that influence a child's overall health and nutritional status. Even in the absence of a causal association between food insecurity and nutritional deficiencies, determining a child's food security status may assist health care providers in targeting and monitoring those children most vulnerable to poor health outcomes.

Iron depletion was highly prevalent (33%) in this sample of primary school-aged Inuit children. This estimate is similar to the prevalence of iron depletion documented in a 2004 survey of women from Nunavik (30%).³⁰ In a study of 32 low- and middle-income countries, the mother's anaemia status was one of the most important predictors of anaemia in her children, surpassing socio-economic predictors in many countries.³¹ It is likely that a similar relation exists for iron depletion in Inuit children. Anaemia (13%) was much less prevalent than iron depletion; nonetheless, most anaemia (25/37 or 68%) in this population is iron-deficiency anaemia. For the 12 children not recognized as having iron-deficiency anaemia, infection and other nutritional deficiencies may be responsible for the low haemoglobin measures. The prevalence of anaemia and iron-deficiency anaemia observed in this study was similar to that reported in a study of Inuit children aged 3-5 years in Nunavut.¹² In contrast, the prevalence of iron depletion in our sample was nearly double that of the preschoolers from Nunavut. This may reflect geographic and dietary difference between the regions and/or increased time since breastfeeding.

Food-insecure children were significantly shorter in stature, by an average of 2 cm, than their food-secure counterparts. For children of this age group, this is close to half a year's growth.²⁵ Adjusting for current iron-related nutritional deficiencies did not alter the significant association between food insecurity and stature. The observed association between food insecurity and linear growth suggests that the diet quality and quantity of children from food-insecure households has been compromised for a long time. Unfortunately, food security was not measured at birth in this cohort, and this study is limited by the assumption that current food insecurity provides a proxy for past insecurity. Clearly, because of limitations inherent in any cross-sectional study, no causal relation can be ascertained. Nevertheless, there is reason for concern that food-insecure Inuit children are not consuming enough protein, calcium and/or zinc.^{3,29} Zinc, in particular, has been associated with linear growth in children and may be a worthwhile avenue of future investigation in this population.¹³

One limitation of this study was that we did not use the full USDA food security survey module.²¹ Two other surveys, also

conducted in predominantly Inuit-inhabited regions of Canada, that used the full USDA module arrived at very similar prevalence estimates of food insecurity.^{1,32} In Nunavut, Nanatsiavut, and the Inuvialuit settlement, 51.3% of households with children were estimated to be food insecure.¹ In Kangiqsujuaq, Nunavik, 43% of women surveyed were considered food insecure.³² Further, the directions of association between food security, nutritional deficiencies, and growth suggest content validity. Finally, because the participants were Inuit children, the generalizability of the observed associations may be limited to similar populations.

CONCLUSION

Food insecurity is a considerable public health problem among Inuit in Nunavik, Canada. Health care providers in the region should be aware that there is a significant association between food insecurity and shorter stature and that the burden of iron-related nutritional deficiencies appears to be greater among food-insecure children. The results of this study raise concerns about the long-term implications of food insecurity for children in Nunavik and highlight the need for health care providers to carefully monitor children suspected to be food insecure.

REFERENCES

- Huet C, Rosol R, Egeland GM. The prevalence of food insecurity is high and the diet quality poor in Inuit communities. *J Nutr* 2012;142(3):541-47.
- WHO. Food Security. Available at: <http://www.who.int/trade/glossary/story028/en/> (Accessed October 17, 2013).
- Blanchet C, Rochette L. Nutrition and food consumption among the Inuit of Nunavik, In: Nunavik Inuit Health Survey 2005. Qanuippitaa? How are we? Institut national de santé publique du Québec (INSPQ), Nunavik Regional Board of Health and Social Services (NRBHSS), 2008.
- Tarasuk V, Mitchell A, Dachner N. Research to identify policy options to reduce food insecurity (PROOF). Household food insecurity in Canada 2011. 2013. Available at: <http://nutritionalsciences.lamp.utoronto.ca/> (Accessed August 15, 2013).
- Campbell AA, Akhter N, Sun K, De Pee S, Kraemer K, Moench-Pfanner R, et al. Relationship of household food insecurity to anaemia in children aged 6-59 months among families in rural Indonesia. *Ann Trop Paediatr* 2011;31(4):321-30.
- Eicher-Miller HA, Mason AC, Weaver CM, McCabe GP, Boushey CJ. Food insecurity is associated with iron deficiency anemia in US adolescents. *Am J Clin Nutr* 2009;90(5):1358-71.
- Jamieson J, Weiler H, Kuhnlein H, Egeland GM. Traditional food intake is correlated with iron stores in Canadian Inuit men. *J Nutr* 2012;142:764-70.
- Park K, Kersey M, Geppert J, Story M, Cutts D, Himes JH. Household food insecurity is a risk factor for iron-deficiency anaemia in a multi-ethnic, low-income sample of infants and toddlers. *Public Health Nutr* 2009;12(11):2120-28.
- Pasricha SR, Black J, Muthayya S, Shet A, Bhat V, Nagaraj S, et al. Determinants of anemia among young children in rural India. *Pediatrics* 2010;126(1):e140-49.
- Skalicky A, Meyers AF, Adams WG, Yang Z, Cook JT, Frank DA. Child food insecurity and iron deficiency anemia in low-income infants and toddlers in the United States. *Matern Child Health J* 2006;10(2):177-85.
- Kirkpatrick SI, Tarasuk V. Food insecurity is associated with nutrient inadequacies among Canadian adults and adolescents. *J Nutr* 2008;138(3):604-12.
- Pacey A, Weiler H, Egeland GM. Low prevalence of iron-deficiency anaemia among Inuit preschool children: Nunavut Inuit Child Health Survey, 2007-2008. *Public Health Nutr* 2011;14(8):1415-23.
- Ramakrishnan U, Nguyen P, Martorell R. Effects of micronutrients on growth of children under 5 y of age: Meta-analyses of single and multiple nutrient interventions. *Am J Clin Nutr* 2009;89(1):191-203.
- Chwang LC, Soemantri AG, Pollitt E. Iron supplementation and physical growth of rural Indonesian children. *Am J Clin Nutr* 1988;47(3):496-501.
- Baig-Ansari N, Rahbar MH, Bhutta ZA, Badruddin SH. Child's gender and household food insecurity are associated with stunting among young Pakistani children residing in urban squatter settlements. *Food Nutr Bull* 2006;27(2):114-27.
- Psaki S, Bhutta ZA, Ahmed T, Ahmed S, Bessong P, Islam M, et al. Household food access and child malnutrition: Results from the eight-country MAL-ED study. *Population Health Metrics* 2012;10(1):24.
- Saha KK, Frongillo EA, Alam DS, Arifeen SE, Persson LA, Rasmussen KM. Household food security is associated with growth of infants and young children in rural Bangladesh. *Public Health Nutr* 2009;12(9):1556-62.
- Dallaire F, Dewailly E, Muckle G, Ayotte P. Time trends of persistent organic pollutants and heavy metals in umbilical cord blood of Inuit infants born in Nunavik (Québec, Canada) between 1994 and 2001. *Environ Health Perspect* 2003;111(13):1660-64.
- Muckle G, Ayotte P, Dewailly EE, Jacobson SW, Jacobson JL. Prenatal exposure of the northern Québec Inuit infants to environmental contaminants. *Environ Health Perspect* 2001;109(12):1291-99.
- Jacobson JL, Jacobson SW, Muckle G, Kaplan-Estrin M, Ayotte P, Dewailly E. Beneficial effects of a polyunsaturated fatty acid on infant development: Evidence from the Inuit of Arctic Quebec. *J Pediatr* 2008;152:356-64.
- Bickel G, Nord M, Hamilton W, Cook J. Guide to Measuring Household Food Security, Revised 2000. Alexandria, VA: Office of Analysis, Nutrition, and Evaluation, Food and Nutrition Service, USDA.
- WHO. Serum ferritin concentrations for the assessment of iron status and iron deficiency in populations. Geneva, Switzerland: World Health Organization, 2011.
- WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Geneva: WHO, 2011.
- Maire B, Delpeuch F. Nutrition indicators for development: Reference Guide. Montpellier, France: Institut de Recherche pour le Développement, 2005.
- Dietitians of Canada. WHO Growth Charts for Canada. 2010. Available at: <http://www.dietitians.ca/secondary-pages/public/who-growth-charts.aspx> (Accessed October 17, 2013).
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: International survey. *BMJ* 2000;320(7244):1240-43.
- Hammond PB. Exposure of humans to lead. *Ann Rev Pharmacol Toxicol* 1977;17:197-214.
- Engle-Stone R, Nankap M, Ndjebayi AO, Erhardt JG, Brown KH. Plasma ferritin and soluble transferrin receptor concentrations and body iron stores identify similar risk factors for iron deficiency but result in different estimates of the national prevalence of iron deficiency and iron-deficiency anemia among women and children in Cameroon. *J Nutr* 2013;143(3):369-77.
- Prentice A, Schoenmakers I, Laskey MA, de Bono S, Ginty F, Goldberg GR. Nutrition and bone growth and development. *Proc Nutr Soc* 2006;65(4):348-60.
- Plante C, Blanchet C, Rochette L, O'Brien HT. Prevalence of anemia among Inuit women in Nunavik, Canada. *Int J Circumpolar Health* 2011;70(2):154-65.
- Balarajan Y, Ramakrishnan U, Ozaltin E, Shankar A, Subramanian S. Anaemia in low-income and middle-income countries. *Lancet* 2012;378(9809):2123-35.
- Lawn J, Harvey D. Nutrition and Food Security in Kangiqsujuaq, Nunavik: Baseline survey for the food mail pilot project. Ottawa, ON: Minister of Indian Affairs and Northern Development, 2004.

Received: March 21, 2014

Accepted: July 4, 2014

RÉSUMÉ

OBJECTIFS : Les Inuits du Canada présentent des niveaux d'insécurité alimentaire alarmants dont les conséquences nutritionnelles et physiologiques sont méconnues, surtout chez les enfants d'âge scolaire. L'objectif de notre étude était d'évaluer la relation entre l'insécurité alimentaire, d'une part, et la carence en fer et la taille, d'autre part, chez les enfants inuits du Nunavik (Nord-du-Québec) en âge de fréquenter l'école.

MÉTHODE : Nous avons évalué l'insécurité alimentaire, la carence en fer et la taille dans une cohorte d'enfants. L'insécurité alimentaire a été déterminée en interviewant les mères des enfants. Par régression logistique multiple, nous avons évalué l'association entre l'insécurité alimentaire et la carence en fer et la petite taille. Nous avons défini la petite taille comme étant la taille dans le tertile inférieur pour l'âge et le sexe, selon les fiches de croissance canadiennes. Le rapport entre l'insécurité alimentaire et la taille (en centimètres) a été analysé selon un modèle linéaire général. Des modèles statistiques ont permis de rajuster les données selon l'âge, le sexe, le poids normal/le surpoids/l'obésité, l'exposition prénatale au plomb et l'exposition postnatale aux biphényles polychlorés.

RÉSULTATS : La moitié des enfants (49,7 %, n=145) étaient confrontés à l'insécurité alimentaire, le tiers des enfants étaient carencés en fer, 12,6 % faisaient de l'anémie, et 8,7 % présentaient une anémie ferriprive. Le

FOOD INSECURITY IN NUNAVIK CHILDREN

rapport de cotes multivarié de l'anémie était de 1,82 (IC de 95 % : 0,97, 3,42, $p=0,06$) chez les enfants confrontés à l'insécurité alimentaire. La prévalence de la petite taille était de 18,7 %. Les enfants touchés par l'insécurité alimentaire mesuraient en moyenne 2 cm de moins (IC de 95 % : -0,48, -3,17) que les enfants épargnés par l'insécurité alimentaire ($p<0,01$).

CONCLUSION : Dans cette population, les enfants confrontés à l'insécurité alimentaire avaient un plus lourd fardeau de carences alimentaires et une croissance linéaire plus lente. Étant donné la forte prévalence de l'insécurité alimentaire chez les enfants inuits du Nunavik, les carences alimentaires et leurs effets indésirables sur le développement devraient être suivis attentivement.

MOTS CLÉS : insécurité alimentaire; carences alimentaires; fer; croissance; enfant; santé autochtone