Youth Smoking in Ontario 1981-1997: A Cause for Concern

Frances M. Hobbs, MA,¹ William Pickett, PhD,^{1,2} Roberta G. Ferrence, PhD,^{1,3}
K. Stephen Brown, PhD,^{1,4} Cheryl Madill, MSc,⁵ Edward M. Adlaf, PhD³

Smoking among youth has widespread and long-term public health implications. Adolescent patterns of smoking often continue into adulthood,¹ and cessation programs aimed at youth are frequently unsuccessful.² Moreover, it is estimated that cigarette smoking will account for over 50% of deaths in Canada before the age of 70 for 15-year-old current smokers if they continue to smoke.^{3,4} The Ontario Tobacco Strategy, introduced in 1992 as part of the Ontario government's health reform agenda,⁵ set a goal to reduce the prevalence of smoking among 12-19 year olds in the province to 10% by the year 2000. This report presents trend data from two major school surveys on smoking among Ontario youth, discusses potential causal factors and outlines strategies to address this public health problem.

METHODS

The descriptive analyses, derived from the Ontario Student Drug Use Survey⁶ and

the Waterloo Smoking Prevention Projects,⁷ span the years 1981-1997. The Ontario Student Drug Use Survey is a biennially repeated cross-sectional survey of Ontario students enrolled in grades 7, 9, 11 and 13, conducted by the Addiction Research Foundation (now a division of the Centre for Addiction and Mental Health) every two years since 1977. The sample design, comprised of 16 strata, is based on a single-stage cluster sample of classrooms stratified equally by grade level (grades 7, 9, 11 and 13) and proportionally by region (Toronto, Western, Eastern and Northern Ontario).

The 1997 sample consists of 3,990 students from 168 schools (response rate = 77%). Earlier cycles are comparable (number of students: 3,270-4,737; number of schools: 137-227; response rates: 76%-84%). *Past year smoking* means smoking more than one cigarette in the previous 12 months. *Daily smoking* is one or more cigarettes smoked each day in the previous 12 months. *Cigarettes per day* is the number of cigarettes smoked daily in the previous 12 months by daily smokers. Calculation of confidence intervals incorporates clustering effects.

In addition to conducting longitudinal surveys for smoking intervention trials, the Health Behaviour Research Group at the University of Waterloo carried out cross-sectional surveys of 9th-graders from 24 schools in Southwestern Ontario in 1995, 1996 and 1997. Youth smoking, measured in the spring of each year, was defined as *regular use* if the respondent usually smoked every week and *experimental* if the student had not quit but smoked less frequently than once a week. Prevalence rates are provided for regular and experimental smokers combined (current smokers).

RESULTS

Ontario Student Drug Use Survey

In 1997, 28% of Ontario students reported cigarette use, a rate that has remained unchanged since 1995 (Figure 1). Between 1981 and 1991, past year smoking among females declined from 35% to 22% ($p < 0.05$), but increased between 1991 and 1997 to 29% ($p < 0.05$). Smoking among males also declined between 1981 and 1991, from 26% to 22% ($p < 0.05$), and increased to 26% ($p < 0.05$) between 1991 and 1997. Significant increases also occurred in rates of daily smoking, with 20% of students reporting daily smoking in 1997 compared to 16% in 1991 ($p < 0.05$).

Among daily smokers, the number of cigarettes smoked has remained stable since 1981 (range = 7.7-8.4, n.s.). Males consistently smoked about one cigarette more per day than females, although these differences were not statistically significant within years.

Waterloo Smoking Prevention Projects

Smoking rates among southwestern Ontario 9th graders for the years 1995 through 1997 are substantially higher than in the provincial survey but show a similar pattern. Rates did not differ significantly by sex. Rates of current smoking for females and males were 34% and 32% respectively in 1995, 38% and 35% in 1996, and 36% and 32% in 1997. Comparable rates for 9th graders in the Ontario survey are 29% and 26% for females and males respectively in 1995 and 30% and 21% in 1997. Despite differences in the methodology and geographic areas sampled, smoking rates within Ontario and the southwestern region did

1. Ontario Tobacco Research Unit, University of Toronto, Toronto, ON
 2. Department of Community Health and Epidemiology, Queen's University, Kingston, ON
 3. Centre for Addiction and Mental Health, Addiction Research Foundation Division, Toronto, ON
 4. Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, ON
 5. Department of Health Studies and Gerontology, University of Waterloo, Waterloo, ON
- Correspondence and reprint requests: Dr. Roberta Ferrence, Director, Ontario Tobacco Research Unit, 33 Russell Street, Toronto, Ontario M5S 2S1, Tel: 416-595-6889, Fax: 416-595-6068, E-mail: rferrence@ar.f.org

The Ontario Tobacco Research Unit is funded by the Ontario Ministry of Health. Funding for data collection was provided by the Centre for Addiction and Mental Health, Addiction Research Foundation Division, and the Heart and Stroke Foundation of Ontario.

not change significantly between 1995 and 1997.

DISCUSSION

The data presented clearly show cause for concern. Rates of smoking among Ontario youth fell substantially during the 1980s, but rose during the 1990s and have recently plateaued. Similar patterns were observed in Ontario among adults during the 1990s, but were statistically significant only for females.⁸ Daily consumption of cigarettes by youth did not change during this time period. Available data suggest that between one quarter and one third of youth are current smokers.⁷ Clearly, the prospects for reducing the prevalence of youth smoking to the Ontario Tobacco Strategy goal of 10% by the year 2000 are poor.

Potential factors in the general increase in youth smoking during the 1990s, which have also been observed in Nova Scotia,⁹ Prince Edward Island,¹⁰ Great Britain,¹¹ and the United States,¹² include the increased access to cigarettes brought about by price cuts, since consumption is inversely related to price,^{13,14} promotions and other marketing efforts,¹⁵ and larger social changes related to use of other drugs and attitudes toward risk and acceptability.¹⁶ Cigarettes are available to youth at about the same price they were after the 1994 tax rollback in which a federal and provincial tax cut reduced the price of a pack of cigarettes by about 50%. Furthermore, the price of a pack of cigarettes in early 1999 in Ontario is the lowest of all ten provinces and is substantially lower than prices in three border states,¹⁷ eliminating any potential for smuggling.

Retailer compliance with legislation forbidding sales to minors did not improve from 1996 (74%) to 1997 (69%).¹⁸ This pattern was evident for all types of establishments, with corner stores the most popular source of cigarettes for minors in 1997, followed by gas stations. The proportion of youth asked for photo identification increased from 39% in 1995 to 45% in 1997,⁶ but still remains relatively low. Finally, research evidence for an effect of enforcement and retailer compliance on youth smoking is not conclusive.¹⁹

Efforts to reduce youth smoking in Ontario should focus on evidence-based

Figure 1. Past Year Smoking and Daily Smoking among Ontario Students 1981-1997

Source: Ontario Student Drug Use Survey 1981-1997 (grades 7, 9, 11, 13), Centre for Addiction and Mental Health, Addiction Research Foundation Division
Note: Past year smoking = more than 1 cigarette in past year

policy initiatives that reinforce existing programming and public education efforts. Young people are particularly responsive to price increases, and a substantial increase in provincial tobacco taxes would reduce both prevalence and consumption.²⁰ Very high levels of vendor compliance may have some impact on prevalence,²¹ but restricting tobacco sales to government outlets would provide more effective control. Expanding bans on smoking in public places and workplaces to include those heavily frequented by young people, such as restaurants and bars, would reduce the visibility of smoking, in addition to protecting youth from second hand smoke. Since smokers report that relapse is more likely when others around them are smoking,²² such bans are critical to support cessation efforts by young people. These measures, along with support for federal elimination of tobacco sponsorship and plain packaging of cigarettes, would constitute a broad public health strategy that would appropriately address youth smoking in this province.

REFERENCES

- Chassin L, Presson C. The natural history of cigarette smoking: Predicting young adult smoking outcomes from adolescent smoking patterns. *Health Psychol* 1990;9:701-16.
- Stone C, Kristeller J. Attitudes of adolescents toward smoking cessation. *Am J Prev Med* 1992;8:221-24.
- Peto R, Lopez AD, Boreham J, et al. Mortality from Smoking in Developed Countries 1950-2000, Indirect estimates from National Vital Statistics. World Health Organization, Imperial Cancer Research Fund, Oxford University Press, 1994.
- Villeneuve P, Morrison H. Health consequences of smoking in Canada: An update. *Chron Dis Can* 1994;15:102-4.
- Ontario Ministry of Health. The Ontario Tobacco Strategy Overview. Toronto: Ontario Ministry of Health, 1993.
- Adlaf E, Ivis F, Smart R. The Ontario Student Drug Use Survey 1977-1997. Toronto: Addiction Research Foundation, 1997.
- Manske SR, Brown KS, Cameron AJR. School-based cancer control: A research agenda. *Cancer Prevention and Control* 1997;1(3):196-212.
- Bondy SJ, Ialomiteanu AR. Smoking in Ontario, 1991 to 1996. *Can J Public Health* 1997;88(4):225-29.
- Poulin C, Elliott D. Alcohol, tobacco and cannabis use among Nova Scotia adolescents: Implications for prevention and harm reduction. *CMAJ* 1997;156(10):1387-93.
- Department of Health and Social Services. Prince Edward Island Student Drug Use 1996: Technical Report. Prepared in collaboration with Dalhousie University, PEI Health and Community Services Agency and PEI Department of Education, November 1996.
- Hall C. Smoking by young men rises to 39%. Based on results from the 1996 Health Survey for England. *UK Electronic Telegraph*. Tuesday January 13, 1998, Telegraph Group Limited 1998.
- University of Michigan. Drug use among American teens shows some signs of levelling after a long rise. Ann Arbor: University of Michigan, 1997. [Available at: www.isr.umich.edu/src/mtf/mfdat97.html].

13. U.S. Department of Health and Human Services. Preventing Tobacco Use Among Young People, A Report of the Surgeon General. Atlanta, Georgia: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1994.
14. Galbraith JW, Kaiserman M. Taxation, smuggling and demand for cigarettes in Canada: Evidence from time-series data. *J Health Economics* 1997;16:287-301.
15. Gilpin EA, Pierce JP. Trends in adolescent smoking initiation in the United States: Is tobacco marketing an influence? *Tobacco Control* 1997;6:122-27.
16. Johnston LD, O'Malley PM, Bachman JG. National Survey Results on Drug Use from the Monitoring the Future Study, 1975-1997. Secondary School Students, Volume I. National Institute on Drug Abuse, US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1998.
17. Sweanor D. Tobacco Taxes and Prices in Canada. Toronto: Smoking and Health Action Foundation, 1998.
18. Nielsen AC. Measurement of retailer compliance with respect to the Tobacco Act & Provincial Sales-to-Minors legislation. Final Results Wave I. Report prepared for Health Canada's Health Protection Branch, Office of Tobacco Control. Toronto: ACNielsen, 1997.
19. DiFranza JR, Rigotti NA. Policies to Reduce Youth Access to Tobacco. Health Science Analysis Project, Policy Analysis 16. Washington, DC: Advocacy Institute, June 1998.
20. Ferrence RG, Garcia JM, Sykora K, et al. Effects of pricing on cigarette use among teenagers and adults in Canada 1980-1989. Toronto, ON: Addiction Research Foundation, 1991.
21. Forster JL, Wolfson M. Youth access to tobacco: Policies and politics. *Annu Rev Public Health* 1998;19:203-35.
22. Ashley MJ, Cohen J, Bull S, et al. Smoking in Ontario, Analysis of Data from the "Q&Q" Study. Prepared for Health Canada and the Ontario Ministry of Health. Toronto, ON: Ontario Tobacco Research Unit, 1997.

Received: September 4, 1998
Accepted: February 4, 1999

C O M I N G E V E N T S A C T I V I T É S À V E N I R

To be assured of publication in the next issue, announcements should be received by May 15, 1999 and valid as of June 30, 1999. Announcements received after May 15, 1999 will be inserted as time and space permit.
Pour être publiés dans le prochain numéro, les avis doivent parvenir à la rédaction avant le 15 mai 1999 et être valables à compter du 30 juin 1999. Les avis reçus après le 15 mai 1999 seront insérés si le temps et l'espace le permettent.

Healthy People and Healthy Communities: A Canada - United States Dialogue on Best Practices in Public Health

Hosted by the Canada West Foundation
8-9 May 1999

Royal York Hotel, Toronto, ON

Contact:

Carey Hill, Conference Organizer
Healthy People and Healthy Communities
The Canada West Foundation
550, 630 3rd Avenue SW
Calgary, AB T2P 4L4
Tel: 403-264-9535 Fax: 403-269-4776
E-mail : hill@cwf.ca

Evaluation: An Essential Contribution

Conference '99, Canadian Evaluation Society
16-19 May 1999 Toronto, ON

Contact:

CES Conference '99 Secretariat
c/o The Willow Group
582 Somerset Street West
Ottawa, ON K1R 5K2
Tel: 613-230-1007 Fax: 613-237-9900

Public Health in the New Millennium/ La santé publique au prochain millénaire

CPHA's 90th Annual Conference/90^e conférence annuelle de l'ACSP
6-9 June/juin 1999 Winnipeg, Manitoba

Contact/Contacter :

CPHA Conference Department
Service des conférences ACSP
400 - 1565 avenue Carling Avenue
Ottawa, ON K1Z 8R1
Tel: 613-725-3769
Fax/Téléc. : 613-725-9826
E-mail/C. électr. : conferences@cpha.ca
www.cpha.ca

The Dawn of a New Era: Challenges for Baby Boomers

Canadian Association on Gerontology Annual Scientific and Educational Meeting
4-7 November 1999

Ottawa, ON

Contact:

CAG Conference Secretariat
100 - 824 Meath Street
Ottawa, ON K1Z 6E8
Tel: 613-728-9347 Fax: 613-728-8913
E-mail: cagacg@magi.com

Celebrating a Century of Progress in Public Health

1999 Annual Meeting of the American Public Health Association
7-11 November 1999 Chicago, Illinois

Contact:

APHA Convention Services
1015 Fifteenth Street, NW
Washington, DC 20005
Tel: 202-789-5620 www.apha.org

Canada's Fourth National Conference on Asthma and Education

Presented by the Canadian Network for Asthma Care
11-13 November 1999 Halifax, NS

Contact:

A. Les McDonald, Executive Director
Canadian Network for Asthma Care
6 Forest Laneway, Suite 1607
North York, ON M2N 5X9
Tel: 416-224-9221 Fax: 416-224-9220
E-mail: ased@cnac.net

CALL FOR ABSTRACTS

Celebrating Our Past, Building Our Future
50th Annual Ontario Public Health Association Conference

15-17 November 1999 Toronto, ON
Co-hosted by the Centre for Health Promotion and Toronto Public Health

Contact:

OPHA Conference Committee Secretary,
Vanessa Tang
Tel: 416-395-7653
or OPHA, Tel: 416-367-3313

Abstract deadline: 14 May 1999